

WP
rA881d
1851

A T L E E



OPERATIONS OF OVARIOTOMY, 1701-1851

No.	Date.	Operators.	Age.	ISCISION.			RECOVERED.			DIED.			Time of Death.	Other Diseases.	Adhesions.	Character of Tumor.	Cause of Death.	REMARKS.	REFERENCES.
				Minor.	Major.	Unknown.	Minor.	Major.	Unknown.	Minor.	Major.	Unknown.							
101	1843	F. Bird	35	1			1								None	Cyst		Sack punctured, emptied and excised.	London Med. Gazette, March 22d, 1844, p. 632.
102	1843	do	21	1	1	1									None	Cyst and solid matter		Incision 5 inches; cyst emptied, withdrawn and excised.	Ibid., December 29, 1843, p. 409.
103	1844	do	21	1			1								None	Cysts		Followed by peritonitis, slight.	Ibid., December 8, 1843.
104	1844	do	35	1			1								Adhesions	Cysts		Many adhesions, sack emptied of firm gelatinous matter, and excised; incision 8 inches, tumor 35 lbs.	Ibid., August 18, 1843, p. 732.
105	1846	do	52	1			1								Strong pelvic ad.	Cysts		Cyst very thick and filled with deotestrum, tapped and extracted. There was no pedicle, and a small section of the uterus was removed with it. See p. 269.	London Lancet, Amer. Ed., April, 1846, p. 315.
106		do		1			1								Adhesions	Cystiform		Tumor adherent to pelvis and uterus. A segment of the cyst left attached.	Ibid.; also, Amer. Jour. Med. Science, Jan., 1851, p. 254.
107	1847	do	32			1			1						Adhesions	Cystiform		Multilocular; both ovaries extirpated. One of compound character weighed 20 lbs.; the other consisted of several cysts, weighed 4 lbs. Both adhered to abdominal walls, the larger one firmly to the omentum, an artery of which was tied. Rapid recovery without an untoward symptom. Menstruation both before and after operation uninterrupted.*	London Lancet, Oct. 30, 1847, p. 467; also, Ranking's Abstract, No. 6, 1848, p. 246.
<p>* We apprehend there has been an error in diagnosis of this case. Otherwise it presents a fac. which contradicts our best physiologists, who maintain that menstruation is a phenomena dependent upon the physiological action of the ovaria. If both ovaries were extirpated, and menstruation continued uninterrupted, then indeed are all our notions wrong as to the functions of these bodies.—B.</p>																			
108		do		1					1			3d day			Adhesions	Large sessile tumor			Ibid., Nov. 23, 1850; also, Amer. Jour. Med. Science, Jan., 1851, p. 234.
109		do		1								3d day			Adhesions	Cystiform		Tumor bound down in the pelvis, causing unceasing suffering. Tapping required every 10 or 12 days.	Ibid.
110		do		1					1			5th day			None			Small tumor.	Ibid.
111		do		1					1			7th day						Large compound tumor.	Ibid.
112		do		1	1				1	1		2d day			Adhesions			Very large malignant mass, inseparably adherent posteriorly. Extreme suffering from distention by solid matter, and rapidly approaching death rendered the attempted operation justifiable.	Ibid., p. 235.
113		do		1			1											Small incision, then tapped.	Ibid.
114		do		1			1											Small incision, then tapped.	Ibid.
115		do		1			1											Incision of rather large size.	Ibid.
116		do		1			1					6 weeks						Small incision to ascertain the character of the adhesions, and of small tumor attached to the cyst, as well as to evacuate the contents. Patient advanced in life and exhausted by suffering. Lived six weeks.	Ibid.
117		do		1			1											Incision.	Ibid.
118		do		1			1											Incision, then tapped.	Ibid.
119		do		1			1											Incision, then tapped.	Ibid.
120		do		1			1											Incision, then tapped.	Ibid.
121		do		1			1											Incision, then tapped. Tapped several times afterwards.	Ibid.
122		do		1			1											Incision, then tapped. Tapped afterwards.	Ibid.
123		do		1			1											Incision, not tapped. Tapped afterwards and died.	Ibid.
124		do		1					1			2d day					Bursting of hepatic abscess.	Incision to ascertain adhesions and to remove very viscid contents; died next day from bursting of hepatic abscess into the peritoneal cavity, as proved by autopsy.	Ibid.
125		do		1			1											Small incision, then tapped. Afterwards tapped.	Ibid.
126		do		1			1											Incision collic.	Ibid.
127		do		1			1											Small incision. Tapped many times afterwards.	Ibid.
128		do		1			1											Incision, then tapped.	Ibid.
129		do		1			1											Incision, then tapped.	Ibid.
130	1846	J. L. Atlee	33						1			15th day			Adhesions	Cystiform	Inflam. of the lungs	Multilocular, weighed 45 lbs. Patient got along very well for several days, and took cold on slight exposure, which produced pneumonia. An autopsy was made and no abdominal or pelvic inflammation was discovered.	Not published.
131	1843	do	29				1								Adhesions	Cyst and hydatids		Both ovaries removed.	Amer. Med. Journal, January, 1844, p. 44.
132	1844	do	42						1			5th day	Dis. of uterus	Adhesions	Fibrous tumors of the uterus.	Hemorrhage		Four uterine tumors, with thick vascular pedicles; extensive adhesions. Hemorrhage from slipping of ligature.	Not yet reported.
133	1843	Mr. Heath	46				1					17 hours	Dis. of uterus		Uterine tumor	Hemorrhage		The uterus, tumor and all were removed.	London Medical Gazette, December 8, 1843, p. 309.
134	1843	Mr. Lane	28				1											Phlegmasia dolens followed operation.	London Medical Gazette, 1844-5, p. 84.
135	1843	do	45				1											Two tumors; cyst emptied before removal.	Med. Chi. Trans., Phillip's Table.
136	1844	do	33				1											Cysts emptied before removal.	London Med. Gaz., 1844-5, p. 84, Mr. Jeaffreson's Table.
137		do					1												Phillip's Table.
138		do					1												Med. Chi. Trans., Phillip's Table.
139		do					1												Ibid.
140	1841	Mr. Key	19				1					9th day			None	Multilocular cysts	Peritonitis	Large vessels on tumor.	From Gay's Hospital Rep., Oct., 1843, p. 473.
141	1843	Mr. Greenhow	29				1					7th day	Dis. of stom.	Adhesions	Dense and vascular cellular tumor.	Peritonitis		For four years previously frequent uterine hemorrhage.	Med. Chi. Trans., v. 27, p. 88; and Amer. Jour. Med. Sci., July, 1844, p. 251.
142	1843	Mr. B. Cooper	32				1					7th day	Mal. dis. uterus	Adhesions	Solid and cysts	Peritonitis		A portion of omentum had been included in ligature.	Ibid., v. 27, p. 76; & Am. Jr. Md. Sci., April, 1844, p. 462.
143	1844	W. L. Atlee	61				1					6th day			None	Bilocular cyst	Insidious peritonitis	Colon involved in a broad pedicle; feeble constitution; both ovaries diseased.	Amer. Jour. Med. Sci., July, 1844, p. 43.
144	1844	do	24				1								None	Fibrous tumor of the uterus.		Thick fleshy pedicle, followed by violent peritonitis; intestines troublesome.	Ibid., April, 1845, p. 300.
145	1849	do	29				1								Procidentia uteri.	Adhesions	Fibrous tumor of ovary.	Extensive and very firm adhesions to the bones of pelvis, and to the iliac vessels. Pouparts ligament was imbedded in the tumor and stretched across it. Complete procidentia uteri. Chloroform, 1 part, ether 2 parts, administered. Recovery without an unpleasant symptom. Pregnant twice since.	Ibid., October, 1849, p.
146	1849	do	33				1						Dis'd uterus.	None	Uterine tumor			The uterus was large, and in consequence the tumor was not removed. The right ovary was as large as an orange, and the left one was also diseased. Chloroform mixture was administered. Rapid recovery. Operation exploratory.	Ibid., April, 1850, p. 318.
147	1849	do	25				1								Adhesions	Cystiform		One large cyst extensively adherent to the omentum by its anterior surface; four gallons of chocolate-colored fluid removed one week before. Weight of tumor 40 lbs. Chloroform mixture given. Rapid recovery; has been married since, and given birth to a fine healthy daughter, after an easy parturition of three hours.	Ibid., p. 328.
148	1849	do	43				1						Dis. of uterus.	None	Uterine tumor			The uterus was enlarged, and had large cystiform bodies incorporated with it; therefore not removed. Chloroform mixture; rapid recovery. Operation exploratory.	Not yet published.
149	1849	do	39				1								None	Fibrous tumor of uterus		Fibrous tumor attached to the fundus of the uterus with a dense pedicle. Weight, 6 lbs. Chloroform mixture. Rapid recovery. Menses on at the time of the operation, and continued without interruption.	Not yet published.
150	1850	do	39				1					6th day			Adhesions	Cystiform	Peritonitis	Extensive adhesions. Two round strong cords; vascular; bound the cyst to the recto-vaginal cul-de-sac; requiring ligatures. Patient ate an orange on the fourth day, up to which time she was doing well, and soon after, inflammation set in. Chloroform mixture.	Not yet published.
151	1850	do	48				1					3d day	Obs. bowels, death impending.		Adhesions	Cystiform	Exhaustion	Multilocular. The cyst was inseparably adhering to the intestines and uterus. The adherent portions of the cyst were detached from the main tumor, and permitted to remain; spermatic artery cut and tied. Weight of tumor, 28 lbs. Chloroform mixture. The bowels had been obstructed for several days before the operation, and it was performed only with the view of arresting the progress of approaching death.	Not yet published.
152	1850	do	40				1					3d day	Ascites and anasarca—death impending.		Adhesions	Cystiform	Exhaustion	The cyst weighed 25 lbs., exclusive of about 2 gallons serum from the peritoneal cavity. Had been tapped for a cetes five times before. The lower limbs were anasarca and ulcerated, weeping away immense quantities of water. The omentum was much thickened oedematous and spread over the cyst, and almost inseparably adherent. Chloroform mixture. The operation was performed with the same view as the case above.	Not yet published.
153	1850	do	41				1						Dis. of uterus.	None	Uterine tumor			The uterus was enlarged; chloroform mixture, which produces a state of catalepsy, rendering it extremely troublesome with the intestines; they were forced out, and could scarcely be returned. A favorable recovery and improved health, since the operation.	Not yet published.
154	1850	do	37				1								Adhesions	Cystiform		Multilocular; adhesion to the whole front of the abdomen and to the omentum. A part of the cyst, as large as the palm of the hand, was left adhering to the walls of the abdomen at the umbilicus. Pedicle three or four inches long; round; as thick as the little finger. First cut the pedicle, and then picked out the vessels and tied them separately. Weight of tumor, 25 lbs; chloroform mixture; rapid recovery.	Not yet published.
155	1850	do	42				1						Anaemia		Adhesions	Cystiform		Multilocular; patient greatly enfeebled by previous disease. Adhesions to the intestines, bladder, rectum, and whole basin of the pelvis. A part of the cyst was left attached to the colon. Several cysts broke while removing them, and flooded the intestines. Pedicle about 4 inches long, and slender. Applied no ligature; used torsion. Pulse 130 at time of operation. Tumor 15 lbs; chloroform mixture; rapid recovery.	Not yet published.
156	1850	do	28				1					80 days	Pregnancy		Adhesions	Cystiform	Inanition from pregnancy.	Multilocular; had been tapped 16 times—56 lbs. fluid removed at one tapping. The mass of the tumor was composed of two immense cysts. It weighed 81 lbs.; chloroform mixture. She was two months gone in pregnancy, at the time of the operation. She recovered from the operation, but this was followed by great irritability of stomach, in consequence of the stage of pregnancy, that she could not be nourished. She died 80 days after the operation. No miscarriage.	Not yet published.

No.	Date.	Operators.	Age.	INCISION.			RECOVERED.			DIED.			Time of Death.	Other Diseases.	Adhesions.	Character of Tumor.	Cause of Death.	REMARKS.	REFERENCES.	
				Minor.	Major.	Unknown.	Minor.	Major.	Unknown.	Minor.	Major.	Unknown.								
157	1851	W. L. Atlee	29		1			1					3d day		Adhesions	Cystiform		Multilocular; incision from the sternum to pubis, nearly. Very firm and extensive adhesion; some peritoneal effusion. Pedicle about 6 inches broad, filled with varicose veins; one of the cysts, a large one, was filled with partially washed crassamentum. Weight of tumor, 35½ lbs.; chloroform mixture.	Not yet published.	
158	1844	F. Bird	21	1		1										Cyst		Contents of cyst, a light amber color, and contained a very small quantity of albumen.	London Med. Gaz., August 16th, 1844.	
159	1844	Ehrhartstein	36		1										None			Tumor tapped before extraction.	Med. Chi. Trans., vol. 27, p. 472. Phillip's Table.	
160	1844	do													None			Serum continued to drain for some time from the aperture; afterwards changed to milky fluid, flowed for 9 weeks; wound cicatrized, and patient cured.	Arch. Generale de Medicene, v. 31, p. 427; also, Amer. Jour. Med. Sci., July, 1847, p. 233.	
161		Dr. Hooper	47		1								30 hours		Adhesions	Solid tumor		Ascites.	Ibid.	
162		do	38		1										Adhesions	Solid		Malignant tumor.	Ibid.	
163		McDonald			1										None				Ibid.	
164		Clorissmann			1										None				Ibid.	
165		Dr. Groth			1										None				Ibid.	
166		Morgan		1														Portion of fluid removed before extraction.	Ibid.	
167		A. B.		1															Ibid.	
168		C. D.	22	1		1									Adhesions				Med. Chir. Trans., v. 27, p. 474.	
169		E. F.		1															Ibid.	
170		G. H.		1											Adhesions				Ibid.	
171		Cascin Gooch		1															Ibid.	
172	1839	Guy's Hospital		1															Ibid.	
173	1844	Dr. Bowles	26		1										Adhesions	Cysts		Not removed on account of adhesion.	Ibid.	
174	1844	Prof. Webster	37		1										Adhesions	Solid			London Medical Gazette, 1844-5.	
															Adhesions	Cyst		Not removed on account of adhesions. The patient rapidly recovered from the operation, after the peritoneal cavity had been exposed for 2 hours, but the disease progressed, and she died from the disease in about 2 months.	Western Lancet, October, 1844.	
180	1844	W. B. Page, Esq.	33		1										None	Cystiform		Cyst opened and withdrawn; hand introduced into abdomen.	London Lancet, April 5, 1844, p. 397; Lond. Ed., p. 84.	
181	1846	do	39	1									12 hours		Adhesions	Cystiform	Hemorrhage	Cyst first tapped and withdrawn, adhering to several inches of intestine; but previous to the operation supposed to be non-adherent. A portion of the pedicle escaped from the ligature, whence arose the hemorrhage.	Ibid., December 12th, 1849; also, Ranking's Abstract, Amer. Ed., No. 5, p. 251, 1847.	
182	1845	Mr. Jno. Dickin	18		1										Adhesions	Cystiform		Multilocular; weight 28 lbs.; recovered without a bad symptom. Ligature came away in 3 weeks; each artery tied separately. The whole pedicle was not included; left ovary healthy. Lee on tumors of the uterus, p. 271.	Provincial Med. and Surg. Journal, Oct. 7th, 1845; also, Ranking's Abstract, No. 2, p. 188.	
183	1845	Mr. G. Southam	38		1										None	Cystiform		Exploratory incision; cyst punctured, and removed 16 or 18 pints clear lemon-colored fluid. Hand introduced, &c. Ligature came away on the 49th day. Left ovary tapped and extracted. Lee, p. 269.	Ibid., Sept. 10th, 1845, p. 561; also, Braithwait's Retrospect, No. 12, p. 247.	
184		do			1								6th day					Multilocular cyst cerebiform solid matter; operation performed under apparently favorable circumstances.	Arch. Gen. de Medicene, v. 20, p. 92; also, Amer. Jour. Med. Science, July, 1847, p. 234.	
185	1845	Dr. Handyside	20		1								70 days					Both ovaries diseased and extracted. Lee, p. 271.	Edinburgh Med. and Surg. Jour., 1846; also, Am. Jour. Med. Science, April, 1846, p. 582.	
186		Dr. Huynz			1								40th day		Adhesions			Exhaustion	Removal impossible from adhesion. No regard paid to temperature during the operation.	Brit. and For. Med. Rev., Forbes' Amer. Ed., Jan., 1847
187		do			1								6 weeks		Adhesions			Peritonitis	A portion of omentum removed with the tumor. No regard paid to temperature during the operation.	Ibid.
188	1846	Mr. Solly	20	1									11 hours		None	Cystiform	Hemorrhage internal	Unilocular cyst of right ovary. Hemorrhage from slipping of the ligature. A portion of the pedicle, containing the fallopian tube, slipped from the ligature, and gave rise to hemorrhage. Lee, p. 271.	Ibid.; also, Lond. Med. Gaz., July 10th, 1846; also, Braithwait's No. 14, p. 317.	
189		Mr. W. (B. K. S. W.)			1								6th day	Exhaustion before operation.	None			Peritonitis	Exhaustion after 3d day. Great tympanitis.	Lee's Tables.
190	1846	Dr. Pr. Smith	39		1									Chm. peritonis	None			Shock of Operation	Multilocular; tumor weighed 20 lbs.; fluid 10 lbs. Lee, p. 271.	Phillip's Table.
191	1846	Mr. Arrowsmith	22	1									4 hours		Adhesions	Cystiform		Multilocular. An exploratory incision was made, of a few inches in extent, but the adhesions were so strong and extensive that the operation was considered unjustifiable. The patient recovered without any untoward symptom. Lee, p. 271.	Ibid.	
192	1846	H. E. Burd	26		1										None	Uterine tumor		Multilocular. The largest cyst was tapped after the incision, and over 8 gallons of glairy fluid were withdrawn; after which the tumor was drawn out. The pedicle was very thick, requiring three strong twine ligatures in different compartments, and each vessel was separately tied besides. Much alarm was excited subsequently, by the frequent occurrence of collapse, which was relieved by the administration of opium, ammonia and brandy. Tumor weighed 50 lbs. The patient was 3 or 4 months gone in pregnancy, and aborted 40 hours after the operation.	London Medical Gazette, April, 1847; also, Ranking's Abstract, No. 5, p. 249.	
193	1846	Mr. W.			1															Lee's Table.
194	1846	Cæsar Hawkins	18	1											None	Cystiform		Unilocular; ligature came away from the 22d to the 25th day. The wound entirely healed on the 29th day. Lee, 271.	Medical Gazette, October 30, 1846.	
195		R. D. Mussey			1										Adhesions			Operation not completed in consequence of numerous and extensive adhesions. Wound kept open by a tent. A clear liquid came away for several days, then became purulent; diminished gradually; ceased in three weeks; closed and the tumor disappeared. A year after, she was delivered of her 14th child, and no return of disease.	Amer. Jour. of Medical Science, February, 1838, p. 380.	
196		do			1										None	Cyst		Unilocular; incised the abdomen, and drew off the fluid from the sack, which was secured to the edges of the wound by suture; wound kept open by tent. The discharge gradually diminished, being first serum, then purulent. The patient's health improved, and for nearly a year she was able to see to her domestic affairs. The wound finally closed, the disease returned and the patient died, about 13 months after the operation.	Case not reported.	
197	1850	do			1								14 hours	Uterine				Tumor in the walls of the uterus.	From Dr. Mussey, himself.	
198		Mr. Bainbridge	31	1														Unilocular; incision 3 inches long; a portion of cyst drawn out, contents evacuated; a portion of cyst then removed; its edges fixed to outer wound, and suppuration induced. On the 5th day the discharge became purulent, and was maintained so by stimulant injections. In three months the discharge gradually diminished, and the general health completely restored.	Prov. Med. Jour., December 16, 1846; also, Braithwait, No. 5, p. 250.	
199	1844	Dr. Waseikowski	40		1													Tumor, weight 6½ lbs. Woman in labor; had gone 3 months beyond her time. Uterus forced out of the vagina. Trocar removed 35 litres of yellowish fluid; day after removed tumor, by incision from 3 inches above the umbilicus to the pubis, and 30 litres more of fluid. No bad symptoms followed. She walked home to a neighboring house, the 24th day afterwards. She was delivered of a boy 13 months afterwards, and of another boy in December, 1846.	Jour. de Medicene, v. 20, p. 92; also, Amer. Jour. Med. Sci., July, 1847, p. 233.	
200	1835	Dr. Bellinger	35		1									Hydrop. uteri				Incision extended nearly to the enciform cartilage. The tumor was removed by cutting the round and broad ligaments. Two arteries of considerable size were tied with animal ligature, both ends of which were cut off close to the knot. A tumor in the pelvis was formed by the uterus distended with a large quantity of dark watery fluid, which escaped through a rent made by the finger, through its posterior walls. The patient was still living and in good health, May, 1847. Menstruation never returned.	South. Jour. Med. and Phar., May, 1847; also, Wood's Quarterly Retrospect, July, 1847, p. 21.	
201		do			1										Adhesions	Several tumors		Operation abandoned, from finding the peritoneum adherent along the line of the incision.	Ibid., p. 22.	
202	1846	Anonymous	30 } 35 }		1								5th day	Dis. of uterus	None	Uterine	Peritonitis	The lateral ligaments were divided; were very vascular, and many arteries were tied with animal ligatures; the neck of the uterus was cut across about ¾ of an inch above the os tincæ, and the entire tumor (the uterus) removed. There had been no internal hemorrhage.	Ibid.	
203	1849	Dr. S. Parkman	27		1								12 hours			Uterine	Hemorrhage	Fibrous tumor developed in the fundus of the uterus; weighed nearly 9 lbs. Pedicle formed by the body of the uterus, about two inches in diameter. The left fallopian tube removed with the tumor. Both ovaries were left in the pelvis; supposed to have contained fluid; attempts at tapping; wrong diagnosis; contained only a clear serous fluid in its interstices or meshes. Half of the pedicle was not firmly constricted by the ligature, and patient sunk from hemorrhage.	Amer. Jour. Medical Science, April, 1847, p. 371.	
204	1701	Dr. R. Houstown	58		1													With a strong fir stick wrapped with lint, and thrust into the cyst, twining and winding it, above two yards long of a substance like glue, was drawn out, which was followed by 9 quarts of such matter, and several hydatids larger than an orange. Dressed by the interrupted suture. Lived in perfect health until 1714.	Philosop. Trans., Abridged, vol. 7, p. 541.	
205	1848	Dr. H. Miller	37		1										Adhesions	Cystiform		Multilocular; extensive adhesions to omentum and parietes; easily loosened. Two cysts punctured to diminish the size, from which flowed a whitish ropy albuminous looking fluid, from which tumor weighed 9½ lbs. Chloroform employed.	Lond. Med. Gaz., June 23d, 1845; also, Amer. Jour. Med. Science, July, 1849, p. 203.	
206	1843	Dr. J. Deane	46		1													Fibrous tumor of the uterus. The tumor, whose length was about 8 or 9 inches, and breadth 5 or 6 inches, embraced the entire left half of the uterus, from which it was considered impossible to separate it, and therefore the further prosecution of the operation was deemed impracticable. Chloroform was used, but its effects were considered very deleterious. The symptoms for several days were very alarming, but she finally recovered from the greatest perils.	Boston Med. and Surg. Jour., Oct. 11th, 1848; also, Amer. Jour. Med. Science, Jan., 1849, p. 253.	

Cases No. 175, 176, 177, 178 and 179 not found in copy.



No.	Date.	Operators.	Age.	INCISION.			RECOVERED.			DIED.			Operation unfinished.	No. Tumor.	Time of death.	Other diseases.	Adhesions.	Character of Tumor.	Cause of Death.	REMARKS.	REFERENCES.
				Minor.	Major.	Unknown.	Minor.	Major.	Unknown.	Minor.	Major.	Unknown.									
207	1849	Dr. J. Deane	45		1						1			12th day		None	Cystiform	Inflam. and gang. of the intestines.	Incision from pubis to sternum, nearly, which was completed after tapping the cyst, which collapsed into the pelvis. The small intestines were brightly injected, congested or inflamed. In other respects all the viscera healthy. Things went on well for a week, and on the 4th day union was accomplished. Afterwards, inflammation of the intestines, constipation and vomiting supervened, and death. A post mortem examination revealed gangrene of the intestines, and parts adjacent to the pedicle. Weight of tumor, 40 lbs.	Dr. James Deane's letter to me, (Prof. W. L. Atlee,) dated, Greenfield, Mass., April 8th, 1851.	
208	1848	P. J. Buckner	32		1		1										Cystiform		Multilocular. The large sack being extensively adherent to the peritoneum, it was unavoidably punctured, and 18 pints of a dirty, opaque cream-colored fluid, of the consistence of pus, discharged. After this, the posterior wall of the cyst was divided, and several bands of organized lymph, severed by the knife. The sack was now torn from its adhesions, which extended above and below the umbilicus, and across the linea semilunaris. The patient remarked, that the act of separating the sack from the walls of the belly, reminded her of tearing the leaf-lard from pork. From another cyst, 6 pints of fluid were removed; this was another division of the tumor adhering extensively to the peritoneum, which was ruptured and bled freely, requiring ligatures. The pedicle was on the right side, embracing the fallopian tube and ovary, and about two inches in diameter, and highly vascular. Patient did not lose 4 ounces of blood. Operation lasted 1 1/2 hours. Tumor weighed 28 1/2 lbs. Solid part, 4 1/2 lbs.	Western Lancet and Hosp. Report, October, 1848.	
209	1848	do			1		1										Fibrous tumor		Weight of tumor, 4 1/2 lbs.	Ibid., p. 213.	
210		do			1						1			6th day				Peritonitis		Amer. Jour. Med. Science, October, 1849, p. 563.	
211	1850	do	39		1		1										Adhesions	Cystiform		Ibid., from Ohio Med. and Surg. Jour., September, 1850.	
212	1850	Dunlap & Buckner			1		1										Adhesions	Cystiform		Not yet published.	
213	1848	Dr. Meeker	32		1						1			6 hours		Adhesions	Cystiform	Hemorrhage	Multilocular. Incision 22 inches, adhesions to omentum parietes, and small intestines, readily separated. Pedicle to the right ovary, 2 1/2 inches broad, and two inches in length. Operation lasted 45 minutes. The ligature slipped from half the pedicle, the cavity filled with fluid blood, and the patient was exhausted by the hemorrhage. The tumor weighed 40 lbs., 8 ounces.	Supplement to Boston Med. and Surg. Journal, September 6, 1848, p. 116.	
214		M. H. Larry				1		1										Pilous ovarium			
215		Mr. Arnott			1									74 hours		Adhesions	Cystiform		Multilocular. Extensive adhesions, on which account the operation was not completed. Dr. Clay attributes the failure, to the small incision.	Mem. del' Acad. Roy. de Md. Tr. xii, & xii, Paris, 1646-48. Med. Gaz., Abstract Rec. No. 12; also, Ranking's Abstract, December, 1848, p. 246.	
216	1847	Dr. Vangirard	25		1		1											Cystiform		Tapped 50 times in 3 1/2 years. The cyst was first tapped and then removed. The tumor was cartilaginous, gelatinous and cerebriiform. Weight, 9 lbs.; ether inhaled. Incision to the left of, and parallel with the linea alba. The pedicle was attached to the left side. The patient was well at the end of a month.	Jour. de Scien. Medico—Chireng; also, Rankin's Abstract, December, 1848, p. 246, and No. 9, p. 275.
217	1849	Prof. Bayless			1		1											Extra uterine		Tumor weighed 18 lbs.	Trans. of Amer. Med. Association, vol. 5, p. 379.
218		Anonymous			1						1									Tumor weighed 26 lbs.	Baltimore paper.
219		M. Sneffard	40		1		1									Adhesions			The tumor of an extra uterine fetus. This was the 6th pregnancy. The fetal movements were perceptible up to the 6th month; then ceased, and were followed by excessive pain and hemorrhage. Six months after this, the tumor was discovered in the right groin. At the operation, two cysts were removed; one was filled with adipocier, closely adherent, and forming one cavity into the uterus, its fundus being destroyed. The other cyst contained the limbs and trunk of a fetus, the head and upper extremities having escaped into the general cavity of the peritoneum, and adhering firmly; both were removed. Ether was administered; patient recovered.	Rankin's Abstract, vol. 9, p. 279; also, Amer. Jour. Med. Science, October, 1849, p. 522.	
220	1849	Jno. Crouch, Esq.	24		1		1									Adhesions	Cystiform		Multilocular; left ovary. Weight of tumor 14 lbs. Rapid recovery. The cysts tapped before their removal.	Prov. Med. and Surg. Jour., Sept. 5th, 1849; also, Amer. Jour. Med. Science, January, 1850, p. 221.	
221	1849	F. Elkington	31		1		1									Adhesions	Cystiform		Multilocular. Adhesions slight; pedicle very broad and vascular. Weight of tumor, 40 lbs. Rapid recovery.	Ibid., Sept. 19th, 1849; also, Amer. Jour. Med. Science, January, 1850, p. 220.	
222	1849	W. H. Vanburen	21		1		1								Uteri proidentia.	Adhesions	Fib. tumor of the ovary		Complete proidentia uteri. Three vessels of the omentum tied, and ligatures cut off close to the knot. About 20 ounces of serum in the peritoneal cavity. Weight of tumor, 7 lbs.; circumference, 23 inches. Chloroform mixture.	New York Jour. Med. and Collat. Science, March, 1850, p. 159.	
223	1849	J. H. Bigelow	22		1		1											Cystiform		Multilocular; had been previously tapped. A multilocular tumor was first removed, weighing 8 lbs. Afterwards, a fibrous tumor was removed from deep in the pelvis—weight, 1/2 lb.	Boston Medical and Surg. Jour., Jan. 23, 1850, p. 503.
224	1849	A. March	49		1		1										None	Cystiform		The pedicle was severed half an inch from the ligature, which slipped off while sponging the wound, and alarming hemorrhage ensued. The vessel was immediately secured by transfixing the pedicle with an armed ligature. "Weight of tumor, 18 lbs. Chloroform mixture given.	Dr. March's Pamphlet.
225	1847	D. Prince	25	1			1									Adhesions	Solid ovarian tumor		The tumor was adherent to the anterior walls of the abdomen; incision 3 inches. A free incision was made into the tumor, and a portion of its interior removed. The wound was left open, and a test introduced. A large amount of pus was discharged, for a long time, and the tumor entirely disappeared. April 10th, 1849, a healthy daughter born.	Amer. Jour. Med. Science, July, 1850, p. 217.	
226	1850	A. H. Grimshaw	37		1						1			5 hours		Adhesions	Cystiform	Hemorrhage	Had been tapped 9 times—last time, 10 days before the operation; 5 gallons of fluid removed; walls of the abdomen.	London Lancet, Nov. 23, 1850; also, Amer. Jour. Med. Science, January, 1851, p. 226.	
227	1850	W. E. Duffin	38	1			1									None	Cystiform		Incision 3 inches; cyst tapped, and drawn out 180 ounces of brown colored,ropy viscid fluid. Woman as large as the 8th month of pregnancy.		
228	1850	C. R. Peaslee	25		1		1									Adhesions	Cystiform		Both ovaries removed; weight of left ovary, with the contents, 29 lbs. of right, 10 drachms. Menses returned 72 hours after operation, and continued three days; 4 ligatures—3 came away by matting themselves, and one with a loop. Unmarried; anaesthesia.	Amer. Jour. Med. Science, April, 1851, p. 371.	
229	1850	Geo. Norman	23		1		1									Adhesions	Solid ovarian tumor		Operation unfinished, in consequence of adhesions. Tumor diminished to half its size, rapidly, after the operation.	Prov. Med. and Surg. Jour., January, 5, 1851; also, Amer. Jour. Med. Science, April, 1851, p. 509.	
230	1850	J. Farrel	28		1						1					Adhesions	Cystiform		Tapped five times previous to operation. Incision 8 1/2 inches. Extensive adhesions existed, to the intestines, uterus, and walls of the abdomen; also, at one of the sites of a previous tapping. It was deemed improper to attempt its removal. The wound healed, except where a broad meche was introduced into the sack. The discharge, which was at first serous, became purulent and foetid, continued until her death, which took place 22 months from first tapping. Left ovary diseased.	Reported to committee of Ohio State Medical Society, 1851, on Ovarian Disease—Not yet published.	
231	1850	J. Farrel			1		1									None	Multicular		Cyst opened and emptied before removal; with it, many smaller ones were situated in the right ovarium. The pedicle and fallopian tubes were secured by ligature, and the tumor removed. Incision 10 inches. Ligature came away 32d day; wound healed very slowly by granulation.	Ibid.	
232		A. Dunlap			1		1							Sev'l weeks after oper'n				Multicular	Diabetic and disease of kidneys.	This case is given by P. J. Buckner, from the statement of a medical friend. The particulars of the case not known to Reporter.	Case has never been reported.
233		E. K. Chamberlain				1		1			1			Immed'tly							From Dr. Chamberlain, himself.
234	1848	H. G. Potter	36		1									17th day		Adhesions	Cystiform	Diarrhoea and peritonitis.	Both ovaries diseased. Incision from 20 to 24 inches long. Both cysts emptied before being tied; only two-thirds of the right cyst removed. Autopsy discovered that this tumor was firmly united to the bladder, uterus and walls of the pelvis. Peritonitic symptoms were almost entirely absent, although the bowels were found agglutinated together, and several ounces of pus were found in the pelvis. Chloroform was administered, and with entire relief from pain.	Lond. Med. Gaz., new series, vol. 6, 1848, p. 1066; also, Amer. Jour. Med. Science, July, 1849, p. 205.	
235		Franckenau			1		1														"Each report a case of extirpation of the ovary, which did not prevent the females from enjoying excellent health afterwards."
236		Pott			1		1														The Lancet, 1828, vol. 1, p. 878; also, Velpeau's Surgery, p. 538, by Mott.
237		Lassus			1		1														
238		M. Deneux			1		1														
239		M. Grath			1		1							16 hours							L'Experience, T. I, p. 631; also, Velpeau's Operative Surg. p. 538; by Mott.
240		do			1		1				1			8 hours	Epip. cribbled with tumors.						Ibid. p. 632; Ibid. p. 538.

