

NATIONAL LIBRARY OF MEDICINE



NLM 00107175 2

Surgeon General's Office


LIBRARY

ANNEX

Section, _____

No. _____

107290.



THE DRUGGIST'S
GENERAL RECEIPT BOOK

COMPRISING A COPIOUS

VETERINARY FORMULARY

NUMEROUS RECIPES IN

PATENT AND PROPRIETARY MEDICINES

DRUGGISTS' NOSTRUMS, ETC.

PERFUMERY AND COSMETICS

BEVERAGES, DIETETIC ARTICLES, AND CONDIMENTS

Photographic Chemicals and Formulæ, Trade Chemicals,
Scientific Processes

AND AN APPENDIX OF USEFUL TABLES

BY

HENRY BEASLEY,

AUTHOR OF "THE POCKET FORMULARY AND SYNOPSIS OF THE BRITISH AND
FOREIGN PHARMACOPŒIAS, AND OF "THE BOOK OF PRESCRIPTIONS,
COLLECTED FROM THE PRACTICE OF THE MOST EMINENT
PHYSICIANS AND SURGEONS."

Ninth Edition

PHILADELPHIA
P. BLAKISTON, SON & CO.

1012, WALNUT STREET

1886



QV

B368d

1886

film no. 10930

Uem 2

P R E F A C E

THE contents of the present volume have been carefully revised. They have also been enlarged by the addition of useful matter, obtained from various sources, and published since the issue of the preceding edition.

The section on Patent Medicines and Druggists' Nostrums has been amplified.

In short, the endeavour has been made to carry out the aim of the originator of the work, by rendering it a useful, convenient, and trustworthy Manual for the Chemist and Druggist.

London, March, 1886.



CONTENTS

	PAGE
VETERINARY MATERIA MEDICA	1
VETERINARY FORMULARY	43
Medicines for Horses	43
Balls and Ball Masses	44
Electuaries, Confections, &c.	60
Powders	61
Liquid Medicines—Drenches	65
Tinctures, Solutions, &c.	77
Clysters	78
External Applications—Liniments	80
Embrocations, various	85
Lotions, or Washes	86
Caustics	89
Fomentations	91
Eye Waters	92
Sundry Solutions	93
Poultices	94
Ointments	96
Charges	106
Powders (for outward use)	107
Medicines for Cattle	109
Drenches, &c.	109
————— for the Foot-and-Mouth Disease	118
————— for the Cattle Plague	121
————— for Calves	123

	PAGE
External Applications—Lotions	125
Eye Waters	126
Embrocations and Liniments	127
Ointments, &c.	128
Medicines for Sheep and Lambs	131
Drenches, &c.	131
External Applications—Eye Waters	136
Washes	136
Powders and Ointments	137
Medicines for Swine	141
Medicines for Dogs	143
External Applications	151
Medicines for Poultry and Rabbits	155
PATENT AND PROPRIETARY MEDICINES, DRUGGISTS'	
NOSTRUMS, &c.	158
Factitious Mineral Waters, and Salts for producing them	198
PERFUMERY	206
Distilled Waters	206
Spirituous Waters, Eaux, Esprits	207
Tinctures used in Perfumery	214
<i>Extraits</i> , Essences	215
Common French Essences, Spirits and Waters	219
Ammoniated Perfumes	219
Acetic Perfumes	220
Pot Pourri, Scented Powders, &c.	222
Pastiles	224, 225
Cosmetics	227
Skin Cosmetics—Lotions	227
Pastes, Pomades, Cold Cream, &c.	229
Face Paints	232
Toilet Soaps	233
Shaving Compounds	236

	PAGE
Hair Cosmetics	237
Compounds to promote the Growth of the Hair	237
Liquid ditto	239
Pomatus, Lotions, for embellishing the Hair, &c.	241
Huiles Antiques	244
Washes for Cleansing the Hair	246
Compounds for Stiffening the Hair	247
Hair Dyes	248
Depilatories	251
Teeth and Mouth Cosmetics	252
Tooth Powders	252
Tooth Pastes	256
Astringent Tinctures—Mouth Washes	258
Tinctures, &c., for Toothache	260
Pastes for Toothache	262
Cements for Stopping Teeth, &c.	262
BEVERAGES, DIETETIC ARTICLES, AND CONDIMENTS	266
Beverages, and Powders for preparing them	266
Dietetic Articles—Directions for using	274
Preparations of Cod-liver	286
Condiments, &c.	288
Curry Powders, Mixed Spices, &c.	288
Culinary Essences, Tinctures, &c.	290
Culinary Vinegars, Sauces, &c.	291
Pickles	294
TRADE CHEMICALS ; Miscellaneous Preparations, Processes and Compounds used in the Arts, in Domestic Economy, Chemical Research and Recreation, &c.	298
APPENDIX.	
Tables of Weights and Measures	459
Comparison of Thermometric Scales	462
Effects of Temperature	463
Specific Gravities	464

APPENDIX.

Table of Chemical Equivalents	466
Table of Solubilities	467
Table of Composition, Equivalent numbers, and Formulæ of some of the more important Com- pounds employed in Pharmacy and the Arts	474
Table of Neutralizing Proportions, &c.	478
Specific Gravities corresponding to degrees of Baumé's hydrometer for liquids heavier than water	479
Specific Gravities corresponding to degrees of Baumé's hydrometer for liquids lighter than water	480

INDEX	481
-----------------	-----

VETERINARY MATERIA MEDICA

Table of the Properties and Doses of the Principal Medicinal Substances used in Veterinary Practice.

N.B.—Where the doses are given without mentioning the animal intended, it must always be understood to refer to the Horse.

ACACIA. See GUM ARABIC.

ACETATE OF AMMONIA SOLUTION. *Spirit of Mindererus.* Diaphoretic and diuretic. It is also regarded as antiseptic. Dose, for horses and cattle, from 5 to 10 oz. For smaller animals, from 5 to 10 drs. Externally, in strains, ophthalmia, and as a discutient.

ACETATE OF COPPER. See VERDIGRIS and COPPER.

ACETATE OF LEAD. Astringent and sedative; in doses of 30 to 40 grains with opium; in internal hæmorrhage, chronic diabetes, and diarrhœa; but chiefly used externally, in cooling lotions, eye-waters, ointments, &c. For DIACETATE OF LEAD, see GOULARD'S EXTRACT. As antidotes for an overdose, give Epsom or Glauber's Salts, with opiates if required.

ACETATE OF POTASH. Diuretic and cooling; dose for horses and cattle, 2 oz. In much larger doses it is a laxative, but not to be depended on.

ACETATE OF SODA. Similar in properties and uses to Acetate of Potash.

ACETIC ACID. Strong acetic (or pyroligneous) acid acts as a rubefacient and caustic, but is rarely employed for this purpose. In the weaker forms of common or distilled vinegar, or diluted wood vinegar, it is frequently used. See VINEGAR.

ACIDS. See MURIATIC ACID, NITRIC ACID, PRUSSIC ACID, SULPHURIC ACID, &c.

ACONITE. Generally given in the form of tincture prepared according to the formula of Dr FLEMING. Dose for the horse, about 10 minims; for cattle, 10 to 20 minims; for sheep and dogs, 1 to 2 minims. Aconite is a most useful and effective sedative and antiphlogistic in febrile attacks and acute local inflammation; hence it is very serviceable in pneumonia, pleurisy, bronchitis, &c. In over-doses it is a most active and *potent poison*.

Antidotes.—Charcoal. *Emetics.* Tannic Acid.

ACUPUNCTURATION. Used in some spasmodic and paralytic affections.

ÆGYPTIACUM. A preparation of verdigris and honey. A mild caustic, used as a local application to ulcers of the mouth, running thrush, grease, &c. Internally *poisonous*.

ETHER. See ETHER.

ALCOHOL. *Poisonous* to all animals—2 drachms will kill a dog. See SPIRITS, ARDENT.

ALLSPICE. *Pimento.* A useful stimulant and carminative; used in cordial balls and drinks, and to correct the action of purgatives. Dose for horses, 2 to 4 drs.; cattle, $\frac{1}{2}$ oz. to 1 oz. Dose of the Tincture, 4 oz., in gripes.

ALOES. The most valuable purgative for the horse, but not to be depended on for cattle and sheep. A horse requires from 4 to 8 drs. of Barbadoes aloes, from 5 to 9 drs. of Socotrine, and from 6 to 10 of Cape. Mr YOUATT says 3 dr. of Barbadoes are equal to 4 of Cape aloes. Mr MORTON considers a mixture of equal parts of Cape and Barbadoes aloes to be quite as efficacious as the latter alone. But the fine gourd Barbadoes aloes are the most certain in their operation. If the animal be prepared by previous mashes, 5 drs. are generally, and 6 drs. almost always, sufficient. Mr BLAINE recommends 2 drs. every 6 hours, till 8 drs. have been taken, as a nauseant and purgative; but Mr YOUATT strongly disapproves of this plan, particularly in inflammation of the lungs. Aloes require from 18 to 36 hours to produce their effect, during which time the horse should not be ridden far or fast. Though not to be depended on for cattle, 4 to 6 drs. are sometimes added to the purgative salts. Large doses (in some cases sufficient to destroy life) have been given to sheep without purging. Small dogs require from 15 to 30 grs.; medium-sized ones

a dr. ; some larger ones require 2 drs., or more. Hogs can bear but a few grains. Externally, in the form of tincture, aloes is used as a stimulating application to wounds. Mr FINLAY DUN says: "The administration of aloes should be avoided in cases of irritation or inflammation of any part of the alimentary canal, and in piles or hæmorrhage from the rectum. In bronchitis and other inflammatory affections of the mucous membranes, and in inflammation of the kidneys, it must be used with great caution, and in very small doses, for in such cases the intestinal mucous membrane is usually irritable, and superpurgation and inflammation are readily induced. During pregnancy, both in the mare and the bitch, the violent operation of aloes must be carefully avoided. Some practitioners give it both to foals and calves, but for young animals, linseed oil or castor oil is more suitable."

ALUM. Astringent and styptic. Given in doses of 2 to 4 drs. to horses in diabetes and diarrhœa ; but BOURGELAT says that its too frequent use induces a phthisical condition. A dose of alum whey, consisting of 2 drs. of the powder in a pint of hot milk, may be given after excessive purging. Cattle require from 2 to 6 or 8 drs. in diabetes and red water ; and from 2 to 4 drs. are given to cows to dry their milk. To calves and lambs it is given in dr. doses, in warm milk, for diarrhœa, &c. Dogs, 10 to 15 grs. Externally it is applied to cracked and greasy heels, joint wounds, sore mouths, inflammation of the eye, chronic discharges from the nostrils, and to arrest bleeding from wounds. Burnt alum is more powerful, and is used as a mild caustic, mixed with honey, to fungous growths, and sore mouths.

ALTERATIVES. Medicines which, without producing any considerable or immediate sensible operation, and without interfering with food or work, effect a slow change in the diseased action of certain parts, so as gradually to restore a healthy state.

AMMONIA, CARBONATE OR SESQUICARBONATE OF. *Volatile Salts.* Stimulant and antacid. Dose, 1 dr. to 2 drs. [MOIROUD says from 2 to 8 drs.] to horses in tympanitis, and the last stage of pneumonia. To cattle, in hoven (distension from the fermentation of green food), 1 to 4 drs. [MOIROUD says to 12 drs.]. The solution of carbonate of ammonia has the same properties as the spirit of hartshorn.

AMMONIA, AROMATIC SPIRIT OF. Properties as the last.

Dose, $\frac{1}{2}$ oz. to 1 oz.; or to cattle in hoven, 2 to 4 oz.

AMMONIA, LIQUID. Water of ammonia is more pungent and stimulant than the carbonate, and is used for the same purposes, particularly in tympanitis and hoven, largely diluted with water or some aromatic infusion; but it is chiefly used externally in stimulating liniments; also both internally and outwardly as an antidote to the bite of vipers. The dose of common water of ammonia may be from 2 to 6 drs.; or for cattle to 2 oz. diluted. For small animals from $\frac{1}{2}$ dr. to $1\frac{1}{2}$ drs. diluted. The vapour from the liquid ammonia (applied by holding an open bottle containing it to the eye) is used for the relief of amaurosis and other chronic affections of the eye.

AMMONIA, ACETATE OF. See ACETATE OF AMMONIA.

AMMONIA, MURIATE OF. See MURIATE OF AMMONIA.

AMMONIACUM, GUM. In properties this gum resin is very similar to Assafoetida, but not quite so active. It is occasionally used for the same purposes, and as a constituent of various charges and plasters. Dose for a horse, 2 to 4 drs.; cattle, 2 to 4 drs.; sheep, $\frac{1}{2}$ to $1\frac{1}{2}$ drs.; pig, $\frac{1}{2}$ to $1\frac{1}{2}$ drs.; dog, 10 to 20 grs.

ANALEPTICS. Medicines or food which restore exhausted strength.

ANGELICA. The root in powder or infusion is a warm tonic.

Dose, $\frac{1}{2}$ oz. to 2 oz.

ANISEED. This warm seed is used as a cordial, carminative, and pectoral. Dose for the horse, $\frac{1}{2}$ oz. to 1 oz.; or $\frac{1}{2}$ dr. of the essential oil. The latter is often added to purgatives to prevent griping. Cattle take 1 to 2 oz. of the powdered seeds. The oil is said to be poisonous to pigeons.

ANODYNES. Medicines which alleviate pain. Opium is chiefly employed for this purpose.

ANTIMONIALS. The preparations of antimony (besides their effect in producing vomiting in carnivorous animals) are considered to have a special action on the skin and lungs. They are also termed resolvent, and purifiers of the blood, and are supposed to be useful in visceral and glandular obstructions, farcy, &c. Mr BLAINE says "they lessen arterial action without operating very sensibly either in nauseating the stomach or greatly relaxing the skin." Some writers attribute diuretic effects to them. They are

also said to promote condition. Pigs are supposed to fatten under their use. The principal antimonial preparations employed in veterinary practice are mentioned below.

ANTIMONY, CRUDE. *Black (or sesqui-) Sulphuret of Antimony. Sulphide of Antimony.* Diaphoretic and alterative. The levigated (prepared antimony) is to be preferred. Given to horses in doses of from 2 to 6 drs., with nitre and sulphur, in surfeit, hide-bound, and other skin diseases; and to improve the coat. Mr YOUATT says the dose should not exceed 4 drs. For cattle, the dose is sometimes increased to 2 or 3 oz.; dogs take from 10 to 30 gr.; hogs, a drachm or more, daily. In consequence of its irregular and uncertain action, black antimony is now rarely employed in veterinary practice.

ANTIMONY, LIVER OF. *Hepar Antimonii and Crocus of Antimony (Crocus Metallorum)* are occasionally used in veterinary practice; but are uncertain in their composition and action. Dose, 1 to 2 drs. Mr CLARK says these compounds, and the glass of antimony, derange the stomach, but that it is doubtful if they have any other effect.

ANTIMONY, CALX OF. Diaphoretic. Dose, 2 to 4 drs.

ANTIMONIAL POWDER. Similar to James's powder. Diaphoretic. In colds, fevers, inflammations, &c. Dose for horses or cattle, 1 to 2 drs.; swine, 6 grs.; dogs, from 2 to 5 grs. Less efficient than Emetic Tartar.

ANTIMONY, PRECIPITATED SULPHURET OF. *Oxysulphuret of Antimony.* Dose, $\frac{1}{2}$ dr. to 2 drs. in obstinate skin diseases. See ANTIMONY, CRUDE.

ANTIMONY, TARTARIZED. *Emetic Tartar, Tartrated Antimony.* Diaphoretic, expectorant, and reduces arterial action. It is also regarded as diuretic and febrifuge. Dose, $\frac{1}{2}$ dr. to 1 $\frac{1}{2}$ drs. in gruel, 3 times a day, in fevers, in inflammation of the lungs, and catarrhal affections. To destroy worms, 2 dr. may be given with powdered tin, or some other mechanical vermifuge, fasting, and followed by aloe; or 1 dr. for 6 mornings, followed on the 7th by a dose of physic. Mr WHITE says he has not seen any good effect from it as a vermifuge. Cattle require from $\frac{1}{2}$ dr. to 1 dr.; sheep from 10 to 20 grs. To swine and dogs it is emetic: the former require from 2 to 5 grs.; the latter from 1 to 3 grs. Externally it produces an eruption on the skin. Formed into an ointment with lard, it has

been rubbed on externally in chest affections, but is dangerously irritant.

ANTIMONY, BUTTER OF. *Chloride or Perchloride of Antimony.* Used externally only, as a caustic in canker, &c.

ANTIPERIODICS. Remedies against those diseases which return at regular intervals, as agues.

ANTISEPTICS. Remedies which counteract putrefaction.

ANTISPASMODICS. Medicines which relieve spasm, as opium, ether, camphor, ammonia, ardent spirits, &c.

ARECA SEEDS. *Areca Nut. Betel Nut.* The seeds, or kernels of the fruit of the catechu or betel-nut palm. Astringent; given for worms, especially in dogs. For horses, 4 to 6 drs.; cattle, 4 to 8 drs.; dogs, 30 grs. to 2 drs. Should be made into a bolus. More effectual in coarse than in fine powder.

ARNICA. Nervine, sedative, and diaphoretic. 40 to 60 grs. of the powdered plant (the flowers in preference) have been given twice a day for paralysis, amaurosis, rheumatism, blows, or falls, &c. A decoction may be used outwardly as a fomentation to bruises, wounds, &c.

ARISTOLOCHIA. See BIRTHWORT.

ARSENIC. *White Arsenic, or Arsenious Acid.* Very poisonous call animals. In small doses, tonic and alterative—but its operation requires to be carefully watched. It has been given, in doses of 2 grs., gradually increased to 20, in farcy and glanders. Externally, as a caustic, but dangerous and unmanageable. Used in solution to destroy vermin in cattle and sheep; but it is not free from danger. Mr YOUATT remarks—"We have better and safer tonics, and better and safer caustics." The best antidotes are, moist hydrated oxide of iron, and calcined magnesia, in very large quantities, or a mixture of lime water and linseed oil.

ASSAFETIDA. Stimulant, antispasmodic, and expectorant. It is prescribed in nervous affections and chronic coughs; also in farcy and worms; and to increase the appetite and digestion. The dose is $\frac{1}{2}$ dr. to 2 drs.; but, according to MOIROUD, may be carried to 2 oz. for the horse, and 2 or 3 oz. for horned cattle. Externally, it is applied to indolent tumours, &c.

ASTRINGENTS. Medicines which produce a more obvious and decided constriction of the muscular fibres than the simple tonics.

- ATROPIA.** A crystallizable alkaloid obtained from *Belladonna* root. *A most potent poison.* When taken internally it excites the capillary circulation, and acts as a general anodyne, and also as a diuretic. The sulphate is the most convenient form for use. The dose, in bolus or solution for horses or cattle is 1 to 2 grs.; for sheep, about 1-10th gr.; for dogs, 1-30th to 1-20th gr. One tenth of these quantities suffice when the medicine is used subcutaneously.
- BALSAMS.** Natural balsams appear to act on the mucous membrane generally; but are chiefly given as diuretics and expectorants. See **BALSAM OF CANADA**, **COPAIVA**, **PERU**, &c.
- BALSAM OF CANADA.** Diuretic. Dose, $\frac{1}{2}$ oz. to 1 oz.
- BALSAM OF COPAIVA.** As a diuretic to horses, $\frac{1}{2}$ oz. to 1 oz.; as an expectorant in chronic coughs, 1 or 2 dr. For dogs, $\frac{1}{2}$ dr. to 1 dr.
- BALSAM, FRYAR'S.** *Comp. Tincture of Benzoin.* It is sometimes given in $\frac{1}{2}$ -oz. doses to horses, in chronic cough, mixed with yolk of egg, gruel, or linseed tea. But more frequently applied to wounds, indolent ulcers, &c.
- BALSAM OF LOCATELLI.** Dose, $\frac{1}{2}$ oz. in old coughs.
- BALSAM OF TOLU, AND OF PERU.** 2 drs. in old coughs; but too expensive.
- BALSAM OF SULPHUR.** A stimulating expectorant in old coughs, in doses of $\frac{1}{2}$ oz. to 1 oz. Sometimes used as an outward application.
- BARBADOES TAR.** Stimulant, diuretic, and expectorant. Dose, 1 to 4 dr. or more [2 to 4 oz.—MORTON], in old coughs and chronic chest affections. Externally in skin diseases, wounds, grease, &c.
- BARK, PERUVIAN.** Tonic, astringent, antiseptic, and anti-periodic. Dose for a horse, 6 or 8 drs. [to 2 or 3 oz.—MOIROUD] in diabetes, general weakness, a tendency to gangrene, &c. To small animals, 1 or 2 drs. Applied also to indolent and foul ulcers.
- BARLEY.** The decoction (of Scotch or pearled barley in preference) is given as an emollient, demulcent, or diluent drink in inflammatory diseases; more frequently as a vehicle for more active remedies.
- BARYTA.** All its compounds are poisonous. The following doses have been given in farcy and glanders:—Chloride of barium, 20 grs., gradually increased to 60 grs.; pure

- baryta, 10 to 20 grs.; carbonate, 1 to 4 grs. A dog was killed by 15 grs.
- BASILICON, YELLOW AND BLACK.** *Resin Cerate.* See *Veterinary Formulary* (DIGESTIVE OINTMENTS).
- BAY BERRIES.** Stomachic and carminative. An ingredient in diapente, but rarely given alone. Dose, of the powdered berries, $\frac{1}{2}$ oz., or of the oil of bays, $\frac{1}{2}$ dr. to 1 dr. The leaves are used in fomentations.
- BELLADONNA.** *Deadly Nightshade.* Narcotic and sedative. Dose of the extract, from $\frac{1}{2}$ dr. to 1 dr., for horses and cattle, in diseases where there is undue action of the nervous and vascular systems [MAVOR]. M. MOIROUD directs from 6 to 8 drs. of the powder. For dogs, from 2 to 8 grs. of the powder. The extract is also applied to the eye, to dilate the pupil.
Antidote.—Opium.
- BENNET, HERB.** *Arens.* Tonic and astringent. Dose, of the powdered root, $\frac{1}{2}$ oz. to 1 oz. or more.
- BENZOIN.** Stimulant and expectorant. Dose, 1 to 3 drs. But seldom used. Externally, it is applied, in balsamic tinctures, to wounds, ulcers, &c. See TINCTURE OF BENZOIN.
- BIRTHWORT.** A gentle stimulant, supposed to act especially on the uterine system. Dose, $\frac{1}{2}$ oz. It is given to cows in cleansing drinks, but it is of doubtful utility.
- BISTORT ROOT.** Astringent. Dose, 4 to 8 drs. [or 2 oz., MOIROUD]. The decoction is used also as an astringent and cleansing lotion.
- BITTER-SWEET.** *Dulcamara.* Diuretic, narcotic, and alterative. Dose, $\frac{1}{2}$ oz. in decoction.
- BITTER APPLE.** See COLOCYNTH.
- BLEEDING.** The quantity of blood usually extracted from the horse is from 2 to 4, or, in some cases, 5 to 6 quarts; or until faintness is produced. From cattle, from 2 to 6 quarts, or till faint. Sheep, 16 oz. Lambs, 4 oz. Dogs, in the proportion of 1 oz. for every 3 lb. weight. [Or 1 or 2 oz. from a very small dog; 7 or 8 oz. from a larger one.—Mr YOUATT.] “Although bloodletting is assuredly the best remedy in the early stages of acute inflammation in vigorous animals, it must not be applied in all cases or in all stages of inflammation. It is always injurious in young weakly subjects, in the later stages of disease, in epizootic affections and eruptive fevers, and,

indeed, whenever the pulse is small, quick and weak."—
FINLAY DUN.

BLISTERING FLY. See CANTHARIDES. Blisters are applied in the form of ointments, or liniments, to excite superficial inflammation, followed by vesication; and are intended to draw away inflammatory action from more deeply seated but not distant parts. Also to excite the action of the absorbents, and to promote suppuration. See BLISTERING OINTMENT, and LIQUID BLISTER, in *Veterinary Formulary*.

BLUE VITRIOL. See COPPER, SULPHATE OF.

BOLE, ARMENIAN. Slightly astringent and absorbent. Dose, $\frac{1}{2}$ oz. to 2 oz., in diarrhœa, bloody urine, &c. A common ingredient in drenches to dry the milk of cows. Dose, 1 to 3 oz. It is also used outwardly as an astringent and desiccative.

BORAGE. A decoction of the plant is pectoral and demulcent.

BORAX. Detergent. Applied to sore mouths, mixed with honey. It is supposed to be a uterine stimulant, but is not often used in veterinary practice as an internal remedy. It is a useful antiseptic.

BOX LEAVES. They are given, chopped with corn, as a vermifuge. They were formerly used as a preventive in hydrophobia. The rasped wood is considered sudorific, and prescribed in rheumatic and skin diseases, and even in farcy and glanders.

BRAN. Mucilaginous, and slightly laxative: given in mashes.

BRANDY. See SPIRITS, ARDENT.

BRIONY. White briony root is *poisonous*. $\frac{1}{2}$ oz. killed a dog.

BROMINE. *Poisonous*. 5 grs. killed a dog. Its medicinal use is not well ascertained, but appears analogous to that of iodine.

BROOM. The Spanish broom, and particularly the seeds, are supposed to produce inflammation of the bladder in sheep and cattle.

BUCKBEAN. A bitter tonic and purgative. The powdered plant has been given to sheep for rot, in 1-dr. doses.—
Dr PARIS.

BUCKWHEAT. Slightly laxative, but chiefly used to fatten poultry.

- BUCKTHORN.** Purgative; principally administered to dogs. Dose, of the juice, 2 or 3 drs.; but it is usually given in the form of syrup. The berries are more active, but seldom employed. See SYRUP OF BUCKTHORN.
- BURDOCK.** Diuretic and sudorific. Used, but rarely, in rheumatism and skin diseases.
- BURGUNDY PITCH.** Similar to resin in its properties. It is chiefly used outwardly, in charges, &c.
- BUTTER OF ANTIMONY.** *Chloride, or Perchloride of Antimony.* See ANTIMONY, BUTTER OF.
- CABBAGE TREE BARK.** Vermifuge. Dose for a horse, 2 to 4 drs. But rarely used.
- CALAMINE, OR LAPIS CALAMINARIS.** *Native Carbonate of Zinc.* Slightly astringent, drying and healing. Sprinkled on excoriations and sores; and used in ointments, lotions, eye waters, &c.
- CALAMUS AROMATICUS.** *Sweet Flag.* A warm stomachic. Dose, from 1 oz. to 2 or 3 oz., in infusion.
- CALOMEL.** Alterative, vermifuge, sialogogue, purgative; it also increases the action of diuretics and diaphoretics. "It is employed in almost all animals in reducing and controlling acute inflammations, and appears especially serviceable in those affecting the serous membranes, as in pleurisy, common and puerperal peritonitis, iritis, and rheumatism."—FINLAY DUN. In doses of 1 to 2 drs. [20 to 60 grs., YOUATT], combined with, or followed by aloes, it is given to horses for worms; or from 10 to 20 grs. as an alterative, in skin diseases, grease, farcy, constitutional affections, &c. If too often repeated it salivates. It does not agree with cattle (see MERCURY), but is sometimes given, in doses of from 10 to 20 grs., in inflammation of the liver and jaundice. Some writers mention much larger doses. On dogs it acts as a purgative, and often as an emetic, and it is very apt to salivate. The same applies to swine. Dose for dogs, 1 to 2 grs. [Never exceeding 3.—YOUATT.] Many dogs are killed by calomel. Hogs require 3 to 5 grs. Poultry should not have more than a grain, in divided doses, in the day.
- CALUMBA.** Tonic. Dose, of the powdered root, from 2 to 4 drs.
- CAMPHOR** is reputed antispasmodic, narcotic, and diuretic. It assists the action of diaphoretics; is frequently added

to fever medicines to allay irritation; and is used as an antiseptic in malignant epidemics, &c. Mr SPOONER combines it with opium in cases of lock-jaw. Dose, 1 or 2 drs. Its use is questionable where active inflammation exists. Externally it is used as a discutient and anodyne, in embrocations, eye-waters, &c. Its vapours are thought to act favorably on old coughs.

CANELLA BARK. A warm tonic. Dose for horses, 2 to 4 drs.; for cattle, 2 to 6 drs.

CANTHARIDES. Stimulant and diuretic. Mr VINES says, "Of all medicines given for farcy and glanders none equal cantharides;" but they should not be given too early, nor without due caution. [Mr BLAINE.] Dose, in debility, 3 to 5 grs.; in dropsy, farcy, and glanders, 5 to 8 grs. daily, gradually increasing the dose to 15 grs.; suspending their use for a time when their diuretic effect is manifest. Of the tincture, 2 to 3 drs. in incontinence of urine; and from $\frac{1}{2}$ oz. to 2 oz. in red water. The practice of giving cantharides as a venereal stimulant is reprobated by the best authorities. Externally, it is used in blistering and stimulating ointments and liniments. It does not permanently blemish, but this effect is often produced by other ingredients combined with it in blistering ointments.

CAPSICUM. *Cayenne Pepper.* A hot stimulant. From 10 to 20 grs. may be given in weakness of the stomach, and from 20 to 60 grs. in flatulent colic, or in severe colds. It is also used externally as a stimulant.

CARBOLIC ACID. *Internally,* in excessive doses, it is an *irritant poison.* Occasionally it is administered in medicinal doses for worms; also given to dogs to stop vomiting. For horses, of the fluid acid, 20 to 40 minims; cattle, the same; sheep, 5 to 15 minims; pig, 5 to 15 minims; dog, 1 to 2 minims, dissolved in dilute spirit of wine, or made into a bolus with linseed meal or common mass. *Externally.*—Caustic, disinfectant, astringent, and styptic. Used in canker, and thrush in the foot of the horse, and for foot-rot in sheep. Said to render inert the virus of cholera, cattle plague, and other contagious diseases. When used externally as a caustic it is applied undiluted; for other purposes it may be used in the form of lotion, liniment, or ointment.

CARBONATE OF AMMONIA; CARBONATE OF POTASH; and

- CARBONATE OF SODA. See AMMONIA, CARBONATE; POTASH, CARBONATE; SODA, CARBONATE.
- CARAWAY SEEDS. Carminative and stomachic. Dose, $\frac{1}{2}$ oz. to 1 oz.; or double that quantity to cattle. Used in cordial balls and drenches; and often added to purgatives, to prevent griping. The essential oil is used for the same purposes, in doses of 10 to 30 drops. Mr YOUATT considers caraway and ginger the only cordials required for the horse.
- CARDAMOM SEEDS. Carminative. Dose, 1 to 4 drs.
- CARMINATIVES are stimulants which by their rapid impression on the stomach, occasion the expulsion of wind, and cause relief from pain.
- CARROTS. Restorative and alterative. Given to horses as food after severe illness; and in coughs, grease, foul humours, &c. Externally in poultices.
- CASCARILLA. A warm, bitter tonic. Dose, 2 or 3 drs.
- CASSIA. A warm stimulant. Dose, 1 to 2 drs.
- CASTOR. Antispasmodic. $\frac{1}{2}$ oz. has been given in locked jaw. Rarely used.
- CASTOR OIL. Laxative. It is uncertain as a purgative for the horse, and sometimes produces much irritation in large doses. $\frac{1}{2}$ pint may be given, with watery solution of aloes, every six hours till it operates. Cattle require a pound, or pint; calves, 2 to 4 oz.; sheep and swine, 1 to 2 oz.; dogs, 2 to 4 drs., with syrup of buckthorn. The seeds are more active; from 2 to 6 are sometimes given to swine and dogs, crushed and mixed with food; but from their effects on man their use would seem to require great caution. They are much used by the native Indian farriers for the cure of mange.
- CATECHU. *Terra Japonica*. Astringent. Dose for a horse, in diabetes, diarrhoea, 1 or 2 drs. [YOUATT], or to 1 oz. [BLAINE]; cattle, 2 to 4 drs. in gruel. [It is usually combined with chalk, opium, and gum.—YOUATT.] Dogs require from 10 to 40 grs. In India it is said to be given in doses of 2 oz. for the purpose of taming vicious horses. The tincture is useful in promoting the healing of wounds.
- CAUSTICS. Solid or liquid substances which burn or destroy the part to which they are applied. The *actual caustery* consists in burning with an iron heated to whiteness.
- CHALK. Antacid and astringent. Horses require from $\frac{1}{2}$ oz. to 1 oz.; cattle, 1 or 2 oz.; sheep and swine, 1 dr.;

dogs, 10 to 20 grs. It is often combined with catechu. Externally it is sprinkled on sores.

CHAMOMILE. A mild tonic, stomachic, and febrifuge. Dose, 1 to 4 dr. of the powdered flowers, or an infusion of $\frac{1}{2}$ oz. of the flowers in a quart of water in debility of the stomach, flatulence, and in the last stage of fevers, and influenza. It is the first tonic that should be used in convalescence. Ginger, or some other aromatic, is usually joined with it.

CHARCOAL. Antiseptic. Used as an application to foul ulcers, either sprinkled on them or mixed with poultices.

CHARGES. Compositions of an adhesive nature, usually mixed with tow, which adhere to the part to which they are applied, for some time. See *Veterinary Formulary*.

CHLORAL HYDRATE. *Hydrate of Chloral.* In excessive doses it acts as a narcotic poison. In medicinal doses it is sedative and antispasmodic. It is said to possess the good, but none of the objectionable properties of opium. The dose for the horse is from $\frac{1}{4}$ to $\frac{1}{2}$ oz.; cattle, $\frac{1}{4}$ to 1 oz.; sheep, 1 to 2 drs.; pig, 1 to 2 drs.; dog, 10 to 30 grs.

CHLORATE OF POTASH. Mr MORTON states that Mr SYMONDS found it useful in hoven and tympanitis. Dose, 1 to 2 drs.

CHLORIDE OF LIME. Antiseptic and disinfectant. From 2 to 4 dr. in a quart of water, given to horses in flatulent colic, and to cattle in hoven; and in putrescent diseases. Externally, as a wash for mange, foul ulcers, &c., and as a disinfectant, $\frac{1}{2}$ oz. to be well mixed with a pint of water, and after a time decanted or strained. Mixed with linseed meal it is applied in the form of a poultice to unhealthy wounds and ulcers.

CHLORIDE OF POTASH. *Eau de Javelle.* Recommended by the French for the same purposes as the chlorides of lime and soda. Dose for hoven or tympanitis, $\frac{1}{2}$ oz. to 1 oz.; for sheep, $\frac{1}{4}$ oz., in water, with or without the addition of ether.

CHLORIDE OF SODA. *Labarraque's Disinfectant Solution.* The properties and uses are the same as of chloride of lime; it is perhaps better adapted for internal use. Dose, 2 to 4 drs. of the solution, gradually increased to 1 oz. or more, largely diluted. It has been tried in glanders. As a lotion, about 1 oz. to a pint of water.

- CHLORINE.** Antiseptic. A strong watery solution of chlorine gas is antiseptic—in large doses *poisonous*. It is used for the same purposes as the chlorides of lime, potash, and soda, but the latter are preferable.
- CHLORIDE OF ANTIMONY.** See **ANTIMONY, BUTTER OF.**
- CHLORIDE OF ZINC.** It is a powerful caustic. A diluted solution is used as a disinfectant.
- CHLOROFORM.** Used to produce insensibility to pain in the same manner as ether; and as a remedy for tetanus. Mixed with spirit it forms the spirit of chloroform of the B. P., and as such is given as an antispasmodic.
- CINCHONA.** See **BARK, PERUVIAN.**
- CINNABAR** *Vermillion. Native, and Factitious Red Sulphuret, or Sulphide of Mercury.* Alterative and vermifuge (?). Dose, $\frac{1}{2}$ oz. daily to horses, in skin diseases and obstinate coughs. Formerly given in large doses, as a vermifuge; but now fallen into discredit as a veterinary medicine. Care must be taken to get pure vermilion, as this compound being used as a pigment, is sometimes adulterated with red lead and other poisonous matters.
- CINNABAR OF ANTIMONY,** so called from the mode of preparation, does not differ from common vermilion in its properties.
- CINNAMON.** Stimulant and carminative. Dose, 2 drs. Cassia is usually substituted for it.
- CLOVES.** A hot stimulant, cordial, and carminative. Dose, 1 to 3 drs. in powder; or from 10 to 20 drops of the oil. The latter is a frequent adjunct to purging balls, to prevent griping. Cloves are also an ingredient in masticatories.
- CLYSTERS.** These are injected into the rectum by a proper syringe, or a bladder and a pipe, either to unload the bowels, abate inflammation and pain, or to act on the system generally, when medicines cannot be given by the mouth. See *Veterinary Formulary*.
- COLCHICUM.** Poisonous to most animals. A diuretic and drastic purgative, chiefly used in rheumatic affections. Dr LEMANN found it useful in constitutional ophthalmia, and in pneumonia, in doses of 1 dr. twice a day, with nitre. According to M. MOIBOUD, the dose for larger animals is from 1 to 2 drs.; for smaller, $\frac{1}{6}$ or 8 grs.
- COLOCYNTH.** *Bitter Apple.* It has little effect on the horse. It is purgative to dogs; and in large doses poisonous.

- CONFECTION OF OPIUM. Anodyne and carminative. Dose, 4 to 6 drs., in flatulent colic.
- CONFECTION OF ROSES. Slightly astringent; but only used to form astringent powders, &c., into balls. Masses formed with it retain their consistence well.
- COPAIVA. See BALSAM OF COPAIVA.
- COPPER. All the compounds of this metal are *poisonous*. In small doses they are tonic. The antidotes are white of egg, milk, iron filings, or hydrated sulphuret (sulfide) of iron.
- COPPER, ACETATE OF. *Crystallized*, commonly called *Distilled, Verdigris*. Caustic and cleansing. Stronger than common verdigris.
- COPPER, DIACETATE OF. See VERDIGRIS.
- COPPER, DINIODIDE OF. Tonic, and promotes absorption. Dose, 1 or 2 drs. daily, in farcy, glanders, swelled legs, &c., and topically, to ill-conditioned ulcers.
- COPPER, NITRATE OF. Sometimes used as a caustic.
- COPPER, SULPHATE OF. *Blue Vitriol*. Tonic and styptic. In doses of $\frac{1}{2}$ dr. gradually increased to 2 drs. or more, daily, it is given in diabetes, farcy, &c. Small doses may be given in balls with gentian and ginger; larger doses in gruel. It has been thought useful in glanders; but Mr YOUATT says it is only proper in nasal discharges without fever. Dose for cattle, 1 to 2 drs.; sheep 20 to 40 grs.; rabbits (in sniffles), 1 or 2 grs. twice a day. Externally the solution is used for the foot-rot of sheep; and as a cleansing wash for foul ulcers in horses and cattle. Used also in the solid state to destroy proud flesh.
- COPPER, AMMONIO-SULPHATE OF. Tonic and astringent. Dose, 1 to 3 drs. twice or thrice a day.
- CORDIALS. Warm stimulating medicines, such as spices, and the aromatic seeds, fermented liquors and spirits, &c., which temporarily restore exhausted strength, revive the spirits, and rouse the system generally. The best modern practitioners condemn their indiscriminate employment as the source of much mischief. For cordial balls, &c., see *Veterinary Formulary*.
- CORIANDER SEEDS. A mild aromatic stimulant and carminative, used in cordial balls and drinks. Dose, $\frac{1}{2}$ oz. to 1 oz.
- CORROSIVE SUBLIMATE. *Perchloride, or Bichloride of Mer-*

cury. One of the most virulent of poisons. In small doses it is alterative and diuretic. It has been tried in doses of 2 to 5 grs., gradually increased to 10 or 20, in farcy and glanders, but rarely with lasting benefit. Externally it is used as a powerful caustic. A dilute solution is employed as a wash for scab and lice in sheep, but the practice is not free from danger. Applied to wounds in cattle it has proved as fatal a poison as when swallowed. The antidote for an overdose is white of egg, or milk, or the hydrated sulphuret (sulphide) of iron; with demulcent drinks.

COTTON WOOL. Applied to blistered surfaces, and in burns and scalds; also to wounds, to protect them from the irritating effects of the atmosphere.

COWHAGE. Vermifuge; but has little effect on the horse.

CREAM OF TARTAR. Cooling, laxative, and diuretic. Seldom given alone; but combined with antimonials, mercurials, or sulphur, as an alterative in skin diseases; and used as an adjunct to aloes in purging balls. Cattle require 2 to 3 oz.; when given in larger doses it should be given in plenty of warm water; sheep require $\frac{1}{2}$ oz. to 1 oz.; dogs, 5 to 20 grs.

CREASOTE. Tonic, stimulant, and antiseptic. Dose, 20 to 30 drops daily, in gruel or linseed tea, in glanders. Externally in lotions and ointments, to fistulous wounds, unhealthy ulcers, &c.

CROCUS OF ANTIMONY. See ANTIMONY, LIVER OF.

CROTON SEEDS AND OIL. Purgative. The oil produces great irritation in the horse. Dose, about 20 drops: 30 drops have proved fatal. The powdered seeds and the meal or ground cake left after expressing the oil, are also used; 3 grs. of the former and 5 of the latter being considered equivalent to 1 dr. of aloes. It operates with less certainty, and produces more debility, than aloes, but is sometimes preferred on account of its more speedy action. It is usually given in the form of a ball, 20 or 30 grs. being mixed with 1 oz. linseed meal. Mr NORTON gives from 12 to 24 grs. of the seed. Mr YOUATT prescribes 30 grs. of the powdered seeds in a drink, in tetanus and brain fever, followed by smaller doses (10 grs.) every 6 hours. It will purge rapidly when placed upon the tongue, but is then likely to inflame the mouth. From 10 to 20 grs. are sometimes added to salts in purging

drenches for cattle, in extreme cases. One drop of the oil purges a dog freely.

CUMIN SEED. A warm carminative. Dose, from 1 to 4 drs. of the powdered seeds; or from 6 to 20 drops of the oil.

CUSPARIA. *Angustura Bark.* An aromatic bitter tonic. Dose, 1 to 4 drs. in debility, diabetes, diarrhœa, &c.

CUSSO, Koussou. The flowering pannicles of *Brayera Anthelmintica*. Given for worms, especially for tapeworm. The dose for a horse is from $\frac{1}{2}$ to 1 lb.; for a dog, from 2 to 6 drs.

CYANIDE OF POTASSIUM. It possesses the same poisonous and medicinal properties as prussic acid. Mr LAFORE has given it with success in a case of idiopathic tetanus of the horse; but it failed to cure traumatic tetanus. Dose, $\frac{1}{4}$ grs.

DAFFY'S ELIXIR. Sometimes given in colic or gripes.

DALBY'S CARMINATIVE. Given to calves in diarrhœa. Dose, $\frac{1}{2}$ a bottle.

DETERGENTS OR DETERSIVES. Remedies which cleanse foul ulcers.

DIAPENTE. A compound powder, reputed cordial and stomachic. Too much of what is sold in the shops is almost worthless. Dose, $\frac{1}{2}$ oz. to 1 oz.

DIAPHORETICS. Medicines which promote perspiration.

DIGESTIVES. Mildly stimulating applications, which excite healthy action in indolent ulcers, wounds, &c.

DIGITALIS. *Foxglove.* Sedative and diuretic. It reduces the frequency of the pulse and diminishes irritability. It is poisonous to animals generally; 6 grs. will kill a dog. It is asserted, however, that it produces no effect on poultry. The common dose of the powdered leaves for a horse is from 10 to 20 grs. Mr YOUATT prescribes 60 grs., with emetic tartar and nitre, in inflammation of the chest; but its effects on the pulse must be carefully watched. To cattle, $\frac{1}{2}$ dr. to 1 drs.; sheep, 5 to 15 grs.; dogs, 1 to 2 grs. An infusion of the leaves is applied to inflamed eyes.

DIURETICS. Medicines which increase the flow of urine. Some of them, juniper, capivi, squills, broom, &c., appear to carry off water only; while the alkaline salts remove solid matters also, and thus purify the blood. Diuretics are employed to lessen the quantity of the circulating

fluid in fevers and inflammations. The legs of many horses cannot be kept fine, nor the grease be subdued, without the use of diuretics. Plenty of water should be allowed with them. But their too frequent use is injurious.

DOG-GRASS. It is emetic to dogs.

DOVER'S POWDER. Sudorific to cattle, in rheumatism. Dose, 1 dr.

EGGS. Nutritive and demulcent. Sometimes given in diarrhœa. They constitute the best antidote in poisoning by corrosive sublimate.

ELATERIUM. It has little effect on the horse.

ELDER. An infusion of the flowers is given in catarrhal complaints. The leaves boiled with lard form an emollient ointment, which is a common application to sore udders. The fresh leaves of the dwarf elder are given, according to **BOURGELAT** and **MOIROUD**, with some success as a deobstruent and aperient, in swelled legs, dropsy, and farcy.

ELECAMPANE. The root is reputed stimulant, diaphoretic, diuretic, stomachic, and expectorant. Dose, 4 to 8 drs., in chronic catarrh, dropsical swellings, indigestion, &c.

EMETIC TARTAR. See **ANTIMONY, TARTARIZED.**

EMETICS. Medicines which excite vomiting. It is scarcely possible to produce this effect in herbivorous animals.

EMOLLIENTS. Medicines which soften and relax the tissues of the organs.

EPSOM SALT. A cooling laxative. It is not to be depended on as a purgative for the horse; but in doses of 4 or 5 oz., in a large quantity of water, repeated three times a day, it is useful as a laxative and diuretic in inflammatory diseases. Cattle require from 12 to 20 oz., with ginger, or any of the warm seeds. It may be rendered more active by aloes or gamboge. Calves require from 1 to 2 oz., according to their age and strength. Sheep, $\frac{1}{2}$ oz. to 2 oz. Dogs, from 1 to 3 drs., wrapped in tissue paper. A large elephant takes a pound and a half, preceded by a dr. of calomel.—**YOUATT.**

ERGOT OF RYE. Styptic in hæmorrhage of the lungs, kidneys, and other organs. It promotes parturition. Dose for a mare or cow, $\frac{1}{2}$ oz. to 1 oz., repeated at intervals of half an hour; an ewe, 20 to 40 grs.; bitch, about 1 dr.

- ERRHINES.** Remedies which excite a discharge from the nostrils.
- ESCHAROTICS.** *Caustics.* Substances which destroy the surface to which they are applied.
- ETHER.** A diffusible stimulant and antispasmodic; used chiefly in colic. Dose, $\frac{1}{2}$ oz. to $\frac{3}{4}$ oz.; cattle, $\frac{1}{2}$ oz. to 1 oz.; dogs, 7 to 14 drops. It is used outwardly in cooling lotions and eye-waters. The vapour, inhaled by means of a proper apparatus, produces insensibility to pain; but some of the experiments with this agent have proved most unfortunate. Chloroform has almost universally supplanted it as an anæsthetic.
- ETHIOPS MINERAL.** The mildest of the mercurial compounds. Alterative and vermifuge. Dose, 2 to 4 drs., daily, in farcy, glanders, grease, skin diseases, and worms, given alone, or with cream of tartar. For cattle, 1 dr.; swine, 3 to 10 grs.; dogs, 5 grs., in mange. Latterly, Ethiops mineral has fallen into discredit and consequent disuse. With an equal weight of prepared antimony it forms Antimonial Ethiops—a more efficient preparation.
- EUPHORBIIUM.** Very acrid and poisonous. Used in blisters, chiefly to economise the more expensive flies; but irritates extremely. It is applied in the form of tincture and ointment as a local stimulant.
- EXCITANTS.** Medicines which quicken the circulation, produce warmth, and render the organs more active.
- FENNEL SEEDS.** A weak carminative and diuretic. Dose, $\frac{1}{2}$ oz. to 2 oz.
- FERN.** Powdered male fern is given in doses of 6 drs., followed by a mercurial purgative, for expelling worms. M. MOIROUD carries the dose to 2 oz.; or 5 or 6 drs. for smaller animals.
- FENUGREEK SEEDS.** Emollient, nutritive, and stomachic. Dose, 1 oz., daily, to promote condition in horses, and in diseases of the chest. It is also added to the food of swine to promote their fattening. Used also externally in fomentations.
- FORGE WATER.** The water of the blacksmith's shop was sometimes given as a tonic, or applied as a wash to ulcerated and cankered mouth, by farriers.
- FOXGLOVE.** See DIGITALIS.
- GALANGAL ROOT.** A warm aromatic, similar in properties to ginger. Dose, $\frac{1}{2}$ oz. of the powder, or 1 oz. in infusion.

- GALBANUM.** Stimulant, expectorant, and antispasmodic. Dose, 2 to 4 drs. But rarely used, Assafœtida being stronger and cheaper.
- GALL NUTS.** Astringent; in diarrhœa. Dose, of the powder, for horses and cattle, 2 to 4 drs.; calves, $\frac{1}{2}$ dr. to 1 dr.; dogs, 4 to 8 grs.
- GAMBOGE.** A drastic purgative. The dose for a horse is said to be from 2 to 6 drs., but its purgative effect cannot be depended on, and it gripes. 3 drs. have been known to cause great prostration, and the horse being killed, marks of intense inflammation were found in its stomach and bowels. It is a bad medicine for herbivorous animals. 2 drs. are sometimes added to salts or other purgatives for cattle. Sheep have been killed by 2 drs. A few grains are given to dogs to destroy and expel worms.
- GARLIC.** A stimulating expectorant. Dose, 1 oz., in chronic coughs and asthmatic complaints, made into balls with liquorice powder; or boiled in milk. It is a common remedy for coughs and chest affections in all domestic animals. It is also reputed vermifuge. For the roup in fowls, it is given in doses of 5 grs.
- GENTIAN ROOT.** Tonic and stomachic; in debility, after severe illness, &c. Dose for a horse, 2, 3, or 4 drs. of the powder; or from $\frac{1}{2}$ dr. to 1 dr. of the extract. (See *Veterinary Formulary*, TONIC BALLS.) Cattle, 2 to 4 drs. or more; sheep, 20 to 60 grs. Generally mixed with ginger. An infusion is recommended as a wash to ulcers.
- GINGER.** Stimulant and carminative; a general ingredient in cordial and tonic medicines. Dose, 1 to 3 drs., or in flatulent colic 2 to 6 drs.; cattle, 2 to 6 drs.; calves, 20 to 30 grs.; sheep, 30 to 60 grs. The smaller of the above doses may be added to all aperient medicines. It is also used as a masticatory. Dose of the tincture, $\frac{1}{2}$ oz. to 2 oz.
- GLASS, POWDERED.** Used to destroy worms in dogs. Mr BLAINE recommends as much as will lie on a sixpence with butter. It must be in very fine powder.
- GLAUBER'S SALT.** *Sulphate of Soda.* Aperient and diuretic. Seldom given to horses as a purgative (Mr CLARK says 1 lb. produces scarcely any effect); but is said to be useful in doses of 6 oz. 3 times a day, in epidemic catarrh. To cattle the usual dose is 16 oz., or from 12 to 20 oz., with ginger or caraway. It is considered more diuretic than Epsom salts.

GLYCERIN. Externally its principal employment is as an emollient in skin affections, accompanied by dryness and irritation, to excoriations, cracked heels, burns, scalds, and freshly blistered surfaces. When mixed with an equal bulk of solution of subacetate of lead, it is very serviceable in allaying irritability. Often added to masses to prevent their hardening.

GOULARD'S EXTRACT OF LEAD. *Solution of Subacetate of Lead. Solution of Diacetate of Lead.* Cooling and astringent. Used externally only, in lotions, &c., in the same cases as sugar of lead. (See LEAD, ACETATE OF.) For inflamed eyes, 1 dr. or 1½ to a pint of distilled or boiled water; for other purposes it is made stronger.

GRAINS OF PARADISE. A warm stimulant; chiefly used in cattle medicines. Dose, 3 to 6 drs.

GRUEL. A decoction of oatmeal. Nutritive and demulcent.

GUAIACUM, GUM. Sudorific and expectorant. It has been given to horses, in doses of 4 drs., in chest affections, farcy, rheumatism, &c.; and to cattle in doses of 4 or 6 drs. But its utility is doubted. The guaiacum wood is given to the amount of 4 oz., in decoction, repeated 2 or 3 times in 24 hours.

GUM ARABIC. Emollient and demulcent. Used in inflammatory affections of the bowels, or of the respiratory or urinary organs. Dose for horses and cattle, 1 to 4 oz., dissolved in water. For smaller animals, from ¼ oz. to 1 oz. It is also used to suspend in water insoluble powders and oils. Gum senegal and gum tragacanth are used for the same purposes. The latter will thicken twenty times as much water as Gum Arabic. For GUM AMMONIACUM, BENZOIN, &c., see AMMONIACUM, BENZOIN, &c.

HARTSHORN, SPIRIT OF. See AMMONIA. It is chiefly used as an ingredient in stimulating liniments and for the bites and stings of venomous reptiles and insects.

HELLEBORE, WHITE. *Poisonous to all classes of animals.* In small doses, it has been strongly recommended as a nauseant and diaphoretic, in inflammatory diseases; but it requires to be very carefully watched, otherwise a fatal collapse may be induced. The usual dose is 20 grs. every four or six hours till nausea is produced, or the pulse affected. Externally, it is used in ointments and washes for the mange; but even in this way its use requires caution. It is also blown into the nostrils as a sternutatory.

- HELLEBORE, BLACK.** The root is used as an irritating seton for cattle, and introduced into fistulous sores of the horse.
- HEMLOCK, SPOTTED.** *A narcotic poison.* In doses of an ounce of the powdered leaves, or 1 dr. of the extract, gradually increased; it is sometimes given to quiet obstinate coughs. It is also an ingredient in some old remedies for farcy, scirrhus tumours, and cancer. For dogs, from 1 to 4 grs., in coughs and cancerous diseases. A decoction of the herb is used as a fomentation to painful tumours. Water hemlock is a more virulent poison, and is often fatal to cattle. M. MOIRAUD says that ruminants bear hemlock better than other animals. Mr YOUATT considers both common and water hemlock harmless to the horse, though he admits that cows have been poisoned by the latter.
- HENBANE.** Narcotic and sedative. Dose, 15 to 20 grs. of the powder [1 to 2 dr. of the extract, MORTON] twice or three times a day, to allay arterial action. On dogs it acts as on man; dose, 3 to 5 grs. German horsedealers are said to give a plump appearance to diseased horses by mixing henbane seeds with their corn.
- HONEY.** Demulcent, emollient, and slightly laxative. Used in cough medicines, and to make up balls. Horses are fond of it. Externally, it is detergent, and is, perhaps, useful in protecting ulcers from the air.
- HOPS.** Tonic and slightly anodyne, but chiefly used in fomentations.
- HOREHOUND.** Sometimes given in coughs; a quart of the decoction, or 1 oz. of the powder.
- HORSE-RADISH.** Stimulant and diuretic. Said to be useful in dropsical complaints, and in recent epidemics attended with chronic inflammation. The fresh root is rasped and mixed with barley meal.
- IODIDE OF IRON.** Tonic and alterative, promoting the action of the absorbents. Also given in diabetes. Dose, $\frac{1}{2}$ dr. to 1 dr.
- IODIDE OF POTASSIUM, *Hydriodate of Potash.*** It possesses the same properties as iodine, but irritates less. It is often combined with iodine, which it renders soluble in water. The dose, by itself, is rather larger than of iodine—from 2 to 4 drs.
- IODIDE OF SULPHUR.** Used externally in scabies and other

skin diseases; also applied to farcy ulcers and indolent sores.

IODINE. Alterative, and promotes absorption. Used externally and internally to reduce glandular swellings and scirrhus and other tumours. 20 grs. of iodine, or $1\frac{1}{2}$ to 2 drs. of the compound tincture, may be given twice a day in farcy. Cattle take from 5 to 10 grs., and from 1 to 2 drs. of the compound tincture; dogs, $\frac{1}{4}$ to 1 gr. twice daily. The compound iodine ointment is used to disperse glandular enlargements. It is rapidly superseding cantharides.

IPECACUANHA. Little used in veterinary practice, except as a sudorific, in combination with opium (DOVER'S POWDER). A drachm or two may be given to horses in asthmatic affections. It purges sheep, purges or produces vomiting in the pig and dog. Dose for the latter, 4 to 20 gr. [From 2 to 30 gr.—MOIROUD.] 3 oz. killed a horse [Mr B. CLARKE].

IRON. The preparations of this metal are tonic; some of them, as the sulphate and perchloride, astringent and styptic. The usual doses for a horse are, 2 oz. of iron filings, once or twice a day, with corn, or in a mash; 1 to 3 or 4 drs. of the sulphate; 2 to 6 drs. of the peroxide or carbonate, or rust of iron, or of the powdered scales; 1 to 3 drs. of tartarized iron; and $\frac{1}{2}$ to $1\frac{1}{2}$ drs. of the iodide, as an alterative, astringent, and tonic. Cattle, 2 to 4 drs. of the sulphate in chronic diarrhoea; for sheep, a sixth or eighth of the above doses. M. MOIROUD prescribes much larger doses than the above. The tincture of perchloride of iron is prescribed in doses of 2 or 3 drs. for incontinence of urine. The sulphate is sometimes used externally in astringent lotions.

JALAP. Purgative, but has little or no such effect on the horse or other herbivorous animals. It is sometimes added to other purgatives, but probably without any benefit. Dose for swine, $\frac{1}{2}$ dr. to 2 drs.; dogs, 15 to 40 grs.; cats, 10 to 20 grs.; but it is rather uncertain.

JAMAICA PEPPER. See ALLSPICE.

JAMES'S POWDER. Similar to antimonial powder, but considered more certain and uniform in its operation. Dose, 20 to 30 grs. in fever and inflammatory complaints. It is also given to dogs as a remedy for distemper, 4 grs. twice a day.

- KAMALA.** Minute glands adhering to the capsules of *Rottlera tinctoria*; imported from India. Purgative, and anthelmintic for tapeworm. For horses, 1 to 2 oz.; dog, 1 to 3 drs. Given in the form of bolus.
- JATROPHA SEEDS.** The seeds of the physic nut (*Jatropha curcas*) are given as croton seeds, doubling the dose.
- JUNIPER BERRIES.** Diuretic and slightly stimulant. Dose for a horse, 1 to 2 oz., or 1 to 2 drs. of the essential oil; for cows, 2 or 3 oz.; sheep, $\frac{1}{4}$ to $\frac{1}{2}$ oz. An extract from the berries (prepared by evaporating a clear decoction, or rather a cold infusion of the berries, to the consistence of treacle) is much used on the Continent as a vehicle for various remedies.
- KERMES MINERAL.** A preparation of antimony, similar to the precipitated sulphuret (sulphide), not much used in this country, but highly esteemed in France. Dose for horses, 1, 2, or 3 drs.; for cattle, 4 drs. or more; for a good-sized dog, $1\frac{1}{2}$ grs. gradually increased. Uncertain and unsatisfactory in its effects.
- LARD.** Half a pound, with warm water, is laxative and emollient. It is also used to make up balls, and is thought to prevent griping, as well as to preserve the consistence of the balls. It forms a common basis for ointments.
- LAUDANUM.** *Tincture of Opium.* See OPIUM.
- LAVENDER.** The compound spirit is carminative and cordial. Dose, 1 oz. in peppermint water.
- LEAD.** The preparations of this metal are poisonous. See ACETATE OF LEAD, and GOULARD'S EXTRACT OF LEAD.
- LEAD, WHITE AND RED.** Common ingredients in ointments and plasters. Also sprinkled on sores as desiccatives. They are likewise used for dusting sheep for the fly.
- LIME.** Quicklime is sometimes used as a caustic; the powder is dusted over foul ulcers, greasy heels, &c.
- LIME WATER.** Antacid and tonic. Sometimes given in diabetes, from 2 to 4 quarts. Used also as a wash for sores, and as an injection into the nostrils for glanders and chronic discharges. Mixed with linseed oil, it forms a liniment for burns.
- LIME, CHLORIDE OF.** See CHLORIDE OF LIME.
- LINSEED.** Demulcent and pectoral. A decoction of the seed is very mucilaginous, and is used in colds, sore throats, and internal inflammations; also to counteract the effects

of corrosive and irritant poisons, and as a vehicle for more active medicines. Linseed meal is used for poultices. Linseed oil is laxative. Dose for a horse, a pint or a pint and a half; for cattle, 1 or 2 pints; sheep, 2 or 3 oz.

LIQUORICE. Demulcent and pectoral, in coughs, &c. Dose, $\frac{1}{2}$ oz. to 2 oz. of the powdered root; or $\frac{1}{2}$ oz. of the foreign extract (Spanish or Italian juice).

LIVER OF SULPHUR. See **SULPHURET OF POTASH.**

LOBELIA INFLATA. It is poisonous to horses, and produces salivation in cattle; but its remedial powers have not been ascertained.

LOGWOOD. Astringent. 2 or 3 drs. of the extract, or a decoction of 3 or 4 oz. of the wood, may be given in diarrhœa, &c.

LOTIONS. *Washes.* Liquid applications, with which external parts are bathed.

MADDER. Madder was formerly used in glanders and farcy, and as a preventive of the effects of the bites of venomous reptiles; but it is now discarded from modern practice. It is sometimes given to pigs, but with what specific intention it is difficult to say. It colours the bones of animals fed with it.

MAGNESIA. Antacid and laxative. From $\frac{1}{2}$ oz. to 3 oz. to horses and cattle, with some warm carminative, in flatulent distension. To calves in diarrhœa, $\frac{1}{2}$ oz. Either the common or the calcined magnesia may be used.

MAGNESIA, SULPHATE OF. See **EPSOM SALT.**

MALLOW. Demulcent. A handful of the leaves is boiled in a quart of water. More frequently used as a lavement. The root of the marshmallow is preferred; a decoction of 2 or 4 oz. is given as a drink in both coughs and internal inflammation, and used as a clyster, and as a fomentation.

MALT. Nutritive, pectoral, and alterative. It is given in the form of mashes, in chest affections, when no inflammation is present, and in grease, farcy, and mange.

MANNA. Slightly laxative and pectoral. Dose, 2 oz. with honey, or dissolved in water, in inflammatory diseases and chronic coughs.

MASHES. See **BRAN MASH, &c.**, in *Veterinary Formulary.*

MERCURY, Quicksilver. The preparations of this metal are alterative, most of them purgative, and all apt to

produce salivation. Dogs may be easily salivated, but graminivorous animals with greater difficulty. The Editor of 'Clater's Cattle Doctor' says, "Mercury does not seem to agree with herbivorous animals, in any form or in any disease." The preparations in use are indicated below.

MERCURIAL LOTION, YELLOW. } Stimulant to unhealthy sores
MERCURIAL LOTION, BLACK. } and ulcers.

MERCURIAL OINTMENT. Applied to callous swellings, enlarged joints, mange, scab in sheep, &c. The weaker ointment is generally sufficiently strong.

MERCURY WITH CHALK. *Alkalized Mercury*. A mild preparation. Dose, 1 to 3 drs., in farcy, glanders, &c.

MERCURY, SUBCHLORIDE OF. See CALOMEL.

MERCURY, BICHLORIDE (PERCHLORIDE, B. P.). See CORROSIVE SUBLIMATE.

MERCURY WITH SULPHUR. *Black Sulphuret, or Sulphide of Mercury*. See ETHIOPS MINERAL.

MERCURY, RED SULPHURET, OR SULPHIDE OF. See CINNABAR.

MERCURY, NITRIC OXIDE. See PRECIPITATE, RED.

MERCURY, BLACK OXIDE. Dose, 1 to 2 drs. MORTON.

MERCURY, AMMONIO-CHLORIDE. See PRECIPITATE, WHITE.

MERCURY, NITRATED OINTMENT OF. See OINTMENT, CITRINE, in *Veterinary Formulary*.

MERCURY, ACID NITRATE OF. Used as a caustic. See CAUSTICS, *Veterinary Formulary*.

MILK. Sometimes given in quantities of 1 to 3 quarts, in acute inflammation, coughs, and all internal irritation, especially those occasioned by acrid and corrosive poisons. It is a convenient vehicle for administering medicines to the dog or cat.

MINDERERUS SPIRIT. See ACETATE OF AMMONIA.

MINT, AND PEPPERMINT. Carminative, cordial, and sudorific. A strong infusion of the plant, or the distilled water, may be given in flatulent colic. Dose, 1 or 2 pints; used chiefly as vehicles for more active remedies. Dose of the oil of peppermint, 20 to 30 drops, or to 60 drops of oil of spearmint. A few drops of the oil are added to purgative medicines, to prevent griping. The other mints have similar properties.

MITHRIDATE. Cordial and anodyne. $\frac{1}{2}$ oz. to 1 oz. may be

given in flatulent colic, but would be injurious in inflammation.

MULLEIN. An infusion of the flowers is given as a demulcent for the same purpose as linseed tea. A decoction of the leaves is used in emollient fomentations and cataplasms.

MURIATIC ACID. *Hydrochloric Acid. Spirit of Salt.* Tonic and antiseptic; but principally used to dissolve calcareous concretions in the bladder. It has been used in the pestilient epidemics of cattle. Dose for a horse, $1\frac{1}{2}$ to 2 or 3 drs., in plenty of water, twice a day. Externally as a caustic, strongly recommended by YOUATT.

MURIATE OF AMMONIA. *Chloride of Ammonium. Sal Ammoniac. Hydrochlorate of Ammonia.* Formerly used in influenza or epidemic catarrh. It is said also to have proved useful in farcy, and perhaps deserves trial in other chronic diseases. It renders the blood more fluid. Its use requires caution. 2 oz. produced inflammation of the mucous membrane of a horse; 2 drs. killed a dog, and $\frac{1}{2}$ dr. a rabbit. M. MOIBOUD states the dose to be from 2 to 8 drs. for horses and cattle, and for small animals from a scruple to a drachm, largely diluted. Externally it is a frequent ingredient in discutient lotions to splints, old strains, bruises, indolent tumours, &c., in horses and cattle. It is also employed as an embrocation to sore teats.

MURIATE OF ANTIMONY. *Chloride, or Perchloride of Antimony.* See ANTIMONY, BUTTER OF.

MURIATE OF BARYTES. *Muriate of Baryta, Chloride of Barium.* Poisonous; in small doses, alterative. It has been given in glanders and farcy, with the usual ill success. Dose, $\frac{1}{2}$ dr. in milk.

MURIATE OF COPPER. *Chloride of Copper.* Used externally only, as a mild caustic.

MURIATE OF LIME. *Chloride of Calcium.* Alterative and resolvent, in glandular diseases; but rarely used in veterinary practice. It has been proposed in glanders and farcy. In an overdose it is poisonous. We have not met with any specific statement of doses. $3\frac{1}{2}$ dr. killed a dog.

MURIATE OF SODA. *Chloride of Sodium.* See SALT, COMMON.

MUSTARD. Stimulant; but little used as an internal remedy. Flour of mustard, with or without vinegar, is applied

externally as a rubefacient, to relieve internal inflammation.

MYLABRIS. *Chinese Blistering Fly.* Its vesicant properties are due to the presence of cantharadin. Physiological effects the same as cantharides, except that it is said not to affect the kidneys when topically applied.

MYRRH. Tonic, expectorant, antiseptic, and balsamic. From 1 to 2 drs. to a horse, in chronic cough; to cattle, the same. The tincture is used for ulcers of the mouth in all animals, and to indolent sores.

NAPHTHA. Rectified wood naphtha may be given in $\frac{1}{2}$ -oz. doses internally, it being an admirable stimulant and diaphoretic.

NAPHTHALIN. A stimulating expectorant. It possesses many of the properties of camphor, and a solution of it in spirit may be substituted for camphorated spirit. With oils and lard it may be used in the form of liniment and ointment. The ointment is substituted for tar ointment.

NARCOTICS. Medicines which induce stupor or sleep, and ease pain.

NAUSEANTS. Medicines which produce nausea, diminish arterial action, and thus abate inflammation.

NETTLE SEED. It is said, when given with the horse's corn, to produce a smooth coat, and to impart an appearance of condition and liveliness.

NITRE. *Nitrate of Potash.* Alterative, febrifuge, refrigerant, and diuretic. In colds, fevers, and inflammatory complaints of the horse, from 2 to 4 drs. may be given daily, in plenty of water, or linseed tea, till the desired effect is produced. An ounce is often given, but repeated smaller doses are better. Cattle, 2 to 4 drs. [1 oz. in 24 hours for some days—*MOIROUD*]; swine and sheep, 30 to 40 grs.; dogs, 4 to 10 grs. A strong solution is applied to gangrenous wounds.

NITRIC ACID. Used externally only, as a strong caustic; or largely diluted (2 dr. to a pint of water) as an antiseptic wash to foul ulcers.

NITRATE OF SILVER. *Lunar Caustic.* Tonic; but rarely given to animals, except to dogs in chorea, in doses of $\frac{1}{2}$ to $\frac{1}{4}$ of a gr. Externally caustic. It is the best caustic that can be applied to the bites of rabid animals. A weak solution (10 gr. to 1 oz. rain water) is used to excite

sluggish wounds, and to remove opacity from the cornea of the eye.

NUX VOMICA. *Poisonous to all animals.* Given in doses of 8 to 10 grs., gradually increased to 30 grs., in paralysis of the horse; but its effect requires to be carefully watched. It has been given in glanders and farcy, but without much success. In small doses it invigorates the digestive functions. The French veterinarians are stated to have sometimes given Nux Vomica with good effect in amaurosis and stringhalt in horses, and chorea in dogs, particularly when accompanied by debility. A few grains proved fatal to a dog. A drachm has killed a horse. See STRYCHNIA.

NUTGALLS. See GALL-NUTS.

NUTMEGS. Stimulant, and perhaps narcotic. Sometimes given in colic, but not much in use.

OAK BARK. Astringent and tonic. Dose, $\frac{1}{2}$ oz. to 2 oz., in powder, or boiled in water, for diarrhœa, diabetes, and debility in horses. To cattle, in dysentery, and in red-water, after purgatives, $\frac{1}{2}$ oz. to 1 oz. The powdered bark and the decoction are applied to unhealthy wounds. In France, a mixture of oak bark, gentian, and chamomile, is used as a substitute for Peruvian bark.

OIL, FISH. Common whale oil is a good preventive of the fly, and does not injure the wool.

OIL OF SPIKE. As sold for veterinary purposes, it consists of turpentine, coloured, and merely scented with foreign oil of lavender. It is used in warm liniments.

OIL OF TURPENTINE. See TURPENTINES.

OIL OF TAR. See TAR.

OILS, EXPRESSED. Olive, almond, and linseed oils are laxative, demulcent, and emollient. Dose, 3 to 16 oz., or a pint. In the latter dose they are given (especially linseed oil) as a substitute for castor oil; they are harmless, but rather uncertain in their operation. See CASTOR OIL. Expressed oils are useful in cases of poisoning by acrid and corrosive substances. Olive oil is used, both inwardly and outwardly, as a remedy for the bites of reptiles and stings of insects. Externally, the expressed oils are used in liniments and ointments. Oil of bay is gently stimulant and antispasmodic, but chiefly used outwardly.

OILS, ESSENTIAL OR VOLATILE. The essential oils of pep-

permint, cloves, aniseed, caraway, &c., possess in a concentrated state the warm carminative properties of the drugs from which they are distilled. They are frequently added to purgative medicines to prevent griping. Oil of juniper is diuretic, in doses of 1 to 3 drs. Oil of organum is almost exclusively used outwardly in stimulating liniments. Oil of bitter almonds is *poisonous*.

OILS, EMPYREUMATIC. Oil of amber and other empyreumatic oils are antispasmodic; but mostly used in outward applications. The fetid oil, called **DIPPEL'S ANIMAL OIL** (or oil of hartshorn), is sometimes given as a warm medicine, in doses of $\frac{1}{2}$ oz., to horses, or a dr. to small animals. As an outward application, it is sometimes added to powders against the fly in sheep; but injures the wool. Oils of paper or rag are empyreumatic fluids obtained by burning these substances. Mixed with water, they are used in inflammation of the eyes, mouth, &c.

OILS, COMPOUND. See **OILS** and **LINIMENTS**, in the *Veterinary Formulary*.

OINTMENTS. See *Veterinary Formulary*.

OLIVE OIL. See **OILS, EXPRESSED**.

ONIONS. Stimulant and diuretic. They are said to be useful in colic and gripes. Externally used in poultices to promote suppuration.

OPIUM. Anodyne, antispasmodic, sedative, indirectly astringent, and in large doses narcotic and stupefactive, and capable of destroying life. In combination with ipecacuanha and tartarized antimony, opium is sudorific. The dose for horses in ordinary cases is from $\frac{1}{2}$ dr. to 1 dr. But in locked-jaw, spasmodic colic, and other urgent cases, it may safely be given in doses of 2 drs., and even (according to **MOIROUD**) to 4 drs. **YOUATT** states the dose as 1 dr. to 3 drs. In inflammation of the bowels, after bleeding, it is recommended to give 2 drs. at once, and 1 dr. every hour afterwards until it takes effect. To cattle, the dose is from 2 to 4 drs.; calves, 10 grs.; sheep, 10 to 30 grs. Much larger doses have been given with impunity. Dogs require from $\frac{1}{2}$ gr. to 2 grs., according to size and case. **M. MOIROUD** says the dose for dogs should not exceed that prescribed for man. **Mr BLAINE** thinks they are much less affected by it than man. The dose of tincture of opium (for horses) is from 1 to 2 oz.; of the extract, 20

- to 30 grs. Externally, opium is used in anodyne liniments, and is useful in inflammation of the eye. See EYE-WATERS, LINIMENTS, &c., in *Veterinary Formulary*.
- OPODEDOC. Soap liniment. Used externally only in stimulating liniments.
- ORIGANUM. *Wild Marjoram*. Stimulant. The essential oil is hot and pungent, and a frequent ingredient in liniments for old strains, and in blisters.
- ORPIMENT. *Yellow Arsenic*. *Poisonous*. Used, mixed with lard, for fistulous sores, warts, &c., but is not free from danger.
- OXYMEL. Cooling and pectoral. Used in catarrhal affections. Dose, 3 to 4 oz.
- OXYMEL OF SQUILLS. A stimulating expectorant. Seldom used in veterinary practice.
- PALM OIL. Emollient. Used in compounding ointments and liniments; and much commended as a basis of aloetic and other balls. It has also been given as a laxative; dose, 12 oz. or more.
- PEPPERMINT. Carminative. The distilled water and the essential oil are chiefly used. See MINT.
- PEPPER, BLACK, WHITE, AND LONG. Warm stimulant, cordial; the latter kind is chiefly used in veterinary practice. It must be carefully avoided in inflammatory complaints. Dose for horses and cattle, 2 to 4 drs. For JAMAICA PEPPER, see ALLSPICE.
- PEPPER, CAYENNE. The ground pods of some species of capsicum. See CAPSICUM.
- PERIWINKLE. The plant, in decoction or chopped up in a mash, is said to relieve quinsy. Pulverised and mixed with Ethiops Mineral, it was once vaunted as a remedy for glanders.
- PERUVIAN BARK. See BARK, PERUVIAN.
- PETROLEUM. See BARBADOES TAR.
- PEWTER. The scrapings are given to dogs for worms. Dose, $\frac{1}{2}$ dr. to 1 dr. Tin filings are safer. See TIN.
- PHELLANDRIUM. The seeds of water-fennel (*Phellandrium Aquaticum*) are used in Germany in chronic catarrhal affections. Dose, $\frac{1}{2}$ oz. to 1 oz.
- PHYSIC. In veterinary practice this term is applied to purgatives. See PHYSIC OR PURGING BALLS, *Veterinary Formulary*.
- PIMENTO. See ALLSPICE.

PITCH. Stimulant, balsamic, probably diuretic; but rarely given internally. It is more frequently used externally in charges and warm plasters. For liquid pitch, see **TAR**.

PITCH, BURGUNDY. Stimulant. Used in charges, and warm and strengthening plasters.

POMEGRANATE. The rind of the fruit is given, in decoction or powder, as an astringent to cattle in diarrhœa. Dose, from $\frac{1}{2}$ oz. to 1 oz. The bark of the root is used to destroy worms. **MOIROUD** directs 5 or 6 oz. to be boiled in water for some hours, and the decoction given in divided doses.

POPPY HEADS. Anodyne; principally used in fomentations. (See also **SYRUP OF POPPIES**.) A watery extract prepared from the ripe poppy heads, freed from seeds, is said to be nearly half the strength of opium. Of the ordinary extract (from the decoction) 5 grs. are said to equal 2 of opium.—**LEBAS**.

POTASH CAUSTIC. *Fused Hydrate of Potash.* A powerful caustic.

POTASH, CARBONATE. *Salt of Tartar or Prepared Kali.* Antacid and diuretic. Dose for a horse, from 2 to 4 drs. or more largely diluted. It is seldom given alone, but sometimes joined with tonics, stomachics, purgatives, and with other diuretics. The bicarbonate is milder, and may be given in larger doses.

POTASH, PERMANGANATE OF. 4 grs. dissolved in 1 oz. of distilled water, forms a useful cleansing wash for diseased surfaces.

POULTICES. Are useful in relieving inflammation and pain. See *Veterinary Formulary*.

PRECIPITATE, RED. A mild caustic, and detergent to indolent and foul ulcers.

PRECIPITATE, WHITE. Principally used to destroy vermin in the horse and other animals.

PRUSSIAN OF POTASH. *Ferro-cyanide of potassium* may probably be found useful in veterinary practice; but its properties and uses are not yet properly ascertained. It must not be confounded with cyanide of potassium, which is an *energetic poison*. See **CYANIDE OF POTASSIUM**.

PRUSSIC ACID. *Hydrocyanic Acid. The Diluted Acid of the B. P.* (Medicinal.) *A strong poison to all animals.* Rarely used in veterinary practice. May be given by enema in tetanus. In medicinal doses, sedative, anti-

spasmodic, and anodyne. Used in chronic cough, chorea, epilepsy, chronic vomiting, palpitation of the heart, and rheumatism. PROFESSOR TUSON prescribes from 20 to 30 minims for horses. Mr YOUATT recommends a lotion composed of a drachm of the medicinal acid in a pint of water, to allay cuticular irritation in dogs.

Antidotes.—*Fresh air, affusions of cold water; inhalation of ammonia; moist peroxide of iron mixed with an alkaline carbonate.*

PULSE. The following table, from VATEL, is inserted as a useful remembrancer to the veterinary practitioner:

Table of the Number of Pulsations in a Minute in various Animals.—In the horse, 32 to 38 [36 to 40—WHITE]; ox or cow, 35 to 42 [42 to 45—CLATER; 50 to 55—SPOONER]; ass, 48 to 54; sheep, 70 to 79; goat, 72 to 76; dog, 90 to 100; cat, 110 to 120; rabbit, 120; guinea-pig, 140; duck, 136; hen, 140; heron, 200.

PURGATIVES. *Cathartics or Laxatives.* Medicines which more or less strictly promote evacuations from the bowels. Aloes is almost the only purgative for the horse that is at once certain and safe. For cattle, Epsom or Glauber's salt is the most preferable. Aloes, gamboge, or linseed or castor oil is sometimes combined with them. Sulphur is used when a very strong purgative is not required; yet this demands some caution. Refer to these various articles.

QUASSIA. A bitter tonic. Dose, 1 or 2 drs., with a little ginger, in debility of the stomach. Its poisonous effects on insects and small animals suggest caution in its use.

QUICKSILVER. See MERCURY.

QUININE, SULPHATE OF. Tonic. Dose, $\frac{1}{2}$ dr. to 1 dr. recommended by Mr MORTON in the prostration which follows influenza. But too expensive for general use. It is given to dogs in chorea and distemper, in doses of 2 to 5 grs.

RAGWORT. The herb is said to produce a kind of lethargy or staggers in horses and cattle. Externally it is used as a poultice in quinsy.

RAKING. Removing hardened fæces from the lower bowel by the hand.

RANUNCULUS REPENS. Acrid stimulant. It is poisonous to sheep.

REED. The great reed (*Arundo donax*), and the Bankside reed (*Arundo phragmitis*), are reputed diuretic. The

former is supposed to have the property of diminishing the secretion of milk.

REFRIGERANTS. *Cooling Medicines.* See TEMPERANTS.

RESIN OR ROSIN. Diuretic. $\frac{1}{2}$ oz. to 1 oz. may be given daily to horses in their corn, for swelled legs. The yellow or amber resin is preferable. Externally, it is adhesive and gently stimulating, and is a common ingredient in digestive ointments, and in plasters and charges.

RHODODENDRON. Supposed to be useful in the rheumatism of cattle. Dose, $\frac{1}{2}$ oz. to 1 oz., boiled in water.

RHUBARB. Tonic and stomachic. Scarcely laxative to large animals. From $\frac{1}{2}$ to 1 oz. is given in jaundice, to horses and cattle. On dogs it acts as a purgative, but is an uncertain one, in doses of $\frac{1}{2}$ dr. to 1 dr.

ROSEMARY. A mild stimulant and carminative. The essential oil is chiefly used in warm liniments and ointments; but is sometimes given in doses of $\frac{1}{2}$ dr. to 1 or 2 drs. in colic.

RUE. Stimulant, uterine, antispasmodic, and vermifuge. It is also supposed to resist contagion and poisons. A decoction or infusion of 2 to 4 oz. of the fresh herb in water or beer is given for worms; as an antidote to the bite of vipers; with diuretics in farcy; (formerly) with box leaves as a preventive of hydrophobia; and with camphor and opium in locked-jaw. The bruised leaves are put into horses' ears for the staggers. It is given to poultry for the cure of roup. Externally, rue is used in fomentations as a stimulant, antiseptic, and discutient.

SAFFRON. Cordial, antispasmodic and uterine; but too weak and expensive for veterinary use.

SAGE. Stimulant and tonic. In habitual relaxation of the bowels. The powder may be given in a ball, or the herb infused. The infusion is used as a mouth-wash.

SAGO. Nutritive and demulcent. Used in the form of gruel.

ST JOHN'S WORT. Vulnerary. The flowers were formerly an ingredient in FRYAR'S Balsam, and other similar compounds. An infused oil of the plant is sometimes used in liniments.

SAL AMMONIAC. See MURIATE OF AMMONIA.

SAL PRUNELLA. *Fused Nitre.* Its uses and doses are the same as nitre.

SALICIN. Tonic. Dose, 10 to 30 grs.

SALT, COMMON, or CULINARY. In small doses it is tonic, digestive, and alterative; in large doses purgative and vermifuge. As a digestive, 1 oz. may be sprinkled on the horse's corn. As a purgative, or to expel worms, the dose may be from 4 to 6 oz. "Its cathartic action is uncertain, often violent, and usually accompanied by considerable irritation of the kidneys."—FINLAY DUN. It is also a common ingredient in laxative clysters. For cattle, an ounce or more may be sprinkled on the hay, to assist digestion; as a purgative, 4 to 8 oz. may be given, but it is not suitable in inflammatory or febrile diseases. Sheep require 2 oz. as a purgative; or smaller doses daily as a preventive of the rot. To dogs, a teaspoonful or one and a half teaspoonfuls will act as an emetic; smaller doses as a vermifuge. Half a teaspoonful of a solution of salt, as strong as it can be made, is given to poultry as an emetic in roup. Externally, salt dissolved in water is used as a discutient, as a stimulant to old strains, and as a collyrium in chronic ophthalmia.

SALTS, EPSOM AND GLAUBER'S. See EPSOM SALTS, and GLAUBER'S SALTS.

SARSAPARILLA AND CHINA ROOT are diaphoretic and alterative; but seldom used in veterinary practice.

SAVIN. An acrid stimulant. The powder is given in doses of 1 to 2 drs. (with or followed by aloes) for worms, but its efficacy is doubtful. Long-continued use of savin is reported to have occasioned the hair to fall off. Externally it is applied, in powder or ointment, to warts.

SCAMMONY. An uncertain as well as expensive purgative, far inferior to aloes.

SCUTELLARIA. *Skull-cap.* By Mr YOUATT and others this plant was formerly employed as a preventive of hydrophobia. Dose, 40 grs. daily, gradually increased.

SEA WATER. Laxative. Dose, 2 or 3 pints.

SEDATIVES. Medicines which produce quiet, and relieve pain.

SENA. Purgative; but rarely used in veterinary practice. 5 or 6 oz. are required to purge a horse.

SERPENTARY. Stimulant, tonic, diaphoretic, and antiseptic. It was once supposed to counteract the effects of the bites of serpents. Dose, from $\frac{1}{2}$ oz. or 1 oz. or more; but rarely used.

SETONS. These consist of cord, tape, or a mixture of horse-

hair and hemp twisted together; they are inserted through a portion of the skin to excite irritation and discharge. Mr MORTON uses cotton cord soaked in a cantharidal liquid. See No. 16, BLISTERING LINIMENTS, *Veterinary Formulary*.

SIMARUBA. Tonic and stomachic, for the same purposes as gentian. Seldom used.

SOAP. Antacid and diuretic. Dose, $\frac{1}{2}$ oz. to 2 oz.

SODA. *Prepared Natron*. Carbonate, or Subcarbonate of Soda. The common washing soda is generally sufficiently pure. Antacid and diuretic. Dose, 2 to 4 drs. It is sometimes added to aloes as a corrective of acidity, and to tonics in weakness of the stomach. The bicarbonate of soda is milder, and may be given in larger doses.

SODA, CHLORIDE OF. See CHLORIDE OF SODA.

SODA, HYPOSULPHITE OF. Action, uses, and doses, as SULPHITE OF SODA.

SODA, SULPHATE OF. See GLAUBER'S SALTS.

SODA, SULPHITE OF. *Sodium Sulphite*. Internally:—Antiseptic, disinfectant, and alterative. Given in indigestion, tympanitis, and in so-called blood diseases. Externally:—Antiseptic, deodorising, and disinfectant. Employed in ulcerated sore-throat, and phagedænic wounds and ulcers. Dose for the horse, $\frac{1}{2}$ to 1 oz.; cattle, the same; sheep, 2 to 6 drs.; pig, 2 to 6 drs.; dog, 20 to 60 grs. Frequently repeated.

SODIUM, CHLORIDE OF. See SALT, COMMON.

SOOT. Some French veterinarians prescribe from 2 to 3 oz. of soot as a vermifuge. Also used externally in mange, &c. We presume wood-soot is intended.

SPERMACEI. Demulcent and pectoral. Dose, $\frac{1}{2}$ oz. to horses in cough; and to cows after calving. Externally emollient, in ointments.

SPIDER'S WEB. Externally, styptic. Internally, has been given to dogs in convulsive fits in $\frac{1}{2}$ -gr. doses.

SPIRIT OF HARTSHORN. This ammoniacal liquor is stimulant, antacid, and antispasmodic. Dose, $\frac{1}{2}$ oz. diluted. But more frequently used in stimulating liniments, and as an application to the bites and stings of venomous reptiles and insects. A diluted solution of Caustic Ammonia is mostly used as a substitute.

SPIRIT OF SAL VOLATILE. This also owes its pungency to ammonia. Dose, 1 oz. diluted.

SPIRITS, ARDENT. Brandy, gin, and rum are given as stimulants and antispasmodics, especially in colic. Dose, from 2 to 4 or 5 oz. with warm water. Rectified spirit of wine may be given in the same way, in smaller doses (1 to 2 oz.); but is more commonly employed for making tinctures; and externally in lotions.

SPIRIT OF MINDERERUS. See ACETATE OF AMMONIA, SOLUTION.

SPIRIT OF NITRE, SWEET. *Spirit of Nitrous Ether.* Diuretic, diaphoretic, and antispasmodic. Dose for horses, in fever, 1 oz., 3 times a day; in colic, from $\frac{1}{2}$ oz. to 2 oz.; cattle, $\frac{1}{2}$ oz. to 1 oz. in low fevers; sheep, 1 dr.; dog, from 10 to 20 drops. In all cases diluted.

SQUILL. A stimulating expectorant. Dose for a horse, 1 dr.; for cattle, $1\frac{1}{2}$ to 2 drs. It is also applied in frictions to the abdomen. MOIROUD states that it removes ascites.

STARCH. Demulcent. Chiefly used in clysters, but sometimes also in drinks. Dose, 1 to 2 oz., rubbed smooth with a little cold water, and then boiled with 3 or 4 pints of water. It is occasionally used in fomentations.

STAVESACRE SEEDS. Poisonous. 2 drs. will destroy a horse. Only used *outwardly* to destroy vermin, either powdered and mixed with grease, or infused in vinegar.

STEEL, SALT OF. See IRON, SULPHATE OF. For the other (so-called) preparations of steel, see IRON.

STIMULANTS. See EXCITANTS. Diffusible stimulants are those which produce a sudden and temporary excitement of the circulation and of the nervous system.

STOMACHICS. Medicines which invigorate the stomach and promote digestion.

STOPPINGS. Compositions employed to keep the feet moist and supple. The term is also applied to mechanical plugs for the feet when they are dry and diseased, as cow-dung, clay, tar, &c.

STORAX. Balsamic and expectorant. Dose, $\frac{1}{4}$ oz. Rarely used.

STRYCHNIA. The active principle of *nux vomica*; chiefly used in paralysis. Dose, 1 to 3 grs.; to be very cautiously increased if necessary: 15 grs. have proved fatal. Dose for the dog, 1-16th to 1-8th gr.

STYPTICS. Astringent applications employed locally to stop bleeding.

SUBLIMATE, CORROSIVE. *Perchloride of Mercury*. See CORROSIVE SUBLIMATE.

SUGAR, SYRUP, AND TREACLE. These are used to sweeten drinks; and to give form to balls and other compounds.

SUGAR OF LEAD. See ACETATE OF LEAD.

SULPHATE OF COPPER. *Blue Stone*. See COPPER, SULPHATE OF.

SULPHATE OF IRON. See IRON, SULPHATE OF.

SULPHATE OF MAGNESIA. See EPSOM SALTS.

SULPHATE OF POTASH. Purgative; but seldom used. Dose, 2 to 4 oz., in colic, &c.

SULPHATE OF QUININE. See QUININE, SULPHATE OF.

SULPHATE OF SODA. See GLAUBER'S SALTS.

SULPHATE OF ZINC. *White Vitriol*. See ZINC, SULPHATE OF.

SULPHUR. *Brimstone*. It is in three forms—roll brimstone, flowers of sulphur, and black brimstone or sulphur vivum. The flowers are generally used. The black is very impure, and sometimes contains arsenic. Sulphur is laxative, alterative, and pectoral. Dose to horses, as an alterative in skin diseases, grease, want of condition, &c., 1 oz.; as a laxative, 4 or 5 oz., but it is rarely employed with this view, and very large doses are not always safe; to cattle, as a laxative, 6 or 8 oz.; sheep, 2 or 3 oz.; dogs, 1 dr. in milk; swine, 2 drs. It is used *outwardly* in ointments, for mange in all animals. As an alterative it is usually combined with antimonials and nitre.

SULPHURET OF IRON. *Sulphide of Iron*. It has been used in hæmorrhage dysentery, and worms. The hydrated persulphuret (see FERRI PERSULPHURETUM HYDRATUM, *Pocket Formulary*) is strongly recommended by BOUCHARDAT as an antidote for metallic poisons; also as a remedy for incipient farcy. Dose, 1½ oz. to 8 oz.

SULPHURET OF MERCURY. *Sulphide of Mercury*. See ETHIOPS MINERAL, and CINNABAR.

SULPHURET OF POTASH. *Potassii Sulphidum*. *Sulphuret of Potassium*. Dose, 1 dr., with astringents, in diabetes and skin diseases. In large doses it is poisonous.

SULPHURIC ACID. *Poisonous*. The strong acid (*Oil of Vitriol*) is used as a powerful caustic. It is also used in ointments, or mixed with tar to form an external application. In small doses, about 1 to 2 drs., plentifully diluted;

- it is rarely given as a tonic. The diluted acid (1 oz. to a pint) is used as a lotion in grease and foul ulcers.
- SUPER TARTRATE (BITARTRATE) OF POTASH.** See CREAM OF TARTAR.
- SYRUP OF BUCKTHORN.** Purgative. Seldom given to horses, except when used in forming powders into balls. Dose for cattle, 2 to 4 oz. with castor oil. A common physic for dogs; dose, 2 to 4 drs.
- SYRUP OF POPPIES.** A mild anodyne and sedative. Dose for dogs, 1 dr. Seldom given to large animals.
- TANNIN.** *Tannic Acid.* The astringent principle of nut-galls. A powerful astringent in diarrhœa. Dose for horses and cattle, 20 grs. to 1 dr. Sheep and dogs, 1 to 3 grs. Catechu is more generally used.
- TANSY.** Tonic and vermifuge. *Externally* in fomentations.
- TAR.** *Internally* in old coughs, from 2 to 4 drs., but now seldom used. *Externally* it is cleansing and gently stimulating. It is particularly useful in thrush and all diseases and wounds of the feet, both of the horse and cattle, to punctured wounds, and for the cure of mange and other skin diseases. Mixed with fish oil, it is applied with a brush to hard, brittle feet. Tar water (see *Pocket Formulary*) is also given in chronic coughs. Oil or spirit of tar is used in mange ointments, and as a dressing for sheep. The latter requires some caution, sheep having been killed by it.
- TAR, BARBADOES.** See BARBADOES TAR.
- TARTAR EMETIC.** See ANTIMONY, TARTARIZED.
- TEA.** "Tonic, in simple indigestion, or when connected with staggers" (DELA FOND). Dose, 4 to 6 drs. infused in 3 or 4 pints of water.
- TEMPERANTS.** Medicines which moderate the circulation, and reduce animal heat.
- TIN.** Vermifuge. A drachm of the filings daily to dogs. A horse requires from 1 to 3 oz.
- TOBACCO.** *An acro-narcotic poison.* In small doses, diuretic and emetic, and of great service in colic, tympanitis and hernia. For *external application*, or for an enema, the infusion should be made with about 1 dr. of tobacco to a pint of hot water. Used as a wash for the mange, and to destroy lice and fly in sheep. But it is not altogether safe, as it is apt to be absorbed. It produces vomiting in

the dog, pig, and cat; but there are safer emetics. Herbivorous animals are less readily affected by it, but instances of its having proved fatal to them are recorded. In some parts of France, jockeys are said to stupefy vicious horses for sale, by tobacco infused in spirits.

TONICS. Medicines which give tone to the fibres, and invigorate the system when relaxed and debilitated. The principal tonics used in veterinary medicines are gentian, Peruvian bark, chamomile and other vegetable bitters and astringents; and the preparations of iron, copper, arsenic, zinc, &c. The over-free use of them, particularly when fever and inflammation are present, is a frequent source of mischief.

TORMENTIL ROOT. Astringent. Dose, 1 to 1½ oz. Its presence in pastures is supposed to prevent the rot in sheep.

TURPETH'S MINERAL. *Subsulphate of Mercury.* An irritating purgative, and in large doses *poisonous.* Dose, ½ dr. in farcy. Given to dogs as an emetic; dose, 1 to 3 grs., but going out of use.

TURMERIC. A weak aromatic stimulant. Supposed to be useful in jaundice or yellows. Dose, 1 oz.

TURPENTINES. They are all stimulant, diuretic, and expectorant; and in larger doses, vermifuge and purgative. Dose of common turpentine, ½ oz. to 1 oz. They are used in digestive ointments. Oil or spirit of turpentine is a more stimulating diuretic, in doses of 2 to 4 drs.; it is also considered efficacious as an antispasmodic in colic (gripes), and as a remedy for worms. Dose for the latter purposes, from 2 to 4 oz., or sometimes still larger doses. To cattle (in hooze, from worms in the bronchial passages), about 2 oz. To sheep, in rot, 1 dr. It is not a safe medicine for dogs, but is sometimes given in doses of 2 drs. with olive oil. *Externally* it is used in stimulating liniments, embrocations, ointments, &c. It is very irritating to the skin of the horse, and also of the dog, instantly producing great excitement. Like the common and Venice turpentine, it enters into the composition of some digestive ointments.

UVA URSI. *Bearberry.* Astringent. Dose, 4 to 6 drs. in diabetes. GIRARDE says it inflames the stomach.

VALERIAN. A stimulant acting chiefly on the nervous system. Dose for horses and cattle, 1 to 4 oz. in powder.

2 oz. twice a day have been given to a horse without any observable effect. In dogs it is said to act as a vermifuge. Dose, 1 to 4 drs.

VERDIGRIS. *Subacetate (or Diacetate) of Copper.* Tonic, caustic, *Poisonous.* It has been given in doses of 1 to 2 drs. daily, in farcy and glanders. *Externally* detergent and caustic, in ointment, and in the form of *Ægyptiacum.* The crystallized acetate of copper is more powerful in its action.

VERJUICE. Properties and uses the same as of vinegar; but preferred by some for outward use.

VERMILION. See CINNABAR.

VINEGAR. Diaphoretic, cooling, and antiseptic. In combination with honey, it is used in coughs. In large quantities, it irritates the stomach; a pint is said to have destroyed a horse. It should always be plentifully diluted. Vinegar which contains much sulphuric acid should be avoided. It is chiefly used as an external application, as a lotion for strains, bruises, sprains, and inflammation; and hot as a revulsive. The vapours are thought to possess disinfecting properties, but are less effectual than chlorine.

VITRIOLATED ZINC. *White Vitriol.* See ZINC, SULPHATE OF.

VITRIOL, BLUE. See SULPHATE OF COPPER.

WALNUT. The green shells are astringent, and sometimes applied, bruised, as a cataplasm, or in a decoction as a lotion.

WATER. Besides its use as a drink, and as a vehicle for medicine, water is used remedially, on the hydropathic system. Rags wetted with cold water, and well covered with dry ones, are used to produce perspiration, their operation being assisted by copious draughts of cold water, adding 4 oz. of sweet spirit of nitre to each pailful. This treatment is said to have succeeded in epidemics of pleuro-pneumonia.

WAX. Chiefly used in making cerates, plasters, charges, &c.

WHEY. A cooling and nutritive drink in inflammatory diseases, and during convalescence from them.

WILLOW BARK. Possesses in some degree the same properties as Peruvian bark. Dose, in powder or decoction, 1 to 4 oz.

WINE. Stimulant. In wine countries it is frequently given

- as a restorative. Port wine has been given as an astringent in obstinate diarrhœa. Dose for horses and cattle, $\frac{1}{2}$ pint to a bottle.
- WINTER BARK.** A warm tonic and stomachic. Dose, 2 to 6 drs.
- WOLFSBANE.** *Aconite, Monkshood.* A virulent poison. See **ACONITE.**
- WORMWOOD.** A bitter tonic and vermifuge. An infusion of from 2 to 4 oz. of the dry, or twice as much of the fresh herb, may be given in dropsy, and diseases of general debility; or from 2 to 4 drs. of the powder may be given in a ball. A few drops of the essential oil are often added to aloes, for worms.
- WORT.** See **MALT.**
- YEW.** It is not used medicinally. The leaves are poisonous to horses and cattle, producing symptoms which resemble those of apoplexy. To counteract its effect, it is recommended to give 10 grs. of croton meal, and afterwards drenches of gruel and vinegar. The croton to be repeated in 6 hours if it has not operated.
- ZEDOARY.** A weak aromatic stimulant, formerly prescribed in jaundice, but now rarely employed. It is weaker than ginger.
- ZINC, CARBONATE OF.** Applied externally as a desiccant, stimulant and astringent.
- ZINC, CHLORIDE OF.** In solution, this constitutes **SIR W. BURNETT'S** disinfecting fluid. Much diluted it is applied as a detergent lotion to foul ulcers. The dry salt is a powerful caustic.
- ZINC, OXIDE OF.** *Flowers of Zinc.* A mild astringent and tonic; dose $\frac{1}{2}$ oz.; but chiefly used in dusting ulcers and excoriations, to promote skinning.
- ZINC, SULPHATE OF.** *White Vitriol.* Tonic and astringent. Dose for the horse, 1 to 4 drs., frequently combined with cantharides. *Externally,* astringent, deterrent, styptic, and healing; in lotions and ointments, to indolent ulcers, and grease; and it is a frequent ingredient in eye-waters,—about 3 grs. to an ounce of water. A saturated solution is used as an injection for quitters.

VETERINARY FORMULARY*

Medicines for Horses

BALLS AND BALL MASSES

THE roots, seeds, and other dry substances are to be reduced to powder; and it is of importance that the aromatic seeds, especially, should have been recently powdered. The drugs should be of good quality. It is hoped that the trash sold as horse-powders will not much longer be known in establishments which have any pretensions to respectability. After this general notice it will be unnecessary to occupy space by repeating the words "powdered," "freshly powdered," "genuine," &c. Balls should not be too hard, but merely stiff enough to retain their form, and should be wrapped in soft paper.

Mode of Administering Balls.—The horse should be backed into the stall, the tongue drawn gently out with the left hand on the off side of the mouth, and then fixed by pressing the fingers against the side of the lower jaw. The ball, being now taken between the tips of the fingers of the right hand, must be passed rapidly up the mouth, as near the palate as possible, until it reaches the root of the tongue; it must then be delivered with a slight jerk, so that, the hand being immediately withdrawn, and the tongue liberated, the ball may be forced through the pharynx into the œsophagus. A slight tap under the chin may then be given, or a draught of water, to assist in carrying it down.

* See also VETERINARY MATERIA MEDICA.

COMMON MASS, as a basis for balls in general. Mix with the hand equal weights of linseed meal and treacle, and add a little palm oil.—CHERRY.

ALTERATIVE BALLS. The term *alterative* is applied to medicines which, without any sensible operation, or with a laxative or diuretic operation so gradual as not to interfere with the usual work or diet, produce a favorable change in the system, and, in common language, "purify the blood." Alterative balls are given in skin diseases, swelled legs, grease, foul humours, &c.; usually 1 daily, or every other day.

Diuretic Alterative Balls. 1. Dried common soda 1 oz., Castile soap 6 drs., resin 2 oz., liquorice powder $\frac{1}{2}$ oz., Barbadoes tar to form 6 balls; 1 daily.—WHITE.

2. Acetate of potash $\frac{1}{2}$ oz., resin $\frac{1}{2}$ oz., fenugreek 1 oz., treacle enough to form a mass for 2 balls; 1 daily.

Laxative Alterative Balls. 1. Aloes 4 oz., soft soap 4 oz., common mass 24 oz.; mix; dose, 1 oz.—V. C.

2. Socotrine aloes 8 oz., soft soap 8 oz., common mass, 16 oz.; mix; dose, 1 oz.—V. C.

3. Aloes 10 drs., soap 12 drs., caraways 12 drs., ginger 4 drs., treacle q. s. for 4 balls; 1 daily.—WHITE.

4. Aloes 1 dr., diuretic mass (see balls, No. 1 or 3) 9 drs.

5. Antimonial powder 1 dr., aloes 1 or 2 drs., diuretic mass (see balls, No. 1 or 3) 1 oz.

Antimonial or Diaphoretic Alterative Balls. 1. Levigated antimony 2 or 4 drs., caraway seeds 4 drs., treacle q. s. to form a ball.—WHITE.

2. Prepared antimony 2 drs., nitre 3 drs., sulphur 2 drs., linseed meal 2 drs., palm oil to form a mass; one every night, in megrims.—CLATER.

3. Tartarized antimony 2 drs., elecampane 2 oz., guaiacum 6 drs., sulphur 1 oz., treacle and flour to form 6 balls; one daily.

4. Tartarized antimony 3 drs., ginger a scruple, soap 1 oz.; for 3 balls; 1 every other morning.—VINES.

5. Emetic tartar 5 oz., ginger 3 oz., opium 1 oz., and syrup to make 16 balls.

Mercurial Alterative Balls. 1. Ethiops mineral 4 oz., sulphur, prepared antimony, cream of tartar, cinnabar, of

each 5 oz., honey to form a mass for 18 balls; 1 every morning for a month, in farcy.—TAPLIN.

2. Calomel $\frac{1}{2}$ dr., aloes 1 dr., Castille soap 2 drs., oil of juniper 30 drops, syrup to form a ball.—WHITE.

3. Blue pill 1 dr., black antimony 2 drs., diuretic mass 4 drs., aloes 1 dr.; for a ball daily.

4. In grease: prepared antimony, sulphur, nitre, Ethiops mineral, of each 3 oz., Castile soap 10 oz., oil of juniper 3 drs., syrup of honey q. s. for 18 balls; 1 every morning for 2 or 3 weeks.—TAPLIN.

5. Dr COLLIER'S Blue Pill.—Quicksilver 2 parts, peroxide of iron 1 part, confection of roses 3 parts. Rub together till the quicksilver disappears. Dose, $\frac{1}{2}$ dr. to 2 drs., with common or other mass q. s.

6. Strong mercurial ointment $\frac{1}{4}$ lb., powdered ginger 3 oz., liquorice powder 10 oz., treacle to mix for 12 balls.—FRANCIS.

Alterative Tonic Balls. See TONIC BALLS.

ASTRINGENT BALLS. These are given in diarrhoea and diabetes.

1. (V. C. Astringent Mass.) Catechu 1 oz., cinnamon 1 oz., common mass 6 oz.; mix; dose, 1 oz.

2. Peruvian bark 12 oz., grains of paradise 2 oz., gentian 3 oz., honey q. s. for 16 balls; 1 every morning; for diabetes.—RYDING.

3. Catechu $\frac{1}{2}$ oz., alum 3 drs., cascarilla 2 drs., flour 2 drs., treacle q. s.—WHITE.

4. Catchu 2 drs., opium $\frac{1}{2}$ dr., linseed meal 2 drs., treacle to form a ball. For profuse staling, 1 night and morning; if they confine the bowels, add 1 dr. of aloes.—CLATER.

5. Peruvian bark $\frac{1}{2}$ oz., alum $\frac{1}{2}$ oz., treacle q. s. For the same purpose.—LAWRENCE.

6. Oak bark 1 oz. (or Peruvian bark $\frac{1}{2}$ oz.), opium 1 dr., ginger 2 drs., syrup to form a ball; for diarrhoea.—WHITE.

7. Opium $\frac{1}{2}$ dr., prepared chalk 6 drs., cassia $1\frac{1}{2}$ drs., tartarized antimony $\frac{1}{2}$ dr., syrup to form a ball; for the same.—WHITE.

8. Nut-galls 2 drs., cassia $\frac{1}{2}$ dr., conserve of roses to form a ball.

9. Burnt rhubarb 1 dr., compound powder of chalk 3 drs., common mass 6 drs.; for diarrhoea.

10. Tormentil or bistort $1\frac{1}{2}$ drs., marshmallow root, $\frac{1}{2}$ oz., chalk 2 dr., syrup to form a ball.

11. For bloody urine. Acetate of lead 10 grs., sulphate of zinc 40 grs., catechu $\frac{1}{4}$ drs., conserve of roses to form a ball; once daily.—BLAINE.

12. Powdered opium $\frac{1}{2}$ dr., soda 1 dr., powdered cassia or ginger $1\frac{1}{2}$ drs., flour and syrup to form a ball.

13. For diabetes. Catechu $\frac{1}{2}$ oz., alum $\frac{1}{2}$ dr., sugar of lead 10 grs., with conserve of roses to form a ball. See also TONIC BALLS.

COUGH BALLS; Expectorant Balls. The following formulæ are chiefly intended for chronic coughs and thickness of wind. The bowels should be kept open by mashes and an occasional laxative. Coughs occasioned by worms require a different treatment. In coughs connected with inflammation of the chest, and epidemic catarrh, see BALLS FOR INFLAMMATION OF THE LUNGS, and BRONCHITIS.

1. Aloes 2 oz., digitalis (powdered) 1 oz., common mass 13 oz.; dose, 1 oz. twice a day.—MORTON.

2. Emetic tartar $\frac{1}{2}$ dr., digitalis $\frac{1}{2}$ dr., nitre $1\frac{1}{2}$ drs., tar enough to form a ball; every night.—YOUATT.

3. Powdered squill 1 dr., gum ammoniac 3 drs., opium $\frac{1}{2}$ dr., syrup to form a ball.—WHITE.

4. Ipecacuanha 1 dr., camphor 2 drs., liquorice powder 1 dr., honey to form a ball; to be given every morning.—BLAINE.

5. Sulphur $\frac{1}{2}$ oz., assafœtida 1 oz., liquorice powder 1 oz., Venice turpentine 1 oz., for 4 balls; one every night for 4 times.—HINDS.

6. Calomel 26 grs., gum ammoniacum 2 drs., balsam of Peru 1 dr., p. squill 1 dr., honey to form a ball; one every morning.—BLAINE.

7. Powd. marshmallow root and liquorice of each 1 dr., elecampane, sulphur, and kermes mineral, of each $\frac{1}{2}$ dr., honey to form a ball; twice a day.—LEBAS.

8. Squill 2 drs., gum ammoniac 4 drs., ipecacuanha 4 drs., opium 4 drs., pimento 1 oz., balsam of sulphur 4 oz., Castile soap 2 oz., treacle to form a mass for 6 balls; one twice a day.—HINDS.

9. Spermæti 1 oz., balsam of copaiva 1 oz., benzoin 2 drs., sulphur 2 oz., elecampane 2 oz., p. squill 4 drs., emetic

tartar 2 drs., syrup of poppies to form a mass for 8 balls.
—B. CLARKE.

10. Liquorice powder $\frac{1}{2}$ oz., linseed or barley meal 1 oz., tar 1 dr., honey to form a ball.

11. Castile soap, aniseed, liquorice, of each 5 oz., Barbadoes tar 6 oz., ammoniacum 3 oz., balsam of tolu 1 oz., honey q. s. to make a mass for 12 balls; one every morning for a fortnight.—TAPLIN.

12. Digitalis 1 dr., nitre 2 drs., liquorice 4 drs., tar enough to form a ball.—CLATER.

13. Digitalis $\frac{1}{2}$ dr., camphor 1 dr., emetic tartar 1 dr., nitre 3 drs., linseed meal 1 dr., make up with Barbadoes tar, and give one daily.—SPOONER. See also MIXED BALLS (PECTORAL CORDIAL).

14. For chronic cough. Ext. Belladonna $\frac{1}{2}$ to 1 dr., aloes Barbadoes in powder 1 dr., nitre 2 drs. Common mass to form into a ball.—TUSON.

BALLS FOR INFLAMMATION OF THE LUNGS, BRONCHITIS, &c.

1. Antimonial powder 2 drs., digitalis 1 dr., nitre 3 drs., cream of tartar 3 drs., honey to form a ball; one every 4, 6, or 8 hours, in inflammation of the lungs.—BLAINE.

2. Digitalis 1 dr., emetic tartar $1\frac{1}{2}$ drs., nitre 3 drs., honey q. s.; when the pulse intermits, reduce the dose to half.—YOUATT.

3. Nitre 6 drs., emetic tartar $\frac{1}{2}$ dr., flour and syrup to form a ball; twice a day.—WHITE.

4. Digitalis 1 dr., emetic tartar 1 dr., nitre 3 drs., sulphur 1 dr., linseed meal 2 drs.; beat together with palm oil.—CLATER.

5. *Epidemic catarrh*. To the last add 2 drs. of the Physic Mass No. 10 (see PHYSIC OR PURGING BALLS); repeat this twice.

6. For pneumonia. White Hellebore $\frac{1}{2}$ dr. (or extract of belladonna 1 dr., or digitalis 1 dr., or calomel 1 dr., with opium $\frac{1}{2}$ dr.), emetic tartar 1 dr., nitre and linseed meal each 2 drs.; one twice a day.—SPOONER.

7. In the advanced stage, when suppuration has taken place. Carbonate of ammonia $1\frac{1}{2}$ drs., opium 1 dr., aniseed $\frac{1}{2}$ oz., syrup to form a ball.—SPOONER.

8. *Cough Ball*. Digitalis $\frac{1}{2}$ dr., camphor and emetic

tartar, each 1 dr., nitre 3 drs., and linseed meal 1 dr., to be made up with Barbadoes tar.

9. A scruple of calomel and a drachm of opium made into a ball, every three hours.

CORDIAL BALLS. For exhaustion, from over-exertion and as a stimulant to weak stomachs. Their frequent and unnecessary use is hurtful.

1. Ginger and gentian, equal parts, treacle to form a mass; dose, 1 oz. to 1½ oz.—V. C.

2. Caraway, bruised raisins, of each 4 parts, ginger and palm oil, of each 2 parts.—YOUATT.

3. Aniseed, caraway, cardamom, each 1 oz., saffron 2 drs., sugar candy 4 oz., liquorice powder 1½ oz., Spanish juice (softened with water) 2 oz., oil of aniseed ½ oz., wheat flour q. s.; dose, 1 oz. to 1½ oz.—BRACKEN.

4. Aniseed, caraway, sweet fennel, liquorice, of each 4 oz., of ginger and cassia, each 1½ oz., honey to form a mass; dose, 1 oz.—WHITE.

5. Ginger, caraway, each 4 lbs., gentian 1 lb., palm oil 4½ lbs., beat together; dose, 1 oz. to 1½ oz.—CLATER.

6. Gentian 8 oz., ginger 4 oz., coriander 8 oz., caraway 8 oz., oil of aniseed ½ oz., treacle q. s.; dose, 1½ oz.—BLAINE.

7. Aniseed, caraway, ginger, each 8 oz., gentian, grains of paradise, cumin, and turmeric, each 4 oz., cassia 2 oz., oil of caraway 2 drs., treacle to form a mass; dose, 1½ oz. To keep it moist add 2 oz. of acetate of potash.

8. Cumin, aniseed, caraway, each 4 oz., ginger 2 oz., treacle q. s.; dose, 1½ oz. to 2 oz.—WHITE.

9. Pimento 1 lb., sifted barley meal 2 lbs., treacle q. s.; dose, 1½ oz.—B. CLARKE.

MIXED BALLS. *Cordial Astringent Ball.* Cordial ball (No. 2) 1 oz., catechu 1 dr., opium 10 grs.; to washy horses, before or after a journey.—YOUATT.

Cordial Anodyne Balls. 1. Cordial mass (No. 6) 10 drs., camphor 1 dr., opium 20 grs.—BLAINE.

2. Opium ½ dr. to 2 scruples, soap 2 drs., ginger 1 dr., aniseed 4 drs., oil of caraway ½ dr., treacle q. s.—WHITE.

Balsamic Cordial Ball. Cordial mass (No. 6) 1 oz., myrrh 1 dr., balsam of tolu 1 dr.—BLAINE.

Pectoral Cordial Balls. 1. For old coughs. Fenugreek,

aniseed, cumin, safflower, elecampane, coltsfoot, sulphur, of each 3 oz., liquorice juice 1 oz., olive oil 8 oz., honey 8 oz., Genoa treacle 12 oz., oil of aniseed 1 oz., wheat meal 1½ lbs., or q. s.; one ball, or 2 oz. (dissolved in water or warm wort), every day for 12 or 15 days if required.—**QUINCE.**

2. Elecampane ½ oz., ginger 1½ drs., squill 1 dr., oil of aniseed 20 drops, syrup of tolu q. s.—**WHITE.**

Diuretic Cordial Balls, to fine the legs of debilitated and overworked horses, and sometimes given in old coughs, &c.

1. Resin 2 oz., soap, nitre, caraway, of each 2 oz., ginger 1½ oz., sulphur 2 oz., oil of caraway ½ dr., oil of juniper ½ dr., syrup to form a mass. Dose 1 oz.

2. Soap and common turpentine, each 4 drs., ginger 1 dr., opium ½ dr., caraway seed q. s. for 1 dose.—**WHITE.**

3. Strained turpentine 8 oz., resin 4 oz., olive oil 2 oz., soap 8 oz.; melt together and add powdered ginger 6 oz., pimento 6 oz., liquorice powder q. s. to form a mass; dose, 1 oz. to 1½ oz.

4. Resin 4 drs., nitre 2 drs., and ginger 1 dr., with sufficient soap to form a ball.—**SPOONER.**

DIURETIC BALLS. For swelled legs, grease, &c., for carrying off bad humours, and in many chronic diseases. The too frequent use of diuretics injures the kidneys and weakens the system. See **ALTERATIVE BALLS.**

1. Resin, soap, nitre, of equal parts, beaten together into a mass; dose, 1 oz. to 1½ oz.—**V. C.**

2. Common turpentine 4 oz., Castile soap 4 oz., caraways 8 oz., ginger 1 oz., flour q. s.; dose 1 oz. to 1½ oz.—**WHITE.**

3. Resin 16 oz., white soap 16 oz., nitre 8 oz., dried common soda 2 oz., oil of juniper 4 oz.; beat together, adding flour if required; dose, 1 oz. to 1½ oz.

4. Nitre 1 lb., Castile soap ½ lb., common turpentine 1 lb., barley meal 2½ lbs., or sufficient; dose, about 1 oz.—**B. CLARKE.**

5. White soap 8 oz., nitre 3 oz., resin 3 oz., camphor 3 drs., oil of juniper 3 drs. For 6 balls; 1 every, or every other morning.—**TAPLIN.**

6. Common turpentine 16 oz., sulphur 24 oz., nitre 8 oz., honey 8 oz., flour or linseed meal q. s.; dose, 1½ oz.

7. Camphor 2 drs., nitre 1 oz., flour and syrup to form a ball; for stoppage of water.—WHITE.

8. Yellow resin 4 lbs., common turpentine 2 lbs., yellow soap 2 lbs., melt together, and add nitre 1 lb. Dose, 1 to 1½ oz.—BLAINE.

9. Common turpentine (or powdered resin) ½ oz., linseed meal ¼ oz., ginger ½ dr., palm oil q. s.—YOUATT.

10. Yellow resin 2 oz., common turpentine 4 oz., soap 3 oz., melt together, stir in 1 oz. sweet oil, add oil of aniseed ½ oz., oil of juniper ½ oz., ginger 2 drs., linseed meal q. s.; mix, and divide into 8 balls; 1 a day till the water is affected.—HINDS.

11. Resin 2½ lbs., cream of tartar ½ lb., sulphur ½ lb., linseed meal 1 lb., palm oil 1 lb.; dose, 1 to 2 oz.—CLATER.

12. Nitre 1 oz., vermilion ½ oz., resin 1 oz., camphor ½ oz., honey q. s. for 4 balls.—LEBAS.

13. Powdered yellow resin 4 drs., nitrate of potash 2 drs., ginger 1 dr.; beat up with soap.—SPOONER.

14. Nitre 8 oz.; oxysulphuret of antimony 1 oz., sulphur 8 oz., resin 8 oz., oil of juniper 1 oz., yellow soap 8 oz., treacle to form a mass; dose, 1½ oz.

15. White soap 1 oz., extract of juniper berries q. s. for 2 balls.—BOURGELAT.

Tonic Diuretic Ball. Gentian 1 dr., ginger ½ dr., sulphate of iron 2 drs., diuretic mass (No. 11) ½ oz., oil of juniper 10 drops, syrup of squills ½ oz.; twice a day in dropsy of chest; less frequently in swelled legs.—CLATER.

Tonic and Diuretic Ball for Pleurisy. Sulphate of copper 1½ drs., ginger and gentian 2 drs. each, with Venice turpentine.—SPOONER. See also LEICESTER RED BALLS (MISCELLANEOUS BALLS).

FEVER BALLS.

1. Emetic tartar ½ dr., camphor ½ dr., nitre 2 drs., common mass 6 drs., or q. s. for 1 ball; to be given once or twice a day.—MORTON.

2. Camphor 1 dr., nitre 6 drs., antimonial powder 2 drs., flour and soap to form a ball.—WHITE.

3. Antimonial powder 2 drs., nitre 3 drs., cream of tartar 2 drs., honey to form a ball; in influenza twice a day, after a mild laxative.—BLAINE.

4. See BALLS FOR INFLAMMATION OF LUNGS, No. 4.—CLATER.

BALLS FOR FARCY AND GLANDERS. Mr COLEMAN says he has tried the various preparations of arsenic, antimony, copper, mercury, zinc, aconite, digitalis, hemlock, henbane, hellebore, nightshade, &c., in glanders, without any specific or curative effect. Mr YOUATT considers it useless to attempt the cure of glandered horses, but that farcy in its early stages and mild form may be successfully treated. Mr BLAINE says, "All the mercurials have been used with benefit in farcy; but they must be discontinued as soon as the mouth is affected, or sickness, loss of appetite, &c., produced." Mr FINLAY DUN says glanders is incurable, and recommends immediate slaughter. He afterwards adds that life may be prolonged by generous diet; and further recommends sulphate of copper and arsenic.

1. Ethiops mineral 2 drs., blue pill 1 dr., prepared antimony 3 drs., diuretic mass 4 drs. One every morning.

2. Strong mercurial ointment 2 to 3 drs., guaiacum 3 drs., soap 4 drs., fenugreek 12 drs., treacle to form a mass, for 6 balls. See MERCURIAL ALTERATIVE BALLS (under ALTERATIVE BALLS).

3. Sulphate of copper 1 dr., corrosive sublimate 8 grs., linseed powder $\frac{1}{2}$ oz.—WHITE.

4. Corrosive sublimate 10 grs., gentian 2 drs., ginger 1 dr., syrup to form a ball; to be given night and morning till some effect is produced; when the mouth is affected, the sublimate may be exchanged for 1 dr. sulphate of copper.—YOUATT.

5. Corrosive sublimate 10 grs., opium $\frac{1}{2}$ to 1 dr., powdered aniseed $\frac{1}{2}$ oz., with syrup to make a ball.

6. Sulphate of copper 1 dr., calomel 20 grs., common turpentine 3 drs., liquorice powder and syrup q. s. for one ball.—COLEMAN.

7. Sulphate of copper 1 dr., white arsenic 8 grs., corrosive sublimate 8 grs., linseed powder $\frac{1}{2}$ oz., syrup to form a ball.—WHITE.

8. Ethiops mineral 2 drs., opium 10 grs., liquorice powder and mucilage to form a ball; to be given twice a day till the breath or urine is affected.—HINDS.

9. Sulphate of iron 2 drs., Peruvian bark 1 oz., opium $\frac{1}{2}$ dr., syrup to form a ball.—SMITH.

10. Cantharides 4 grs., gradually increased to 6 or 8 grs., gentian, ginger, and caraway, each 1 dr.; syrup q. s.; every, or every other day.—VINES.

11. Sulphate of iron 2 drs., iodide of potassium 10 grs., ginger 1 dr., gentian 2 drs., made into a ball with treacle.—SPOONER.

12. Diniodide of copper 1 dr., gentian $1\frac{1}{2}$ drs., pimento 1 dr., cantharides 5 grs.; for one ball.—MORTON.

13. Sulphate of zinc 15 grs., cantharides 7 grs., pimento or ginger 15 grs., treacle and oatmeal to form a ball; 1 daily.—BRACY CLARK.

14. Sublimate, arsenic, verdigris, each 8 grs., sulphate of copper 20 grs. for one ball (with common mass q. s.); the dose may be gradually increased, carefully watching its effects, but should never exceed 15 grs. of sublimate and arsenic.—BLAINE.

15. Sublimate 10 grs., gentian 2 drs., ginger 1 dr., linseed meal $\frac{1}{2}$ oz., palm oil to form a ball; night and morning for a fortnight; for farcy.—CLATER.

16. Sulphate of copper $\frac{1}{2}$ dr. to 1 dr., ginger and gentian, each 1 dr., linseed meal and palm oil to form a ball; morning and night for a fortnight, then daily as long as necessary: in glanders.—CLATER.

17. Strong mercurial ointment 3 oz., white soap 2 oz., starch 2 oz., form a mass and divide into 12 balls; 1 every morning.—MOIROUD.

18. Assafœtida 3 oz., vermilion 2 oz., muriate of lime 3 drs., galangal 1 oz., strong mercurial ointment 2 oz.; beat together into a uniform mass, and divide into 6 balls; one every other morning.—LEBAS.

19. Ethiops mineral 8 oz., powdered burdock root 16 oz., treacle q. s.; make into 32 balls.—MOIROUD.

20. Antihecticum Poterii 2 drs., with 6 drs. of cordial ball; every other day.—LAWRENCE.

21. Calomel 1 oz., assafœtida 4 oz., galangal powder 1 oz., mercurial ointment 2 oz. Mix, and form 6 balls. One every other morning.—LEBAS.

22. Hydrargyro-iodide of potassium (see HYDRARGYRI ET POTASSII IODIDUM, *Pocket Formulary*) 3 oz., powdered althæa root, and honey q. s. to make 100 balls. Give

from one to eight daily, gradually increasing the dose to the latter number.—BOUCHARDAT.

BALLS FOR GREASE. See DIURETIC BALLS, and ALTERATIVE BALLS.

BALLS FOR YELLOWS OR JAUNDICE, AND INFLAMMATION OF LIVER (HEPATITIS).

1. For *hepatitis without purging** calomel 1 dr., antimonial powder 2 drs., aloes 3 drs., syrup to form a ball; one every four or five hours, till the bowels are opened.—BLAINE.

2. Calomel $\frac{1}{2}$ dr., aloes 1 dr., soap 2 drs., rhubarb $\frac{1}{2}$ oz., syrup to form a ball; to be given every 12 hours, till it purges moderately.—WHITE.

3. Aloes 2 drs., calomel 1 dr., syrup to make a ball, twice a day.—YOUATT.

4. Opium 1 dr., calomel 1 dr. emetic tartar 1 dr., liquorice powder 3 drs., syrup to form a ball; one every 12 hours.—WHITE.

5. Opium $\frac{1}{2}$ dr., calomel 1 dr., resin 3 drs., carbonate of potash 2 drs., with soft soap. To be preceded by blisters to the side, and purgative draughts.—SPOONER.

Yellows (Jaundice) without Fever. 1. Calomel 1 dr., aloes 2 drs., soap 2 drs.; for one ball; night and morning till purged, then so as to keep the bowels lax.—BLAINE.

2. Calomel $\frac{1}{2}$ dr., aloes $1\frac{1}{2}$ drs., Castile soap 2 drs., rhubarb 3 drs., syrup to form a ball.—WHITE.

3. In the latter stage, when not costive, calomel 12 grs., sulphate of copper 1 dr., gentian 3 drs., oak bark 3 drs., chamomile 3 drs., syrup to form a ball; once or twice a day.—BLAINE.

PHYSIC OR PURGING BALLS. The animal should be prepared by bran mashes for two days, and the ball given fasting in the morning. Gentle exercise with a ball is useful, but not after it begins to operate. Barbadoes aloes alone should be used (from the gourd, not melted), and the dose seldom need exceed 6 drs. A week should be allowed after the operation of one ball before

* "Nothing can be done without purgatives in jaundice."—FINLAY DUN.

another is given. See ALOES, in the *Veterinary Materia Medica*.

1. (V. C. Cathartic Mass.) Bruised Barbadoes aloes 8 oz., olive oil 1 oz.; melt together in a vessel placed in hot water; remove it from the fire, add 3 oz. of treacle, and stir all together; dose 6 to 12 drs., equal to 4 to 8 drs. of aloes.

2. (V. C. Stronger.) To each dose of the last add from 4 to 8 drops of croton oil.

3. Aloes Barbadoes in small pieces 8 parts, glycerine 2 parts, ginger in powder 1 part. Melt together in a water-bath, and thoroughly incorporate by constant stirring. If desirable, gentian may be substituted for ginger. Dose, from 6 to 8 drs.—TUSON.

4. Barbadoes aloes 4 to 8 drs., soap 3 to 4 drs., ginger 1 dr., oil of cloves 10 drops (or oil of caraway or aniseed 20 drops), water 1 dr. or q. s.; beat together into a mass.—WHITE. Mr WHITE says it is the best that can be employed.

5. Barbadoes aloes 15 oz., ginger 1 oz.; mix and beat up with 8 oz. of palm oil. Dose, 1 to 1½ oz.—YOUATT.

6. Barbadoes aloes 2½ drs. Cape aloes 12 drs., olive oil 4 drs.; treacle 12 drs.; dose, 7 to 14 drs.; mix as No. 1.—MORTON.

7. Barbadoes aloes 5 drs., 7½ drs., or 9 drs., oil of caraway 10 drops; made up with palm oil or lard.—Mr BLAINE'S Nos. 1, 2, and 3.

8. Melt Barbadoes aloes (in a tin vessel immersed in boiling water) with a fifth of its weight of treacle, and, while soft, pour it into paper moulds; 1 oz. is a full dose for a large-sized saddle or coach horse.—B. CLARK. [For a convenient apparatus for melting and casting these balls, see Mr BRACY CLARK'S *Pharmacopœia Equina*; or Vol. V (1st Series) of the *Pharmaceutical Journal*.

9. Barbadoes aloes 5 to 8 drs., cream of tartar 2 drs., oil of cloves 10 drops, treacle to form a ball.—PEALL.

10. Aloes 7 drs., Castile soap 4 drs., aromatic powder 1 dr., oil of caraway 6 drops; mucilage to form a ball.—HINDS.

11. Barbadoes aloes 7½ parts, Socotrine aloes 7½ parts, ginger 1 part; mix the powders, add 7½ parts of palm oil, and beat to a mass; keep it in a jar closely covered; dose, 1½ to 1¾ oz.—CLATER.

12. Barbadoes aloes $13\frac{1}{2}$ oz., lard 6 oz., treacle $1\frac{1}{2}$ oz., water $1\frac{1}{2}$ oz.; put them in an earthen vessel, placed in boiling water; mix, and form the mass into 18 balls.—MCEWEN.

13. Aloes and hard soap each 5 oz., pearl ashes 1 oz., powdered ginger 1 oz. Melt in a ladle and divide, while warm, into 8 balls.

Mercurial Physic Balls. 1. Physic mass (No. 10, page 53) 10 to 14 drs., calomel 1 to $1\frac{1}{2}$ drs.; mix.—CLATER.

2. For stomach staggers: aloes 1 oz., calomel $\frac{1}{2}$ dr., cascarilla 3 drs., syrup to form a ball.—WHITE.

LAXATIVE BALLS.

1. Ipecacuanha 1 dr., aloes 3 to 4 drs., liquorice powder and mucilage to form a ball.—HINDS.

2. Aloes 3 to 4 drs., soap 3 drs., oil of caraway 20 drops, syrup q. s.—WHITE.

3. Aloes 3 to 4 drs., soap 4 drs., emetic tartar 1 dr., mucilage to form a ball.—HINDS.

For other Formula see ALTERATIVE BALLS (LAXATIVE).

NAUSEATING BALLS. These are given in inflammatory diseases.

1. Powdered white hellebore $\frac{1}{2}$ dr., linseed meal 4 drs., treacle to form a ball; one night and morning till some effect is produced: in inflammation of the kidneys.—CLATER.

2. White hellebore 20 grs., common mass or other proper material to form a ball; give one every 4, 6, or 8 hours, till symptoms of nausea appear, taking care not to carry it too far.—PERCIVAL. See FEVER BALLS. See HELLEBORE, in *Veterinary Materia Medica*.

STOMACHIC BALLS. For indigestion, and during recovery from debilitating diseases which have impaired the appetite. A mild purge should be previously given.

1. Gentian, quassia, grains of paradise, of each 3 drs., Venice turpentine q. s. for 1 ball.—BLAINE.

2. Gentian 2 or 3 drs., carbonate of soda 1 dr., ginger 1 dr., treacle to form a ball.—WHITE.

3. Chamomiles 2 drs., calumba 2 dr., common salt 1 dr., fenugreek 2 drs., syrup to form a ball.

4. Myrrh $1\frac{1}{2}$ drs., cascarilla 2 drs., Castile soap 1 dr., syrup to form a ball.—WHITE.

5. *Laxative Stomachic Ball.* Aloes 3 drs., rhubarb 3 drs., carbonate of soda 2 drs., ginger $1\frac{1}{2}$ drs., treacle to form a ball.—WHITE.

6. Calumba and chamomile in powder, each 2 drs., Venice treacle $\frac{1}{2}$ oz., oil of caraway 25 drops, honey, q. s.—LAWRENCE. See TONIC BALLS, for other formulæ.

TONIC BALLS. In diseases attended with general debility, and to restore strength after a tedious illness.

Vegetable Tonics. 1. Peruvian bark $\frac{1}{2}$ oz., opium $\frac{1}{2}$ dr., ginger $1\frac{1}{2}$ drs., oil of caraway 20 drops, treacle to form a ball.—WHITE.

2. Sulphate of quinine 1 dr., gentian, oak bark, and honey, to form a ball.—MOIROUD.

3. Gentian 1 dr., ginger $\frac{1}{2}$ dr., cascarilla 1 dr., treacle and linseed meal to form a ball.—CLATER.

4. Myrrh 2 drs., mustard flour 1 dr., cantharides 5 grs., chamomile 4 drs., Venice turpentine q. s. for one ball.—BLAINE.

5. Gentian 4 drs., chamomile 2 drs., carbonate of iron 1 dr., ginger 1 dr., syrup q. s. for one ball.—YOUATT.

6. Quassia 2 drs., canella 2 drs., opium $\frac{1}{2}$ dr., ginger 1 dr., treacle q. s.—WHITE.

Mineral Tonics. 1. Sulphate of iron 4 oz., ginger 4 oz., common mass 10 oz.; beat together to form a mass; dose, 1 to $1\frac{1}{2}$ oz.—V. C.

2. Sulphate of iron $\frac{1}{4}$ oz., aromatic powder 2 drs., mucilage q. s. to form a ball.—WHITE.

3. Scales of iron 12 oz., gentian 8 oz., honey to form a mass.—MOIROUD.

4. Myrrh 3 drs., sulphate of iron 2 drs., chamomile 3 drs., ginger 1 dr., Venice turpentine or palm oil to form a ball.—BLAINE.

5. Gentian 4 drs., chamomile 2 drs., carbonate of iron 1 dr., ginger 1 dr., syrup for 1 ball.—YOUATT.

6. Sulphate of iron 2 drs., carbonate of potash 2 drs., cascarilla 2 drs., caraway 4 drs., treacle q. s.—WHITE.

7. Sulphate of iron 1 dr., carbonate of soda 2 drs., myrrh 1 dr., ginger 1 dr., cantharides 6 grs., caraway $\frac{1}{2}$ oz., treacle q. s.—WHITE.

8. Tonic mass. Sulphate of copper 2 oz., ginger 2 oz.,

common mass 12 oz., beat together; dose, 1 oz. to 1½ oz.—V. C.

9. Sulphate of copper and ginger, of each 1 dr., canella 4 drs., conserve of roses q. s. for one ball.—BLAINE.

10. White arsenic 5 to 10 grs., aniseed ½ oz., opium ¼ dr., treacle q. s.; sometimes 2 drs. of sulphate of zinc may be added.—WHITE.

11. Arsenic 10 grs., gentian and cascarilla, of each 3 drs., conserve of roses q. s.—BLAINE.

Mild Alterative Tonics. To promote condition; a mild dose of physic should be previously given.

1. Aloes 1 dr., winter's bark 2 drs., verdigris 1 dr., treacle or honey q. s.

2. Arsenic 8 grs., pimento 1 dr., extract of gentian 4 drs.; daily.

3. Nitre 1 oz., sulphur 6 drs., physic mass ½ oz., gentian 6 drs., ginger ½ oz., palm oil q. s. for 4 balls. One daily, after an attack of stomach staggers.—CLATER.

Tonic Condition Balls. Ginger and camphor, of each 1 dr., gentian and sulphate of iron, of each 2 drs.; make up with linseed meal.—SPOONER.†

WORM BALLS.

1. Calomel 1 or 2 drs. at night, and an aloetic ball in the morning.—CLATER.

2. Emetic tartar 1 dr., ginger 1 scruple, linseed meal and treacle to form a ball; 1 every morning an hour before feeding.—YOUATT.

3. Calomel 8 grs., arsenic 8 grs., tin filings 1 oz., Venice turpentine ½ oz.; mix, and give every morning fasting, for a fortnight.—BLAINE.

4. Common salt ½ oz., gentian 2 drs., rust of iron 2 drs., savin 1 dr., treacle to form a ball; to be given every morning for a week; then a purging ball.

5. Barbadoes aloes 6 drs., ginger 1½ drs., oil of wormwood 20 drops, carbonate of soda 2 drs., syrup to form a ball; ½ dr. or 1 dr. of calomel may be added, or given the previous night; to be repeated at intervals of 10 days if required.—WHITE.

6. Emetic tartar 1 dr., common mass 6 drs.; to be given for 6 mornings, and a purging ball on the seventh.

7. Assafoetida 2 drs., calomel 1 or 2 drs., savin 1½ drs., oil

of wormwood 20 drops, syrup q. s.; at night, and a physic ball in the morning.

8. Emetic tartar 1 dr., sulphur 1 dr., linseed meal 4 drs., palm oil to form a ball; 1 every morning after a mercurial physic ball.—CLATER.

9. *For long round worms.* Emetic tartar 1 dr., ginger $\frac{1}{2}$ dr., tin filings 6 drs., linseed meal 1 dr., palm oil to form a ball.

10. Assafoetida $\frac{1}{2}$ oz., gentian 2 oz., strong mercurial ointment 1 oz., honey to form a mass for 16 balls; 1 or more every morning.—LEBAS.

MISCELLANEOUS BALLS.

Garlic Ball. Beat garlic to a paste with enough linseed or liquorice to form a mass; dose, 10 drs.

Camphor Ball. Mix into a ball 2 drs. of camphor with liquorice powder and syrup enough to give it a proper consistence.

Iodine Ball. Iodine 5 grs., linseed meal 5 drs., palm oil to form a ball.

Ball to prevent Rabies. Skull-cap 2 scruples, belladonna $2\frac{1}{2}$ grs., form them into a ball, to be given night and morning; the second week 2 balls, the third 3 balls, and this continued for 6 weeks.—YOUATT.

Leicester Red Balls. Nitre 1 lb., resin 1 lb., common soda 2 oz., Castile soap $\frac{1}{2}$ lb., ginger 2 oz., oil of juniper 2 drs., cinnabar $\frac{1}{2}$ oz.; dose, $1\frac{1}{2}$ oz.

Ball for Appetite. Equal weights of assafoetida, saffron, bay-berries, and aloes, made into a mass with extract of gentian; dose, 1 oz.—LEBAS.

Anodyne Ball. Opium $\frac{1}{2}$ dr. to 1 dr., camphor 1 dr., aniseed $\frac{1}{2}$ oz., soft extract of liquorice q. s.—WHITE.

Antispasmodic Ball. Opium 1 dr., powdered belladonna 10 grs., linseed meal 3 drs., palm oil or treacle q. s.; twice or thrice a day, in spasm of the neck of the bladder.—CLATER.

Ball for Roaring. The Cough Ball, No. 12. may be tried; and the compound iodine ointment rubbed on the throttle for some weeks or months.

Stimulating Diaphoretic Ball. Emetic tartar 1 dr., ginger 2 drs., camphor $\frac{1}{2}$ dr., opium 2 scruples, oil of caraway 15 drops, honey to form a ball; for hide-bound and unhealthy coat without any other disease.—WHITE.

BARTLETT'S Perspirative Ball. Dover's powder 3 drs., camphor 1 dr., treacle q. s.

HIND'S Sweating Ball. Emetic tartar, 1 dr., assafœtida 1 dr., liquorice powder and syrup to form a ball; repeat in 12 hours if required.

Grease Ball. Liver of antimony 16 oz., salt of tartar 8 oz., gum guaiacum, fenugreek, parsley seed, of each 4 oz., treacle to form a mass; dose, 1½ oz.

Sedative Ball. In slight colic. Assafœtida 4 drs., opium 4 drs., syrup and liquorice powder to form 4 balls.—HINDS.

Cordial and Anodyne Ball. Castile soap 3 drs., camphor 2 drs., ginger 1¼ drs., and Venice turpentine 6 drs. into 1 ball.

Stimulating Diuretic Balls. Cantharides ½ dr., aloes 2 drs., strained turpentine 1 oz., honey q. s.; make 4 balls, and roll in elecampane powder.—M. GOHIER, in dropsy.

Stimulating Expectorant Ball. Assafœtida, 3 drs., galbanum 1 dr., carbonate of ammonia ½ dr., ginger 1½ drs. honey q. s.—WHITE.

Sedative Aperient Ball. In epidemic catarrh or distemper. Balls for Inflammation of the Lungs (No. 4) 6 drs., physic ball (No. 10) 2 drs.; one at night and another in the morning.—CLATER.

Zinc and Valerian Ball. Oxide of zinc 1 oz., valerian 2 oz., oil of hartshorn 1 oz., soft extract of juniper berries q. s. to make 4 balls; one, twice a day.—ECKEL.

CHEWING BALLS, or MASTICATORIES.* The ingredients are to be tied in a piece of rag, and fixed by a string so that they may be kept in the mouth and chewed.

1. *Emollient Masticatory.* Marshmallow root, liquorice, gum arabic, of each (in powder) 1 oz., honey 1 oz., or q. s.—LEBAS.

2. *To promote Appetite.* Assafœtida, liver of antimony, juniper berries, bay-wood, pellitory, made into a mass with verjuice, tied as above.—SOLLEYSSELL.

3. Assafœtida, common salt, mastic, galangal, each 1 oz.—LEBAS.

4. Assafœtida 2 oz., salt 1 oz.—BOURGELAT.

5. Angelica ½ oz., assafœtida 1 oz., vinegar 2 drs.—SOLLEYSSEL.

* Rarely, if ever, employed in modern veterinary practice.

6. Flour of mustard $\frac{1}{2}$ oz., sal ammoniac 2 drs., powdered pellitory 1 oz.—MOIROUD.

ELECTUARIES & CONFECTIONS

Electuaries are compound medicines in the state of a soft paste. When the paste is hard enough to be formed into balls, the compound resembles ball masses or balls, under which we have placed them. French Pharmaciens often use the term *opiates* as nearly synonymous with electuaries; but we only apply the name *opiates* to compounds containing opium.

OPIATE CONFECTION. (*Veterinary.*) Opium $1\frac{1}{2}$ oz., mace-rate in a little hot water till soft, and rub it to a paste; then add ginger 3 oz., caraway 6 oz., treacle $1\frac{1}{2}$ lbs.; dose, $1\frac{1}{2}$ to 2 oz.—WHITE.

DEMULCENT AND PECTORAL ELECTUARIES.

1. Marshmallow root and liquorice (in powder) of each 2 oz., honey 10 oz.; mix; to be given at twice, with a spatula.—MOIROUD.

2. Melt $\frac{1}{2}$ oz. spermaceti with 2 oz. of olive oil, add 6 oz. of honey, and mix with $1\frac{1}{2}$ oz. p. marshmallow root; to be given daily.—MOIROUD.

3. (*With Opium.*) Powdered gum 2 oz., marshmallow 1 oz., extract of opium 2 drs., honey 3 oz.; for 2 doses.

4. *Cough Electuary with Manna.* Manna 2 oz., honey 6 oz.; in the morning; said to have cured acute bronchitis.

5. Powdered liquorice 8 oz., elecampane 4 oz., sulphur 2 oz., honey of squill 32 oz.; mix; for 8 doses.

STIMULANT AND CORDIAL ELECTUARIES. M.

LEBAS gives a form for an electuary (*Thériaque*) of many ingredients, the first of which (cordial powder) itself contains 26 different substances. We only insert here the simple formulæ of the French veterinarians.

1. Powdered angelica root 2 oz., masterwort 1 oz., muriate of ammonia $\frac{1}{2}$ oz., honey 8 oz.—MOIROUD.

2. *Stimulant and Expectorant.* Assafœtida 4 oz., elecampane 8 oz., honey 32 oz.; for 6 doses.—MOIROUD.

3. Powdered cassia and ginger, each 1 oz., honey 6 oz. Dose 1 oz. to $1\frac{1}{2}$ oz.—MOIROUD.

TONIC AND ASTRINGENT ELECTUARIES.

1. Red oxide of iron 8 oz., gentian 12 oz., extract of juniper berries 32 oz. MOIROUD prescribes 6 oz. for a horse, or 1 oz. for a sheep; but these are larger doses than are customary in England.

2. Peruvian bark 6 oz., nitre 1 oz., camphor $\frac{1}{2}$ oz., honey 16 oz. Dose 1 oz. to $1\frac{1}{2}$ oz.—LEBAS.

3. Powdered bistort 1 oz., calcined magnesia 4 drs., honey 4 oz. Dose 1 oz. to $1\frac{1}{2}$ oz.—MOIROUD.

PURGATIVE AND LAXATIVE ELECTUARIES. Aloetic compounds are usually made stiff enough to form into balls. See PHYSIC BALLS.

1. Oil of croton 20 drops, powdered senna 4 drs., honey q. s.—MOIROUD.

2. Sulphate of magnesia 4 oz., honey 16 oz., bran a quart; infuse the bran in sufficient hot water, and add the salt and honey; twice a day till the bowels are relaxed.—BOURGELAT.

3. Sulphate of soda or magnesia 5 oz., manna 4 oz., bran 1 quart; as the last.—MOIROUD.

DIURETIC ELECTUARIES.

1. Acetate of potash 2 oz., oxymel of squills 4 oz.; oatmeal or flour, to give a soft consistence.—MOIROUD.

2. Nitre 1 oz., camphor 2 drs. (rubbed with yolks of 2 eggs), oxymel 4 oz.; flour or liquorice powder, to give a suitable consistence.—MOIROUD.

DIAPHORETIC ELECTUARIES. (Dose 1 oz. to $1\frac{1}{2}$ oz.)

1. Sulphur 1 oz., powdered angelica $1\frac{1}{2}$ oz., honey 5 oz., —MOIROUD.

2. Prepared antimony $1\frac{1}{2}$ oz., elecampane 2 oz., treacle 4 oz.

3. Kermes mineral 1 oz., powdered sassafras and elecampane, each 2 drs., honey 6 oz.

VETERINARY POWDERS.

MR B. CLARK'S PULVIS UTILIS, *as a vehicle for other powders.* Turmeric $\frac{1}{2}$ lb., oatmeal or sifted barley-meal 4 lbs.; mix.

AROMATIC POWDER, OR HORSE SPICE.

1. WHITE'S *Aromatic Powder.* Caraway 6 oz., pimento 4 oz., ginger 2 oz., liquorice 2 oz.; mix; dose, 6 to 8 drs.

2. *Common Horse Spice.* Caraway, aniseed, coriander seeds, of each 16 oz., turmeric 32 oz., cumin seeds, liquorice, and ginger, of each 8 oz.; mix.

3. This is inserted, not as a desirable form, but as a specimen of what is used in the trade. Cayenne 2 oz., bean flower 45 lbs., mustard hulls 45 lbs., cumin seed 15 lbs., caraway 15 lbs., turmeric 9 lbs., bay-berries 3 lbs., ivory-black 1 lb.—GRAY'S SUPPLEMENT. The cordial powder of LEBAS contains 26 ingredients.

ABSORBENT POWDERS.

1. Carbonate of soda 2 to 4 drs., ginger 1 dr., calumba 2 to 4 drs.—WHITE.

2. Prepared chalk 4 drs., gentian 2 to 4 drs., aromatic powder (above) 1 or 2 drs.

ALTERATIVE, DIURETIC, AND DIAPHORETIC POWDERS. For swelled legs, grease, foul humours, hidebound, mange, surfeit, old coughs, and to render the skin fine. They are usually given with moistened corn. Too free use of these powders may prove injurious.

1. Sulphur 2 parts, black antimony 1, nitre 1; mix; dose, $\frac{1}{2}$ oz. to 1 oz.—V. C.

2. Sulphur 4 drs., levigated antimony 2 drs., nitre 3 drs.; mix; in *hide-bound* and unthrifty coat, every night.—YOUATT.

3. Ethiops mineral $\frac{1}{2}$ oz., cream of tartar 1 oz.; mix; give every night in a mash; for grease.—BLAINE.

4. Sulphur 12 oz., black antimony 12 oz.; mix, and divide into 24 powders; for mange, &c.—TAPLIN.

5. Nitre 16 oz., resin 16 oz., prepared antimony 4 oz., flowers of sulphur 24 oz.; mix; dose, 1 oz. every evening, with moistened corn, for 6 or 8 times.

6. Equal weights of antimony, nitre, and cream of tartar; dose, 6 to 9 drs.—BLAINE.

7. Nitre 6 oz., vermilion $\frac{1}{2}$ oz., resin 6 oz., tartarized antimony 2 drs.; for 12 doses.

8. Sulphur $\frac{1}{2}$ oz., prepared antimony 1 dr.; once a day, in the food, for 10 or 14 days.—CLATER.

9. TAPLIN'S *Alterative Powders.* Levigated antimony 8 oz., sulphur 8 oz., Ethiops mineral 4 oz., cream of tartar 4 oz.; in 12 doses.

10. Cream of tartar 2 drs., nitre 2 drs., sulphur 4 drs.; for 1 dose.—BLAINE.

11. Nitre 1 oz., resin 1 oz., rust of iron 1 oz., emetic tartar 15 grs.; dose, 1 oz.—LEBAS.

12. *In Farcy.* Prepared antimony 12 oz., sulphur 12 oz., cream of tartar 8 oz., cinnabar 6 oz.; mix, and divide into 20 doses; one every night, in corn.—TAPLIN.

CONDITION POWDERS. A want of condition is generally indicated by, and connected with, the unthrifty state of the coat, which the above alterative powders are supposed to remedy. Sometimes warm and bitter tonics are added to those ingredients which promote the action of the skin and kidneys, to increase the appetite and promote nutrition; but the most scientific practitioners condemn these additions; and particularly when the animal is changing its coat.

1. Black antimony 4 oz., flowers of sulphur 2 oz., bean flour or barley-meal $\frac{1}{2}$ lb.; a tablespoonful with corn.—B. CLARK.

2. Sulphur 2 lbs., fenugreek 4 lbs., cream of tartar 1 lb., liquorice 1 lb., nitre 1 lb., black antimony $\frac{1}{2}$ lb., gentian $\frac{1}{4}$ lb., aniseed $\frac{1}{4}$ lb., common salt 1 lb.; dose, 1 oz., daily for 2 or 3 weeks.

3. Gentian 4 oz., liquorice 4 oz., fenugreek 16 oz., diapente 6 oz., nitre 4 oz., salt 4 oz.; to promote appetite.

4. Aromatic powder 2 oz., assafœtida $\frac{1}{4}$ oz., cream of tartar $\frac{3}{4}$ oz., crocus metallorum $\frac{1}{2}$ oz.; for 2 doses.—LEBAS.

DIAPENTE. This should be made with equal parts of myrrh, gentian, ivory-dust, bay-berries, and birthwort. A worthless compound is frequently sold for it. The following is one of the least objectionable substitutes:—Equal parts of gentian, turmeric, bay-berries, and mustard. Another form in use is—Bay-berries 2 $\frac{1}{2}$ lbs., guaiacum wood 2 lbs., gentian 14 lbs., bole 2 lbs., bark which has been used for the tincture 2 lbs. Never employed in modern veterinary practice.

Fever Powders. 1. Nitre 1 oz., camphor 2 drs., tartarized antimony 1 dr.—WHITE.

2. Nitre 6 drs., camphor 2 drs., calx of antimony 1 $\frac{1}{2}$ drs.—HINDS.

3. Nitre 1 oz., unwashed calx of antimony 2 drs., antimonial powder 3 drs., camphor 1 dr.—WHITE.

Pectoral Powder. Powder of gum tragacanth 6 oz., nitre 1 oz.; give a tablespoonful in the mashes or food. In coughs.

Purgative Powder. Epsom salts 8 oz., aloes 10 oz., aniseed 3 oz.; dose, 2 oz.—LEBAS.

Powder for the Gripes. Aloes, senna, ginger, cream of tartar, of each 1 lb.; mix. This was formerly honoured with the title of *Pulvis Sanctus*.

Worm Powders. 1. Sulphur 12 parts, quicksilver 4 parts; triturate together till the mercury is extinct; then add male fern, rhubarb, tansy, gentian, of each 4 parts, wormwood, savin, aloes, castor seeds, of each 1 part; dose, $1\frac{1}{2}$ to 2 oz.—LEBAS.

2. Fern root 4 parts, tansy 2 parts, assafœtida and aloes, each 1 part; dose, as the last.—MOIROUD.

3. Sulphur 1 oz., emetic tartar 4 drs., common salt 8 oz., liver of antimony 1 oz.; mix; for 6 doses; one daily in wetted corn.—HINDS.

4. For Tapeworm. $\frac{1}{2}$ lb. Kosso in a drench.

MR WHITE'S *Compound Arsenical Powder.* White arsenic 1 dr., cream of tartar 9 drs.; mix carefully; give 10 grs. 3 times a day.

HAYNE'S *Bitter Powder*, for loss of appetite. Sulphate of potash 2 oz., gentian 1 oz., flour q. s. To be given twice a day.

MEDICATED PROVENDER. Bruised oats 4 lbs., bruised juniper berries 2 oz., common salt 1 oz.; mix. Nourishing and stimulant.—DELAFOND.

Liquid Medicines for Horses

DRINKS, DRENCHES, MIXTURES, MASHES,

ETC.

Drinks, properly speaking, are liquids which the horse will take willingly; *Drenches* are those liquid medicines which must be administered by a horn, bottle or funnel. This distinction is not always observed.

MILD DRINKS. *Demulcent, pectoral, cooling, and diuretic.*

Barley Water. Barley 1 lb., water 2 gallons; boil to 6 quarts, strain, and add 1 lb. of honey. If common barley is used, it should be first boiled with a little water, and this thrown away. If pearl barley is used, this will be less necessary. In inflammatory and catarrhal complaints.

Oatmeal Gruel. 1. Mix gradually 4 oz. of fresh oatmeal with as much cold water as will form a smooth mixture. Put 2 quarts of water in a saucepan over a clear fire, and before it gets very hot, add the mixture of oatmeal and water; stir the whole till it boils, and let it simmer a little while. Take care not to smoke it.

2. Mix half a pint of oatmeal with the same measure of water; triturate them in a marble mortar with a wooden pestle, for some time; then add 1 gallon of boiling water, and boil for a few minutes.—B. CLARK.

Blanche Water. Wet 3 or 4 handfuls of bran with scalding water, and work it with the hands till it becomes clammy; then add as much more water as may be desired. A mixture of oatmeal and cold water is also called white water, and in France, potato or other starch is used for the same purpose.

Linseed Tea. 1. Infuse 4 oz. of linseed in 3 pints of boiling water for several hours near the fire, stirring occasionally; then strain off, and add 4 oz. of honey; for 2 doses; in coughs, &c.

2. Pour 1 gallon of boiling water on $\frac{1}{2}$ lb. of linseed,

let the infusion stand till nearly cold, then pour off the clear liquid.—YOUATT.

Compound Decoction of Linseed. Linseed 4 oz., liquorice root 4 oz., mallows 2 handfuls; boil in six quarts of water for half an hour. Let the horse drink it freely.—BLAINE.

Cooling and Refreshing Drink. Barley water, linseed tea, or blanche water, 8 quarts, simple oxymel 16 oz.—MOIROUD.

Cooling and Diuretic Drink. Dissolve 1 oz. of nitre in a pail of water.

Camphorated Diuretic Drink. Water 10 quarts, nitre 1 oz., camphor (rubbed with yolks of 2 or 3 eggs), $\frac{1}{2}$ oz.; mix, and let the animal drink when thirsty.—MOIROUD.

MASHES

Bran Mash. Bran or pollard $\frac{1}{2}$ peck; put it in a bucket, and pour on it enough scalding water to wet it thoroughly; let it be well stirred with a stick, or worked with the hands, and let it stand, covered up, till new-milk warm. Emollient and slightly laxative. When intended to be nutritive, oats should be scalded with the bran.—B. CLARK.

Malt Mash. Upon a peck of ground malt pour a gallon and a half of water at 140° F. Stir frequently, and give when new-milk warm. Nutritive, in diseases attended with great debility.—MARKHAM.

Linseed Mash. HIND'S *Cooling Decoction.* Linseed 2 quarts, coarse sugar 2 oz., boiling water 6 quarts; simmer for three or four hours.

DRENCHES

DRENCHES FOR DIARRHŒA, DYSENTERY, AND DIABETES.

For Diarrhœa. 1. *Restraining Draught.* Opium 1 dr., prepared chalk 1 oz., compound powder of tragacanth 1 oz., mint water 1 pint.—WHITE.

2. Laudanum 1 oz., ether 1 oz., tannin 1 scruple, given in gruel or ale. This is to be given when the purging has gone on for some days, or when laxatives have been given without the desired effect.—FINLAY DUN.

3. Prepared chalk 8 oz., gum arabic, $\frac{1}{2}$ oz., catechu 2 drs., thin starch $\frac{1}{2}$ pint.—BLAINE.

4. Prepared chalk 1 oz., catechu 2 drs., opium 1 dr., ginger 1 dr.; rub together with the contents of 1 egg and add $\frac{1}{2}$ pint of thin gruel.—CLATER.

5. *For purging from corrosive sublimate.* Powdered opium 2 drs.; rub down with the yolk and white of one egg, and the contents of two more eggs, and gradually stir in $\frac{1}{2}$ pint of thin gruel.—CLATER.

Draught for Enteritis. Opium $1\frac{1}{2}$ drs., tartar emetic 1 dr., spirit of nitrous ether 1 oz.; mix, and add $1\frac{1}{2}$ pints of linseed oil.—SPOONER.

For dysentery or Mottled Grease. 1. Castor oil 8 oz., ipecacuanha 1 dr., opium 20 grs., liquid arrowroot 8 oz. Repeat once or twice at intervals of 6 hours; then substitute boiled starch for the castor oil.—BLAINE.

2. Opium 2 drs., nux vomica $\frac{1}{2}$ dr., ipecacuanha 1 dr., red wine 1 quart; mix. To be given morning and evening.

For Diabetes. 1. Opium 1 dr., ginger 2 drs., p. oak bark, 1 oz., decoction of oak bark 1 pint.—WHITE.

2. Sulphuret of potash 2 drs., p. uva ursi 4 drs., oak bark 1 oz., catechu 2 drs., opium $\frac{1}{2}$ dr. In strong chamomile tea.—BLAINE.

3. Calomel 3 drs., cascarilla 2 drs., salt of steel $2\frac{1}{2}$ drs., salt of tartar $1\frac{1}{2}$ drs., tincture of opium $\frac{1}{2}$ oz., strong beer q. s.—WHITE.

CARMINATIVE AND ANTISPASMODIC DRENCHES for Spasmodic and Flatulent Colic, or Gripes.

N.B. As most of these drenches would be injurious in enteritis, care should be taken to distinguish between this disease and colic. Inflammation is known by the quick but small pulse, redness of the inside of the eyelids, coldness of the ears and legs, and scanty and high-coloured urine. In colic, the attacks and remissions of pain alternate; in inflammation, the pain and distress continue. In colic, the pain is relieved by friction and motion; in inflammation, it is increased. Cold is sudden in its attack; inflammation more gradual in its approach.

1. Brandy, rum, or gin from 4 to 6 oz., hot water 12 oz. Mr CLARK directs a wineglassful of spirits to half a pint

of warm water. A pint of ale is sometimes substituted.
—WHITE.

2. Half a large bottle of Daffy's elixir, with hot water.

3. Tincture of pimento 4 oz., warm water half a pint.—
B. CLARK.

4. WHITE's *Anodyne Carminative Tincture* (see further on) 2 to 4 oz., hot water half a pint.—WHITE.

5. *Antispasmodic Draught*. Spirit of nitric ether 2 oz., tincture of opium 1 oz., solution of aloes (see further on) 4 oz.—V. C.

6. Spirit of nitric ether $\frac{1}{2}$ oz., tincture of opium $\frac{1}{2}$ oz., oil of turpentine 3 oz., gruel 1 pint.—BLAINE.

7. Rectified oil of turpentine 3 oz., tincture of opium 1 oz., warm ale 1 pint. If it does not relieve, repeat half the quantity with 1 oz. aloes dissolved in warm water.—
YOUATT.

8. Ether 1 oz., laudanum 2 oz., oil of peppermint 1 dr., ale and gin, each a $\frac{1}{2}$ of a pint.—BLAINE.

9. Camphor 2 drs., tincture of opium 1 oz., oil of peppermint 30 drops, warm water 1 pint. In a violent attack, add 1 oz. of spirit of turpentine.—PEALL.

10. The juice of 3 or 4 onions, with $\frac{1}{2}$ a pint of sound ale.

11. Pepper $\frac{1}{2}$ oz., oil of turpentine 3 oz., laudanum 1 oz., ale $\frac{1}{2}$ of a pint.—BLAINE.

12. Pepper a teaspoonful, juice of 2 or 3 large onions, gin $\frac{1}{2}$ of a pint.—BLAINE.

13. Laudanum 1 oz., sweet spirit of nitre 4 oz., oil of juniper 1 oz., tincture of benzoin 2 oz., spirit of sal volatile $1\frac{1}{2}$ oz., oil of peppermint 1 dr.; mix; give a fourth part in warm water or gruel, and repeat in 2 or 3 hours, if necessary.—HINDS.

14. *For Flatulent Colic*. Tincture of opium 1 oz., Tinct. of myrrh 1 oz., sulphuric ether 6 drs., tepid water a pint. Repeat in an hour if relief is not obtained.—SPOONER.

15. Heat $\frac{1}{2}$ lb. of common salt, and quench it in a quart of good ale. Give it new-milk warm.—DOWNING.

16. *In flatulent colic, when there is an evident distension of the abdomen with gas*: chloride of lime $\frac{1}{2}$ oz. (or solution of chlorinated soda 1 oz.), water 1 quart; repeat in half an hour if necessary.

17. Ginger, caraway, nutmeg, pimento, of each $\frac{1}{2}$ oz.,

bruise, and boil them in $\frac{3}{4}$ of a pint of ale for a few minutes, and add a gill of any spirit.—TAPLIN.

18. *In colic, accompanied with tympanitis*: Sol. of ammonia, Sp. nitrous ether, compound tincture of gentian, of each, equal parts. Dose, 1 to 2 fluid oz.

19. *Cordial Antispasmodic Drink*, for Spasm of the Diaphragm. Ginger 1 dr., caraway 2 drs., laudanum 1 oz., sweet spirit of nitre 1 oz., warm ale $\frac{1}{2}$ a pint.—CLATER.

20. *Antispasmodic Drench for Suppression of Urine*. Nitre 1 oz., camphor 2 drs., linseed tea 1 pint.—WHITE.

21. *Diuretic Camphor Drink*. Camphor 2 to 3 drs., olive oil 1 oz., carbonate of soda 1 dr.; rub together, and add tincture of opium 1 dr., warm water 2 pints.

22. *Antispasmodic Drenches for Locked Jaw*. 1. Opium $1\frac{1}{2}$ drs., camphor 2 drs., ginger 3 drs., brandy-and-water 8 oz.—WHITE.

2. Ether $\frac{1}{2}$ oz., tincture of opium 2 oz., camphor 1 dr., peppermint water $\frac{1}{2}$ pint.

CORDIAL AND STIMULANT DRENCHES. These are used in the same cases as the cordial balls, but are preferred where a more quick and powerful operation is required. Some of them are used in indigestion and slight attacks of colic.

1. Cloves and black pepper (bruised) $\frac{1}{2}$ oz., boiling water a quart; infuse and give warm.—MOIROUD.

2. Any of the cordial balls may be dissolved in warm ale or water, or peppermint water, and given as a drench.

3. A bottle of wine, 1 oz. of extract of juniper berries, and $\frac{1}{2}$ oz. of cinnamon in powder.—M. LEBAS.

4. Peppermint 2 oz., chamomiles $\frac{1}{2}$ oz.; infuse in $2\frac{1}{2}$ pints of water, and give it before it is cold; in slight colic and indigestion.

PECTORAL AND EMOLLIENT (or DEMULCENT) DRENCHES, for Coughs, Epidemics, Catarrh, &c. (For LINSEED TEA, COMPOUND INFUSION OF LINSEED, BARLEY-WATER, &c., see DRENCHES, further back.)

1. *Simple Emulsion*. Olive oil 2 oz., honey 3 oz., soft water 1 pint, subcarbonate of potash 2 drs.; mix.—WHITE.

2. Linseed tea 1 pint, honey 2 oz., syrup of poppies 2 oz., linseed oil 4 oz.

3. *B. CLARK'S Cough Drench.* Linseed oil 2 oz., solution of potash 40 drops, treacle 1 oz., soft water 10 oz.; mix.

4. Powdered gum 2 oz., warm water a quart; dissolve, and add honey 4 oz.—MOIROUD.

5. Marshmallow root 2 oz., water 2½ or 3 pints; boil to a quart, and add 4 oz. of treacle.—MOIROUD.

6. Liquorice and marshmallow roots, of each 2 oz., water a quart; boil, strain, and add honey 4 oz.—LEBAS.

7. Marshmallow root 2 oz., 4 poppy heads, water a quart; boil for ten minutes, strain, and add to the liquor before quite cold, 4 oz. of olive oil, 6 oz. of honey, and the yolks of 4 eggs, previously well beaten together.—MOIROUD.

8. Compound decoction of linseed (see LIQUID MEDICINES FOR HORSES, further back) 1 quart, oxymel 3 oz.

9. Spermaceti ½ oz., olive oil 3 oz.; melt together, and add honey 4 oz., water (by a little at a time) to make up a quart; repeat it twice a day.—LEBAS.

10. *Camphorated Emulsion.* Reduce to powder with a few drops of spirit, 1 or 2 drs. of camphor, add 12 drops of oil of aniseed, and 1½ oz. of simple emulsion.—WHITE.

11. Oxymel of squills 2 oz., opium ½ dr. to 1 dr., linseed oil 2 oz.; mix the opium with 8 oz. of water, and add the other ingredients; for one dose.—WHITE.

12. *For chronic coughs.* Tar water ½ pint, lime-water ½ pint, powdered squill 1 dr.; every morning.—BLAINE.

13. *In inflammation of the lungs, or catarrhal fever.* Tartarized antimony 2 drs., digitalis 1½ drs., nitre 3 drs., simple oxymel 4 oz., compound decoction of linseed 8 oz.—BLAINE.

14. The same, omitting the digitalis, and substituting 6 oz. of warm water for the dec. linseed. In influenza, when soreness of throat prevents swallowing balls.—BLAINE.

15. *In inflammation of the lungs:* Ipecacuanha 2 drs., laudanum 4 drs., powdered camphor 2 drs., Mindererus spirit 4 oz., linseed tea ½ pint.—BLAINE.

16. *In pleurisy.* Boil pearl barley, split figs, and raisins, each 6 oz., and liquorice root 2 oz., in 4 quarts of water down to 3; strain, and add honey 1 lb., vinegar 1

pint; give 1 oz. of nitre in a pint of this decoction every 6 hours.—TAPLIN.

17. *In epidemic (epizootic) catarrh*: Spirit of nitrous ether 1 oz., Mindererus spirit 6 oz., with linseed tea.—BLAINE.

18. GIBSON'S *Drink for Catarrhal Epidemic*. Colts-foot, hyssop, chamomile, of each a handful, linseed and garlic, each 1 oz., liquorice root sliced 1 oz., saffron $\frac{1}{2}$ oz.; infuse in 2 quarts of boiling water; give half in the morning and the rest in the afternoon.

19. *In influenza (after bleeding)*. Oil of croton 5 drops, nitre 4 to 6 drs., tartarised antimony 1 dr., spirit of nitric ether $\frac{1}{2}$ oz. to 1 oz., solution of acetate of ammonia (B. P.) 2 to 4 oz., warm water q. s. Once or twice daily. Sometimes $\frac{1}{2}$ oz. of cream of tartar is added.—SPOONER.

20. *For malignant epidemic*. Oxymel $\frac{1}{2}$ oz., spirit of Mindererus $\frac{1}{2}$ oz., beer yeast 4 oz., sweet spirit of nitre 1 oz.

DIURETIC DRENCHES for Dropsical Complaints, &c.

The use of stimulating diuretics in retention of urine from inflammation of the neck of the bladder, is dangerous.

1. MARKHAM'S *Dropsy Drench*. Decoction of wormwood in ale 2 quarts, soap 1 oz., grains of paradise 2 drs., long pepper 2 drs., treacle 3 oz.; for one dose, fasting.

2. *For dropsy of the belly*. Castile soap 2 oz., strong beer 1 pint; dissolve, and add cascarilla 2 drs., ginger 3 drs., oil of juniper 2 drs. (or balsam of copaivi 1 oz.); mix, for one dose.—WHITE.

3. White soap 1 oz., spirit of turpentine 1 oz., honey 4 oz., decoction of linseed 2 quarts; for two doses.—MOIROUD.

4. Strained turpentine 2 oz., yolks of 6 eggs; triturate together till incorporated, and add gradually 2 quarts of linseed tea; for two doses.—MOIROUD.

5. White wine and water 4 quarts, nitre 3 oz., honey 4 oz.; for 3 doses.—LEBAS.

6. Acetate of potash 2 or 3 oz., honey 6 drs., decoction of hemp or linseed 2 quarts; for 1 dose.—MOIROUD.

7. Acetate of potash 2 oz., camphor (rubbed with yolks of 2 eggs) 2 drs., decoction of linseed 2 quarts; for 2 doses, at an interval of some hours; in irritation of the urinary

passages, especially arising from cantharides or resinous irritants.—MOIROUD.

8. *Squill Drench.* Decoction of pellitory of the wall 1 quart, oxymel of squills 4 oz.—MOIROUD.

9. *Colchicum Drench.* Colchicum wine 2 oz., simple oxymel 4 oz., barley water 1 quart.

10. *In dysuria not arising from mechanical obstruction, or inflammation of the neck of the bladder.* Sweet spirit of nitre 4 oz., white wine 1 quart, water 2 quarts; for 3 doses.—LEBAS.

11. *Saline Diuretic Drink.* Glauber's salt 2 oz., nitre 6 drs., warm water 1 pint, sweet spirit of nitre 1 dr.—CLARK.

DRENCHES FOR FARCY AND GLANDERS.

1. Expressed juice of cleavers 6 oz., strong decoction of hempseed 6 oz., essence of spruce 6 oz.; mix; give every evening; and a mercurial or arsenical ball in the morning.—BLAINE. See Balls for FARCY AND GLANDERS.

2. *For Glanders.* Sulphate of copper 3 to 6 drs., gum arabic 2 or 3 oz., dissolved in 2 or 3 pints of water.—SEWELL.

FEVER DRENCHES.

1. Nitre 2 drs., tartar emetic $\frac{1}{2}$ dr., warm water or thin gruel 12 oz.; once or twice a day.—B. CLARK.

2. Sweet spirit of nitre 1 oz., spirit of Mindererus 6 oz., water 4 oz.

3. *Draught for Influenza.* Spirit of nitrous ether 2 oz., tartarised antimony 1 dr., nitre 4 drs., warm water $\frac{1}{2}$ pint.—SPOONER.

LAXATIVE AND PURGATIVE DRENCHES.

1. Castor oil 6 oz., linseed oil 8 oz., gruel q. s.—BLAINE.

2. Glauber's or Epsom salts 6 or 8 oz., whey or gruel 1 quart, castor oil 6 or 8 oz.—WHITE.

3. Barbadoes aloes 2 drs., tartarized antimony 1 dr., warm water 4 oz.; mix, and add castor oil 4 oz.—WHITE.

4. *Laxative Febrifuge in Influenza.* Linseed oil 12 oz., nitre 3 drs., camphor powdered 1 dr., sweet spirit of nitre 1 oz., warm water $\frac{1}{2}$ pint.—CLATER.

5. *Laxative Anodyne Drink.* In inflammation of the bowels. Linseed oil 1 pint, opium 2 scruples, sweet spirit of nitre 6 drs., warm water 4 oz.—CLATER.

6. Aloes 2 or 3 drs., salt of tartar 1 dr., water or mint water $\frac{1}{2}$ pint; mix, and add castor oil 4 to 6 oz.—WHITE.

7. *A Cooling Purging Drink.* Infuse 2 oz. senna with 3 drs. salt of tartar in a quart of boiling water for 2 hours; strain, and add 4 oz. Glauber's salts, and 2 or 3 drs. of cream of tartar.—BARTLET.

8. *Draught for Hepatitis.* Carbonate of potash and aloes, each 2 drs.; dissolve in hot water, shake with 12 oz. of linseed oil, and calomel 1 dr. Repeat twice a day without the aloes until relief is obtained. When in pain, add 1 oz. of tincture of opium.—SPOONER.

STRONGER PURGATIVE DRENCHES.

1. *In Inflammation of the Brain.* Barbadoes aloes 2 oz., gum arabic 1 oz.; powder and mix them, and pour on them a pint of boiling water. Take 10 grs. of farine of croton, and add to it gradually 4 oz. of the above solution. Repeat this dose every 6 hours till it operates.—CLATER.

2. Aloes 1 oz., soap 2 drs., salt of tartar 1 dr., water 1 pint; in apoplexy or staggers.—WHITE.

3. Infuse 1 oz. of senna in a quart of boiling water, strain, and add 1 oz. of aloes in powder.—BOURGELAT.

4. Aloes 1 oz., sulphate of magnesia 2 oz., aniseed powder $\frac{1}{2}$ oz., water a quart.—LEBAS.

5. Aloes 1 oz., syrup of buckthorn 4 oz., warm water a quart.—LEBAS.

DRENCHES FOR STOMACH STAGGERS, or Staggers from Indigestion.

1. *After a ball of aloes and calomel, and clyster of salt water.* Spirit of sal volatile $\frac{1}{2}$ oz., cascarilla powder 2 drs., warm water $\frac{1}{2}$ pint, twice a day; and the same without the cascarilla every hour.—WHITE.

2. Aloes 3 drs., pimento 2 drs., ginger 1 dr.; infuse in a quart of hot water, and, when cold, add 2 oz. spirit of turpentine, and 1 oz. of spirit of hartshorn. Repeat in an hour if required.—BLAINE.

3. *Laxative Tonic Drinks.* Linseed oil 1 pint, pow-

dered gentian 2 drs.; every 6 hours till the bowels are properly opened.—CLATER.

4. Common salt $\frac{4}{8}$ oz., ginger 2 drs., magnesia 1 oz., warm water 1 quart.—WHITE.

5. Valerian 1 oz., serpentary $\frac{1}{2}$ oz., saffron 2 drs., infuse in a pint of boiling water, and when nearly cold, strain off, and add 1 oz. tincture of assafœtida and 2 drs. of laudanum.—TAPLIN.

6. *After a Purgative.* Volatile tincture of valerian 1 oz., powdered valerian $1\frac{1}{2}$ oz., peppermint water 8 oz.; mix, for a dose.—WHITE.

TONIC DRENCHES. Tonics are more generally administered in the form of balls.

1. *Mild Tonic in later stages of Epidemic Catarrh or Distemper.* Gentian 1 dr., powdered ginger $\frac{1}{2}$ dr., cascarrilla 1 dr., warm water $\frac{1}{2}$ pint, sweet spirit of nitre $\frac{1}{2}$ oz. to 1 oz.; to be repeated night and morning unless the pulse is quickened.—CLATER.

2. Gentian root 2 oz., smaller centaury 1 oz., wormwood $\frac{1}{2}$ oz.; boil in 3 pints of water to a quart.—VATEL.

3. *CLARK'S Bitter Drench.* Quassia chips 2 oz., water 3 pints; boil to 2 pints; for 3 doses.

4. Quassia 1 oz., ginger 2 drs., water 2 pints; boil for 10 minutes; for 2 doses.

5. *Metallic Tonic.* Sulphate of zinc $\frac{1}{2}$ dr., ginger or pimento 1 dr., treacle 1 oz.; mix, and add gradually 12 oz. of warm water.—B. CLARK.

6. *Egyptian Tonic Drink.* In farcy and nasal gleet. Ægyptiacum $\frac{1}{2}$ oz., pimento or ginger $\frac{1}{2}$ dr., warm water 12 oz.—CLARK.

7. *Cantharides Tonic Drench* (for the same). Sulphate of zinc 15 grs., cantharides 7 grs., pimento 15 grs., treacle 1 oz., warm water to form a drench.

DEOBSTRUENT DRENCHES.

1. Guaiacum wood 2 oz., sassafras 1 oz., linseed $\frac{1}{2}$ oz., water q. s. to yield a quart of decoction; boil, strain, and add of corrosive sublimate 10 grs., sal ammoniac 2 drs.—LEBAS.

2. Iodide of potassium 40 grs., iodine 10 grs., water a quart.—MOIROUD.

3. Muriate of lime $\frac{1}{4}$ oz., water a quart.—MOIROUD.

WORM DRENCHES.

1. Common salt 2 oz., infusion of wormwood a quart. Repeat it for some days.—MOIROUD.

2. A quart of linseed oil.—CLATER.

3. Oil of turpentine 4 oz., linseed or castor oil 8 oz., gruel a pint; preceded by a mild dose of aloes, and bran mash.

4. Fern root 2 oz., valerian 1 oz., DIPPÉL'S animal oil (empyreumatic oil of hartshorn) 1 oz., yolks of 2 eggs, honey 2 oz.; boil the roots in 2 quarts of water to half, incorporate the oil with the egg, and then the honey, and mix the whole with the decoction.—VATEL.

5. Animal oil 1 oz., yolks of 2 eggs, honey 1 oz., water or some bitter infusion a quart. CHABERT recommends infusion of savory as a vehicle for the oil.—LEBAS.

6. Soot (wood soot?) in fine powder 2 oz., spirit of wine 2 oz.; mix, and add a quart of infusion of rue, or of tansy. Some practitioners prefer milk as a vehicle for worm medicines.—MOIROUD. For other worm remedies, see WORM BALLS.

. Worms may be prevented by a properly regulated dietary, and access to rock salt.—FINLAY DUN.

DRENCHES for the MALIGNANT EPIDEMIC, or DISTEMPER.—Pestilential or Putrid Fever.

1. Gentian 1 dr., calumba 2 drs., ginger 1 dr., laudanum $\frac{1}{2}$ oz., spirit of nitrous ether $\frac{1}{2}$ oz., peppermint water 3 oz.—CLATER.

2. Gentian 2 oz., willow bark 6 oz., water 3 pints; boil to a quart, and add solution of acetate of ammonia (B. P.) 6 oz.—MOIROUD.

3. Dissolve $\frac{1}{2}$ dr. of chloride of lime in 8 oz. water, and add spirit of nitric ether $\frac{1}{2}$ oz., laudanum $\frac{1}{2}$ oz., tincture of calumba 1 oz.; twice a day.—CLATER.

4. Bruised bark 1 oz., acetate of ammonia (B. P.) 4 oz., camphor 1 dr.; boil the bark in 2 quarts of water in a covered vessel for a quarter of an hour; strain, and when cool add the camphor (rubbed with yolk of egg or honey) and the acetate of ammonia.—LEBAS.

5. Spirit of nitric ether 1 oz., Mindererus spirit 4 oz., infusion of chamomile 6 oz., beer yeast 6 oz., tincture of opium 3 drs.—BLAINE. See also ANTISEPTIC DRENCHES (page 76).

ANTISEPTIC DRENCHES, *to check mortification.*

1. Peruvian bark 1 oz., ginger 2 drs., opium 1 dr., fresh beer q. s.—WHITE.

2. Opium 1 dr., carbonate of ammonia 1 dr., aromatic powder 2 drs., camphor $1\frac{1}{2}$ drs., good ale or porter, a pint.

3. Chloride of lime or soda 2 to 3 drs., serpentry in powder 1 oz., fresh beer or sweet wort 1 quart.—WHITE.

DRENCH FOR POISONING BY YEW. Stronger purgative drink (No. 1) 4 oz., vinegar 4 oz., thick gruel 4 oz.; repeat it every 6 hours, without the croton, till purging is produced.—CLATER.

DRENCH FOR PREVENTING RABIES.

1. Box leaves 8 oz., rue 8 oz.; cut them very fine and boil in 3 pints of milk, in a close vessel, for an hour, and strain; boil the ingredients another hour in 3 pints of water, and strain; mix the decoction; give a third part every morning fasting.—BLAINE. (Exploded.)

2. Excise and cauterise the bitten part at once. Chloral hydrate, belladonna, and ammonia, the latter subcutaneously.—FINLAY DUN.

3. Curara, the potent arrow-poison of the South American Indians, has been proposed as a remedy for rabies.

DRENCHES TO PROMOTE PARTURITION.

1. Ergot of rye in fine powder 2 or 3 drs., pennyroyal water, or infusion of rue, 1 quart.

2. Saffron 6 drs., chamomile 2 oz., boiling water a quart; make an infusion, to be given warm. 1 oz. of dried savin, with 1 oz. of cassia, may be substituted for the saffron.—MOIROUD.

ANODYNE DRENCHES.

1. Opium 1 dr. dissolved in warm water, $\frac{1}{2}$ pint; add 1 quart of starch gruel.

2. Oil of peppermint 50 drops, added to a pint of warm water, with 2 oz. of gum arabic; add tincture of opium $\frac{1}{2}$ oz.

3. Mix tincture of opium $\frac{1}{2}$ oz., with sweet spirits of nitre $1\frac{1}{2}$ oz., essence of peppermint 1 dr., and water 1 pint.

IODINE DRENCH. Iodide of potassium 2 scruples, iodine 12 grs.; triturate together, and add gradually a quart of water.—MOIROUD,

MISCELLANEOUS LIQUID MEDICINES

TINCTURES, SOLUTIONS, &c.

Solution of Aloes. Aloes 1 part, water 7 parts, proof spirit 1 part; dissolve the aloes in water by means of a water-bath, and when removed, add the spirit.—MORTON.

Anodyne Carminative Tincture. Opium 1 oz., cloves 1 oz., ginger 1 oz., old brandy (rum or gin) 1 quart; digest in a corked bottle, shaking daily. Dose, a fourth or sixth part.—WHITE.

Ethereal Tincture of Opium. Turkey opium 1 lb., spirit of nitric ether 8 lbs.; macerate for a month. Dose, $\frac{1}{2}$ oz. to 1 oz., in spasmodic colic.—DICKENS.

Gripe Tincture. Tincture of Pimento. Ground pimento 1 lb., rectified spirit, and soft water, of each 3 pints; digest for some days and strain: give 4 fl. oz. at once, and repeat every hour till relieved.—B. CLARK.

Tincture of Foxglove. Digest 3 oz. of dried foxglove in a quart of any spirit.—YOUATT.

Infusion of Foxglove. Infuse 1 oz. of powdered foxglove in a quart of boiling water till cold.—YOUATT.

Tincture of Myrrh. Myrrh 2 oz., sand 2 oz., rectified spirit and soft water, of each $\frac{1}{2}$ pint.—B. CLARK.

Tincture of Aloes and Myrrh. Aloes 12 oz., myrrh 6 oz., rectified spirit 1 gallon, water $\frac{1}{2}$ gallon; digest 14 days, frequently shaking, and filter. For outward use, rectified wood naphtha may be substituted for the spirit.—V. C.

Tincture for Colic. Opium 1 dr., horseradish 2 oz., capsicum 1 oz., spirit of nitric ether 1 lb.; macerate 14 days; dose, 1 oz., with 2 oz. of spirit of nitric ether, every 2 hours as long as necessary.—GREGORY.

Tincture of Croton. Bruised croton seeds 1 oz.; rectified spirit 16 oz. Digest for 7 days, and filter. Dose, $\frac{1}{2}$ oz. to 1 oz. in water.

Tincture of Iodine. Iodine 1 part, rectified spirit 8 parts; dose, 1 to 2 drs.—V. C. The following is preferable:

Compound Tincture of Iodine. Iodine 1 oz., iodide of potassium 2 oz., spirit of wine 12 oz.

Solution of Chloride of Lime. 1. Chloride of lime 1 dr., water 8 oz.; mix in a mortar, and filter.—CLATER.

2. Chloride of lime 1 part, water 10 parts.—CHEVALIER.

3. Chloride of lime 1 part, water 48 parts.—LARBARAQUE. See LOTIONS, under MEDICINES FOR HORSES.

Solution of Nitre. Nitre 1 part, water 7 parts.—V. C.

Solution of Ammonio-Sulphate of Copper. Dissolve 1 oz. of sulphate of copper in 16 oz. of water, and add ammonia until it begins to precipitate. 4 ounces every 8 hours as a tonic.—JECKYLL.

Solution of Henbane. Extract of henbane 4 drs., spirit of nitric ether 4 oz. Antispasmodic; dose, 2 oz., with or without solution of aloes.—WRIGHT.

For SOLUTIONS and TINCTURES for outward use, see EXTERNAL APPLICATIONS (*Veterinary Formulary*), further on.

CLYSTERS

Laxative. 1. Aloes 1 oz., water 2 or 3 quarts.—YOUATT.

2. Water gruel 1 gallon, olive oil 1 pint.—WHITE.

3. Epsom salts 6 oz. (or common salt 6 oz., or soap 2 oz.), thin gruel or broth 5 quarts.—BLAINE.

4. Soft soap 2 oz., warm water $\frac{3}{4}$ of a pailful.—B. CLARK.

5. Infuse 3 oz. senna in 2 quarts of water, and add Epsom salts 4 oz., honey 6 oz.—MOIROUD.

6. Chamomiles, fennel seed, coriander seed, of each 1 oz., caraways $\frac{1}{2}$ oz.; boil in 2 quarts of water to 3 pints; strain, add 2 oz. Epsom salts, and when nearly cool, $\frac{1}{4}$ pint of olive oil and $\frac{1}{4}$ pint of tincture of senna.—TAPLIN.

Purgative. 1. Aloes 8 to 12 drs., salt 8 oz., water 1 gallon: in staggers.—WHITE.

2. Senna 2 oz., tobacco 2 oz.; boil for a quarter of an hour in 2 quarts of water, strain, and add common salt 4 oz., emetic tartar 1 dr.; for 2 doses. Very irritating.—LEBAS.

Emollient. Dried mallow leaves, or marshmallow root, 1 $\frac{1}{2}$ oz., linseed $\frac{1}{2}$ oz., water 2 quarts; boil and strain; to be used warm.—MOIROUD.

Emollient and Anodyne. 1. Mix 6 drs. of starch in powder with a little cold water, and add it to a decoction of 6

poppy-heads in 2 quarts of water; boil for an instant and strain: in intestinal irritation.—MOIROUD.

2. Gruel 2 pints, liquid starch or arrowroot 1 pint, powdered opium 1 to $1\frac{1}{2}$ drs.—WHITE.

3. Boil 6 poppy-heads in 4 quarts of water till reduced to 2 quarts; add prepared chalk 2 oz., boiled starch 2 quarts; once or twice a day *in diarrhæa*.—BLAINE.

4. A double handful of coarse bran, 6 poppy-heads, 2 quarts of water; boil and strain.—MOIROUD.

5. Tripe liquor (or suet boiled in milk) 3 pints, thin starch a quart, laudanum $\frac{1}{2}$ oz.; *in diarrhæa*.

Cooling. Butter-milk or whey, barley-water, of each a quart.—MOIROUD.

Carminative and Stimulant. 1. Chamomiles 3 oz., aniseed or fennel seed $1\frac{1}{2}$ oz., 4 poppy-heads; boil the poppies in sufficient water, and infuse the flowers and seeds in the hot decoction.—VATEL.

2. *To expel wind.* Boil 1 lb. of figs in 3 quarts of water for half an hour, then add 2 handfuls of chopped rue; boil a few minutes, strain, and add 8 oz. of olive oil.—SOLLEYSSELL.

For Gripes. Mash 2 onions, pour over them 2 oz. of oil of turpentine, and 4 quarts of thin gruel.—BLAINE.

Astringent. 1. Alum whey 1 quart, thin starch a quart.

2. Suet milk 3 pints, starch gruel 2 pints, laudanum $\frac{1}{2}$ oz.

Vermifuge. 1. *For Thread-worms.* Powdered aloes $\frac{1}{2}$ oz., powdered gum arabic $\frac{1}{2}$ oz.; mix with half a pint of boiling water; then mix the white of an egg with a quart of linseed oil, and gradually add the solution of aloes.—CLATER.

2. Infuse 4 oz., of tansy in 2 quarts of water; strain and add 2 oz. of animal oil (empyreumatic oil of harts-horn); also the worm drenches (Nos. 4 and 6) may be used in this method.—MOIROUD.

Uterine Stimulants. 1. Infuse a handful of rue in 2 quarts of water, and add 2 oz. of common salt.

2. Savine 2 oz., sal ammoniac 4 drs.: as the last.

Diuretic. 1. Nitre 1 oz., decoction of linseed 3 pints.

2. *Camphorated.* Incorporate 4 drs. of camphor with the yolks of 2 eggs, and add it to the last.—MOIROUD.

For Irritable Bladder. 1. Belladonna leaves 3 oz., water 3 pints; boil and administer warm.—MOIROUD.

2. Extract of belladonna 2 drs., boiling water $1\frac{1}{2}$ pints.
- Nourishing.* 1. Thick gruel 3 quarts, ale 1 quart.—BLAINE.
2. Milk 2 quarts, yolks of 4 eggs; mix, and give warm.—BOURGELAT.
3. Strong broth 2 quarts, thickened milk 2 quarts.—BLAINE.
4. Tripe liquor or broth 3 quarts, flour 4 oz.; mix the flour in the hot broth; repeat frequently.—MOIROUD.

External Applications

LINIMENTS AND EMBROCATIONS

BLISTERING LINIMENTS, OR LIQUID BLISTERS, AND SWEATING OILS.

1. Powdered Spanish flies 1 oz., spirit of wine 6 oz., water of ammonia 2 oz.; let it stand for a week, shaking frequently, and strain. (See No. 11.)—WHITE.

2. Flies 1 oz., euphorbium $\frac{1}{2}$ oz., oil of turpentine 4 oz.; digest for 2 or 3 days, and pour off the liquid; digest the flies, &c., in 4 oz. of spirits of wine and 2 oz. of water of ammonia for 3 or 4 days, shaking frequently; strain off this liquid, and mix it with the former. This is more active than the last.—WHITE.

3. BLAINE'S *Liquid Blister*. Spanish flies, coarsely powdered, 8 oz., oil of turpentine 2 quarts; steep for 3 weeks, strain, and add a quart of olive oil.

4. BLAINE'S *Milder or Sweating Liquid*. Mix 4 oz. of the last with 6 oz. of oil.

5. CLATER'S *Strong Liquid Blister*. Spirits of turpentine coloured with alkanet, 1 gallon, powdered flies 1 lb.; macerate for a month, shaking daily, then pour off the clear fluid for use.

6. *Common or Sweating Liquid*. Mix the last with equal part of spermaceti oil.

7. Powdered flies 3 oz., spirit of turpentine a pint; digest for a few days.—YOUATT.

8. *Blistering Liquid*. The vinegar of cantharides (ACETUM CANTHARIDES. EPISPASTICUM) of the British Pharmacopœia.

9. *Blistering Liniment for immediate use.* Spanish flies in fine powder 1 oz., oil of turpentine 6 oz. To be rubbed on the belly in inflammation of the bowels.—**WHITE.** Cantharides 3 drs. in hartshorn 4 oz; in a fortnight, strain, and add olive oil 4 oz.—**SPOONER.**

10. *Croton Liniment.* A tincture of croton nuts with oil of turpentine is used as a blister, but is not so efficacious as cantharides.—**YOUATT.**

11. *WHITE'S Mustard Blister.* Best flour of mustard 8 oz., water enough to form a paste, oil of turpentine 2 oz.; water of ammonia 1 oz.

12. *Blistering Tincture.* Flies 1 oz., proof spirit 8 oz.; macerate 2 or 3 weeks; mix and filter. To be rubbed in, and repeated next day if necessary.—**WHITE.**

13. Saturated tincture of cantharides 1 oz., bichloride of mercury 6 grs.—**KENT.**

14. Powdered cantharides 1 dr., olive oil 2 oz. To be applied every 48 hours for a week in old spavin.—**TAPLIN.**

15. *Oil of Cantharides (by infusion).* Digest 1 oz. of powdered cantharides in 8 oz. of olive oil in a water-bath, for 2 hours, and strain.

16. *Cantharides Solution (for setons).* Digest 1 oz. of p. flies with 8 of oil of turpentine, with a gentle heat, for 14 days; strain, and add to the clear liquid an equal weight of Canada balsam. Soak the cotton cord in the solution, draw it between the finger and thumb, and dry it.—**MORTON.**

STIMULATING LINIMENTS.

1. Soft soap 4 oz., camphor 1 oz., proof spirit 2 pints, water of ammonia $\frac{1}{2}$ pint.—**V. C.**

2. Sweet oil 2 oz., spirit of hartshorn 1 oz., oil of turpentine $\frac{1}{2}$ oz.—**WHITE.**

3. *In chronic sprains.* Common oil 6 oz., liquid blister 2 or 3 oz.—**BLAINE.**

4. *Soap Liniment.* Soft soap 6 oz., water 8 oz.; dissolve, and add 1 pint of rectified spirit, in which is dissolved 2 oz. camphor, 1 oz. oil of rosemary, and 2 to 4 oz. strong water of ammonia.—**WHITE.**

5. *For splints.* Oil of origanum 1 oz., spirit of turpentine 1 oz., spirit of wine $\frac{1}{2}$ oz. To be applied night and

morning for a few days, discontinuing it as often as any moisture appears.—LANCET.

6. *For the same purpose.* Oil of origanum $\frac{1}{2}$ oz., oil of turpentine $\frac{1}{2}$ oz., camphorated spirit of wine 2 oz.—TAPLIN.

7. *For sprains, old swellings, rheumatism, &c.* Spirit of hartshorn 2 oz., camphorated spirit 2 oz., oil of turpentine 1 oz., laudanum $\frac{1}{2}$ oz., oil of origanum 1 dr.

8. Camphorated oil 4 oz., oil of turpentine 1 oz., oil of origanum 1 dr.

9. *For callous swellings after bruises.* Soap liniment 4 oz., camphor 2 drs., water of ammonia 1 oz.

10. *For indolent tumours.* Mercurial ointment 2 oz., olive oil 2 drs., camphor 2 drs.

11. Olive oil 4 oz., water of ammonia 2 oz., oil of turpentine 2 oz.

12. *For strains.* Barbadoes tar 2 oz., spirit of turpentine 2 oz., opodeldoc 4 oz.—TAPLIN.

13. Oil of turpentine 2 parts, muriatic acid 1 part.—POTT.

14. Camphorated oil 4 parts, oil of turpentine and tincture of cantharides, of each 2 parts, acetic acid 1 oz.—LEBAS.

15. *Turpentine Liniment.* Equal parts of oil of turpentine and oil. Digestive and rubefacient.—V. C.

16. *Compound Turpentine Liniment.* Soft soap 4 oz., camphor 1 oz., oil of turpentine 16 oz.; mix.—V. C.

LINIMENT FOR BOG SPAVIN. Mercurial ointment 2 oz., oil of cantharides 4 drs.—MORTON.

LINIMENT FOR SORE BACKS. Extract of lead $\frac{1}{2}$ oz., vinegar 1 oz., olive oil 2 oz.—WHITE.

LINIMENTS FOR ITCHING HUMOURS, MANGE, LICE, &c.

1. Equal parts of oil of tar, oil of turpentine, and seal oil. Apply every second day for 2 or 3 times, then wash.—V. C.

2. Sulphur 4 oz., turpentine 4 oz., oil of tar and train oil 6 or 8 oz. The parts to be first washed with soft soap, and dried.

3. *For lice.* Sublimate 1 dr., muriatic acid 3 drs., tobacco water 2 pints, oil of turpentine 4 oz.—WHITE.

4. *Liniment for Mange.* Goulard's extract of lead 2 oz., olive or rape oil 2 oz., sulphur 1 oz.

LINIMENTS FOR CANKER OF THE FOOT AND BAD THRUSHES. See also CAUSTICS and LOTIONS.

1. Barbadoes tar 1 oz., oil of turpentine 1½ oz.; mix carefully, and add oil of vitriol 1 dr.—WHITE.

2. Butter of antimony alone.

3. Crystallized verdigris in fine powder 1 oz., honey 2 oz., bole and alum, of each ½ oz., vinegar to form a liniment; to be mixed over a gentle fire. Greasy applications are to be avoided.—WHITE.

4. Carbolic acid.

See also DETERGENT LINIMENTS, below.

DETERGENT LINIMENTS.

1. Oil of turpentine 1 oz., oil of vitriol 2 drs. by measure; mix in a large gallipot, and when cool, add 2 oz. of linseed oil.—WHITE.

2. *Ægyptiacum.* Bruised sulphate of copper 12 oz., vinegar 4 lbs., treacle 3 lbs.: place over a clear fire, and let it boil up.—B. CLARK.

3. *Wash for Grease.* Sulphate of copper 2 drs., and alum 2 drs. in water 1 pint.

MISCELLANEOUS LINIMENTS AND MIXED OILS.

See also EMBROCATIONS.

Creasote Liniment. Creasote 2 oz., oil of turpentine 4 oz., olive oil 4 oz.; mix; in fistulous sores, unhealthy wounds, &c.—V. C.

Oil of Cantharides. Powdered flies 1 oz., olive oil 8 oz.; digest in a water-bath for 2 or 3 hours, and filter.—V. C.

Goulard Liniment. Extract of lead 1 oz., olive oil 4 oz.—MORTON. For excoriated surfaces.

Saturnine Balsam. Acetate of lead 1 oz., oil of turpentine 2 oz.; digest with a gentle heat.—MOIRAUD.

Drying Liniment. Linseed oil and spirit of wine, of each equal parts.—SOLLEYSSELL.

Marshallow Liniment. Olive oil and marshallow ointment, of each 4 oz.; melt the ointment and add the oil.—BOURGELAT.

Emollient and Anodyne Liniment. Neatsfoot oil 4 oz., poplar ointment, marshmallow ointment, of each 2 oz.—**MOIROUD.**

Lime water Liniment. Lime water 8 oz., olive or linseed oil 2 oz.

Narcotic Liniment. Olive oil 4 oz., laudanum 2 oz.—**MOIROUD.**

Liniment for confirmed Grease. Verdigris, sugar of lead, of each $\frac{1}{4}$ oz., honey 1 oz; mix.—**CLATER.**

Compound Iodine Liniment. Iodine 1 oz., soap liniment 8 oz.—**V. C.**

Turpentine Liniment. Equal parts of turpentine and olive oil.—**V. C.**

Resolvent Liniment. Olive oil 2 oz., strong mercurial ointment 2 drs., water of ammonia 2 drs.

Black Oils. Olive (or rape) oil 1 pint, oil of turpentine 2 oz., mix, and add gradually 6 drs. of sulphuric acid; leave the bottle open till cold.—**PERCIVALL.**

Oils for Mange. Oil of turpentine 1 pint; add to it, very gradually and cautiously, 2 oz. of oil of vitriol, stirring the mixture constantly, then add a quart of linseed oil; from 4 to 8 oz. to be rubbed in with a brush every second day, for 3 or 4 times.—**CLATER.**

WARD'S White Oils. Spirit of wine, oil of turpentine, rape oil, beef brine, camphor, of each equal parts.

White Oils or Egg Oils. 1. Yolks of 2 eggs, 3 oz. solution of ammonia, 1 oz. oil of origanum, $\frac{1}{2}$ oz. oil of turpentine, a pint of vinegar; mix s. a.—**PHARMACEUTICAL JOURNAL.**

2. Distilled vinegar $1\frac{1}{2}$ pints, oil of turpentine $1\frac{1}{2}$ drs., spirit of wine $1\frac{1}{2}$ oz., GOULARD'S extract of lead $\frac{1}{2}$ oz., whites and yolks of 2 eggs; mix the turpentine and Goulard with the eggs, gradually add the vinegar, and lastly the spirit.—**REDWOOD'S GRAY'S SUPPLEMENT.**

Liniment of Ammonia. This is sometimes termed *White Oils.* Olive or rape oil $\frac{1}{2}$ oz., water of ammonia 1 oz. Sometimes 1 oz. of oil of turpentine is added to increase its activity.

Darby's Oils. Equal parts of oil of amber, Barbadoes tar, and balsam of sulphur.

MARSHALL'S Oils. Linseed oil 1 lb., olive or rape oil 1 lb., green oil $\frac{1}{2}$ lb., oil of turpentine $\frac{1}{2}$ lb., oil of vitriol $1\frac{1}{2}$ drs.

Newmarket Oils. Linseed oil, oil of turpentine, green oil, of each 3 lbs., oil of vitriol 1 oz.

Nine Oils (so-called). Train oil 23 lbs., oil of turpentine 6 lbs., oil of bricks 1 lb., oil of amber 1 lb., spirit of camphor 2 lbs., Barbadoes tar 7 lbs., oil of vitriol 2 oz.—GRAY'S SUPPLEMENT.

RADLEY'S Oils. Barbadoes tar 8 oz., linseed oil 4 oz., oil of turpentine 4 oz.

LORD STAMFORD'S Mixed Oils. Oil of origanum 6 oz., oil of turpentine 24 oz., spirit of wine 16 oz., green oil 6 lbs., camphor 3 oz.

EMBROCATIONS, VARIOUS

(See also LINIMENTS and LOTIONS)

**Embrocation for Strains.* 1. Soft soap 1 oz., spirit of wine 4 oz., oil of rosemary 2 drs., camphor 2 drs.

2. *For strains in the shoulder.* Oil of turpentine 1 oz., camphorated spirit 2 oz.—BLACKER.

3. Equal quantities of soft soap, oil of turpentine, spirit of wine, and elder ointment.—WHITE.

4. Soft soap 2 oz., oil of bays 1 oz., water of ammonia 1½ oz., oil of origanum ½ oz.

5. Barbadoes tar 2 oz., spirit of turpentine 2 oz., opodeldoc 4 oz. After fomenting with hot vinegar and Goulard.—TAPLIN.

Mustard Embrocations. 1. Mustard flour 4 oz., water of ammonia 1½ oz., oil of turpentine 1 oz., water enough to bring it to the consistence of cream.

2. Camphor 1 oz., oil of turpentine 1 oz., water of ammonia 2 oz., flour of mustard 8 oz., water to form a thin paste.—WHITE.

Embrocations for Poll Evil. Spirit of wine ½ pint, camphor 2 drs., Goulard's extract of lead 1 dr.; mix.—HINDS. See DISCUTIENT LOTIONS.

Embrocations for Saddle-galls or Warbles. 1. Goulard's extract of lead 2 drs., distilled vinegar 3 oz., spirit of wine 4 oz.—WHITE.

2. Soap liniment and Mindererus spirit, equal parts.

* These embrocations should only be applied after the subsidence of the inflammation.

3. Sal ammoniac $\frac{1}{2}$ oz., muriatic acid 2 drs., water 8 to 12 oz.

4. White vinegar 3 oz., spirit of wine 3 oz., sugar of lead 2 drs., water 6 oz.; mix.—HINDS.

CLARK'S *Embrocatio Frigorifera*. Vinegar 4 oz., camphor (dissolved in spirit) $\frac{1}{2}$ oz., water to fill up a wine bottle.

CLARK'S *Embrocatio Excitans*. Olive oil 3 oz., camphor $\frac{1}{2}$ dr., spirit of turpentine $\frac{1}{2}$ oz., water of ammonia 3 drs.

TAPLIN'S *Embrocation for Windgalls*. Oil of origanum, spirit of turpentine, of each $\frac{1}{2}$ oz., camphorated spirit 1 oz. Applied with tow, and covered with a piece of lead bound on.

LOTIONS AND WASHES

COOLING LOTIONS, for external inflammation.

1. Sal ammoniac 1 oz., nitre 2 oz., water 16 oz. To be used as soon as made.—MORTON.

2. Goulard's extract of lead 1 oz., vinegar 2 oz., camphorated spirit 3 oz., water 16 oz.; for recent spavin.—TAPLIN.

3. V. C. *Goulard Water*. Goulard's extract 2 drs., spirit 2 drs., soft water 1 pint.

4. WHITE'S *Saturnine Lotion*. Sugar of lead 1 oz., vinegar and water of each 1 pint.

5. B. CLARK'S *Lotio Refrigerans*. Liquor of diacetate of lead 1 dr., spirit of nitric ether 1 dr., water 2 pints. In slight rubs and bruises.

DISCUTIENT LOTIONS for dispersing indolent tumours and saddle-galls and for chronic strains.

1. Mindererus spirit 4 oz., camphorated spirit 4 oz., water 16 oz.—PERCIVALL.

2. Sal ammoniac 1 oz., vinegar 8 oz., camphorated spirit 1 oz.—MORTON.

3. For saddle-galls and warbles. Goulard's extract 2 drs., distilled vinegar 3 oz., spirit of wine 4 oz.—WHITE.

4. For saddle-galls and windgalls. Muriate of ammonia $\frac{1}{2}$ oz., muriatic acid 2 drs., water 8 to 12 oz.—WHITE.

5. BLAINE'S *Saline Embrocation*. Sal ammoniac 8 oz., vinegar 3 pints.

6. Mindererus spirit 2 oz., soap liniment 2 oz.—WHITE.

7. *For warbles.* White vinegar 3 oz., spirit of wine 3 oz., sugar of lead 2 drs., water 6 oz.—HINDS.

8. Strong solution of salt 1 oz., tincture of myrrh $\frac{1}{4}$ oz., for saddle-galls.—YOUATT.

9. *In incipient poll-evil.* Common salt 4 oz., vinegar $\frac{1}{2}$ pint, cold water 1 quart, spirit of wine and laudanum, each 1 oz.—CLATER.

10. White vinegar 1 pint, extract of lead 2 oz., camphorated spirit $\frac{1}{4}$ oz., soft water 1 pint.—TAPLIN.

11. *For strains.* Bay salt $\frac{1}{2}$ lb., sal ammoniac 2 oz., sugar of lead $\frac{1}{4}$ oz., vinegar $1\frac{1}{2}$ pints, water 1 pint.

12. *For sitfasts.* Blisters and ointment of biniodide of mercury (see RED IODIDE OF MERCURY OINTMENT).

ASTRINGENT LOTIONS, for drying up sores and diminishing their discharge (especially in grease and scratched heels), after the inflammation has been subdued by linseed or carrot poultices.

1. *For grease and cracked heel.* Alum 4 oz., boiling water 1 pint.—TAPLIN.

2. Alum 2 drs., sulphate of zinc 1 scruple, water 1 pint.—YOUATT.

3. *Mild, for cracks.* Sugar of lead 2 drs., sulphate of zinc 1 dr., infusion of oak bark 1 pint.—BLAINE.

4. *For confirmed grease.* Nitric acid 1 oz., water 8 oz.—BLAINE.

5. *Strong.* Blue vitriol $\frac{1}{4}$ oz., alum 3 drs., water 1 pint.—SPOONER.

6. Sugar of lead 1 oz., blue vitriol 1 oz., water 1 quart.—WHITE.

7. Lime-water 16 oz., spirit of camphor $\frac{1}{4}$ oz., sugar of lead 1 dr.—BOURGELAT.

8. Sulphate of iron 2 oz., alum 2 oz., vinegar 8 oz., water $3\frac{1}{2}$ pints.—MOIROUD.

9. Tincture of myrrh 1 oz., camphorated spirit 1 oz., distilled vinegar and water, each 2 oz.—TAPLIN.

10. *For anburies.* Alum 2 oz., water 1 pint, sulphuric acid 1 dr.

11. *To wash farcy buds after they have been opened.* Sulphate of iron 1 oz., water 1 quart; dissolve, and add $\frac{1}{4}$ oz. (by weight) of oil of vitriol.—CLATER.

12. One fluid dr. of carbolic acid in from 6 to 12 oz. of water.

DETERGENT LOTIONS, for foul ulcers.

1. Sulphate of copper 1 oz., nitric acid $\frac{1}{2}$ oz., water 6 oz.—WHITE.

2. Sulphate of copper 1 oz., sulphuric acid 12 drops, water 4 oz.

3. *For stimulating old ulcers.* Sulphate of copper 2 drs., water 1 pint.—YOUATT.

4. *To remove fungous granulations.* Sulphate of copper 1 oz., water 1 oz.

5. Nitrous acid 1 oz., quicksilver $\frac{1}{2}$ oz.; dissolve, and add water 8 oz.

LOTIONS FOR MANGE.

1. White hellebore 2 oz., tobacco 2 oz., water 3 pints; boil, strain, and add, when cold, a pint of fresh lime-water.—BLAINE.

2. Boil 4 oz. of white hellebore in 3 pints of water to 2 pints, and add corrosive sublimate 2 drs., previously dissolved in 3 drs. of muriatic acid.—WHITE.

3. Boil 2 oz. of tobacco in a quart of water, strain, and add common salt 3 oz., soap 2 oz.—LEBAS.

4. Liver of sulphur $\frac{1}{2}$ oz., water 1 quart.—MOIROUD.

5. Liver of sulphur 4 oz., soft soap 16 oz., water 2 gallons.—LEBAS.

6. Acid nitrate of mercury 2 drs., distilled water 16 oz.—MOIROUD.

7. *Mercurial Wash.* Sublimate 2 drs., spirit of wine 2 oz., water 2 pints.—CLARK.

8. Chloride of lime 1 lb., water a gallon. Mix.—LUCAS.

9. One fluid dr. of carbolic acid in from 6 to 12 oz. of water.

VARIOUS LOTIONS

Conglutinum. Sulphate of zinc 4 oz., water a pint.—BRACY CLARK.

Black Wash. *For sluggish ulcers.* Calomel 2 drs., lime-water 1 pint.

Yellow Wash. Sublimate 8 grs., lime-water 4 oz.

Nitric Acid Lotion. Nitric acid 2 or 3 drs., water 1 pint; for exciting sluggish ulcers.—MORTON.

Lotion of Nitrate of Silver. For the same. Nitrate of silver 10 grs., distilled water 1 oz.—YOUATT.

Lotion for Farcy. Dissolve 1 oz. of sulphate of iron in a quart of water, and add $\frac{1}{2}$ oz. of oil of vitriol.—CLATER.

Styptic Lotion, for Stopping Bleeding. Alum 2 oz., sulphate of zinc 2 drs., water 1 quart.

Catechu Lotion for Ulcers of the Mouth. Infuse 2 oz. of catechu in a quart of boiling water for an hour; strain, and add 1 oz. of spirit of wine. (For saddle-galls, add 4 oz. of tincture of catechu and 8 oz. of common salt.)—CLATER.

Lotion of Chloride of Lime. For mange, and as a stimulant to unhealthy wounds and fistulous sores. To chloride of lime 1 lb. add gradually 1 gallon of water; mix, and filter or decant. Diluted with 10 or 15 parts of water it is used as a lotion for grease, exfoliated bones, &c., and is a disinfectant for foul stables. For ulcers of the tongue, mix 1 dr. of chloride of lime with a pint of water; for mange, 4 drs. to a pint.

Wash for destroying Lice about the Legs. Corrosive sublimate 1 dr., muriatic acid 3 drs., tobacco water 1 quart, oil of turpentine 4 oz.—WHITE.

Alum Mouth Wash. Alum 2 drs., sage tea a quart.—ECKEL.

For Bruised Gums. Alum 2 drs., tincture of myrrh 1 oz., honey 1 oz., water 2 oz.—SPOONER.

Acid Collutorium. Infusion of sage a quart, muriatic acid 1 oz., flour 3 oz., honey 8 oz. To be applied to the mouth frequently.—ECKEL.

LOUSE WATER.

1. Tobacco 4 oz., boiling water a quart; infuse for 24 hours.—CLARK.

2. *Mercurial.* Sublimate 2 drs., spirit of wine 2 oz., water 1 quart.

THE FOLLOWING LIQUID CAUSTICS are for canker, thrush, foul, unhealthy wounds, and to remove proud flesh, &c. See also LINIMENTS (DETERGENT).

Milder. 1. Tincture of perchloride of iron.

2. Sulphate of copper 1 oz., water from 4 oz. (V. C.) to a pint.—CLATER.

3. Saturated solution of sulphate of zinc: in quitters.—**WHITE.**

4. Alum $\frac{1}{2}$ oz., borax $\frac{1}{2}$ oz., boiling water 4 oz., styptic tincture 1 oz.—**TAPLIN.**

5. Hydrochloric acid, alone or diluted.

6. Ægyptiacum 2 oz., nitrous acid 20 drops.

7. Any of the stronger caustics (except butter of antimony) diluted with water.

8. Goulard's extract 4 oz., sulphate of zinc 2 oz., sulphate of copper 2 oz.; white vinegar 32 oz.—**VILLATE.**

9. Aloes 5 oz., weak spirit 10 oz.; dissolve, and add 6 oz. of sulphuric acid.—**DUVILLE.**

Stronger. 1. Butter of antimony. This is the safest and most useful caustic in canker.

2. Dissolve 1 oz. of quicksilver, by heat, in 2 oz. of nitric acid, and evaporate till the liquid weighs $2\frac{1}{2}$ oz.

3. Verdigris 1 oz., nitrous (red nitric) acid 1 oz.; dissolve.—**WHITE.**

4. Red precipitate 1 oz., nitrous acid 2 oz.—**WHITE.**

5. Nitrous acid, alone or with a little water.

6. Sulphuric acid, alone or with a little water.

7. Sublimate 1 dr., muriatic acid 2 drs., water q. s.—**WHITE.**

8. Creasote.

9. *For canker.* Dissolve corrosive sublimate $\frac{1}{4}$ oz. in hydrochloric acid 1 oz., then add spirits of wine 4 oz., and water 4 oz.

10. Chloride of zinc with enough water to dissolve it; or **Sir WILLIAM BURNETT'S PATENT SOLUTION.**

Caustic for Poll-Evil. 1. Lunar caustic 1 dr., distilled water $\frac{1}{2}$ oz.—**BLAINE.**

2. Corrosive sublimate 2 drs., water 3 oz.

Scalding Mixture for Poll-Evil. 1. Sublimate 2 drs., verdigris 2 drs., blue vitriol 2 drs., sulphate of iron 4 drs., honey 2 oz., oil of turpentine 8 oz., spirit of wine 4 oz.; to be applied hot, and confined by stitches.—**GIBSON.**

3. Sublimate 1 dr., finely powdered and mixed with 4 oz. of basilicon, and melted to scalding heat.—**BLAINE.**

4. Caustic potash 1 dr., rubbed down with 4 oz. oil of turpentine.—**BLAINE.**

Caustic for Farcy Buds. Sublimate 1 dr., muriatic acid 3 drs., spirit of wine 1 oz., water $\frac{1}{2}$ oz.—**WHITE.**

Solid Caustics. Lunar Caustic, Caustic Potash, and Chloride of Zinc. See ARGENTI NITRAS, POTASSÆ HYDRAS, and ZINCI CHLORIDUM, *Pocket Formulary*.

Canquoin's Caustic is made by mixing chloride of zinc with twice its weight of flour and a little water into a stiff paste, which is to be rolled out to the required thickness, and cut to the size of the part to be destroyed, the skin being previously removed by a blister. Another caustic is made with 2 parts of chloride of zinc, 1 of butter of antimony, and 5 of flour.

Sulphuric Caustic is made by triturating hay saffron with oil of vitriol, so as to form a ductile mass. BOUCHARDAT recommends solidifying the acid by ivory or lamp black.

Solidified Nitric Acid is merely lint soaked with strong nitric acid, squeezed, and formed to the required shape.

Filho's Caustic is made by melting together in an iron ladle 2 parts of caustic potash and 1 of lime over a quick fire, and pouring it into leaden tubes of the desired size. The air must be excluded when not in use, by beeswax, or other means.

FOMENTATIONS.

These should be applied moderately warm (about 120° F.) by means of flannel dipped in the liquid, and frequently renewed from time to time, keeping the parts covered.

Emollient. 1. Coarse bran 2 double handfuls, water 6 quarts, boil and strain.

2. Mallow-leaves 8 oz., water 4 quarts; boil and strain; 6 poppy-heads may be added.

Anodyne. 1. Boil 24 poppy-heads and 2 handfuls of hemlock in 6 quarts of water for 2 hours, and strain.—WHITE.

2. Belladonna 2 handfuls, 6 poppy-heads, water 3 quarts; boil and strain.—MOIROUD.

3. Dried wormwood and chamomile, of each 4 oz., bay-leaves 2 oz., rue 3 oz.; boil in a gallon of water.

4. Take wormwood, chamomile, mallow (or either of them), cut them to pieces, and put 2 handfuls into a bucket, pour scalding water on them, and cover with a cloth.—B. CLARK.

Discutient and Astringent. Vinegar or verjuice 1 quart; make it hot, and add 2 oz. of Goulard's extract of lead; apply warm; in strains of the sinews of the legs.—TAPLIN.

COLLYRIA, OR EYE-WATERS.

1. Acetate of lead and sulphate of zinc, of each $\frac{1}{2}$ dr.; dissolve them separately in $\frac{1}{2}$ pint of boiled water; mix, and filter.

2. Sugar of lead 10 to 20 grs., water 8 oz.—MORTON.

3. Extract of lead 1 dr., spirit 2 drs., water 8 oz.—WHITE.

4. Sol. acetate of ammonia (B. P.) 3 oz., rose-water 6 oz.—BLAINE.

5. Sugar of lead 2 drs., vinegar $\frac{1}{2}$ oz., soft water 16 oz., rose-water $\frac{1}{2}$ oz.—BLAINE.

6. Infuse 1 oz. of foxglove in 2 pints of boiling water, and strain.—YOUATT.

7. Tincture of opium 2 drs., water 8 oz., extract of lead 1 dr.—WHITE.

8. Brandy 1 oz., vinegar 1 oz., tincture of opium 2 drs., rose-water 8 oz.—BLAINE.

9. Extract of henbane 1 dr., water 8 oz.—WHITE.

10. Decoction of poppies 8 oz., saffron $\frac{1}{2}$ dr., infuse the saffron in the hot decoction.—LEBAS.

11. Lapis divinus 3 drs., soft water $\frac{1}{2}$ pint.—CLATER.

12. Common salt $\frac{1}{2}$ dr., water 6 oz.—YOUATT.

13. *For Watery Bloodshot Eyes.* Burnt alum 1 oz., calcined white vitriol 1 oz., boiling water 3 pints.—BRACKEN.

14. *Emollient.* Infusion of marshmallow leaves or flowers 1 quart, starch (rubbed smooth with a little water) $\frac{1}{2}$ oz.; mix and boil. To be used warm.—MOIROUD.

15. *Astringent.* Alum 2 drs., whites of 2 eggs, water $\frac{1}{2}$ pint; mix in a mortar.—BOURGELAT.

16. Tincture of digitalis $\frac{1}{2}$ oz., soft water 8 oz.—CLATER.

17. *To remove Opacity of the Cornea.* Nitrate of silver 10 grs., distilled water 1 oz. 1 or 2 drops to be dropped into the eye.—YOUATT.

18. *For Cloudiness of the Eye.* Sublimate 4 grs., spirit of wine 20 drops; rub together, and add soft water $\frac{1}{2}$ oz. A few drops to be introduced into the eye 3 or 4 times a day.—CLATER.

19. Tincture of aloes 1 oz., rose-water 8 oz.—LEBAS.

20. *Stimulating.* Infusion of elder flowers 16 oz., brandy 2 oz.—MOIROUD.

21. *Lapis mirabilis* $\frac{1}{2}$ oz., water 4 to 8 oz. The *Lapis mirabilis* is thus made: White vitriol 2 lbs., rock alum 3 lbs., fine bole $\frac{1}{2}$ lb., litharge 2 oz., water 3 quarts; boil together to dryness.—SOLLEYSSELL. Another form is given under SUNDRY SOLUTIONS.

22. *Alum Collyrium*. Decoction of marshmallow 16 oz., alum 2 drs., camphorated spirit 1 dr.; mix. To be used towards the decline of inflammation.—STRAUSS.

23. *Tannin Collyrium*. Dissolve 1 dr. of tannin in 13 oz. of water, and add 3 oz. of cherry-laurel water.

24. *In Chronic Inflammation*. Sulphate of zinc 8 grs., water 4 oz.—CLATER.

25. *In Specific Ophthalmia*. Tincture of opium 2 drs., extract of belladonna 1 dr., with distilled water 1 pint.

SUNDRY SOLUTIONS, &c.

Styptic Stone. For saddle-galls, kicks, sprains, bruises, ulcers, and as a collyrium. Sulphate of iron 8 oz., sal ammoniac, sulphate of zinc, and oxide of copper, each 1 oz.; mix, and melt together with a gentle heat. About the size of a nut of this compound to be dissolved in a quart of warm water and applied with compresses renewed every 3 or 4 hours.—KNAUP.

Lapis Divinus. Sulphate of copper, alum, nitre, of each 3 oz.; melt together, and stir in 1 dr. of camphor: used in eye-waters and lotions.

Wound Stone. Alum, sulphate of zinc, of each 3 oz., verdigris and sal ammoniac, of each, 1 dr.: melt together and add $\frac{1}{2}$ dr. of powdered saffron: detergent and drying.

Clark's Conglutinum. Sulphate of zinc 4 oz., water a pint.
Solution of Alum. Alum 1 oz., water 16 oz. Dissolve.—V.C.

Solution of Sulphate of Zinc. Sulphate of zinc 1 oz., water 3 oz.—V. C. In quitters.

Solution of Sulphate of Copper. Sulphate of copper 1 oz., water 4 oz.—V. C.

Compound Solution of the same. Sulphate of copper 3 oz., alum 3 oz., water 2 lbs., sulphuric acid $1\frac{1}{2}$ oz.

Solution of Perchloride of Mercury. Sublimate, and hydrochloric acid, each 1 part, spirit or water 7 parts.—V. C.

Goulard Water. Extract of lead 1 oz., camphorated spirit

- 2 oz., rain water a quart.—TAPLIN. V. C. uses extract of lead and rectified spirit, each 2 drs., soft water 1 pint.
- Tincture of Catechu.* See *Pocket Formulary*. Used externally for wounds.
- Tincture of Euphorbium.*—Euphorbium 1 oz., rectified spirit 6 oz.
- Alkaline Tincture of Euphorbium.* Euphorbium 8 oz., solution of carbonate of potash 3 pints: used as a caustic and stimulant, particularly in curbs after the inflammation has been subdued.
- Compound Tincture of Cantharides.* Powdered flies 4, euphorbium 1, proof spirit 24 parts.—LEBAS.
- Styptic Tincture.* Tincture of myrrh, spirit of camphor, and Friar's balsam, equal parts.—TAPLIN.
- Ægyptiacum.* (Veterinary.) Sulphate of copper in powder 12 oz., vinegar 4 oz., treacle 48 oz.; boil together to a proper consistence.—B. CLARK.
- Ægyptiacum with Turpentine.* Honey 28 oz., pyroligneous acid 14 oz., powdered verdigris 10 oz.; boil together in a copper vessel till the mixture has a reddish-purple colour and is of the consistence of thin honey; add Venice turpentine 28 oz., and keep it on a slow fire, stirring constantly for a quarter of an hour.—LELOUP.
- Liniment of Verdigris.* V. C. Verdigris in fine powder 9 oz., alum 6 oz., treacle 1½ lbs. Boil until the compound assumes a brown colour.
- Liniment of Sulphate of Copper.* Powdered sulphate of copper 1 part, treacle 4 parts. Simmer in a pipkin over a slow fire until the whole assumes a reddish-brown colour. In canker, severe thrush, &c.—MORTON.

POULTICES OR CATAPLASMS

These are useful in reducing inflammation and relieving pain. They should not be used too hot, nor applied too tightly, especially to the feet.

COMMON POULTICES.

1. Bran moistened with hot water, and as much linseed meal added as will give it tenacity.—V. C.
2. Boil a quart of bran for 10 minutes with enough

water to make a thin mash, then add to it 4 oz. of linseed meal; apply it in a flannel bag.—BLAINE.

3. Fine bran 3 parts, linseed meal 1 part, hot water q. s.

CHARCOAL POULTICES.

1. Oatmeal $\frac{1}{2}$ pint, linseed meal $\frac{1}{2}$ pint, charcoal 4 oz., beer grounds q. s.

2. Carrots scraped, or carrots boiled, with charcoal powder q. s. Antiseptic.—BLAINE.

YEAST POULTICES.

1. *In old grease with an offensive smell.* Linseed meal, oatmeal, boiling water q. s.; mix, and ferment with a tablespoonful of yeast.—BLAINE.

2. *In gangrene.* Add 2 oz. of turpentine to the last.—BLAINE.

ANODYNE POULTICES.

1. Boil poppy-heads in water, strain, and add linseed meal to stiffen it.—YOUATT.

2. Sprinkle the surface of a simple poultice with laudanum.

CLEANSING POULTICES. Mashed turnips, not pressed, with enough linseed meal, or oatmeal, to give them consistence; or, the charcoal poultice above.

DRAWING POULTICES.

1. Boil 2 lbs. of chopped onions in water, and add to it the crumb of a 4-lb. loaf.—HINDS.

2. Sorrel boiled and squeezed 4 parts, onions baked in ashes 1 part, basilicon ointment 1 part; mix, and apply warm.—VATEL.

RESOLVENT POULTICES.

1. Rye meal 8 oz., prepared chalk 2 oz., vinegar 10 oz.; mix warm, and stir, till no more gas is disengaged; apply warm.—SOLLEYSSELL.

2. *To indolent glandular tumours.* Linseed meal 12 oz., powdered hemlock 4 oz., muriate of ammonia 4 oz., vinegar q. s.—LEBAS.

GOULARD POULTICES.

1. To a linseed-meal poultice add 1 or 2 drs. of Goulard's extract of lead.—YOUATT.

2. Bread and barley meal equal parts, Goulard water q. s., lard 4 or 6 oz.—TAPLIN.

CHLORINE POULTICE. Chloride of lime $\frac{1}{2}$ oz., water 1 pint, linseed meal q. s.: to grease when offensive.—YOUATT.

POULTICES FOR GREASE.

1. The herb cleavers (or goose-grass) beaten to a paste.
2. Mash bread and boiled turnips with stale beer, and stir in 1 oz. flour of mustard, turpentine 2 oz., linseed meal 2 oz., lard 6 oz.; apply night and morning.—TAPLIN.

MUSTARD POULTICE.

1. Mustard flour and linseed meal, equal parts; mix with sufficient hot vinegar to give a proper consistence.
2. Flour of black mustard 3 lbs., hot vinegar, or water, q. s.—MOIROUD.

RUBEFACIENT POULTICE:

1. Fresh horseradish root, grated, and immediately applied.—MOIROUD.
2. *Stronger.* Old yeast 2 lbs., flour of black mustard 1 lb., euphorbium powder 4 oz., vinegar q. s.; mix, and apply cold.

OINTMENTS, CERATES, CHARGES, &c.

SIMPLE EMOLLIENT OINTMENTS.

1. *Simple Cerate.* Olive oil 1 pint, beeswax 4 oz.
2. *Spermaceti Ointment.* Lard 12 oz., white wax 2 oz., spermaceti 1 oz.

BLISTERING OINTMENTS. The Spanish flies should be finely powdered, and added to the melted substances, at as low a temperature as practicable.

1. Lard 4 oz., common turpentine 1 oz., p. flies 1 oz.; melt the lard and turpentine, and stir in the powdered flies. V. C. Mr YOUATT substitutes resin for the turpentine.

2. Venice turpentine and resin, of each 1 lb., palm oil or lard 2 lbs.; melt together, and gradually stir in 1 lb. of powdered flies.—BLAINE.

3. Palm oil 4 lbs., resin 1 lb.; melt together, and stir in 1 lb. of powdered flies.—CLATER.

4. *Mild.* Lard 4 oz., Venice turpentine 1 oz., p. flies 6 drs.—WHITE.

5. *Stronger.* Mercurial ointment 2 oz., oil of bays 2 oz., Barbadoes tar 1 oz., oil of rosemary 2 drs., p. flies 1 oz.—WHITE.

6. *Strong.* Oil of turpentine 1 oz., oil of vitrol 2 fluid drs.; mix in a basin, and add melted lard 6 oz., oil of origanum 1 oz., powdered flies 1 to 2 oz.—WHITE.

7. *Strongest.* Strong mercurial ointment 4 oz., oil of origanum $\frac{1}{2}$ oz., finely powdered euphorbium 3 drs., p. flies $\frac{1}{2}$ oz.—WHITE.

8. *BLAINE'S Mercurial.* Common blister (No. 2, *ante*) $\frac{1}{2}$ oz.; sublimate in fine powder $\frac{1}{2}$ dr.; for splints, spavins, &c.

9. *For common purposes.* Lard 6 oz., Venice turpentine 4 oz., beeswax 2 oz., yellow resin $\frac{1}{2}$ oz., oil of origanum $\frac{1}{2}$ oz., powdered cantharides 3 oz. It may be softened in winter by rubbing it with a little turpentine.—WHITE.

10. Powdered flies 5 drs., lard 4 oz., oil of turpentine 1 oz.—HINDE.

11. *Mustard Blister.* Best flour of mustard 8 oz., water to form a paste.—YOUATT. Two oz. of oil of turpentine, and 1 oz. of pure water of ammonia are sometimes added.

12. Mylabris in fine powder (see *Veterinary Materia Medica*) 1 dr., prepared lard 4 oz. Digest together over a water-bath for 3 hours, occasionally stirring, while hot filter through paper, and allow to cool.

Note.—In applying vesicants, the hair should be clipped closely, or shaved off, the part fomented with warm water, and the blistering ointment well rubbed in. In inflammation of the lungs, &c., blistering is more successful after bleeding. In 24 hours, a little olive or neatsfoot oil should be applied, and repeated night and morning. The head should be tied up for the first two days, and the litter removed from the stable. If strangury is produced give plenty of linseed tea. The simplest blisters are perhaps the best for common purposes. Sublimate blemishes. Sweating down is effected by milder stimulants; for this purpose, the liquid blister (see LINIMENTS, under "MEDICINES FOR HORSES") is diluted by some mild oil, &c.

DETERGENT OINTMENTS, for cleansing foul and indolent ulcers.

1. Suet 4 oz., Venice turpentine 6 oz., red precipitate, finely powdered, 2 oz.—**WHITE**.

2. Citrine ointment, alone or with $\frac{1}{2}$ its weight of Venice turpentine.

3. Sulphate of zinc 1 dr., sulphate of copper 1 dr., oil of turpentine 2 drs.; grind smooth, and mix it with 4 oz. of melted tallow. See also **DIGESTIVE OINTMENTS**, No. 4.

4. Yellow basilicon 2 oz., black basilicon 1 oz.; melt together, remove from the fire, add 1 oz. of turpentine, and $\frac{1}{2}$ oz. finely powdered red precipitate.—**TAPLIN**.

5. *Verdigris Ointment*.—Verdigris in fine powder 1 part, common turpentine 1 part, lard 12 parts; mix.—**MORTON**.

DIGESTIVE OINTMENTS, to promote a discharge from unhealthy and indolent ulcers.

1. Resin 16 oz., linseed oil 12 oz.; melt together with a gentle heat.—**CLARK**.

2. Strained turpentine, honey, of each 2 oz., yolks of 4 eggs, myrrh $\frac{1}{2}$ oz., aloes 1 oz.; mix.—**SOLLEYSSELL**.

3. Equal parts of common turpentine and lard melted together.—**WHITE**.

4. To 1 lb. of the last add 1 oz. of finely powdered verdigris.—**WHITE**.

5. Yellow wax 3 oz., common turpentine 3 oz., black pitch 1 oz., resin 6 oz., linseed oil 16 oz.; melt together with a gentle heat, then add oil of turpentine 4 oz., and stir till cold.

6. Olive oil 1 pint, yellow wax and black resin, of each 4 oz., Burgundy pitch and turpentine, of each 2 oz.; melt the other ingredients, and add the turpentine when it is removed from the fire.—**TAPLIN**.

7. Common turpentine 1 part, lard 3 parts; melt together.—**V. C.**

8. (*Basilicon*.) Resin 5 oz., yellow wax 2 oz., lard 8 oz.; melt together.

9. (*Black Basilicon*.) Pitch, wax, resin, of each 11 oz., olive (or rape or linseed) oil, a pint.

EYE OINTMENTS. The powder should be very fine, and the whole rubbed smooth.

1. *In chronic ophthalmia.* Nitrate of silver 5 to 10 grs., lard 1 oz.; rub till perfectly smooth. The size of a pea to be introduced between the lids.—MORTON.

2. Calamine $\frac{1}{2}$ oz., tutty $\frac{1}{2}$ oz., sulphate of copper $\frac{1}{2}$ dr., sulphate of zinc $\frac{1}{2}$ oz., alum $\frac{1}{2}$ oz., camphor 2 drs., fresh butter 3 oz.; mix, and apply warm, with a feather, to watery, inflamed eyes.—BRACKEN.

3. Ointment of nitrated quicksilver 1 dr., zinc ointment 1 oz., camphor 1 dr.

4. *In inflammation of the eyelids.* Verdigris 1 part, Venice turpentine 1 part, lard 12 parts.—MORTON.

5. *For wounds in the eye.* Tutty ointment 1 oz., honey of roses 2 drs., calcined white vitriol 20 grs.; apply with a feather night and morning, and sponge daily with warm milk and water.—BRACKEN.

6. *For removing opacity of the cornea.* Iodine 2 grs., iodide of potassium 20 grs., lard or butter $\frac{1}{2}$ oz.

EUPHORBIIUM OINTMENT. Euphorbium 1 part, lard 8 parts; mix.—DELAFOSSÉ.

OINTMENTS FOR SCURVY AND CRACKED HEELS, AND CONFIRMED GREASE. The inflammation should be first subdued by poultices. The milder preparations (which are here placed first) should be employed in the first instance, and afterwards those for confirmed grease.

1. *For scurvy heels.* Goulard's extract $\frac{1}{2}$ dr., lard 1 oz.; mix. The heels should first be gently rubbed with soap and water.—CLATER.

2. *For scurvy or cracked heels.* Sugar of lead $\frac{1}{2}$ oz., oxide of zinc $\frac{1}{2}$ oz., lard or palm oil 4 oz.

3. Melt together 3 oz. of white diachylon, 4 oz. of olive oil; mix, and when nearly cold, add 3 drs. of sugar of lead in fine powder. First wash the heel, then apply the Astringent Lotion No. 9, and afterwards this ointment; or elder ointment 4 oz., camphor 6 drs., laudanum 2 drs., extract of lead 2 drs.; mix.—TAPLIN.

4. Melt yellow wax 2 oz. with sweet oil 8 oz., and add sugar of lead $\frac{1}{2}$ oz.

5. *Healing ointment for cracked heels.* Lard 4 lbs., resin 1 lb.; melt together, and stir in 1 lb. true calamine.—CLATER. See also SOFTENING and COOLING OINTMENTS further on.

6. *For cracked heels and grease.* Alum 1 oz., turpentine 1 oz., lard 3 oz.; melt the turpentine and lard, and stir in the powdered alum.—V. C.

7. *For grease.* Venice turpentine 4 oz., wax 1 oz., lard 4 oz.; melt together, and add sugar of lead 1 oz. (or alum 2 oz.) in fine powder.—WHITE.

8. Lard, honey, common turpentine, of each 8 oz.; melt together, and add powdered alum 6 oz., white vitriol 2 oz.

9. Common turpentine 1 lb.; melt, and add powdered alum 1½ lbs., bole 2 lbs.; stir till cold, spread on brown paper, and tie over with list.

10. Lard ½ lb., honey ½ lb., common turpentine ½ lb.; melt, and add p. alum 1 lb., white vitriol 2 oz.; stir till cold.

11. *For confirmed grease.* Common verdigris ½ oz., alum, sulphate of zinc, sugar of lead, of each ½ oz.; tar 6 oz.—BLAINE.

12. Citrine ointment 3 oz., lard 2 oz., turpentine 2 drs., saturated solution of nitrate of copper 2 drs.—BLAINE.

13. *Ægyptiacum* 8 oz., lard 4 oz., sulphate of zinc in powder 1 oz.; rub together till perfectly mixed.—LASSAIGNE.

14. Prepared verdigris 1 oz., lard 4 oz., honey a sufficient quantity.—DELAFOSSÉ.

15. *To remove the fetor in grease.* Chloride of lime 1 to 2 parts, lard 3 parts; mix.—MORTON.

OINTMENTS TO PROMOTE THE GROWTH OF THE HAIR, and to remove the blemish from broken knees.

1. Camphor ½ dr., oil of rosemary 1 dr., weak mercurial ointment 1 oz., ivory black and bole to colour.—WHITE.

2. Poplar-bud ointment and honey, applied twice a day for 15 or 20 days.—PYE.

3. Calamine 2 drs., prepared charcoal 1 dr., oil of turpentine 1 dr., lard 4 drs.; rub well together with 1 dr. of blister ointment.—CLATER.

4. Liquid blister (No. 3, 6, or 7) 1 dr., ivory black 1 dr., camphor 1 dr., palm oil 1 oz.

5. Citrine ointment 1 oz., camphor 1 dr.; colour as above.

HELLEBORE OINTMENT. Powdered white hellebore 1 part, lard 8 parts; an irritating dressing for rowels and setons.—V. C.

HOOF OINTMENT. Tar and tallow, equal parts, melted together.—WHITE.

IODINE OINTMENT (Simple). Iodine 1 part, lard 8 parts; mix.—V. C.

FARCY OINTMENT. Iodine 1 dr., lard 1 oz., mercuria ointment 1 oz.; mix. Useful when the complaint is confined to one leg; from 5 to 10 grs. of iodide of potassium being given daily, with a mineral tonic.

IODINE OINTMENT (Compound). Iodine 1 dr., iodide of potassium 2 drs., lard 2 oz.—V. C.

IODIDE OF MERCURY OINTMENT. Red iodide of mercury 1 part, lard or palm oil 7 parts; mix; the size of a nut to be rubbed on daily; in thoroughpin.

For strangles. Iodide of mercury 1 dr., lard 1 oz.—**SPOONER.**

OINTMENTS FOR MANGE AND LICE.

1. Sulphur 4 oz., soft soap 4 oz., oil of bays 4 oz., train oil q. s.

2. Sulphur 1 oz., train oil 1 oz., Venice turpentine 2 oz., —**YOUATT.**

3. Train oil 3 oz., sulphur 1 oz., oil of turpentine 6 oz., —**WHITE.**

4. Sulphur 8 oz., common turpentine 2 oz., strong mercurial ointment 2 oz., linseed oil 1 pint; rub the flowers of sulphur with a fourth part of the oil, then rub in the turpentine and ointment, and gradually add the rest of the oil; half to be rubbed in daily for three days; on the sixth day, wash off with soft soap and warm water.—**CLATER.**

5. Oil of turpentine 3 oz., oil of vitriol 1 oz.; mix cautiously, avoiding the fumes, and add melted lard 8 oz., train oil 4 oz., oil of turpentine 2 oz., flowers of sulphur or sulphur vivum 4 oz.; stir till cold; apply daily for 3 or 4 times, and give an alterative powder twice a day.—**WHITE.**

6. Oil of bay 16 oz., strong mercurial ointment 6 oz.,

oil of turpentine 2 oz., soft soap 4 oz.; mix and apply in the sun; but it is not quite safe.—BRACKEN.

7. Oil of turpentine 4 oz., oil of tar 4 oz., train oil 8 oz., sulphur 4 oz.

8. Sulphur vivum 8 oz., powdered stavesacre 1 oz., mercurial ointment 2 oz., turpentine 2 oz., lard or train oil 8 oz.—BLAINE.

9. Sulphur 4 oz., white hellebore $\frac{1}{2}$ oz., oil of tar 3 oz., train or linseed oil 12 oz.—SPOONER.

10. Soft soap and tar, equal parts.

11. Weak mercurial ointment $\frac{1}{2}$ lb., sulphur vivum 4 oz., white hellebore 3 oz., black pepper 3 oz., oil of tar 1 oz., olive oil enough to make it soft; use daily for 7, 10, or 14 days.—TAPLIN.

OINTMENTS FOR MALLENDERS AND SALLENDERS.

(Scurvy eruptions.)

1. Citrine ointment 2 oz., tar ointment 1 oz.; mix.

2. Lard 2 oz., finely powdered red precipitate 2 drs.—WHITE.

3. Sugar of lead 1 part, tar 2 parts, lard 6 parts; mix, give a diuretic ball occasionally.—YOUATT.

4. Lard 2 oz.; melt, and stir in Goulard's extract 1 oz.—WHITE.

5. Quicksilver 1 oz., common turpentine 3 oz.; mix.—BRACKEN.

6. Sublimate 10 grs., mercurial ointment 1 oz.; mix.

7. Iodide of potassium 1 dr., lard 2 oz., Goulard's extract 4 drs.; mix.

8. Camphor 1 dr., sugar of lead $\frac{1}{2}$ dr., mercurial ointment 1 oz.; mix, and apply after washing with soap and water.—BLAINE.

9. Naphthalin 1 dr., cod-liver oil 1 oz., zinc ointment 1 oz.

MARSHMALLOW OINTMENT. The following is often substituted for the old Pharmacopœia preparation:—Rape oil 1 lb., yellow wax 6 oz., palm oil $\frac{1}{2}$ lb., common turpentine 1 oz.

MERCURIAL OINTMENT. This is prepared in the usual way; but Venice turpentine is often used to incorporate the quicksilver more speedily, as it does not interfere with its veterinary uses.

Strong Mercurial Ointment. 1. Quicksilver 16 oz., Venice turpentine 2 oz.; rub together till the metal is incorporated, then add 16 oz. of lard.

2. Quicksilver 16 oz., liquid styrax 5 drs., lard 3 oz.; triturate until the metal is incorporated, and add 12 oz. more lard.—CRESENT.

Weaker Mercurial Ointment. 1. Strong mercurial ointment 1 part, lard 2 parts.

2. Quicksilver 2 oz., balsam of sulphur $\frac{1}{2}$ oz.; rub together till the globules disappear, and add 6 oz. of lard.—TAPLIN.

Compound Mercurial Ointment. Mercurial ointment 1 part, soft soap 2 parts.—V. C.

RESOLVENT OINTMENTS, for indolent tumours of the withers, spavins, windgalls, farcy buttons, splints, &c.

1. Strong mercurial ointment 4 oz., cantharides in powder $\frac{1}{2}$ oz., oil of rosemary 2 drs.—WHITE.

2. *For spavin and thoroughpin.* Binodide of mercury 1 part, lard or palm oil 7 parts; rub together in a mortar; the quantity of a nut to be rubbed on daily till a scurf is produced.—SPOONER.

3. Blister ointment 2 oz., strong mercurial ointment 1 oz., soft soap $\frac{1}{2}$ oz., oil of bays 3 drs., yellow wax 3 drs.; melt the wax by a gentle heat, and add the other ingredients, mix by stirring, remove, and stir till cold.—LEBAS.

4. Common turpentine 12 parts, corrosive sublimate 1 part; mix.—GIRARD.

OINTMENT OF NITRATE OF SILVER. Nitrate of silver 5 to 10 grs., lard 1 oz.—MORTON.

CREASOTE OINTMENT. Creasote 1 part, lard 8 parts.—V. C.

MILD CITRINE OINTMENT. Ointment of nitrate of quicksilver 1 part, lard and oil, of each 2 parts. In tarsal ophthalmia.

SOFTENING AND COOLING OINTMENTS, for cracks, and ulcers on the heel.

1. Spermaceti ointment 4 oz., olive oil 1 oz., sugar of lead 2 drs., oxide of zinc 1 oz.—WHITE.

2. Extract of lead $\frac{1}{2}$ dr., lard 1 oz.; mix.—CLATER.

3. Marshmallow ointment 4 oz., extract of lead 8 drs., elder ointment $\frac{1}{2}$ oz., calamine 1 oz.

OINTMENT FOR SIT-FASTS, and all hard tumours. Strained ammoniacum 4 oz., mercurial ointment 8 oz., oil of turpentine 10 oz.—**HINDS**.

OINTMENTS FOR SORE BACKS AND SADDLE-GALLS. (See *LOTIONS, Veterinary Formulary*.)

1. Camphor 2 drs., oil of rosemary 1 dr., elder ointment or lard 3 oz.

2. Marshmallow ointment 4 oz., extract of lead 1 oz.—**WHITE**.

SULPHURIC ACID OINTMENT.

1. Sulphuric acid 1 dr., lard 1 oz.; mix.

2. Sulphuric acid 1 fluid oz., lard 8 oz., oil of turpentine 1 oz.

OINTMENT FOR SPAVINS AND WINDGALLS. See **RESOLVENT OINTMENTS**.

TAR OINTMENT. Equal parts of tallow and tar, melted together.

OINTMENTS FOR THRUSH AND CANKER.

1. Common verdigris $\frac{1}{2}$ oz., calamine $\frac{1}{2}$ oz., sulphate of zinc 1 dr., tar 3 oz.—**BLAINE**.

2. Blue vitriol 2 oz., white vitriol 1 oz., rubbed down and mixed with lard 2 lbs., tar 1 lb.; a pledget of tow covered with it to be introduced into the cleft of the frog every night, and renewed in the morning.—**YOUATT**.

3. *Thrush Paste.* Alum, blue vitriol, white vitriol, of each 1 oz.; rub them into a fine powder; melt 2 lbs. of tar with 1 lb. of lard, and when getting cool stir in the powder.—**CLATER**.

4. Verdigris $1\frac{1}{2}$ oz. (or burnt alum 8 oz.), red lead 8 oz., treacle 4 lbs.; boil to a proper consistence, and add 1 oz. of nitrous acid.—**FERON**.

5. Barbadoes tar 4 oz., sulphuric acid $\frac{1}{2}$ oz.—**SPOONER**.

6. *In Canker.* Tar 4 parts, nitric acid 1 part; mix.—**MORTON**.

* * * The feet should be kept perfectly clean.

OINTMENTS FOR FARCY BUDS. 1. Sublimate 1 oz., white arsenic $\frac{1}{2}$ oz., yellow arsenic $\frac{1}{2}$ oz., euphorbium $\frac{1}{2}$ oz., oil of bays 4 oz.; mix.—LAMOTTE.

2. (*Topique Terrat.*) Corrosive sublimate 1 oz., white and yellow arsenic, each $\frac{1}{2}$ oz., oil of bays 4 oz.; mix with a gentle heat.

TURPENTINE OINTMENT. Common turpentine 1 part, lard 3 parts; melt together.—V. C.

VERDIGRIS OINTMENT. Verdigris in powder 1 part, common turpentine 1 part, lard 12 parts; for foul ulcers and tarsal ophthalmia.—MORTON.

OINTMENT FOR WARTS AND ANBURIES. 1. Chloride of ammonium 2 drs., powdered savin 1 oz., lard $1\frac{1}{2}$ oz.; to be applied daily.—BLAINE.

2. Arsenic 1 part, lard 8 parts.—FINLAY DUN.

OINTMENTS FOR CHRONIC VIVES.

1. Emetic tartar 2 drs., olive oil 1 dr.; rub together till smooth, and add lard 1 oz.

2. Iodide of potassium 1 dr., palm oil 1 oz.; rub together till quite smooth.—CLATER.

ASTRINGENT PASTE, for broken knees and for wounds after the inflammation has subsided.

1. Powdered alum and pipeclay, mixed with water to the consistence of cream. For broken knees it may be coloured with bole and lamp-black.—WHITE.

2. *Paste for Open Knee-Joint.* Flour and stale beer, boiled to the consistence of paste, and coloured as above. To be spread thick all round the joint, and covered with a pledget of tow and $\frac{1}{2}$ sheet of brown paper; and the leg of a cotton stocking drawn over the whole. The stocking to be covered with the paste, and enveloped with 2 calico bandages regularly applied.—TURNER.

STOPPING FOR THE FEET.

1. Cow-dung beaten with a fourth part of clay.—YOUATT.

2. Soft soap 4 oz., Barbadoes tar 16 oz., linseed meal $2\frac{1}{2}$ lbs.—WHITE.

3. Tallow and tar, equal parts, melted together.

4. Common tar 2 parts, soft soap 1 part, linseed meal

q. s. To be spread over the sole of the foot $\frac{1}{2}$ of an inch thick, covered with a layer of tow, and a leather sole over all.—V. C.

HOOF OINTMENTS.

1. Equal parts of wax, olive oil, lard, veal suet, turpentine, and honey; melt the wax, suet, and lard with the oil by a gentle heat, remove from the fire, and add the honey and turpentine, stirring till cold; when intended to embellish the hoof as well as to soften it, it may be coloured with lamp-black or ivory-black.—BOURGELAT.

2. Tallow 4 lbs., beeswax 4 oz., tar $\frac{1}{2}$ lb.; melt slowly, remove from the fire, and when they begin to cool, stir together. A portion of pitch may be added when intended to fill fissures, &c.—BRACY CLARK.

COMPOSITION FOR SAND CRACKS. Beeswax 4 oz., yellow resin 2 oz., common turpentine 1 oz., tallow $\frac{1}{2}$ oz.; melt together; fill the cracks with the composition, and turn the horse out to grass.

SUPPLING LINIMENT FOR BRITTLE HOOF. Oil of tar 1 pint, fish oil 2 pints.—CLATER.

CHARGES

The usual method of applying charges is to soften the compound by heat, and apply it with a large spatula to the part, as warm as the animal can comfortably bear it, and while warm, to cover it with cut tow. Charges are used for old sprains of the loins, strains of the back sinews, windgalls, &c. Cold charges are spread on cloth or leather, and renewed as they become dry.

1. *Simple Charges.* Pitch 4 oz., turpentine 1 oz.—GASPARIN.

2. *For Strains of the Loins.* Pitch 4 lbs., turpentine 6 oz., olive oil 4 oz.; melt together.—B. CLARK.

3. Burgundy pitch 4 oz., wax 4 oz., yellow resin 4 oz., common turpentine 1 oz.; melt together, and when it begins to thicken, stir in 1 oz. of bole.—WHITE.

4. Burgundy or common pitch 5 oz., tar 6 oz., wax 1 oz.; melt together, and when they are becoming cool, stir in $\frac{1}{2}$ dr. of powdered cantharides.—YOUATT.

5. Pitch 3 lbs., tar 1 lb., beeswax $\frac{1}{2}$ lb.; melt together.—CLATER.
6. Resin 2 oz., Burgundy pitch 4 oz., Barbadoes tar 2 oz., wax 3 oz., red lead 4 oz.—WHITE.
7. Pitch 8 oz., suet 4 oz., oil of turpentine 3 oz.; tincture of cantharides 3 oz.—DELAFOSSÉ and LASSAIGNE.
8. *Cold Charge.* For sprains in the back sinews. Bole $\frac{1}{2}$ lb., white of egg and vinegar to form a soft paste, to be applied on doubled cloth or leather, and removed as it dries.—BRACKEN.
9. Bruised leaves of elder, or cabbage, or mallow.—B. CLARK.
10. *Mercurial Charge.* Burgundy pitch $1\frac{1}{2}$ oz., wax $1\frac{1}{2}$ lbs.; melt, and add, while cooling, 9 oz. of mercurial ointment previously mixed with 6 drs. of iodine.—Mr S. FISHER.
11. *Soot Charge.* Common turpentine $4\frac{1}{2}$ oz., soot 3 oz.; mix.—DELAFOND.

POWDERS FOR OUTWARD USE

ASTRINGENT POWDERS; chiefly used for sprinkling greasy or ulcerated heels, after the inflammation has been subdued by poultices,—and in joint wounds.

1. Calamine (true) 4 parts, alum 1 part; mix.—MORTON.
2. Burnt alum, dried sulphate of iron, and myrrh, equal parts.—V. C. (Comp. powder of alum.)
3. Alum 1 dr., charcoal $\frac{1}{2}$ oz., chalk 2 oz.—BLAINE.
4. Sulphate of zinc, chalk slightly calcined, white pepper, in equal parts.—B. CLARK.
5. Alum 4 oz., bole 1 oz.—WHITE.
6. Oak bark 1 oz., verdigris 2 drs.—BLAINE.
7. White vitriol 2 oz., oxide of zinc 1 oz.—WHITE.
8. Prepared chalk 4 oz., sulphate of zinc 1 oz., charcoal 1 oz., Armenian bole 2 oz.—SPOONER.

DETERGENT AND ESCHAROTIC POWDERS; for cleansing foul ulcers and repressing fungus or proud flesh. They should all be very finely powdered and well mixed.

1. Equal parts of calcined white vitriol and alum.—BRACKEN.

2. Bole 2 drs., blue vitriol or verdigris 1 oz.—WHITE.

3. Red precipitate $\frac{1}{2}$ oz., acetate of copper $\frac{1}{2}$ oz., calamine $\frac{1}{2}$ oz.—BLAINE.

4. Red precipitate $\frac{1}{2}$ oz., burnt alum 2 drs.

5. Blue vitriol 1 oz., alum 1 oz., white lead 1 oz.

6. Equal parts of verdigris and sugar of lead.—CLATER.

7. *In joint wounds.* Alum, dried sulphate of iron, and myrrh, of each equal parts.

8. Alum, sulphates of iron, of zinc, and of copper, of each 1 oz., chloride of ammonium $\frac{1}{2}$ oz., camphor and saffron, of each, $1\frac{1}{2}$ drs.—BOUCHARDAT.

STYPTIC POWDER. Alum, with an equal or double weight of flour.—WHITE.

STYPTIC STONE. See SUNDRY SOLUTIONS, under "MEDICINE FOR HORSES," further back.

SNEEZING POWDERS. The ingredients to be very finely powdered and mixed.

1. Asarabacca 4 drs., white hellebore 1 dr.; mix, and keep in a bottle for use.—BRACKEN.

2. Snuff 1 oz., hellebore 1 dr., euphorbium 10 to 20 grs.—PECK.

3. *In incipient cataract.* Turpeth mineral 2 drs., asarabacca 4 drs.; mix, and apply as much as will lie upon a sixpence, daily.—BRACKEN.

MEDICINES FOR NEAT CATTLE

DRINKS OR DRENCHES

Note.—The peculiar structure of the digestive organs in cattle renders it proper to give their medicines in a liquid form. For the same reason, drenches should be given very slowly, so as to enter at once the third or fourth stomach. It is only in cases of *hoven* or *blown*, that it is desirable to introduce medicine into the first stomach or *rumen*.

For treatment of the Cattle Plague of 1865, see p. 121.

PURGING AND LAXATIVE DRENCHES.

These are given when fever exists, or is threatened; to prevent downfalls of the udder; after calving, to prevent milk-fever; to remove undue accumulations in costiveness; in the first stage of red-water and jaundice; and in all inflammatory complaints.

1. Epsom salts 8 oz., sulphur 4 oz., ginger 2 drs., warm water a pint, linseed oil 12 oz.—SPOONER.

2. Epsom salts 6 or 8 oz., castor oil 8 oz., gruel $1\frac{1}{2}$ pints, ginger $\frac{1}{2}$ oz.

3. Glauber's or Epsom salts 16 oz. (or in bad cases with fever, 24 oz.), caraways 1 oz., warm gruel a quart.—CLATER.

4. Castor oil from 16 to 24 oz., with gruel; but it is not to be depended on.

5. To No. 3, add 2 or 3 drs. of gamboge, or 4 drs. of aloes.

6. Sulphur 8 oz., ginger $\frac{1}{2}$ oz., warm gruel a quart; in rheumatism, or joint-fellon.—CLATER.

7. Common salt 6 oz., flour of mustard a tablespoonful, grated ginger or ground pepper, of either, a teaspoonful gin $\frac{1}{2}$ pint, water 2 pints.

8. Common salt 1 lb., warm water, or guel, q. s. The last three are only proper where there is not much fever.

9. *In red-water.* Sulphate of magnesia 8 to 16 oz., sulphur 2 to 6 oz., carbonate of ammonia $\frac{1}{2}$ oz., ginger $\frac{1}{2}$ oz., warm water q. s.; a fourth of this every 6 hours till the bowels are sufficiently acted on.—SPOONER.

10. *When the last does not operate.* Calomel 20 grs., yeast $\frac{1}{2}$ pint.—HARRIS.

11. Aloes 4 to 6 drs., common salt 4 to 6 oz., ginger 1 to 3 drs., water a quart, anodyne tincture 2 oz., in red-water.—WHITE.

12. *Cordial Purgatives.* Aloes 4 drs., Epsom salts 4 oz., ginger 1 dr., carminative tincture 2 oz., water 1 quart.—WHITE.

13. *In the commencement of puerperal or milk fever.* Epsom salts 6 or 8 oz., powdered croton seeds 20 to 30 grs., ginger 4 drs., in 3 or 4 pints of gruel: repeat in 6 hours, if required, without the croton seeds.—BLAINE.

14. *In locked jaw.* Barbadoes aloes $1\frac{1}{2}$ oz., powdered croton kernel 10 grs., boiling water q. s.; given when cool.—CLATER. See Mr FINLAY DUN's treatment, further on.

15. *Mild laxative and tonic.* Epsom salts $\frac{1}{2}$ lb., sulphur 4 to 6 oz., ginger $\frac{1}{2}$ oz., gentian $\frac{1}{2}$ oz., warm water q. s.—EVESON.

16. *In flatulent colic with costiveness.* Aloes $1\frac{1}{2}$ oz., carbonate of potash 3 drs., ginger $\frac{1}{2}$ oz., warm water a pint, linseed oil 8 oz.—WHITE.

17. Palm oil 16 oz., Glauber's salts 12 oz., boiling water q. s.—PECK.

18. *Laxative drink for cows that are kept on hay.* Aloes 4 drs., ginger $1\frac{1}{2}$ drs., water a quart, Epsom salts 6 oz., carbonate of soda $\frac{1}{2}$ oz.; for one dose.—YOUATT.

FEVER DRENCHES, for fevers, colds, influenza, &c.

1. *In simple cold or catarrh.* Tartar emetic 1 dr., digitalis $\frac{1}{2}$ dr., nitre 3 drs.; mix, and give in a quart of gruel.—CLATER.

2. *After bleeding, in inflammation of the bladder.* Antimonial powder 2 drs., opium a scruple; rub together, and mix with thick gruel.—WHITE.

3. *In influenza, or epidemic (epizootic) colds.* Nitre $\frac{1}{2}$ oz., salt of tartar 1 oz., camphor 2 drs., valerian, liquo-

rice, turmeric, of each 1 oz., mustard 2 oz., juniper berries 1 oz., gruel a quart.—SKERRETT.

4. *For the same.* After bleeding and a laxative, give antimonial powder 2 drs., camphor $1\frac{1}{2}$ drs., ginger 3 drs., laudanum $\frac{1}{2}$ oz., in gruel.—WHITE.

5. *In bad colds, attended with fever.* Nitre 1 oz., camphor $\frac{1}{2}$ dr., tartar emetic $\frac{1}{2}$ dr., in gruel.—PECK.

FEBRIFUGE TONIC DRENCHES.

1. Antimonial powder $\frac{1}{2}$ dr., camphor 1 dr., Peruvian bark 1 oz., gruel, or decoction of arrowroot, or starch, q. s. for two doses. PECK.

2. *In the decline of fevers and influenza.* Emetic tartar $\frac{1}{2}$ dr., nitre 2 drs., gentian 3 drs., chamomile 1 dr., ginger $\frac{1}{2}$ dr.; pour on them a pint of boiling ale, and give when cool.—CLATER.

3. Emetic tartar $\frac{1}{2}$ dr., gentian 2 drs., digitalis $\frac{1}{2}$ dr., nitre $\frac{1}{2}$ oz., spirit of nitric ether 4 drs., gruel q. s.

TONIC DRENCHES.

1. Cascarella 3 drs., ginger 3 drs., carbonate of soda 2 drs., in gruel.—WHITE.

2. Gentian $\frac{1}{2}$ oz., ginger 1 dr., Epsom salts 2 oz., warm gruel a pint.—CLATER.

3. Tartarized iron 1 dr., gentian 2 drs., ginger 1 dr., gruel 1 pint, after laxatives, in indigestion.

DRENCHES FOR INFLAMMATION OF THE LIVER.

After bleeding give—

1. Calomel $1\frac{1}{2}$ drs., opium $\frac{1}{2}$ dr., ginger 2 drs., thick gruel q. s. Six hours afterwards, give Epsom salts 1 lb., sulphur 6 oz., linseed oil $\frac{1}{2}$ pint, gruel q. s.—SPOONER.

2. Epsom salts 1 lb., caraway $\frac{1}{2}$ oz., Barbadoes aloes $\frac{1}{4}$ oz.; in a quart of warm gruel.—CLATER. After the yellowness appears, give—

3. Half of No. 2, with 20 grs. of calomel morning and night.—CLATER.

4. Mr FINLAY DUN's treatment is—Cathartics. Salines. Aconite. Chloride of Ammonium. Laxative diet.

DRENCHES FOR JAUNDICE OR YELLOWS.

1. Opium 10 grs., calomel 10 grs., thick gruel q. s., at night, and the tonic drink (No. 2) in the morning.—CLATER.

2. Mr. SPOONER says salts in $\frac{1}{2}$ -lb. doses, with a little ginger, are generally sufficient. "Nothing can be done without purgatives."—FINLAY DUN.

3. Chloride of sodium $\frac{1}{2}$ oz., carbonate of soda $\frac{1}{2}$ oz., turmeric 2 oz., Glauber's salts 6 oz., powdered gentian and chamomile 2 drs., gruel q. s.

4. Castile soap $\frac{1}{2}$ oz.: Venice turpentine $\frac{1}{2}$ oz., ginger 3 drs., gentian 1 oz.; rub the soap and turpentine in a mortar, and gradually add a pint of water, and afterwards the ginger and gentian.—WHITE.

5. Castile soap 1 oz., salt 1 oz., Venice turpentine 1 oz., yolk of 2 eggs; mix together, and gradually add a strong decoction of barberry bark.

6. Powdered cumin seed, aniseed, and turmeric, each 2 oz., grains of paradise and salt of tartar, each 1 oz., mix. Slice 1 oz. of Castile soap to mix with 2 oz. of treacle. Pour a quart of boiling ale upon all the ingredients, and administer when lukewarm. To be repeated two or three times a day.

CLEANSING DRINKS, for cows after calving. These are often applied for, but are condemned as useless or hurtful by veterinarians of the new school. The following are some of the forms in use; probably a gentle laxative would be in most cases preferable.

1. Spermaceti, Irish slate, and birthwort, in powder, of each 1 oz., powdered aniseed 2 oz., liquorice powder 2 oz.; in linseed tea.

2. Aniseed, myrrh, birthwort, allspice, cumin seed, of each 1 oz., in a quart of gruel.—M'EWEN.

3. Juniper berries 3 oz., birthwort 2 oz., fenugreek 1 oz., spermaceti 2 oz., antimony 1 oz., saffron $\frac{1}{2}$ oz.; in a quart of warm ale.—DOWNING.

4. Resin, soap, of each $\frac{1}{2}$ oz., spermaceti $\frac{1}{2}$ oz., aniseed, caraway seed, of each 1 oz., ginger $\frac{1}{2}$ oz., treacle 4 oz., warm gruel a quart.

5. 1 oz. spermaceti, 1 oz. birthwort, 2 oz. powdered bay-berries, 1 oz. myrrh; in juniper-berry tea.

LAXATIVE DRINK AFTER CALVING. Epsom salts 12 oz., aniseed 1 oz., olive oil 6 oz., gruel a pint, or q. s.

DRENCH FOR STRANGURY. After laxatives and a clyster, give camphor 2 drs., spirit of nitrous ether $\frac{1}{2}$ oz.,

tincture of opium $\frac{1}{2}$ oz., nitre 1 oz., gruel a pint.—**WHITE.**

DRENCHES FOR HOVEN OR BLOWN (flatulent distension of the paunch). It appears doubtful whether any liquid enters the paunch in these cases. More dependence is now placed on the introduction of a tube, constructed for the purpose.

1. Ginger $\frac{1}{2}$ oz., spirit of nitric ether 2 oz., oil of peppermint 30 drops, warm water a pint.—**WHITE.**

2. Liquid ammonia, or spirit of hartshorn, $\frac{1}{2}$ oz. to 1 oz. ($1\frac{1}{2}$ oz.—**WHITE**), cold water 3 pints.—**MOIROUD.**

3. Chloride of potash 4 drs., water 4 oz., ether 3 drs. The solution of chlorinated soda may be substituted for chloride of potash (*Eau de Javelle*).—**CHARLOT.**

4. Aloes 3 drs., pimento 2 drs., oil of turpentine, 2 oz., spirit of hartshorn 1 oz., in gruel or warm water.—**BLAINE.**

5. Chloride of lime 2 drs., water a quart. Administer it by means of a stomach-pump, and repeat in an hour if required.—**YOUATT.**

RHEUMATIC DRENCH.

1. Sulphur 8 oz., ginger $\frac{1}{2}$ oz.; in gruel, every third day if necessary.—**CLATER.**

2. Antimonial powder 2 drs., Dover's powder $\frac{1}{2}$ dr., aniseed 1 oz., thick gruel a pint; night and morning, the bowels having been opened by No. 1.—**CLATER.**

3. Rhododendron leaves 4 drs., water a quart; boil to a pint, strain, and add powdered gum guaiacum 2 drs., caraway-seeds and aniseed, of each 2 drs., warm ale $\frac{1}{2}$ pint.

ANTISPASMODIC DRENCH FOR LOCKED JAW.

1. Camphor 1 dr. (rubbed with spirit), powdered opium 1 dr., thick gruel $\frac{1}{2}$ pint.

2. MR FINLAY DUN'S TREATMENT OF LOCKED JAW.

Purgatives. Belladonna and chloral hydrate. Cool air. Warm clothing. Perfect quiet. Soothe any wound. Vapour-bath.

CORDIAL CARMINATIVE DRENCHES. Drenches for indigestion, and colic without inflammation.

1. *In indigestion.* Salt 3 or 4 oz., carbonate of soda 2 drs., ginger $\frac{1}{2}$ dr., anodyne carminative tincture (see

TINCTURES, SOLUTIONS, *Veterinary Formulary*) 2 oz., water 10 or 12 oz.—WHITE.

2. *The same.* Aloes 4 drs., common salt 4 oz., ginger 2 drs., anodyne carminative tincture (see TINCTURES, SOLUTIONS) 2 oz., gruel q. s.

3. *In Colic.* Salt 4 oz., aloes 3 drs., ginger 1 dr., opium $\frac{1}{2}$ oz., water 1 pint, peppermint water 1 pint.

4. *Carminative.* Oil of turpentine 1 oz., tincture of opium 6 drs., spirit of nitric ether 2 oz., water 1 pint.—WHITE.

5. *Warm Cordial.* A bottle of red wine, extract of juniper 1 oz., powdered cinnamon $\frac{1}{2}$ oz.—LEBAS.

6. *Mild.* Peppermint 2 oz., chamomiles $\frac{1}{2}$ oz., hot water 5 pints; infuse, and give while warm.

7. Chamomile 2 oz., aniseed $1\frac{1}{2}$ oz.; infuse in hot water, and strain; when cold, add ether 2 oz.—VATEL.

DRENCHES FOR BLOODY URINE.

Bloody Urine (*Hæmaturia*) and Red-water, although often confounded, are different diseases, and require a different treatment. *Hæmaturia* is distinguished by the presence of actual blood in the urine, in a state of coagulation, and by great tenderness across the loins. It generally occurs in oxen of good condition. It is to be treated by gentle purgatives, stimulating applications to the loins, emollient drinks and opiates. [SPOONER.] Give one of the following drenches:

1. Epsom salts 6 to 8 oz., water a quart, castor oil 4 to 6 oz.—WHITE. Or,

2. Linseed oil 1 pint, gruel 1 pint, caraways 2 drs., Epsom salts 8 oz. (in warm water $\frac{1}{2}$ pint), tincture of opium 2 drs. Or either of the laxative drenches for red water (further on).

3. After the above, when the pain and difficulty have abated, but the water continues bloody, give—Catechu 2 drs., opium $\frac{1}{2}$ dr., alum 3 drs., gum arabic $\frac{1}{2}$ oz., water $\frac{1}{2}$ pint; simmer for a few minutes, and add $\frac{1}{2}$ pint of ale. Repeat if required.

4. *In obstinate cases.* Oil of juniper $\frac{1}{2}$ oz., oil of turpentine 1 oz., laudanum 1 oz., in a pint of linseed tea; at first twice, and afterwards once a day.—RUSH.

5. Three quarters of a pint of black beer, and 2 oz. Irish slate.—KNOWLSON.

6. Sulphate or sesquichloride of iron. Gallic acid.

DRENCHES FOR RED WATER.

This commonly attacks milch cows, and appears generally to arise from the nature of the pasture. Moderate bleeding is recommended, but is less necessary than in hæmaturia. Laxatives should then be given. Mr FINLAY DUN (on the contrary) says blood-letting is decidedly injurious, and he also discountenances the employment both of diuretics and astringents.

1. Epsom salts 8 to 12 oz., sulphur 2 to 4 oz., carbonate of ammonia $\frac{1}{2}$ oz., ginger $\frac{1}{2}$ oz., warm water 4 pints; give a fourth part every 6 hours till the bowels are acted on.—SPOONER. Or,

2. Glauber's salts 12 oz., carbonate of soda $\frac{1}{2}$ oz., nitre $\frac{1}{4}$ oz., sugar 1 oz., powdered caraways $\frac{1}{2}$ oz., in a quart of gruel. Or the Purging drenches, No. 1, 2, or 3. After the bowels are well open, give astringents or mild stimulants.

3. The laxative drench, No. 11.—WHITE. This is WHITE's drench for red-water, No. 1. To be followed by drenches of whey.

4. *Astringent*. Powdered oak-bark $\frac{1}{2}$ oz., catechu 2 drs., opium 10 grs., gruel 1 pint.—CLATER.

5. Catechu 2 drs., mucilage 4 oz., lime-water 6 oz.—BLAINE.

6. Laudanum $\frac{1}{2}$ oz., sugar of lead $\frac{1}{2}$ dr., catechu 4 drs., gruel 1 quart.—WHITE.

7. *After laxatives*. Ginger, gentian, each 1 dr., spirit of nitrous ether 1 oz., gruel q. s.; twice a day.—SPOONER.

8. Powdered oak-bark 1 oz., charcoal 1 oz., bole 2 oz., in a quart of new milk.—PYE.

9. Catechu $1\frac{1}{2}$ oz., alum $1\frac{1}{2}$ oz., diapente 2 oz., Locatelli balsam 2 oz., warm gruel 3 pints.—M'EWEN.

10. Dragon's blood 2 oz., rust of iron 1 oz., nitre 3 oz., oil of turpentine 2 oz.; mix; for 2 doses, in gruel.—DOWNING.

11. Sulphuric acid 1 dr., tincture of opium $\frac{1}{2}$ oz.; treacle 4 oz., warm gruel 4 quarts; daily, for a week.—BLAINE.

DRINK FOR ACUTE DIARRHŒA. Sulphate of soda, sulphate of magnesia, of each 2 oz., ipecacuanha $\frac{1}{2}$ dr., sulphate of iron 6 grs.—BLAINE.

FOR CHRONIC DIARRHŒA. Calomel $\frac{1}{2}$ dr., aloes 1 dr., gentian 2 drs., opium 5 grs., decoction of chamomiles 1 pint.—BLAINE.

ASTRINGENT DRENCHES, for dysentery (scouring rot) or lax.

1. After purging drenches: prepared chalk 2 oz., oak-bark 1 oz., catechu $\frac{1}{2}$ oz., opium 2 scruples, ginger 2 drs., warm gruel 1 quart.—CLATER.

2. Two quarts of alum whey.

3. First give blue pill 2 or 3 drs., rhubarb 3 drs., castor oil 4 oz., gruel 1 pint, well stirred before giving it; repeat this 3 or 4 mornings; then give—thick starch (made with 4 oz. of starch) 3 or 4 pints, tincture of opium 2 drs., ginger 3 drs., catechu $\frac{1}{2}$ oz.—WHITE.

4. Mutton suet 1 lb., new milk 2 quarts; boil, and add opium $\frac{1}{2}$ dr., ginger 1 dr.—CLATER.

5. *Cordial Astringent Drench.* (After the laxative drench No. 2.) Catechu $\frac{1}{2}$ oz., allspice $\frac{1}{4}$ oz., caraways $\frac{1}{2}$ oz., ale $\frac{1}{2}$ pint, water $\frac{1}{2}$ pint.—WHITE.

6. Decoction of sloes, with prepared chalk.—TUSSEY.

7. Decoction of wormwood a quart, gum Arabic 2 oz., aromatic confection 1 oz., catechu 2 drs.; with linseed tea, repeated every 6 days for three times.—RAWLINGS.

8. Sheep's heart, liver, and lights, all chopped up together.—SUMNER (obsolete).

EXPECTORANT AND COUGH DRENCHES IN HOARSE or CATARRH. [See MEDICINES FOR CALVES, FURTHER ON.]

1. Bruised liquorice 2 oz.; boil in a quart of water to a pint, strain, and add powdered squill 2 drs., gum guaiacum 1 dr., tincture of tolu 4 drs., honey 2 oz.—CLATER.

2. *In chronic coughs.* Balsam of sulphur 2 oz., Barbadoes tar 1 oz., yolks of 2 eggs, honey 4 oz., salt of tartar $\frac{1}{2}$ oz., oil of aniseed 1 dr., elecampane 1 oz., gruel 1 quart.

3. *In chronic cough.* Fresh squill 2 oz., garlic 2 oz., vinegar 24 oz.; digest for a day with a gentle heat, strain

and press, and boil the liquor with 24 oz. of treacle : for 6 doses.

4. *For recent coughs.* Digitalis 20 grs., emetic tartar $\frac{1}{2}$ dr., nitre 3 drs., squill 1 dr., opium 20 grs., gruel 1 pint.

5. Boil 4 oz. Iceland moss and 1 oz. liquorice root in 4 quarts of water for a quarter of an hour, and strain; add to the liquor 1 oz. nitre, cream of tartar 1 oz. In hoose from cold, if inflammation of the lungs and fever be present, bleed before giving the drink, at the commencement of the inflammation only. See also FEVER DRENCH (further back).

If the disease be connected with worms in the air passages, give the following :

WORM DRENCH. For cough from worms: Oil of turpentine 2 oz., sweet spirit of nitre 1 oz., laudanum $\frac{1}{2}$ oz.; mix, and give in a pint of gruel.—CLATER.

DRENCHES FOR THE EPIDEMIC, affecting the feet and mouth, and attended with a low fever.

Mr FINLAY DUN recommends the following treatment: Soft laxative food, brought to the patient; rest; cleanliness; comfortable, soft lodging; hydrochloric acid and treacle, and Condry's fluid for mouth, udder, and feet; lead acetate solution. Milk the cows frequently with syphon.

For treatment of the Cattle Plague of 1865, see further on.

1. Glauber's salts 1 lb., treacle 4 oz., sulphur 4 oz., aniseed $\frac{1}{2}$ oz., cream of tartar 1 oz., warm water 3 pints; give it new-milk warm. The above is for a full-grown beast. The mouth to be washed with a strong solution of blue vitriol, burnt alum, and vinegar. If the feet crack, apply a mixture of equal parts of hydrochloric acid and water.

2. Some cattle-masters give common salt in gruel, with great success.—BLAINE.

3. After a mild dose of salts—sweet spirit of nitre 1 oz., ale yeast 6 or 8 oz.—BLAINE.

4. Epsom salts 8 oz., sulphur 2 oz., nitre $\frac{1}{2}$ oz., ginger $\frac{1}{4}$ oz.; half of this to be given in warm water, with 1 oz. of sweet spirit of nitre. Repeat daily. When the bowels are properly relaxed, and the fever reduced, but much weakness remains, give the following :

5. *Tonic Drench.* Gentian 4 drs., ginger 2 drs., sul-

phate of iron 2 drs., sweet spirits of nitre 1 oz., warm water q. s.; wash the mouth with one of the LOTIONS FOR CANKER (*Veterinary Formulary*); dress the feet, after paring and poulticing, with equal parts of tincture of myrrh and butter of antimony, and afterwards apply the ASTRINGENT POWDER (see *Veterinary Formulary*).—SPOONER.

6. Linseed oil a pint, oil of turpentine 8 oz., aloes $\frac{1}{2}$ oz., ginger $\frac{1}{4}$ oz., laudanum 1 oz.; mix. For 2 doses, to be given with gruel.

MURRAIN DRENCHES.

1. Sweet spirit of nitre $\frac{1}{2}$ oz., laudanum $\frac{1}{2}$ dr., solution of chloride of lime 2 drs., prepared chalk 1 oz., mix, and give in a pint of warm gruel.—CLATER.

2. Cascarella powder 2 oz., spirit of nitrous ether $\frac{1}{2}$ oz., liquid acetate of ammonia 4 oz., beer yeast 8 oz.; every 4 hours.—BLAINE.

3. Opiate confection 1 oz., liquid acetate of ammonia 2 oz., water 1 quart; for one dose.—VATEL.

4. *Tonic*. Calumba 2 drs., canella 2 drs., ginger 1 dr., sweet spirit of nitre $\frac{1}{2}$ oz., thick gruel 1 pint.—SPOONER.

5. *Ceylon Remedy*. A small piece of lard the size of a walnut. Said to be used with perfect success.

6. *To Prevent Murrain*. Myrrh 1 oz., Epsom salts 2 oz., sulphur 1 oz., liver of antimony $\frac{1}{2}$ oz., diapente 1 oz.; in rue tea.—DOWNING.

7. Sulphite of soda 1 oz. in 3 galls. of water. The animals are to be allowed to drink *ad libitum*.—Sir J. TYRELL.

LOTION FOR THE MURRAIN OR "FOOT AND MOUTH DISEASE." Dissolve 1 lb. of sulphate of copper in Cj of soft water, wash the animals' mouths with this, with a sponge attached to a stick, then put 2 or 3 large tablespoonfuls of oatmeal and powdered alum, equal parts, into their mouths, near the root of their tongue. Wash their feet, especially between the claws, with the copper solution, and let them stand on dry straw.—Mr SCOTT.

FOR THE DISTEMPER.

Warm tar-water is much recommended by some authorities. It should be given in doses of 3 quarts 3 or 4 times a day, gradually diminishing the quantity.

DIURETIC DRINKS.

1. Common turpentine $\frac{1}{2}$ oz., ginger 2 drs.; mix with a little treacle, and add gradually spirits of nitrous ether 1 oz., gruel a pint.—CLATER.

2. *Tonic Diuretic.* Common turpentine 4 drs., ginger 2 drs., gentian 2 drs., tartrate of iron 1 dr; rub together with a little treacle, and add gradually 1 oz. of sweet spirit of nitre.

BULLING DRENCHES. These are strongly condemned by modern veterinary writers; those drenches, at least, which contain cantharides.

1. Aniseed, grains of paradise, bay berries, of each 1 oz.; cantharides in fine powder 20 to 30 grs.; to be given in a quart of milk.

2. Black hellebore $\frac{1}{2}$ oz., capsicum 2 drs., birthwort $\frac{1}{2}$ oz., bay berries 1 oz., cantharides 20 grs.; in a quart of warm ale.—DOWNING.

3. A quart of milk from a cow in season.—CLATER.

4. Powdered cantharides 20 grs., aniseed 2 oz., black hellebore $\frac{1}{2}$ oz.; in ale, gruel, or milk.—PECK.

DRYING DRENCHES, for drying a cow's milk. Bleed the night before, and give the drink, warm, in the morning. The bleeding is questionable treatment.

1. Boil 6 drs. of alum in milk, and strain.—WHITE.

2. Alum 6 lbs., bole 2 lbs., cream of tartar or red tartar 1 lb.; mix. Give from 6 to 9 oz., in stale beer; or in gruel with $\frac{1}{2}$ pint of vinegar.

3. Roche or common alum $\frac{1}{2}$ oz., dragon's blood $\frac{1}{2}$ oz., tumeric 1 oz., in a pint of rennet water, and a pint of vinegar.—DOWNING. [These large doses of alum, though often given, are not regarded as necessary or proper by modern veterinary writers.]

DRENCH FOR THE BITES OF VIPERS. Olive oil 2 pints, spirits of hartshorn 1 oz.; mix.

ALUM WHEY. Boil $\frac{1}{2}$ oz. of alum in 2 quarts of milk for ten minutes, and strain.

ANODYNE CARMINATIVE TINCTURE; and tincture of Pimento. See MEDICINES FOR HORSES (TINCTURES).

ALTERATIVE POWDERS.

1. Sulphur 4 oz., black antimony 1 oz., Æthiops mineral

$\frac{1}{2}$ oz., nitre 2 oz.; mix, for 4 doses: to be given daily in gruel.

2. *Alterative Tonic.* Add to the last 2 oz. gentian and 1 oz. ginger; and make 6 doses.

FEVER POWDERS, IN INFLAMMATION, &c.

1. Antimonial powder a scruple, camphor $\frac{1}{2}$ dr., nitre 1 oz.; mix; give twice a day in gruel.—PECK.

2. Peruvian bark 8 oz., nitre 24 oz.; for 16 doses.

CORDIAL POWDERS.

1. Black mustard $\frac{1}{2}$ oz., flowers of sulphur 1 oz., aromatic powder (see VETERINARY POWDERS, further back) 1 oz., fenugreek 4 oz., common salt 16 oz.; give 1 oz. on a slice of bread.—MATTHIEU.

2. *Cow Spice.* As Horse Spice, No. 2 (see VETERINARY POWDERS under MEDICINES FOR HORSES, further back). Or,

3. Powdered turmeric, liquorice, aniseed, and diapente, each 1 oz.

CLYSTERS.

1. Salt 1 lb., warm water a gallon.

2. Linseed oil 8 oz., Epsom salts 8 oz., gruel 3 quarts.

MASTICATORIES.

1. Bruised garlic 4 cloves, salt a tablespoonful, ground pepper 1 oz., honey 4 oz. Boil for a short time in a glass of vinegar, immerse it in a piece of linen, and roll it up. Keep it in the animal's mouth for an hour, night and morning. Antiputrescent; in epizootic maladies, and in ulcers of the mouth.—J. ROBINET.

2. Bruised mustard and pepper, each $\frac{1}{2}$ oz., rolled up in linen, and sprinkled with vinegar; to be kept in the mouth not more than half an hour, morning and evening, in epizootic diseases.

REMEDIES FOR THE EPIZOOTIC PNEUMONIA. In the fatal form of this disease which prevailed some years ago, the following treatment is said to have proved effectual:—Bleed freely; then administer $\frac{1}{2}$ pint of brandy every 2 hours. Mr Jeckyll gives, in pleuro-pneumonia, when a tonic is indicated, $\frac{1}{2}$ oz. of the following solution every 8 hours:—Sulphate of copper 1 part, water 4 parts; dissolve, and add ammonia until it begins to precipitate.

Mr FINLAY DUN says bloodletting is quite inadmissible. He prescribes 25 to 30 drops of Flemming's tincture of aconite four times a day.

TREATMENT OF THE CATTLE PLAGUE OF 1865, OR RINDERPEST.

Whether the plague be Smallpox, or a form of enteric fever, or a disease *sui generis*, the precaution of separating infected animals from the remainder of the herd is equally necessary. The diseased carcass must be buried or destroyed at a distance. The most scrupulous cleanliness must be observed in the yards and sheds, and provision made for ventilation and abundant supply of water, with the counteraction of all noxious smells and emanations by means of carbolic acid. When the cattle have sickened with the disorder, it cannot be checked, but, like other contagious diseases, must run its course. The treatment can only be palliative. Among innumerable recipes we select the following as the most likely to be useful. Their application must be guided by the circumstances of the case.

1. *Vaccination.* This has been recommended on the theory that the Cattle Plague is analogous to or identical with Smallpox. Vaccination with the lymph of Cow Pock may then be resorted to as a prophylactic. It may be practised on the udder or vulva of a cow. It has already been resorted to very extensively, but with doubtful success.

2. *The Laxative Treatment.* Linseed oil 6 to 12 oz., with turpentine 1 to 2 oz.; the dose to be repeated once or twice. This dose, as the others, may be given in warm gruel.

3. *The Astringent Treatment.* Applicable in later stages, when there is often much diarrhœa. Chalk and opium (Dr LETHEBY); or laudanum 2 oz., comp. infusion of catechu 1 pint, in a warm mash, with 2 or 3 tablespoonfuls of charcoal.—MALTON AGRICULTURAL ASSOCIATION.

4. *The Saline Treatment.* Useful in moderating the fever, and generally in mild cases. Table salt $\frac{1}{2}$ lb., Epsom salts $\frac{1}{4}$ lb., brimstone $\frac{1}{4}$ lb., ground ginger 2 oz., nitre 2 oz., old ale 1 quart (Dr ALLNATT); or chlorate of potash 2 drs. to $\frac{1}{2}$ oz., thrice daily.—Mr F. BUCKLAND.

5. *The Mixed Saline Treatment.* Adopted in imitation of Dr STEEVENS' treatment of cholera. Chloride of sodium 3 oz., nitre 2 oz., carbonate of potash 3 oz., chlorate of potash 2 oz., sulphate of magnesia 4 oz., old ale 2 quarts (Dr ALLNATT). It would be as well to divide the dose into 3 or 4. Bicarbonate of soda 1 oz., common salt 1 oz., chlorate of potash $\frac{1}{2}$ oz., Rochelle salt 1 oz., in a gallon of water; 1 pint every second hour or oftener.—Dr TUCKER.

6. *The Diaphoretic Treatment.* Useful as the last, in mild cases. Sweating to be promoted by rubbing, wrapping in blankets, or vapour baths. Sweet spirit of nitre 1 to 2 oz., and spirit of Mindererus 1 to 2 oz., in 2 or 3 quarts of linseed meal gruel. Will also act as a diuretic.—MALTON ASSOCIATION.

7. *The Stimulant Treatment.* (1) 16 oz. of linseed oil and a "mutchkin" of whiskey. (Professor DICK.) Large draughts of old ale are recommended by some. 2 drs. of carbonate of ammonia 3 times a day. (Professor DICK.) (2) 12 to 15 grs. of camphor, with 20 grs. of carbonate of ammonia in gruel, thrice daily.—Dr COPLAND.

8. *The Tonic Treatment.* A dr. each of the tincture of perchloride of iron and dilute hydrochloric acid in a quart of linseed tea or gruel, thrice daily. (Dr COPLAND.) $\frac{1}{2}$ oz. of sulphate of iron, twice daily. (Dr TUCKER.) $1\frac{1}{2}$ oz. of powdered cinchona. (Dr SMART.) Strong hydrochloric acid $2\frac{1}{2}$ oz., strong nitric acid $1\frac{1}{2}$ oz., sulphate or chloride of iron $1\frac{1}{2}$ oz., water to make a quart; 1 oz. of this in a gallon of water for a dose.—MALTON ASSOCIATION.

9. *Arsenical Treatment.* Adopted by the homœopaths, with their usual want of success. 1-10th to 1-5th of a grain of arsenic for a dose, every 2 hours, to alternate with 1-50th gr. of phosphorus. Vapour-baths also recommended, and immense quantities of old ale given.—Dr KIDD.

10. Sloppy food. Chlorate of potash, sulphite of soda, sulphurous acid inhalation and sponging. Carbolic acid, mineral acids and bitters. Separate healthy from sick. Disinfectants.—FINLAY DUN.

N.B.—However useful to the individual, these remedies are of little avail in preventing the spread of the disorder. With regard to this plague, and that of late

years, the foot-and-mouth disease, both eminently contagious, the well-being of the community demands the most ruthless use of the poleaxe in all suspected cases. The stamping out of the disorder by the slaughter of all diseased animals, and the prohibition of importations from infected districts abroad, which is done by an Order of Council, are the securest means of checking these destructive epidemics.

MEDICINES FOR CALVES

PURGATIVE DRENCHES.

1. Epsom salts 1 to 2 oz.,* according to the age and size of the calf; dissolve in $\frac{1}{2}$ pint of gruel, and add 20 grs. of ginger and 3 drops of essence of peppermint.—CLATER.

2. Epsom salts $1\frac{1}{2}$ oz., castor oil 2 oz. ginger 10 grs., caraway 2 drs., gruel $\frac{1}{2}$ pint.

3. *In costiveness and accumulation in the paunch and stomach.* Dissolve 2 oz. of Epsom salts in 2 or 3 quarts of water, or 4 oz. in a gallon, according to the age of the calf, and throw it in gently by means of a stomach-pump.

4. *Laxative.* Epsom salts 2 or 3 oz., carbonate of soda 2 drs., water 6 or 8 oz., ginger 1 dr.; mix. After it has operated, give the cordial, No. 3, just further on.—WHITE.

DRENCHES FOR DIARRHŒA, OR CALVES' CORDIAL.

1. YOUATT'S *Cordial.* Prepared chalk 2 oz., catechu 1 oz., ginger $\frac{1}{2}$ oz., opium 1 dr., peppermint water 1 pint; dose for a calf, from 2 to 4 tablespoonfuls.

2. Prepared chalk 2 drs., opium 10 grs., catechu $\frac{1}{2}$ dr., ginger $\frac{1}{2}$ dr., essence of peppermint 5 drops; mix, and give twice a day in $\frac{1}{2}$ pint of gruel.—CLATER.

3. Caraway $\frac{1}{2}$ oz., ginger $\frac{1}{2}$ dr., subcarbonate of soda 1 dr., brandy or gin 1 oz., water 8 oz.—WHITE'S *Cordial.*

4. Half a bottle of Dalby's carminative.

5. Suet boiled in milk $\frac{1}{2}$ pint, opium 5 grs., alum 5 grs., prepared chalk $\frac{1}{2}$ oz.; mix.

6. If No. 2 fails: Dover's powder 2 scruples, aromatic

powder 1 dr., kino $\frac{1}{2}$ dr.; give it night and morning, with 1 oz. of arrowroot boiled in a pint of water.—CLATER.

INFLAMMATORY DISORDERS. Bleed; give 2 to 6 oz. Epsom salts. [Give to a calf of six months old $\frac{1}{2}$ the dose for cattle; at a year and a half, $\frac{1}{3}$ the dose.—SPOONER.]

MEDICINE FOR PILES IN CALVES.

Oil of vitrol 15 drops, tincture of opium $\frac{1}{2}$ oz.—PECK.

SOLUTION OF POTASH, FOR CORDS, &c. Carbonate of potash 2 oz., fresh lime-water 8 oz. To correct acidity in the stomach, give 1 or 2 teaspoonfuls in gruel; the first dose to be given with an ounce or two of Epsom salts in $\frac{1}{2}$ pint of thin gruel. If the disorder is attended with griping pains, add a teaspoonful of anodyne carminative tincture.—WHITE.

ALUM WHEY. See page 119.

TO PROMOTE THE FATTENING OF CALVES.

Aniseed $\frac{1}{2}$ lb.; fenugreek $\frac{1}{2}$ lb., linseed meal (with the oil in it) 1 lb.; make it into a paste with milk, and cram them with it.

Fattening Powder. Common salt with a little carbonate of soda; a small quantity added to the food promotes fattening, and prevents scouring, &c.

HOOSE, OR COUGH FROM WORMS IN AIR PASSAGES.

1. $\frac{1}{2}$ pint lime-water every morning, and a tablespoonful of salt every afternoon, to each calf.—MAYER.

2. Linseed oil 4 oz., oil of turpentine 1 oz., oil of caraways 20 drops; repeated once or twice at intervals of 10 days. This dose for calves of 6 to 10 months old.—DICKENS.

3. A tablespoonful of oil of turpentine, a little sweet oil, and 6 or 8 oz. of warm water.—WHITE

External Applications for Heat Cattle

LOTIONS OR WASHES. (See also EMBROCATIONS, further on.)

LOTIONS FOR CANKER IN CALVES.

1. Alum 1 oz., water 8 oz., tincture of myrrh 1 oz., honey of roses 1 oz.

2. Equal parts of tincture of myrrh and water.—CLATER.

3. Alum $\frac{1}{2}$ oz., water 1 pint, tincture of myrrh 1 oz.

LOTION FOR COW-POCK. Sal ammoniac $\frac{1}{4}$ oz., white vinegar $\frac{1}{2}$ pint, camphorated spirit 2 oz., Goulard's extract 1 oz.; mix.—CLATER.

LOTIONS FOR SLIGHT BRUISES.

1. Extract of lead $\frac{1}{2}$ oz., vinegar 4 oz., soft water 1 pint.—WHITE.

2. Acetate of ammonia 4 oz., water $\frac{1}{2}$ pint, spirit of camphor $\frac{1}{2}$ oz.

DISCUTIENT LOTION, for dispersing tumours. Bay salt 4 oz., vinegar 1 pint, water 1 quart, oil of origanum 1 dr.; rub the oil with the salt, and gradually add the others.

LOTIONS FOR STRAINS.

1. Bay salt 4 oz., oil of origanum 1 dr.; rub together, and add vinegar $\frac{1}{2}$ pint, spirit of wine 2 oz., water 1 quart.

2. Common salt 1 oz., sal ammoniac 1 oz., water 1 pint.

LOTIONS FOR FOUL IN THE FOOT. After poulticing, and removing loose horn, apply—

1. Butter of antimony, or

2. Strong solution of alum.

3. Solution of sulphate of copper.

4. (When the above are not sufficient.) Dissolve 2 drs. of corrosive sublimate in 12 oz. of water.—WHITE.

5. One fluid drachm of carbolic acid in from 6 to 12 oz of water.—TUSON.

LOTIONS FOR WOUNDS.

1. Tincture of myrrh and aloes.
2. *For proud flesh.* Strong solution of sulphate of copper.
3. *For offensive wounds.* Chloride of lime 1 oz., water 1 pint; mix well, and strain.

LOTION FOR BULL-BURNT. Goulard's extract 1 oz., spirit of wine 2 oz., water $\frac{1}{2}$ pint.

LOTION FOR BLAIN IN THE MOUTH. After lancing the bladder, apply a saturated solution of salt in water.—YOUATT.

LOTION FOR THE MOUTH, in ulceration during the epidemic. Alum 1 oz., sulphate of zinc $\frac{1}{2}$ oz., warm water 1 pint, treacle $\frac{1}{4}$ lb.—SPOONER.

WASHES FOR DESTROYING VERMIN. They are all poisonous.

1. Stavesacre seeds 4 oz., water 4 pints; boil to 2 pints, and apply it daily.—PECK.
2. Sublimate 2 drs., spirit of wine 2 oz., water 1 pint.—CLATER.
3. Stavesacre 4 oz., white hellebore root 2 oz., water 1 gallon; boil to half: apply with a sponge.

LOTION FOR MANGE. Corrosive sublimate 2 drs., hydrochloric acid $\frac{1}{2}$ oz., water 12 to 16 oz.; mix. In obstinate cases only.—WHITE. See LINIMENTS.

EYE WATERS, OR COLLYRIA.

1. White vitriol a scruple, spirit of wine 1 dr., water a pint.—CLATER.
2. Sugar of lead 10 to 20 grs.; soft water 8 oz.—V. C.
3. *Sedative Eye Drops.* Powdered digitalis $1\frac{1}{2}$ oz.; infuse in a pint of Cape wine for a fortnight, and filter; a few drops to be introduced into the eye twice or thrice a day.—SPOONER.
4. Extract of lead 2 drs., wine of digitalis (see above) 2 drs., tincture of opium 2 drs., water a pint.

EMBROCATIONS AND LINIMENTS

STRONG EMBROCATION, for deep-seated strains, &c.

1. Oil of origanum $\frac{1}{2}$ oz., oil of turpentine $\frac{1}{2}$ pint, sweet oil $1\frac{1}{2}$ pints, powdered cantharides 1 oz.—CLATER.

2. Olive oil 4 oz., oil of turpentine 1 oz., water of ammonia 1 oz. For strains and bruises, after the inflammation has subsided.—WHITE.

CROTON LINIMENT. Bruised croton seeds 1 part, oil of turpentine 8 parts. Macerate for 14 days, and strain. It irritates the skin powerfully; for general purposes it requires to be diluted with olive oil.—MORTON.

MUSTARD EMBROCATION. Flour of mustard 4 oz., oil of turpentine 2 oz., water of ammonia 2 oz.—WHITE.

RHEUMATIC EMBROCATIONS.

1. Olive oil 2 oz., strong water of ammonia 1 oz., marsh-mallow ointment 1 oz.

2. Neatsfoot oil $\frac{1}{2}$ oz., camphorated oil 1 oz., oil of turpentine 1 oz., laudanum 1 oz., oil of origanum 1 dr.—CLATER.

3. Sweet oil 4 oz., oil of turpentine 2 oz.; mix.—WHITE.

EMBROCATIONS FOR GARGET, or Downfall of the Udder.

1. Oil of elder 4 oz., water of ammonia $\frac{1}{2}$ oz., Minde-rerus spirit 1 oz., camphorated oil 2 oz.

2. Olive oil 3 oz., oil of turpentine 1 oz., camphor 2 drs.—WHITE.

3. Soft soap 8 oz., oil of bays 8 oz., oil of turpentine 8 oz., spirit of camphor 4 oz. (See also OINTMENTS, just further on.)

DRIFFIELD OILS. Barbadoes tar 1 oz., linseed oil 1 lb., oil of turpentine 3 oz., oil of vitriol 1 oz.

LINIMENT FOR MANGE. Sulphur vivum or flower of sulphur 4 oz., train oil 12 oz., oil of turpentine 4 oz.; mix.

LINIMENT FOR SORE THROATS. Oil of turpentine 1 oz., sweet oil 1 oz., water of ammonia 2 oz.—WHITE.

BLISTERING LINIMENT. Cantharides bruised 1 oz., oil of turpentine 8 oz.; digest 14 days, and strain. To be applied by friction on the skin.—**YOUATT.**

LIQUID CAUSTIC. Butter of antimony alone, or mixed with an equal quantity of tincture of myrrh.

LIQUID SNUFF. Alum, sulphate of zinc, capsicum, of each 1 oz.; camphor 2 drs.; pulverize, and macerate in 32 oz. of strong vinegar and 1 oz. of turpentine; shake up when used, and introduce a teaspoonful into the nostrils, to promote a discharge for the relief of inflammation of the chest.—**MATTHIEU.**

OINTMENTS

BLISTERING OINTMENTS.

1. *For Setons.* Resin cerate 1 oz., cantharides finely powdered 3 drs., oil of turpentine 2 drs.—**CLATER.**

2. Lard 12 oz., resin 4 oz.; melt together, and when sufficiently cool add oil of turpentine 4 oz., powdered cantharides 5 oz., stir till cold: to be rubbed in after removing the hair.—**CLATER.**

OINTMENT FOR MANGE, LICE, &c. (See also LOTIONS, further back.)

1. Sulphur 1 lb., common turpentine 4 oz., mercurial ointment 2 oz., linseed oil a pint. Melt the turpentine with the oil, and when nearly cold stir in the sulphur, and afterwards the mercurial ointment.—**YOUATT.**

Note.—Cattle are easily salivated, and greatly weakened thereby. Mercurials should therefore be used with great caution.

2. Sulphur 1 lb., strong mercurial ointment 2 oz., common turpentine $\frac{1}{2}$ lb., lard 1 $\frac{1}{2}$ lbs.—**CLATER.**

3. *French Liniment.* Olive oil a pint, sulphur 4 oz., heat till the oil becomes coloured by the sulphur; remove from the fire, and when nearly cold add 4 oz. of oil of turpentine; apply with a feather.

4. Lard 2 lbs.; melt and add oil of turpentine 8 oz., sulphuric acid 2 oz., sulphur vivum 8 oz.; stir till cold.

5. *Mange Liniment.* Sulphur vivum finely powdered 4 oz., train oil 12 oz., oil of turpentine 4 oz.; mix.—

WHITE.

IODINE OINTMENT, for Empyema. Rub together $1\frac{1}{2}$ drs. of iodine, and 1 dr. iodide of potassium, with a few drops of water, then add 3 oz. of strong mercurial ointment and $\frac{1}{2}$ oz. of powdered camphor. To be rubbed over the chest every night till it causes an exudation, then occasionally, to keep it up.

GARGET OINTMENT, for Downfall of the Udder.

1. Soft soap 1 lb., mercurial ointment 2 oz., camphor (powdered with spirit) 1 oz.; mix: give first a laxative, then a fever or diuretic drink.

2. Green elder ointment 2 oz., water of ammonia $\frac{1}{2}$ oz.

3. Beat fox-glove leaves with twice their weight of whey butter; to every pound add 1 oz. of sal ammoniac, 1 oz. of turpentine, and $\frac{1}{2}$ oz. of bole; mix and apply 2 or 3 times a day.—**DOWNING**.

4. Spirit of camphor 1 oz., mercurial ointment 1 oz., elder ointment 8 oz.—**YOUATT**.

5. *In obstinate cases.* Iodide of potassium 1 part, lard 7 parts. To be rubbed in once daily.—**SPOONER**.

6. Liquid ammonia 1 oz., olive oil 2 oz., iodine 1 dr.—**FINLAY DUN**.

. Powerful stimulants and blisters are contra-indicated.

FOOT OINTMENT (for all domestic animals). Equal parts of tar, lard, and resin, melted together.

OINTMENT FOR ULCERS ABOUT THE JOINTS.

Equal parts of basilicon and citrine ointments.—**CLATER**.

HEALING AND CLEANSING OINTMENT. Lard 2 lbs., yellow resin $\frac{1}{2}$ lb.; melt together, and when it begins to cool, add calamine in powder $\frac{1}{2}$ lb.

APPLICATION TO WOUNDS. Mix the whites of eggs with flour to a proper consistence. Applied over the part, it soon dries, and shields it from the air.

DIGESTIVE OINTMENT. 1. Lard, common turpentine, of each 4 oz.; melt, and add 1 oz. powdered verdigris.—**WHITE**.

2. Boil leaves of black hellebore with an equal weight of lard, until the leaves are crisp; strain, and add an equal weight of common turpentine. A similar ointment made with ivy leaves is likewise very stimulating.

OINTMENT FOR FOUL IN THE FOOT, OR LOW.

1. Melt 4 oz. of lard with 4 oz. of common turpentine, and add 1 oz. of finely powdered sulphate of copper, stirring until cold.—WHITE.

2. Melt together equal weights of soft soap and common turpentine.—SKERRET.

OINTMENT FOR CANCEROUS TUMOURS. Iodide of potassium $\frac{1}{2}$ oz., hot water $\frac{1}{2}$ oz.; dissolve, and mix with 2 oz. of lard.

CHARGE FOR OLD STRAINS. Burgundy pitch 4 oz., common pitch 4 oz., wax 2 oz., tar 6 oz.; apply hot, and cover with cut tow.

SETONS.

1. *Common.* A piece of cord or coarse tape; or horse-hair and tow, platted together.

2. *Irritating.* Root of common dock; or of black hellebore.

3. Cotton cord soaked in Morton's cantharidal solution (see LINIMENTS AND EMBROCATIONS FOR HORSES. *Veterinary Formulary*).

PASTE FOR STOPPING BLEEDING. Equal quantities of white, green, and blue vitriol, flour, and bole; beaten up with fresh nettles and a little vinegar.

ASTRINGENT POWDER FOR SORE FEET, &c. Sulphate of copper $\frac{1}{2}$ oz., prepared chalk 2 oz., powdered alum $\frac{1}{2}$ oz., bole 1 oz.; rub together.—SPOONER.

MEDICINES

FOR

SHEEP AND LAMBS

THESE are best given in a liquid form, and should be carefully and slowly administered. Sheep generally require one sixth (or from one eighth to one sixth) of the doses given to cattle.

PURGING DRENCHES.

1. Epsom salts 2 oz., powdered caraway $\frac{1}{2}$ oz., warm thin gruel sufficient to dissolve the salts. The editor of CLATER says that this is the best purging drink that can be used. For *Lambs* give a fourth of this, and repeat in 6 hours if necessary.

2. Epsom salts $1\frac{1}{2}$ oz. or 2 oz., ginger 1 dr., treacle 1 oz., hot water 4 oz.

3. Castor oil 2 oz., ginger and salt of tartar, of each 2 scruples, moist sugar a spoonful, gruel q. s.—M'EWEN.

4. Epsom or Glauber's salts from 1 to 2 oz., common salt a teaspoonful, boiling water sufficient to dissolve the salts, and a little gruel. A teaspoonful of tincture of ginger or of pimento, or of anodyne carminative tincture (see MISCELLANEOUS LIQUID MEDICINES, *Veterinary Formulary*, further back), may be added.

5. Sulphur $\frac{1}{2}$ oz., Epsom salts 1 oz., common salt a teaspoonful, thin gruel $\frac{1}{2}$ pint.

6. Linseed oil 2 or 3 oz., croton oil 2 or 3 drops, warm gruel q. s.

7. For *Lambs*. Epsom salts 2 to 4 drs., ginger $\frac{1}{2}$ dr., in gruel.—SPOONER.

8. For *Sheep on the first attack of Smallpox*. Epsom salts 2 oz., ginger $\frac{1}{2}$ dr., in chamomile tea or infusion of gentian (or with 1 dr. of powdered gentian or chamomile).

—WARNECKE.

FEVER DRENCH. Powdered digitalis 20 grs., emetic tartar 10 grs., nitre 2 drs. Twice a day, mixed with gruel.—**CLATER.**

TONIC DRENCHES.

1. *General Tonic.* Gentian 2 drs., calumba 1 dr., ginger $\frac{1}{2}$ dr., all in powder; tincture of orange peel 1 dr., gruel $\frac{1}{4}$ oz.; for one dose.—**CLATER.**

2. *In the last stage of Fever.* Gentian 1 dr., ginger 20 grs., spirit of nitrous ether 1 dr., tincture of cardamom 20 drops, in gruel.

3. *For Debility and Indigestion, after a purgative.* Gentian, caraway, each 1 oz., calumba and ginger, of each $\frac{1}{2}$ oz. (all sliced or bruised), boiling water a quart; infuse till cool, and strain. Give a tablespoonful daily, with the same quantity of gruel.—**CLATER.**

DRENCHES FOR RED-WATER. The pasture should be changed for shorter, the animal bled, and the bowels kept open with the above purging drinks. If these means do not remove the disease, give one of the following drenches:—

1. Epsom salts 6 oz., nitre 2 oz., bole $\frac{1}{2}$ oz., hot water 3 pints, oil of turpentine 4 oz.; mix, and give 3 or 4 tablespoonfuls (from a horn that will measure that quantity), shaking the bottle well before each dose is poured out.

2. Powdered catechu 30 grs., alum 20 grs., ginger 20 grs., decoction of oak-bark 4 oz., for a dose.

3. Olive oil 1 oz., oil of turpentine 1 oz., thick gruel $\frac{1}{2}$ pint.

* * Mr FINLAY DUN condemns bloodletting in Red-water.

FOR EXTERNAL RED-WATER. (Vesicles on the skin, containing a reddish fluid.)

1. Sulphur 2 to 3 drs., in gruel, once or twice a day. If it continue, give—

2. Epsom salts 1 oz., gruel sufficient to dissolve it.—**Sir JAMES MACKENZIE.**

DRENCHES FOR DIARRHŒA, SCOUR, &c., IN LAMBS.

[The Purging Drink, No. 6, or a fourth of No. 1, should be given before the Astringent Drinks.]

1. Prepared chalk 2 oz., catechu 1 oz., ginger $\frac{1}{2}$ oz.,

opium 1 dr., peppermint water a pint. Dose, for lambs, a tablespoonful night and morning.—YOUATT.

2. Prepared chalk $\frac{1}{4}$ oz., ginger $\frac{1}{2}$ dr., catechu $\frac{1}{2}$ dr., opium 2 grs., in gruel; once or twice daily.

3. A tablespoonful of Calves' Cordial (see MEDICINES FOR CALVES, further back).

4. Compound powder of chalk with opium 2 drs., gentian 1 dr., essence of peppermint 3 drops; in a little thin starch morning and night.—CLATER.

5. Ginger 2 drs., caraway 4 drs., prepared chalk 4 drs.; mix: give a teaspoonful in gruel.—WHITE.

6. *In white skit.* A teaspoonful of WHITE'S Alkaline Solution (p. 124) in a little gruel; and afterwards No. 7.

7. Epsom salts 3 drs., common salt a scruple, powdered ginger a scruple, thin gruel 4 oz. Repeat if necessary.

DRENCH FOR THE LATE EPIDEMIC. Epsom salts 1 oz., sulphur 2 drs., nitre $\frac{1}{2}$ dr., ginger 15 grs., in warm water. Repeat half this, with a teaspoonful of sweet spirit of nitre, daily.—SPOONER.

DRENCH FOR COW-POX. Mix 3 parts of flowers of sulphur, 1 of common salt, and 1 of honey, into an electuary; give half of this daily, in gruel. Keep the mouth and nose clean with vinegar and water. See also PURGING DRENCH, No. 7, above.

DRENCH FOR INFLUENZA. Epsom salts $\frac{1}{2}$ oz., chamomile tea 4 oz. Afterwards give half doses of the FEVER DRENCH, further back.—DARBY.

DRENCHES FOR BLOWN OR BLAST.

1. Glauber's salts 1 oz., hot water 1 oz., peppermint-water 4 oz., tincture of ginger 1 dr., tincture of gentian 2 drs.; every six hours till the bowels are opened, and half the quantity the next 4 mornings.—CLATER.

2. Common salt 1 oz., solution of potash (WHITE'S) $1\frac{1}{2}$ drs., castor or olive oil 2 tablespoonfuls, water 3 oz. After letting out the air by a tube or probang.—WHITE.

DRINK TO PREVENT RESP OR MEADOW-SICKNESS. Pearlash $1\frac{1}{2}$ drs., hot water 8 oz. To be given from a flat bottle the second and fourth morning after putting them to keep.—HOLDITCH.

DRENCH FOR STURDY AND APOPLEXY. After bleeding, 2 oz. of Epsom and Glauber's salts, in warm water or thin gruel.

DRENCHES FOR FLUX, OR SCOURING, OR DYSENTERY.

1. Epsom salts 1 oz., hot water or thin gruel to dissolve it; add castor oil 2 oz., laudanum 30 drops. When it has operated, give No. 2.—BLAINE.

2. Ipecacuanha 15 grs., prepared chalk 1 dr., opium 2 grs.; boiled starch or arrowroot 4 oz. Night and morning.

3. Linseed oil 2 oz., powdered opium 2 grs., linseed tea q. s. Afterwards give No. 4.—SAYER.

4. Opium 2 grs., ginger $\frac{1}{2}$ dr., gentian $\frac{1}{2}$ dr., linseed tea gruel q. s.

5. Epsom salts $1\frac{1}{2}$ oz., hot water 4 oz.; dissolve and add castor or olive oil $1\frac{1}{2}$ oz.—WHITE. Afterwards give No. 6.

6. Catechu $\frac{1}{2}$ dr., allspice $\frac{1}{2}$ dr., caraway 1 dr., water or beer 4 oz.; simmer together.

DRENCHES FOR DRY BRAXY, OR INFLAMMATION OF THE BOWELS.

1. After bleeding: Epsom salts $1\frac{1}{2}$ oz., warm water a pint.—STEVENSON.

2. After bleeding: common salt 1 oz., water $\frac{1}{2}$ pint, laudanum a teaspoonful.—WHITE.

3. Laudanum 2 drs., castor oil 3 oz., calomel 12 grs., treacle 3 oz.—FINLAY DUN.

DRENCH FOR BITES OF VENOMOUS REPTILES.

Olive oil 4 oz., spirit of hartshorn $\frac{1}{2}$ oz., gruel or arrowroot $\frac{1}{2}$ pint.—WHITE.

DRENCHES TO PROMOTE PARTURITION IN THE EWE.

1. A decoction of horsemint, or any other kind of mint.

2. Bruised ergot of rye 1 dr., boiling water a pint; infuse for a quarter of an hour, and give a third part. Repeat if necessary.

DRENCHES AND POWDERS FOR THE ROT.

1. Juniper berries 6 oz., gentian 1 oz.; boil in 3 gallons of water for a quarter of an hour, strain, and add

common salt 4 lbs., powdered ginger 4 oz., tartarized iron 2 oz.; stir, and let it stand till cool. Put it into wine-bottles filled two thirds full, and add to each $1\frac{1}{2}$ oz. oil of turpentine and $\frac{1}{2}$ oz. sweet spirit of nitre. Give a tablespoonful night and morning, shaking the bottle before pouring it out.

2. Common salt 8 oz., gentian powder 8 oz., ginger 1 oz., tincture of calumba 4 oz., water to make up a quart.—CLATER. See the next.

3. To a quart of No. 2, add spirit of turpentine 3 oz.; shake well together, and give 2 tablespoonfuls at night, before the night's food is given, and a tablespoonful of No. 2 every morning.

4. Epsom salts 2 oz., oil of turpentine 1 oz., ginger $\frac{1}{2}$ oz. Give in treacle and water.—FINLAY DUN.

Powders for the same.—A French recipe. 1. Dry bran 10 lbs., salt $\frac{1}{2}$ lb., aromatic herbs (as thyme, sage, juniper, rosemary, &c.) cut small, 6 oz., green anise and coriander, of each 5 oz.; mix, and give morning and night every third day. The above quantity is for 30 sheep.

2. Juniper berries 4 oz., bay-berries 1 oz., grains of paradise $\frac{1}{2}$ oz., bay salt $1\frac{1}{2}$ lbs., loaf sugar $\frac{1}{2}$ lb.; powder all together, and keep the powder in a bottle for use. Give the sheep dry and sweet hay, sprinkled with the powder.—LAWRENCE.

DRENCHES FOR INFLAMMATION OF THE LUNGS, CATARRH, HOARSE, AND COUGH. After bleeding from the neck, give Epsom salts 2 oz., gruel or linseed tea q. s.

DRENCH FOR INFLUENZA. Epsom salts $\frac{1}{2}$ oz., chamomile tea 4 oz. Afterwards, small doses of digitalis, opium, tartarized antimony, and vegetable tonics.—DARBY.

External Applications for Sheep

EYE WATERS.

1. *Strong.* For cloudiness of the eye: corrosive sublimate 4 grs., spirit of wine $\frac{1}{2}$ oz.; dissolve, and add water a pint.—CLATER.

2. Tincture or wine of opium a teaspoonful, water $\frac{1}{2}$ pint.

WASHES FOR THE SCAB, LICE, AND TICKS. (The scab ointments will also destroy them, and are less hazardous, and less injurious to the wool.)

1. *Arsenical wash.* White arsenic $\frac{1}{2}$ lb., salt of tartar $\frac{1}{2}$ lb., water 12 gallons; boil for half an hour.—YOUATT.

2. Arsenic 2 lbs., soft soap 4 lbs., water 30 gallons; dissolve. The sheep to be immersed in this liquid (the head only being kept out), and while in it, the fleece to be well rubbed. When taken out, the fluid should be well pressed out of the fleece, and the sheep kept from cold and wet for a few days.—CLATER. Mr SPOONER says 2 lbs. of arsenic should make 48 gallons of the liquid.

3. Arsenic 1 lb., yellow soap 6 lbs., pearlash 12 oz., water 30 gallons.—MATTHEWS.

4. *Mercurial.* Corrosive sublimate 1 oz., spirit of wine 2 oz.; rub together till dissolved, then add cream of tartar 1 oz., bay salt 4 oz., dissolve the whole in 2 quarts of water, and apply it with a sponge wherever lice appear.—CLATER.

5. Tobacco 4 oz., water 1 gallon; boil, and add soft soap 1 lb., sulphur vivum 1 lb.; when cold add a pint of oil of turpentine.

6. Equal parts of decoction of tobacco and lime water.—YOUATT.

* * * "On no account should sheep be returned to their grazings until they are dry, and all risk of dripping over."—FINLAY DUN.

WASH TO KILL MAGGOTS. Shake up in a bottle together, 1 quart of water, spirits of turpentine 1 oz., and corrosive sublimate 10 grs. Stop with a cork in which quill is inserted. When the maggots are observed, a

small quantity of the mixture is to be shaken on the spot through the quill, and the maggots will shortly creep out and die.

SMEARING MIXTURE.

1. One gallon of common tar, and 12 lbs. of any sweet grease, melted together.

2. Oil of tar is used as a preventive of the *fly*; but fish oil is equally serviceable according to Mr HOGG, and is less injurious to the wool. Oil of tar has sometimes destroyed sheep.

FLY POWDER FOR SHEEP.

1. White lead 2 lbs., red lead $\frac{1}{2}$ lb., sulphur $1\frac{1}{2}$ lbs., oil of wormwood, animal oil (empyreumatic), or creasote $\frac{1}{4}$ oz.; mix.

2. White lead 2 lbs., red lead 1 lb.; mix, and apply by sprinkling from a dredger, following a stick drawn through the wool.—CLATER.

3. Powdered colocynth 3 drs., black brimstone 1 lb., tincture of assafoetida $\frac{1}{2}$ oz.; mix.

4. White lead 4 parts, arsenic 1 part, sulphur 6 parts, vermilion 2 parts.—SPOONER.

POWDER FOR THE EYES. Equal parts of sal ammoniac, white sugar, and oxide of zinc, triturated together. It may be mixed either with rose-water or honey.—SPOONER.

ASTRINGENT POWDER FOR THE FEET, in the epidemic affecting them. The same as for cattle. See further back.

OINTMENT FOR THE SCAB OR SHAB.

1. Quicksilver 1 lb., Venice turpentine $\frac{1}{2}$ lb.; rub them together until the globules are no longer visible; then add $\frac{1}{2}$ pint of oil of turpentine, and 4 lbs. of lard. The mode of applying this ointment is as follows.—Begin at the head of the sheep, and proceeding from between the ears along the back to the end of the tail, divide the wool in a furrow till the skin can be touched; and let a finger slightly dipped in the ointment be drawn along the bottom of the furrow. From this furrow similar ones must be drawn along the shoulders and thighs to the legs, as far as

the wool extends. And if much infected, two or more should also be drawn along each side, parallel with that on the back; and one down each side before the hind and fore legs. It kills the sheep-fag, and probably the tick and other vermin. It should not be used in very cold or wet weather.—SIR JOSEPH BANKS.

2. Tar oils. Tobacco juice. Stavesacre.—FINLAY DUN.

3. Strong mercurial ointment 1 part, lard 5 parts; mix.—YOUATT.

4. Quicksilver 1 lb., Venice turpentine $\frac{1}{2}$ lb., spirit of turpentine 2 oz., lard $4\frac{1}{2}$ lbs.; to be made and used as No. 1. In summer 1 lb. of resin may be substituted for a like quantity of lard.—CLATER.

5. Strong mercurial ointment 1 lb., lard 4 lbs., oil of turpentine 8 oz., sulphur 12 oz.—WHITE.

6. *Mild.* Flowers of sulphur 1 lb., Venice turpentine $\frac{1}{2}$ lb., rancid lard 2 lbs., strong mercurial ointment 4 oz.; mix well.—CLATER.

7. Lard or other fat, with an equal quantity of oil of turpentine.—DAUBENTON.

8. *Without Mercury.* Lard 1 lb., oil of turpentine $\frac{1}{2}$ lb., flowers of sulphur 6 oz.—WHITE.

9. Strong mercurial ointment 1 lb., lard 4 lbs., Venice turpentine 8 oz., oil of turpentine 2 oz. If mixed by heat, care must be taken to use no more heat than is necessary, and to add the oil of turpentine when the other ingredients begin to cool, and to stir till cold.—M'EWEN.

10. Corrosive sublimate 2 oz., white hellebore 3 oz., fish oil 6 quarts, resin $\frac{1}{2}$ lb., tallow $\frac{1}{2}$ lb. The sublimate and then the hellebore to be rubbed with a portion of the oil till perfectly smooth, and then mixed with the other ingredients melted together.—STEVENSON.

11. The following once had considerable local celebrity, but it obviously requires to be used with caution. Dissolve $2\frac{1}{2}$ oz. of corrosive sublimate in the same quantity of hydrochloric acid, and beat up the solution with 6 lbs. of strong mercurial ointment; put it in a large pan, and pour on it $19\frac{1}{2}$ lbs. of lard, and $1\frac{1}{2}$ lbs. of common turpentine, melted together and still hot, and stir the whole continually until it becomes solid.

OINTMENT FOR DEEP WOUNDS OR ULCERS FROM FLIES. The Fly Powder, No. 2, mixed with tar.—CLATER.

OINTMENT FOR SORE HEADS. Black pitch 2 lbs., tar 1 lb., flowers of sulphur 1 lb.; melt together, taking care that it does not boil. To be spread thickly on leather while warm, and fitted to the head.

CAUSTIC ASTRINGENTS FOR FOOT-ROT.

1. Blue vitriol 1 oz., white vitriol 1 oz., burnt alum 2 oz., bole $\frac{1}{2}$ oz., honey to form a stiff paste.—M'EWEN.

2. Sulphate of copper 2 oz., water 12 oz., dilute sulphuric acid 2 drs.—WHITE.

3. Butter of antimony, alone, or mixed with tincture of myrrh.

4. Verdigris, bole, and sugar of lead, in equal parts, rubbed together into a fine powder. Sprinkle on the sore, cover with tow, and bind down with tape for 24 hours, using afterwards No. 2, or No. 3.—CLATER.

5. *Strong.* Verdigris 1 oz., nitrous acid 2 oz., water 4 oz.—WHITE.

6. *Strongest.* Red precipitate 1 oz., nitrous acid 2 oz.; dissolve, and add water 2 oz., spirit of wine 1 oz.—WHITE.

7. Aloes 16 oz., weak spirit 32 oz., sulphuric acid 17 oz.; mix.—DUVILLE.

8. Dissolve sulphate of copper $2\frac{1}{2}$ oz. in $1\frac{1}{2}$ pint of water, and add a solution of $3\frac{1}{2}$ drs. of sulphate of iron previously calcined. Diffuse $\frac{3}{4}$ oz. of slaked lime in water, and add to the mixed solutions; then add 7 oz. common salt, 1 oz. wood vinegar, and water to make up a quart. [Nearly the composition of a celebrated French nostrum.]

9. *LELOUP'S Terebinthinated Oxymel of Copper.* Honey 14 oz., pyroligneous acid 7 oz., powdered verdigris 5 oz.; boil it in a large copper pan until it assumes a reddish purple colour; then add (keeping the mixture on a slow fire) 14 oz. Venice turpentine; stir with a wooden spatula for a quarter of an hour, and pour it into jars. To be applied twice, at 12 hours' interval, by means of a small piece of wood, after cleaning the part with an iron blade.

10. White vinegar 78 parts, powdered sulphate of copper 10 parts; dissolve, and add 12 parts of sulphuric acid. Apply it with a feather. (A French remedy.)

11. *Detersive Ointment.* Burnt alum 4 parts, verdigris 1 part, camphor 1 part, green ointment of elder or poplar 16 parts.—LEBAS.

12. Honey 4 oz., burnt alum 2 oz., Armenian bole $\frac{1}{2}$ lb.; mix with as much train oil as will convert these ingredients into a salve. The honey must first be completely dissolved in the oil made hot, then the bole stirred in, and lastly the alum.

13. Carbolic acid 1 fl. dr., water 6 to 12 oz.

BLACKLOCK condemns all caustic applications, using only mild poultices and emollient ointments.

MEDICINES FOR SWINE

ALTERATIVE MEDICINES, given in mange and other skin diseases, and in obstinate costiveness.

1. Sulphur $\frac{1}{4}$ oz., Æthiops mineral 3 grs., nitre $\frac{1}{2}$ dr.; cream of tartar $\frac{1}{2}$ dr.; daily, in thick gruel or wash.—**CLATER**.

2. Black antimony $\frac{1}{2}$ oz., sulphur 2 oz., nitre $\frac{1}{2}$ oz.; mix; for 8 doses.

FEVER MEDICINE. Digitalis 3 grs., antimonial powder 6 grs., nitre $\frac{1}{2}$ dr.; after bleeding, in a little warm swill, milk, or wash, morning, noon, and night.—**CLATER**.

PURGING MEDICINES.

1. Epsom salts 1, 2, or 3 oz., in broth or swill.

2. Sulphur 2 drs., daily; full dose $\frac{1}{2}$ oz., with milk or other food. This may be repeated for 2 or 3 days, in surfeit from overfeeding.

3. Jalap 1 dr.; if insufficient, add 10 or 12 grs. of scammony, or 10 grs. of calomel.—**WHITE**.

4. Jalap $\frac{1}{2}$ dr., sulphur 2 drs., antimony $\frac{1}{2}$ dr.

5. Jalap $\frac{1}{2}$ dr., Epsom salts 1 oz.

6. Castor oil 1 to 2 oz., with gruel.

7. Castor oil 1 oz., gruel q. s., Epsom salts 2 oz., salt $\frac{1}{2}$ oz.; mix.

8. Calomel 5 grs. This must not be repeated more than twice.

CARMINATIVE DRENCH, for flatulent distension, from sour whey, &c. After using the probang, or where it cannot be had, give—

1. **WHITE**'s solution of potash (see p. 124) 2 oz., anodyne carminative tincture 1 tablespoonful, water 8 oz.

2. A tablespoonful of common salt in warm water, a teaspoonful of mustard or powdered ginger, and a glass of gin.

THRIVING POWDER, to promote fattening. Powdered fenugreek, alone, or mixed with a fourth of liquorice powder; an ounce daily with the food. Cleanliness greatly conduces to the same end.

REMEDIES FOR MEASLES. After bleeding by tail, ear, palate, or vein inside the forearm, an inch above the knee, give one of the purging drinks, and turn the animal into the open air.

REMEDIES FOR THE DISTEMPER, affecting the Mouth and Feet. The same drink, and astringent powder as for SHEEP.

DRENCH FOR INFLAMMATION OF THE BRAIN. Castor oil 2 oz., with gruel; afterwards 2 grs. white hellebore powder twice or thrice a day.—**CUPISS.**

HEALING OINTMENT FOR SORE EARS.

1. Lard 1 lb., resin 4 lbs.; melt together, and stir in $\frac{1}{2}$ lb. lapis calaminaris.—**CLATER.**
2. Zinc ointment 1 oz., yellow basilicon 3 oz.
3. Tar ointment mixed with a little soap.

MANGE OINTMENT.

Sulphur 4 oz., Venice turpentine 1 oz., old lard 8 oz., mercurial ointment 1 oz.; the animal to be previously scrubbed all over with a good soap lather. The alterative powders (see further back) should be given at the same time.

OINTMENT FOR SORE TEATS. Soft soap 4 oz., camphor (powdered with spirit) $\frac{1}{2}$ oz., mercurial ointment $\frac{1}{2}$ oz. It must be carefully washed off.

MEDICINES FOR DOGS

N.B.—The doses required vary considerably, according to the strength and size of the dog, which should always be duly considered.

PHYSIC BALLS AND OTHER PURGATIVE MEDICINES.

1. Barbadoes aloes 8 oz., antimonial powder 1 oz., ginger 1 oz., palm oil 5 oz.; beat together into a mass. Dose from $\frac{1}{2}$ dr. to 2 drs., every 4 or 6 hours, till the bowels are relieved.—YOUATT.

2. The same, with the addition of 1 oz. of calomel. From 45 grs. to 2 drs. for a dose.—CLATER.

3. Aloes $\frac{1}{2}$ dr. to 2 drs., made into a ball with syrup of ginger.

4. *In inflammation of the bowels, and in worms.* Aloes $\frac{1}{2}$ dr. to 1 $\frac{1}{2}$ drs., calomel 2 to 5 grs., syrup to form a ball.—BLAINE.

5. Cape aloes $\frac{1}{2}$ dr. to 1 dr., calomel 2 to 3 grs., oil of caraway 6 drops, syrup to form a ball.—M'EWEN.

6. Calomel 12 grs., aloes 3 drs., opium 1 gr., syrup q. s. to form a mass, for 4, 6, or 8 balls; one every 4 or 5 hours till the bowels are relieved.—BLAINE.

7. Croton oil 1 drop, Castile soap 20 grs., conserve to form a ball.

8. Castor oil 3 parts, syrup of buckthorn 2 parts, syrup of poppies 1 part; dose, from 1 to 2 tablespoonfuls.—Mr YOUATT'S purge. Mr CLARK says syrup of buckthorn for dogs should be made with treacle, and the spices omitted.

9. Epsom salts, from 1 to 4 drs., wrapped in tissue paper, dividing the doses into convenient-sized packets.

10. *In costiveness with inflammation.* $\frac{1}{2}$ oz. to 2 oz. castor oil.—Mr SPOONER.

11. Jalap, powdered, 30 grs., calomel 8 grs.; make into a pill with gum water, and administer every morning. In distemper.

ALTERATIVE BALLS AND POWDERS.

1. Sulphur 2½ lbs., nitre ½ lb., Æthiops mineral 4 oz., linseed meal ½ lb., palm oil 1 lb., or as much as may be required; beat together, and keep in a jar for use. Dose, from 2 scruples to 1½ or 2 drs.—CLATER.

2. Æthiops mineral 20 to 40 grs., cream of tartar 20 to 40 grs., nitre 5 to 10 grs.; night and morning, made into a ball with butter.—SPOONER.

3. *Tonic Alterative.* Mercurial pill 1 dr., aloes 2 drs., myrrh, benzoin, balsam of Peru, of each 1½ drs.; to be divided into 10, 15, or 20 pills; one every evening, for the yellows, after aloes and calomel.—BLAINE.

4. *Alterative Powder.* Æthiops mineral 2 to 5 grs., cream of tartar 4 to 10 grs., tartarized iron 1 to 3 grs., once a day.—CLATER.

5. *To give a fine skin.* Give a tablespoonful of tar, made up with oatmeal.—MAYER.

ASTRINGENT BALLS, &c.

1. Catechu 1½ drs., sulphate of quinine 20 grs., opium 5 grs., ginger 1 dr., conserve of roses q. s. to form a mass; to be divided into 4, 6, or 8 balls.—BLAINE.

2. *In the advanced stage of distemper.* Prepared chalk 2 oz., powdered gum arabic ½ oz., powdered catechu ½ oz., powdered oak bark ½ oz., powdered ginger ½ oz., opium 15 grs., palm oil 1 oz.; beat well together. Dose, ½ dr. to 2 drs., morning, noon, and night.—CLATER.

3. *In diarrhœa.* Opium 5 grs., catechu 2 drs., gum arabic 2 drs., ginger ½ dr., syrup of poppies q. s.; divide into 6, 9, or 12 balls.—BLAINE.

4. Myrrh 1 dr., ipecacuanha 1 scruple, opium 3 grs., chalk 2 drs., carbonate of iron 1 dr.; as No. 3.—BLAINE.

5. *In obstinate cases.* Alum 1 dr., chalk 2 drs., opium 6 grs., resin 3 drs.; into 4, 6, or 8 balls.

6. *In diarrhœa,* after 1 to 4 drs. of Epsom salts. Prepared chalk 1 to 3 scruples, catechu 5 to 10 grs., opium ¼ to 2 grs.; twice a day.—SPOONER.

7. *Astringent Drink.* Boil 1 oz. of logwood in a quart of milk to ½ a pint. A teaspoonful every morning, in prolapsus.

COUGH BALLS IN ASTHMA, &c.

1. *After a few Emetics.* Calomel 3 grs., foxglove 3 grs.,

cream of tartar 1 dr., antimonial powder 12 grs., honey to form 6 boluses. One twice a day.—BLAINE.

2. Digitalis 20 grs., antimonial powder 40 grs., nitre 2 drs., sulphur 3 drs., palm oil 3 drs., or q. s. Divide into 10, 15, or 20 balls, according to the size of the dog; give one morning and night, interposing an emetic every third or fourth day.—CLATER.

3. *In old cases.* P. squill $\frac{1}{2}$ gr. to 1 gr., gum ammoniacum 5 grs., balsam of Peru 8 grs., benzoic acid 1 gr., balsam of sulphur to form a ball.

4. Extract of hemlock $\frac{1}{2}$ dr., extract of henbane 10 grs., p. digitalis 20 grs., conserve of roses to form a mass. Divide into 6, 8, or 10 balls. One night and morning.—BLAINE.

DISTEMPER MEDICINES.

1. *Turpeth's mineral 1 to 3 grs., assafœtida $\frac{1}{2}$ dr., aloes 20 grs., soap 10 grs., syrup of poppies to form a ball. To be preceded by an emetic, and given every third day.

2. After bleeding (if required) and an emetic, give a physic ball; and afterwards the following:—2 or 3 times a day: Antimonial powder 2, 3, or 4 grs.; nitre 5, 10, or 15 grs.; ipecacuanha 2, 3, or 4 grs.; form a ball. If the disease proceed to the debilitating stage, give the *Tonic Ball* No. 2; in the putrid or malignant stage give the *Astringent Ball* No. 1.—BLAINE.

3. After the Emetic Powder No. 1 (which should be repeated every 3rd or 4th day) give the *Cough Ball* No. 2, from $\frac{1}{2}$ dr. to 2 drs. in weight. And if the dog lose flesh, give equal parts of the *Cough Ball* and the *Tonic Ball* (No. 1). In the more advanced stages give the tonic alone; or the *Astringent Ball* if diarrhœa comes on.—CLATER.

4. Give a third of a paper of James's powder, mixed with butter, and afterwards warm broth or milk. In 2 hours, another third; and if this neither vomit nor purge, give the other third at the end of 4 hours.—DANIEL.

5. BLAINE'S *Distemper Powders*, which are sold in packets, with directions for use.

6. *In the malignant stage, with diarrhœa.* Camphor 3 to 5 grs., charcoal 10 grs., opium 1 gr., aromatic confection q. s. to form a ball.

7. Antimonial powder 2 to 4 grs., nitre 5 to 10 grs.,

* Discarded by the modern veterinary practitioner.

digitalis $\frac{1}{4}$ to 2 grs. Afterwards the Tonic Pills No. 4.—
SPOONER.

Poudre Kusique: a French nostrum. Mix 45 grs. of nitre, 45 of sulphur, and 1 charcoal. Divide into 3 doses. Give 1 for 2 successive mornings, and the third on the 4th morning, mixed with lard or butter, or in milk. For a large dog a second packet (of 3 powders) may be required.—HABERT.

Another French nostrum, Hemel's Powder, is of a similar kind.

8. A strong solution of salt, to the amount of $\frac{1}{2}$ pint daily.

9. Powdered tin, sulphur, gunpowder, of each 1 oz.; lard sufficient to form a mass. The size of a nutmeg to be given twice or thrice a week.

10. Physic Ball No. 11.

11. $\frac{1}{4}$ oz. to 1 fl. oz. of cod-liver oil, twice a day, according to size.

12. Emetics, gentle laxatives, milk diet, and from 5 to 15 grs. of chlorate of potash, twice a day.—FINLAY DUN.

REMEDIES FOR SPASMODIC COLIC.

1. Castor oil $\frac{1}{2}$ oz., oil of peppermint 1 drop, laudanum 20 drops. If it does not open the bowels, give $\frac{1}{2}$ dr. to $1\frac{1}{2}$ drs. of aloes.—BLAINE.

2. Castor oil 3 oz., syrup of buckthorn 2 oz., syrup of poppies 1 oz. Give from a teaspoonful to a tablespoonful.—YOUATT.

3. Ether $\frac{1}{2}$ dr., laudanum $\frac{1}{2}$ dr., camphor 3 to 6 grs., castor oil (unless purged) 3 to 5 drs.—BLAINE.

CONVULSIONS.

Give Colic Mixture No. 3, and apply warm bath and flannel.

FOR FITS OR EPILEPSY.

1. Calomel 8 grs., carbonate of iron $\frac{1}{2}$ dr., extract of hemlock 20 grs., conserve of roses, or palm oil, to form a mass for 6, 9, or 12 balls.—BLAINE.

2. Give the Alterative Balls No. 1, or the pills of nitrate of silver, as for St. Vitus's dance.—CLATER.

3. *For epilepsy of suckling bitches.* Ether 1 dr., laudanum $\frac{1}{2}$ dr., strong ale 2 oz.; give from a dessert-spoonful to 1 or 2 tablespoonfuls every 2 or 3 hours.—BLAINE.

4. *For epilepsy attending distemper.* The Tonic Balls, or the pills for chorea.—CLATER.

5. *After an emetic.* Gentian 10 to 20 grs., ginger 3 to 6 grs., carbonate of iron 2 to 4 grs., or from an eighth to a fourth of a gr. of nitrate of silver, and $\frac{1}{2}$ gr. of spiders' web once a day.—SPOONER.

6. Ether 1 dr., laudanum $\frac{1}{2}$ dr., camphor 6 grs., spirit of hartshorn 1 dr.; in a spoonful of ale: for small dogs give half the quantity.—BLAINE.

7. When the fit is over, give an active purgative, and 20 to 40 drops of turpentine.—FINLAY DUN.

EMETIC POWDERS.

1. Calomel, emetic tartar, of each 1 oz.; vermilion 10 grs.; rub together; dose, from 1 to 3 grs., dropped on the tongue, or mixed with a teaspoonful of milk.—CLATER.

2. Emetic tartar, from 1 to 3 grs.

3. Turpeth's mineral, from 1 to 3 grs.

4. A teaspoonful of common salt.

MEDICINES FOR INFLAMMATORY DISORDERS.

1. *In inflammation of the lungs.* After bleeding and purging, digitalis 12 grs., emetic tartar 3 grs., nitre 1 dr.; mix, and divide into 6, 9, or 12 powders.—BLAINE.

2. *Ditto, with much cough.* Tincture of digitalis 1 dr., emetic tartar 3 grs., nitre 1 dr., simple oxymel 2 oz.; dose, 1 or 2 drs. every 3 hours.—BLAINE.

3. *In pleurisy, with incipient water in the chest.* Digitalis 6 grs., calomel 6 grs., tartarized iron 18 grs.; into 6, 9, or 12 doses.—YOUATT.

4. *In inflammation of the liver.* Digitalis 8 grs., antimonial powder 16 grs., nitre 1 dr.; divide into 7, 9, or 12 powders, or boluses.—BLAINE.

5. *In chronic inflammation of the liver.* Calomel 20 grs., antimonial powder $\frac{1}{2}$ dr., myrrh, gentian, aloes, of each 2 drs.; mix, and divide into 15, 20, or 25 balls.

6. *In inflammation of the bowels.* After bleeding and a warm bath, give the castor-oil mixture (see PURGATIVES, No. 8).—CLATER.

7. *Bilious inflammation* (with offensive, often black vomiting and purging): Calomel 10 grs., opium 4 grs.: in 4 or 8 pills—one 3 times a day; afterwards the astringent remedies for diarrhœa.—SPOONER.

MEDICINES FOR RHEUMATISM.

1. After warm bath, and friction, give tincture of opium 20 drops, ether 20 drops, castor oil $\frac{1}{2}$ oz. to 1 oz.—BLAINE.

2. Calomel 2 to 4 grs., opium $\frac{1}{4}$ gr., oil of peppermint 1 drop, aloes 1 dr.; form a ball with butter or lard: repeat it every 4 hours till the bowels are well opened; and use the Embrocation No. 3.—CLATER.

3. After warm bath, &c., give 40 drops of laudanum, and a teaspoonful of hartshorn, in warm beer; and rub with the Embrocation No. 1.—MAYER.

TONIC MEDICINES.

1. Gentian 1 oz., chamomile $\frac{1}{2}$ oz., oak-bark $\frac{1}{2}$ oz., ginger $\frac{1}{4}$ oz., carbonate of iron $\frac{1}{4}$ oz., palm oil 1 oz.; beat them together to form a mass; dose, 2 to 6 scruples.—CLATER.

2. *Debilitating stage of distemper.* Sulphate of quinine $\frac{1}{2}$ dr., powdered chamomile 3 drs., balsam of Peru $1\frac{1}{2}$ drs., camphor 1 scruple; form a mass with conserve of roses, and divide into 6, 9, or 12 balls; one every 6 hours.—BLAINE.

3. *In gutta serena.* Chamomile 1 oz., rue $\frac{1}{2}$ oz., ginger $\frac{1}{4}$ oz. (all in powder); beat them into a mass with 7 drs. of palm oil, and divide into 12, 16, or 20 balls; one night and morning.—CLATER.

4. Gentian powder 10 to 20 grs., ginger 5 grs., cascarilla 10 to 20 grs.; conserve of roses, or syrup, to form a ball. One twice a day.

WORM MEDICINES.

1. Carbonate of iron $\frac{1}{2}$ oz., Æthiops mineral 1 dr., gentian 1 oz., ginger $\frac{1}{2}$ oz., levigated glass 1 oz., palm oil 9 drs.; beat well together; dose, from $\frac{3}{4}$ to 2 drs.—CLATER.

2. As much very finely-powdered glass as will lie on a sixpence, mixed with butter.—BLAINE. Mr YOUATT says from $\frac{1}{2}$ dr. to 1 dr., powdered glass, with a little ginger, made into a ball with lard.

3. Aloes, sulphur, prepared hartshorn, and juice of wormwood, made into a mass; the size of a hazel nut to be given three times a week, fasting, wrapped in butter.—DANIEL.

4. Tin filings, or pewter filings, $\frac{1}{2}$ dr. to 1 dr., with butter or lard.

5. Jalap 10 to 15 grs., calomel 2 to 3 grs. mixed with butter; no cold liquid should be allowed.—WHITE.

6. Cowage $\frac{1}{2}$ dr., iron filings 4 drs., conserve of roses q. s. to form a mass, to be divided into 4, 6, or 8 balls; one every night and morning; and afterwards the purgative No. 4.—BLAINE.

7. Epsom salts 1 oz., common salt 1 dr.; give a small or large teaspoonful daily.

8. Give green walnut leaves boiled in milk.—MAYER.

9. From $\frac{1}{2}$ dr. to 2 drs., according to size, of betel nut, in coarse powder, made into a ball; an excellent remedy.

10. *For Tape Worm.* Oil of turpentine $\frac{1}{2}$ dr., mixed with yolk of egg; for very large dogs 2 scruples. Some writers prescribe larger doses (1 or 2 drs.), but these sometimes prove fatal.—BLAINE.

11. From 2 to 6 drs. of cuseo according to size.

12. *For Tape Worm.* Oil of turpentine and olive oil, of each $\frac{1}{2}$ oz.; mix, and give carefully; 3 or 4 hours after, give 1 oz. castor oil.

13. *For Stomach Worms.* Give the emetic powder (see further back) and afterwards a physic ball.

14. *Thread Worms.* These are destroyed by an aloetic clyster.

MEDICINES FOR THE YELLOWS.

1. After bleeding—Calomel 2 to 3 grs., jalap 10 grs., scammony 4 grs.—WHITE.

2. Aloes 20 to 40 grs., calomel 2 to 4 grs.; afterwards the tonic alterative balls. See ALTERATIVE BALLS AND POWDERS.

MEDICINES FOR ST VITUS'S DANCE, OR CHOREA.

1. Nitrate of silver 8 grs., ginger 10 grs., syrup to form a mass; divide into 64 pills, and give one or two, morning and night.—CLATER.

2. Strychnia 1 gr., oxide of zinc 24 grs., assafoetida 24 grs., conserve of roses q. s.; mix very accurately, and divide into 6, 9, or 12 balls.

3. Nitrate of silver 3 grs., carbonate of iron 2 drs., gentian 3 drs., conserve of roses to form a mass, for 6, 9, or 12 balls.—BLAINE.

4. Give 2 grs. of valerianate of quinine or sulphate of zinc, twice a day.—FINLAY DUN.

MEDICINES FOR DROPSICAL COMPLAINTS.

1. Digitalis 9 grs., squill 12 grs., cream of tartar 2 drs., mix, and divide into 9, 12, or 15 powders; one night and morning.

2. Foxglove 12 grs., antimonial powder 15 grs., nitre 1 dr.; as the last.—BLAINE.

3. Foxglove 1 gr., nitre 10 grs., ginger 8 grs.; night and morning; then iodide of potassium $\frac{1}{2}$ gr. to 1 gr.—YOUATT.

BALLS FOR ENLARGED GLANDS AND CANCEROUS DISEASES.

1. Extract of hemlock 1 to 3 grs., burnt sponge 10 to 20 grs.; make a ball, to be given once or twice a day.

2. Iodine 12 grs., powdered gum 40 grs., syrup to form a stiff mass, divide into 48 pills, and give one or two, night and morning.—CLATER.

BALLS TO PROMOTE PARTURITION. Ergot of rye 20 grs.; pulverise, and add ginger 16 grs., syrup q. s.; beat into a mass, and divide into 5 pills; give one every hour, or to a small bitch, half of one.—CLATER.

TO PREVENT RABIES, OR CANINE MADNESS.

1. Powdered leaves of the *scutellaria lateriflora* 40 grs., powdered belladonna $2\frac{1}{2}$ grs.; to be given night and morning for 6 weeks, gradually increasing the dose.—YOUATT.

2. Infuse a teaspoonful and a half of powdered *scutellaria* in a quart of hot water; give half a pint morning and night, omitting the dose every third day, when a mild dose of sulphur must be given.—Dr SPALDING.

3. Fresh leaves of the tree-box 2 oz., rue 2 oz., sage $\frac{1}{2}$ oz., chop them fine, and boil them in a pint of water till reduced to half a pint; strain, and press out the liquid; beat the herbs, and boil them in a pint of new milk to half; strain, press the herbs, and mix the liquids. For a man, give a third of this quantity every other morning fasting; double the above quantity makes 3 doses for a horse or cow; two thirds will suffice for a middle-sized

dog, and a third for smaller dogs. It produces extreme nausea and distress, and has occasionally proved fatal to dogs.—BLAINE. All the above are obsolete.

External Applications

ASTRINGENT & DETERGENT LOTIONS, for Wounds, &c.

1. Bruised oak-bark 2 oz., catechu 1 oz., water 3 pints; boil to a pint, and strain.—CLATER.

2. Tincture of myrrh and aloe 1 oz., alum $\frac{1}{2}$ oz., water 1 pint.

3. *For sore feet.* The Lotion (No. 1), 4 oz., tincture of aloe $\frac{1}{2}$ oz., water 1 pint.

4. *To excite sluggish wounds.* Nitrate of silver 10 grs., water 1 oz.

LOTIONS FOR CANCER IN THE EAR. See OINTMENTS FOR CANCER IN THE EAR, further on.

1. Sulphate of zinc 20 grs., sugar of lead $\frac{1}{2}$ dr., water 4 oz.—WHITE.

2. Sulphate of zinc 20 grs., decoction of oak-bark 4 oz.—BLAINE.

3. Nitrate of silver 1 gr., rain-water 2 oz.

4. Sugar of lead $\frac{1}{2}$ dr., rose-water 4 oz. A teaspoonful to be introduced blood-warm into the ear.

5. *Mild Canker Lotion.* Infusion of foxglove leaves $\frac{1}{2}$ pint, Goulard's extract of lead $\frac{1}{2}$ oz.; mix. To be used as the last.—CLATER.

6. *Strong Canker Lotion.* Goulard's extract 2 drs., white vitriol 1 dr., alum 2 drs., water $\frac{1}{2}$ pint.

7. Chloride of lime 20 to 30 grs., water $\frac{1}{2}$ pint.

LOTION TO ALLAY ITCHING. Dilute hydrocyanic acid 1 dr., water a pint.—YOUATT.

OINTMENTS FOR CANCER OF THE EAR.

1. Equal parts of zinc ointment, and ointment of nitrate of quicksilver.—BLAINE.

2. Sublimate 3 grs., Turner's Cerate 1 dr., sulphur a scruple.

3. White vitriol, alum, each in fine powder, a drachm, lard 4 oz. To be rubbed gently into the crack.—CLATER.

4. *Stronger.* Nitrate of silver 20 grs., lard 1 oz. Rub them well together.—CLATER.

5. Levigated red precipitate $\frac{1}{2}$ oz., lard 2 oz.—MAYER.

EYE WATERS.

1. *Astringent Wash for Weak Eyes.*—White vitriol 4 grs., spirits of wine $\frac{1}{2}$ dr., water 4 oz.—CLATER.

2. Sugar of lead 30 grs., rose-water 6 oz.—BLAINE.

3. Laudanum $\frac{1}{2}$ dr., infusion of green tea 4 oz.—M'EWEN.

4. *For naturally weak eyes.* Laudanum 2 drs., water 8 oz. To be used every morning.—CLATER.

5. Sugar of lead $\frac{1}{2}$ dr., distilled water 6 oz., tincture of opium $\frac{1}{2}$ dr.—SPOONER.

6. *In inflammation after bleeding.* Physic and warm fomentations.

OINTMENTS FOR ULCERATED EYELIDS.

1. Red precipitate, levigated, 10 grs., zinc ointment $\frac{1}{2}$ oz.

2. Ointment of nitrate of quicksilver 1 dr., sugar of lead 20 grs., spermaceti ointment 3 drs.—BLAINE.

3. Dissolve a drachm of quicksilver in a drachm and half of strong nitric acid, and well mix the warm solution with 6 oz. of melted lard.—H. CLAYTER.

OINTMENTS AND LOTIONS FOR THE MANGE.

N.B.—An alterative ball should be given daily and a physic ball occasionally. Bleeding is sometimes prescribed.

1. *For scabby mange.* Sulphur 4 oz., sal ammoniac $\frac{1}{2}$ oz., aloes 1 dr., Venice turpentine $\frac{1}{2}$ oz., lard 6 oz.; mix. After four applications wash well with soap and water.—BLAINE.

2. Horse turpentine and palm oil, of each $\frac{1}{2}$ lb., train oil $\frac{1}{2}$ pint. Melt together, and while cooling, stir in 3 lbs. of flowers of sulphur.—CLATER.

3. Aloes 2 drs., hellebore $\frac{1}{2}$ oz., sulphur 4 oz., lard or train oil 6 oz.—M'EWEN.

4. Sulphate of zinc 1 dr., snuff $\frac{1}{2}$ oz., white hellebore $\frac{1}{2}$ oz., sulphur 4 oz., aloes $\frac{1}{4}$ oz., soft soap 6 oz.—BLAINE.

5. Charcoal powder 2 oz., sulphur 4 oz., salt of tartar 1 dr., Venice turpentine $\frac{1}{2}$ oz., lard 6 oz.

6. *For red mange.* Add 1 oz. of strong mercurial ointment to 6 oz. of either of the above.

7. Charcoal 1 oz., chalk 1 oz., sugar of lead 1 dr., white precipitate 2 drs., sulphur 2 oz., lard 5 oz.—BLAINE.

8. *Wash for red mange.* Corrosive sublimate 20 grs., spirit of wine 2 drs.; dissolve, and add milk of sulphur $\frac{1}{2}$ oz., lime-water $\frac{1}{2}$ pint. Apply by means of a sponge.—CLATER.

9. *For ulcerated mange.* Ointment of nitrated quicksilver 2 drs., sugar of lead 20 grs., flowers of sulphur $\frac{1}{2}$ oz., lard 1 oz.; mix.—BLAINE.

ARSENIC OINTMENT. *In mange and other skin diseases.* Yellow sulphuret (sulphide) of arsenic $1\frac{1}{2}$ grs., cerate or lard $\frac{1}{2}$ oz.—DELAFOND.

SURFEIT OINTMENT. After bleeding and purging, apply sugar of lead 1 dr., spermaceti ointment 2 oz.

OINTMENT AND POWDERS FOR PILES.

Ointment.—Sugar of lead 6 grs., tar $\frac{1}{2}$ dr., elder ointment 3 drs.—BLAINE.

Powders.—Nitre $\frac{1}{2}$ dr., milk of sulphur 3 drs.; mix, and divide into 9, 12, or 15 doses.—BLAINE.

HEALING OINTMENT.

1. Palm oil 3 lbs., resin 1 lb.; melt together, and when they begin to cool add 1 lb. of powdered calamine.—CLATER.

2. Oxide of zinc $\frac{1}{4}$ oz., lard $1\frac{1}{2}$ oz., balsam of Peru 1 dr.

OINTMENT FOR SCIRRHOUS TUMOURS. Iodide of potassium 1 dr., lard 7 drs.; rub together till perfectly smooth.

STIMULATING MERCURIAL OINTMENT. Mercurial ointment 1 oz., simple cerate 1 oz. A small quantity to be rubbed over the region of the liver once a day till the mouth is sore.—BLAINE.

EMBROCATIONS FOR PALSY, RHEUMATISM, &c.

1. Oil of turpentine 2 oz., spirit of hartshorn 2 oz., tincture of opium $\frac{1}{4}$ oz., olive oil 2 oz.—BLAINE.

2. Cajeput oil 1 oz., soap liniment 2 oz.
3. Spirit of turpentine, spirit of hartshorn, camphorated spirit, of each 1 oz., laudanum $\frac{1}{2}$ oz.—CLATER.

LOTIONS FOR STRAINS AND BRUISES.

1. Common salt and cold vinegar.
2. Sal ammoniac $\frac{1}{2}$ oz., vinegar a pint.
3. *For strains.* Oil of turpentine 1 oz., old beer $\frac{1}{2}$ pint, brine $\frac{1}{2}$ pint.—MAYER.
4. Spirit of Mindererus 4 oz., Goulard water 8 oz.

ASTRINGENT LOTION FOR WOUNDS, SORE FEET, &c. Bruised oak-bark 2 oz., catechu 1 oz., water 3 pints; boil to 1 pint and strain.—CLATER.

POWDERED AND LIQUID CAUSTICS FOR WARTS.

1. Equal parts of sal ammoniac and savine, powdered together.
2. Sublimate 1 dr., hydrochloric acid 1 dr., spirit of wine 3 drs., water 2 drs. The warts to be touched with the liquid twice a day.—YOUATT.
3. Salicylic acid $\frac{1}{2}$ dr., collodion $\frac{1}{2}$ oz. Paint over the wart, which will gradually peel away after each application.

FLEAS.

1. Rub the skin with powdered resin and bran.
2. Let the dog sleep on deal shavings.
3. Scotch snuff steeped in gin.—MEYER. (This requires caution.)
4. Oil of aniseed.—FINLAY DUN.
5. Persian insect powder.

CLYSTERS.

- Astringent.* Alum whey.
- Purgative.* The purgative medicine No. 8; with gruel.
- For Worms.* Solution of aloes 2 oz., linseed oil 1 oz.
- Mix.
- Anodyne.* Boiled starch $\frac{1}{2}$ pint, laudanum 5 to 10 drops.—CLATER.

MEDICINES

FOR

POULTRY, RABBITS, & C.

FOR ROUP, POULTRY GLANDERS, AND GARGLE IN GEESE.

1. A saturated solution of common salt. Medium dose, half a teaspoonful.

2. Antimonial powder 1 gr., with sopped bread, twice a day.—CLATER.

3. Garlic, rue, brickdust, and butter, beaten together, and a little crammed down the throat.

4. *For wet roup in pigeons.* Give 3 or 4 pepper-corns in 3 or 4 days.—MOORE.

5. *For dry roup.* Give 2 or 3 pills of garlic every day. Some recommend assafœtida to be mixed with the food of Poultry, whenever they manifest disease by drooping their wings.

FOR RUMP ROUP, OR INFLAMMATION OF THE OIL-VESSEL. Open the tumour, and squeeze out the collected oil.

GAPES (OR PIP), FROM WORMS IN THE AIR-PASSAGES.

1. Pills of flowers of sulphur, turpentine, and wheat flour.

2. Oil of turpentine 2 drs., linseed oil 1 oz.; or oil of turpentine 2 drs., flour enough to make it into 20 pills. For 20 doses, one every other day for 3 or 4 times.

3. Tobacco smoke.

INFLAMMATION OF THE LUNGS AND ASTHMA.

Give a grain each of calomel and antimonial powder, daily.

PURGING FLUX, OR DIARRHŒA.

1. Change the diet and give whole wheat or rice; and

if obstinate, cram down small pieces of the following mass :—Chalk, p. caraway, and syrup of poppies.—CLATER.

2. Put chalk in their water, or give forge-water.

CROPSICK, OR CONSTIPATION.

If the obstruction is in the crop, endeavour to force the contents into the gullet and mouth by gentle pressure. When partially emptied give rue and butter.

When the obstruction is in the bowels, give bran and pollard, mixed with a little greasy hot liquor, to which, if necessary, a little sulphur may be added; or give a teaspoonful of the castor-oil mixture (see No. 8, Medicines for Dogs, Physic Balls).—See CHIPPING.

PIP, OR BLAIN IN THE TONGUE.

1. Wash the mouth two or three times a day, with a mixture of equal parts of tincture of myrrh and water.

2. Rub the sore with common salt.

3. Solution of chloride of soda 1 dr., water 1 oz., honey of roses 3 drs.

CANKER IN PIGEONS. Apply burnt alum, mixed with honey.

SCABS IN BREASTS AND BACKS OF PIGEONS. Dill seed, cumin seed, fennel seed, of each 1 lb., assafœtida 1½ oz., bay salt ¼ lb., common salt ¼ lb.; knead them with fine clay and flour. Bake it in earthen pots, and set it for the birds to peck.—MOORE.

Genuine Salt-Cat.—Sifted gravel, brickmakers' earth, rubbish of old walls, of each a peck, cumin seed 1½ lbs., bay-salt ¼ lb.; mix.

FOR FILANDER WORMS IN HAWKS. Aloes, iron filings, nutmeg, and honey; mix, and give a small piece as often as necessary.

CHIPPING, IN CHICKENS.—Remove the chickens to a warm place. Mix 1 oz. of castor oil with ¼ oz. syrup of ginger; mix a teaspoonful of this with a little thick gruel, and force a little down several times a day, so that it shall get half a teaspoonful of the mixture in the course of the day.

FOR CHILL, IN TURKEY CHICKS. Give ground malt and barley-meal in equal quantities, adding a little powdered caraway or coriander-seed.

PASTE FOR WEAK TURKEY CHICKS. Eggs boiled hard, nettles, and parsley, all chopped up, and moistened with wine or water.

MEGRIMS OR GIDDINESS. Castor oil 1 oz., syrup of ginger $\frac{1}{4}$ oz., syrup of poppies $\frac{1}{4}$ oz.; mix with gruel and force a little down occasionally.—CLATER.

CONVULSIONS OF DUCKS. Give to grown-up ducks 4 grains of pepper, mixed with fresh butter.

FOR BLINDNESS. Foment with warm water, then drop a few drops of the following solution into the eyes :—
laudanum 1 teaspoonful, water a teacupful.

LOTION FOR WOUNDS.

1. Laudanum, a few drops, added to a teacupful of water.

2. Tincture of myrrh and paregoric, each a teaspoonful, water $\frac{1}{2}$ a pint.

TO PROMOTE THE LAYING OF EGGS. A little sulphate of soda, placed within reach of the hens, is said to be useful. Warmth, good feeding, with a little chopped meat in winter, are also recommended. To prevent their laying soft eggs, supply them with old mortar, bruised egg-shells, or chalk.

FUNGUS, OR PROUD FLESH, FROM WOUNDS IN THE HEAD. Burnt alum 2 drs., honey 1 oz.; mix, and apply twice a day.

VERMIN, TO DESTROY. Tobacco smoke, with good food, and cleanliness.

MOULTING. It is usual to put saffron into the water of cage birds when moulting; others recommend a rusty nail.

FOR SNIFFLES IN RABBITS. Sulphate of copper 1 to 2 grs., morning and evening, in bran.—CLATER.

FOR ROT OR POT BELLY. Give them young green broom, and bread well toasted.—MAYER.

PATENT AND PROPRIETARY
MEDICINES,
DRUGGISTS' NOSTRUMS,

ETC.

THIS division consists of those medicinal compounds which are excluded from the *Pocket Formulary*, as belonging rather to empirical than to regular practice. It includes, in addition to those secret and patent remedies which are usually termed QUACK MEDICINES, preparations of various drugs made according to private formulæ; some favourite domestic remedies; and a few compounds, which, though not empirical, are better known by the names of individual practitioners than by any other title. The supposed composition of some of the secret remedies is given on the authority of Dr PARIS, the *Philadelphia College of Pharmacy*, the *Pharmaceutical Journal*, the *Chemist and Druggist*, and other sources; but without vouching for their correctness.

ABERNETHY'S PILLS. The nostrum to which this distinguished surgeon's name has been applied is said to consist of 2 grs. of blue pill, and 3 of compound extract of colocynth.

AGUE DROPS (tasteless). A solution of arsenic, probably similar to the *liquor potassæ arsenitis* (*liquor arsenicalis*, B.) of the Pharmacopœia.

ALCOCK'S POROUS PLASTER. The plaster seems to be made by heating and melting together caoutchouc, Burgundy pitch, olibanum, and myrrh, with the aid of turpentine oil.—HAGER.

ALLEN'S ANTI-FAT. A fluid form of *Fucus Vesiculosus* (bladder wrack).

ALLEN'S WORLD'S HAIR RESTORER. An analysis of the contents of an 8 oz. bottle indicated 75.6 grs. of sulphur; and an amount of lead corresponding to 87 grs. of acetate. (*Chemist and Druggist*.)

ANDERSON'S PILLS. See *PILULÆ ANDERSONIS, Pocket Formulary.* Other published formulæ are—

1. Barbadoes aloes 1 oz., jalap $\frac{1}{4}$ oz., soap 1 dr., oil of aniseed $\frac{1}{2}$ dr., tincture of aloes q. s.; mix, and divide into 4-grain pills.

2. Barbadoes aloes 5 oz., water 1 oz.; soften by the heat of a water-bath, and add powdered jalap, powdered aniseed, and ivory-black, of each 1 oz., oil of aniseed 1 dr.

3. Barbadoes aloes 16 oz., black hellebore, jalap, subcarbonate of potash, of each 1 oz., oil of aniseed $\frac{1}{2}$ oz., syrup of buckthorn q. s. to form a mass. To be divided into 4-grain pills.

4. Barbadoes aloes 24 oz., soap 4 oz., colocynth 1 oz., gamboge 1 oz., oil of aniseed $\frac{1}{2}$ fl. oz.; mix, and divide into pills of 3 grs. each. *Phil. Coll. of Pharmacy.*

ANODYNE NECKLACES. Beads formed of the root of henbane, and used as necklaces, to allay the pain of teething.

ANTIPERTUSSIS. Dr PARIS states that the basis of this nostrum is a salt zinc.

ARQUEBUSADE (acid). 1. Sulphuric acid $\frac{1}{2}$ lb., vinegar and spirit of wine, of each 3 lbs., clarified honey 1 lb.—**SWEDIAUR.**

2. Distilled vinegar and rectified spirit, of each 10 oz., sulphuric acid (by weight) 14 oz., sugar 24 oz.; mix. For the aromatic spirituous arquebusade water, see **ARQUEBUSADE WATER**, under **PERFUMERY**; also **SPIRITUS VULNERARIUS. Pocket Formulary.**

AROMATIC VINEGAR. Acetic acid (sp. gr. 1.044) 1 lb., camphor 1 oz.; dissolve, and add 1 oz. each of oil of lavender, oil of cloves, and oil of lemon.

APERIENT AND ANTIBILIOUS PILLS. See **ANDERSON'S, BAILLIE'S, BARCLAY'S, DIXON'S, &c., Pills.** The following are useful forms—

1. Compound extract of colocynth 60 grs., comp. rhubarb pill 30 grs., soap 10 grs. In 24 pills.

2. Compound extract of colocynth 2 drs., extract of rhubarb $\frac{1}{2}$ dr., compound soap pill 10 grs.; mix, and divide into 40 pills; 1, 2, or 3 for a dose.

3. Compound extract of colocynth 8 oz., soap 1 oz., scammony 2 oz., extract of rhubarb 2 oz., oil of cassia 5 drs., spirit q. s. to form a mass. Divide into 4-grain pills.

4. Blue pill, compound extract of colocynth, of each a

scruple; scammony and Castile soap of each 10 grs., oil of caraway 4 drops. Mix and divide into 15 pills—3 at bedtime.—Sir B. BRODIE.

5. Compound extract of colocynth 4 scruples, scammony a scruple, extract of rhubarb 12 grs., soap 6 grs., oil of cinnamon 4 drops. Mix, and divide into 24 pills.—Mr VANCE.

6. Mr VANCE'S *Stronger Pills with Calomel*. Compound extract of colocynth 4 scruples, scammony 2 scruples, calomel 24 grs., oil of cinnamon 6 drops, in 24 pills.—Dr J. JOHNSON.

7. Compound extract of colocynth 1 dr., calomel 15 grs., emetic tartar 1 gr., oil of cassia 5 drops. In 24 pills.—Dr J. JOHNSON.

8. Scammony 10 to 15 grs., compound extract of colocynth 2 scruples, extract of rhubarb $\frac{1}{2}$ dr., soap 10 grs., oil of caraway 5 drops. In 20 pills. One or two when required.—Sir C. SCUDAMORE.

9. Compound rhubarb pill $\frac{1}{2}$ dr., ipecacuanha 6 grs., compound extract of colocynth 20 grs. In 12 pills. One or more at bedtime occasionally.—Dr BARON.

10. The same as PIL. COLOC. ET HYOSCYAMI, Ed. Ph.—Dr HAMILTON.

11. (Strong.) Compound extract of colocynth 2 drs., aloes and myrrh pill 2 drs., calomel 1 dr.; mix, and divide into 40 pills. Two for a dose.—Dr LYNN.

12. Compound extract of colocynth 2 scruples, ipecacuanha 6 grs., soap 10 grs., extract of henbane 30 grs. Into 18 pills. Two at bedtime.—Dr COPLAND.

13. Dr NELIGAN'S *Purgative Pills for general use*. Comp. colocynth pill and soap of jalap, of each 1 dr. In 24 pills. Two when required.

14. (Without aloes.) Simple extract of colocynth 24 grs., extract of jalap 12 grs., blue pill 12 grs., ipecacuanha 4 grs., oil of peppermint 3 drops. Into 12 pills.

Other formulæ will be found in the *Pocket Formulary*. Those which do not contain calomel should be preferred for general and repeated use.

ATKINSON'S INFANT PRESERVATIVE. Carbonate of magnesia 6 drs., white sugar 2 oz., oil of aniseed 20 drops, spirit of sal volatile $2\frac{1}{2}$ drs., laudanum 1 dr., syrup of saffron 1 oz., caraway water to make a pint.

ATROPINE PAPER. Green tissue paper imbued with a solu-

tion of sulphate of atropia, so that a piece 1-5th of an inch square contains as much as a drop of a solution of 2 grs. to 1 oz. of water. The paper is hung up and turned about while drying. A piece of the size named will dilate the pupil if placed on the sclerotic and the lids closed over it, and tied with a handkerchief.—Mr STREET-FIELD.

Discs of atropine made of gelatine, and containing usually about 1-5000th of a grain of sulphate of atropine in each, are also employed in ophthalmic practice. See *Pocket Formulary*.

BACHER'S TONIC PILLS. Alkaline extract of black hellebore 2 drs., extract of myrrh 2 drs., powder of holy thistle 1 dr.; mix, and divide into 4-grain pills.

Dr BAILLIE'S PILLS. Compound extract of colocynth 1½ drs., extract of aloes 1½ drs., Castile soap ½ dr., oil of cloves 15 drops. In 38 pills; 3 at bedtime occasionally.

Dr BAILLIE'S DINNER PILLS. Aloes 20 grs., ginger ½ dr., ipecacuanha 8 grs., syrup q. s. Mix, and divide into 16 pills. One daily, before dinner.

BAILEY'S ITCH OINTMENT. Olive oil 1 lb., suet 1 lb., alkanet root 2 oz. Melt, and macerate until coloured; then strain, and add 3 oz. each of alum, nitre, and sulphate of zinc, in very fine powder; adding vermilion to colour it, and oil of aniseed, lavender, and thyme, to perfume.

BAKING POWDER. Tartaric acid 8 oz., bicarbonate of soda 9 oz., arrowroot, or rice flour, 10 oz. Mix. DELFORT'S is said to consist of alum 5 oz.,* bicarbonate of soda 2½ oz., bicarbonate of ammonia ½ oz., arrowroot 4 oz.

BALM OF GILEAD (factitious). 4 oz. of gum benzoin may be dissolved by heat in 1 lb. of Canada balsam, and to the mixture, when cold, ¼ oz. each of the oils of rosemary, lemon, and cassia, added.

BALSAM OF RAKASIRI. Oil of rosemary dissolved in common gin.

BALM OF SYRIACUM. PERRY'S. Gin 1 pint, moist sugar ½ lb., water 4 oz. Dissolve the sugar in the water, mix in the gin, then add Tinct. Camph. Co. (P. L., 1836) 1 oz., tincture of tolu ½ oz., aniseed and spearmint oils a few drops. Mix well, let stand, and decant the clear portion.

BALSAM. See FORD'S, HILL'S, FRIAR'S, &c.

* The employment of alum in bread-making ought to be discountenanced.

- BARCLAY'S (REV. D.) ANTIBILIOUS PILLS.** Extract of colocynth 2 drs, soap of jalap $2\frac{1}{2}$ drs., extract of guaiacum wood 3 drs., emetic tartar 8 grs., oil of juniper, caraway, and rosemary, of each 4 drops; into 4-grain pills.
- BAREGIAN BALLS.** Extract of soap-wort (or of artichoke leaves) 3 oz., gelatine $1\frac{1}{2}$ oz., water 3 oz.; heat together till dissolved, pour the solution into a warm iron mortar; add 6 oz. of sulphuret of lime, and 1 oz. of salt, previously powdered and mixed. Stir constantly till a mass is obtained, and divide it into balls of $2\frac{1}{4}$ oz. each. Use one for a general bath, half of one for a footbath.
- BARK, ESSENTIAL SALT OF.** See *EXTRACTUM CINCHONÆ SICCUM*, *Pocket Formulary*.
- BARKER'S TOOTH TINCTURE.** An alcoholic tincture of pyrethrum, coloured with tincture of red cabbage.
- BATEMAN'S PECTORAL DROPS.** 1. Compound spirit of aniseed 16 fluid oz., opium 1 dr., camphor 1 dr., oil of fennel 20 drops, cochineal 2 drs.
2. Proof spirit 4 galls., red sanders 2 oz.; digest 24 hours, filter, and add powdered opium 2 oz., camphor 2 oz., catechu 2 oz., oil of aniseed 4 fluid drs.; digest for 10 days. The old wine gallon is here intended. *Philadelphia College of Pharmacy*.
- BATEMAN'S ITCH OINTMENT.** Carbonate of potash $\frac{1}{2}$ oz., rose-water 1 oz., vermilion 1 dr., sulphur 11 oz., oil of bergamot $\frac{1}{2}$ dr., lard 11 oz.; mix.
- BATE'S ANODYNE BALM.** Soap liniment 2 parts, tincture of opium 1 part.
- BATE'S CAMPHORATED EYE-WATER.** Sulphate of copper 15 grs., French bole 15 grs., camphor 4 grs., boiling water 4 oz.; infuse, strain, and dilute with 4 pints of cold water.
- BATE'S STYPTIC WASH.** See *LIQUOR ALUMINIS CO.*, *Pocket Formulary*.
- BATHING SPIRITS.** These resemble liquid opodeldoc (soap liniment), and are usually coloured by the addition of some dark tincture. See *FREEMAN'S BATHING SPIRITS*.
- BATH DIGESTIVE PILLS.** Rhubarb 2 oz., ipecacuanha $\frac{1}{2}$ oz., cayenne pepper $\frac{1}{2}$ oz., soap $\frac{1}{2}$ oz., ginger $\frac{1}{4}$ oz., gamboge $\frac{1}{4}$ oz.; mix, and divide into 4-gr. pills.
- BATH LOZENGES (in imitation of DAWSON'S).** Pure extract of liquorice 1 oz., powdered gum arabic 1 oz., white sugar

- 1 lb., hot water q. s. to form a mass; to be rolled into pipes.
- BATTLE'S VERMIN POWDER.** Said by Dr MACADAM to contain starch, sugar, and Prussian blue, with 10 per cent. of strychnia.
- BATLEY'S LIQUOR CINCHONÆ and LIQUOR OPII.** See *Pocket Formulary*.
- BATLEY'S SENNA POWDER.** Senna leaves heated until they become light in colour, reduced to powder, and mixed with some finely powdered charcoal.
- BAUME DE VIE.** Socotrine aloes 2 drs., rhubarb 6 drs., saffron 2 drs., liquorice root 1 oz., proof spirit 8 oz.; digest for 8 days, and filter. The original Swedish form is this:—Aloes 9 drs., rhubarb, gentian, zedoary, saffron, theriaca, agaric, of each 1 dr., proof spirit 2 pints.
- BAY RUM.** Prepared from the leaves of *Myrica acris* (known as "Bayberry" in America) by distilling them with rum.—*Chemist and Druggist*.
- BAYNTON'S PLASTER.** Simple litharge plaster 16 oz., yellow resin 6 drs.; melt together, and spread on linen or calico.
- BEDDOW'S PILLS;** for gravel. Carbonate of soda dried without heat, 1 dr., soap 4 scruples, oil of juniper 10 drops, syrup of ginger q. s. for 30 pills.
- BEEBLE WAFERS.** Red lead, sugar, and flour; made into wafers.
- BELLOSTE'S PILLS.** Quicksilver, scammony, and jalap, of each 1 lb., sugar 4 oz.; mixed and made up into a mass with sherry wine.
- BESTUCHEFF'S NERVOUS TINCTURE.** A mixture of a strong solution of perchloride of iron with sulphuric ether and spirit, exposed in long bottles to the rays of the sun until it has quite lost its brown colour.
- BETTON'S BRITISH OIL.** Oil of turpentine 8 oz., Barbadoes tar 4 oz., oil of rosemary 4 drs.; mix. See BRITISH OILS.
- BEWLEY AND EVANS' CHALYBEATE WATER.** Citrate of iron 13 grs., carbonated water 6 oz., syrup of orange peel 1 oz.
- BISCUITS, APERIENT.** An ounce of powdered jalap, mixed with 16 oz. of the materials for gingerbread, or other kind of cake. See GINGERBREAD, PURGATIVE.
- BLACK DROP.** See GUTTÆ NIGÆ, *Pocket Formulary*.

BLACK DRAUGHT. 1. Infusion of senna 10 drs., sulphate of magnesia 3 drs., syrup of ginger 1 dr., aromatic spirit of ammonia 20 drops.

2. Tartrate of potash $1\frac{1}{2}$ drs., manna $\frac{1}{2}$ dr., tincture of jalap $\frac{1}{2}$ dr., aromatic spirit of ammonia 20 drops, extract of liquorice 4 gts., infusion of senna 11 drs. See *MISTURA SENNÆ COMP.*, *HAUSTUS APERIENS*, and *MISTURA APERIENS*, in *Pocket Formulary*, for other formulæ.

BLAINE'S DISTEMPER POWDERS. The basis of these is said to be aurum musivum (sulphide of tin).

BLISTERING PAPER. Melt together cantharadin 1 dr., white wax 1 dr., olive oil 5 drs. Paint it with a brush on white bibulous paper, and hang it up to dry in a current of air. Take a piece of pink paper of form and size required, paint the under coloured side with a weak solution of india rubber, cut the cantharadine paper to the size, less a margin, of the pink paper, and place it on while the india-rubber solution is still sticky. Before applying, the blister must be held over the steam of hot water. See *PAPIER EPISPASTIQUE*.

BLISTERING TISSUE. *Taffetas Vesicant.* Powdered cantharides exhausted by ether, the tincture distilled to recover the principal part of the ether for the same use, and the residue heated in a water-bath till it ceases to boil. The green butyraceous oil which remains is to be melted with twice its weight of wax, and spread on waxed silk, or any convenient and adhesive material. An extract prepared by evaporating a tincture made with 4 parts of flies, 1 of strong acetic acid, and 16 of rectified spirit, is used for the same purpose.

BROWN'S BRONCHIAL TROCHES. See *TROCHISCI CUBEBÆ*, *Pocket Formulary*.

BOCHET'S SYRUP, for scrofulous affections. Compound syrup of sarsparilla, with senna, and 1 per cent of iodide of potassium.

BRANDISH'S ALKALINE SOLUTION, OR CAUSTIC ALKALI. See *LIQ. POTASSÆ BRANDISHII*, *Pocket Formulary*.

BRANDISH'S ALKALINE TINCTURE OF RHUBARB. Coarsely powdered rhubarb 1 oz., alkaline solution (BRANDISH'S) 32 fluid oz. The original formula directs only $\frac{1}{2}$ oz. rhubarb, but as smaller doses of the tincture than were given by Mr B. are usually prescribed, the quantity of rhubarb is here increased. Or an alkaline infusion of

rhubarb may be made by pouring boiling water 38 parts on rhubarb 3 parts, and carbonate of potash 1 part.

BRANDBETH'S PILLS. Root of may apple 10 grs., extract of ditto 10 grs., extract of poke berries 30 grs., powdered cloves 10 grs., gamboge 2 to 5 grs., traces of saffron, and a few drops of oil of peppermint; made into 24 pills.
—HAGER.

BRITISH HERB TOBACCO. The principal ingredient in this compound is dried coltsfoot leaves, to which a smaller portion of thyme, wood-betony, eye-bright, and rosemary, are added.

BRITISH OILS. Oil of turpentine and linseed oil, of each 8 oz.; oil of amber and oil of juniper, of each 4 oz.; true Barbadoes tar 3 oz.; American petroleum (seneca oil) 1 oz.; mix. See **BETTON'S BRITISH OIL**, *Ante*.

BRODUM'S NERVOUS CORDIAL. Iron wine, compound spirits of lavender, tinctures of calumba, gentian, cinchona, and cardamoms, equal parts of each.

BROCCHIERI'S STYPTIC WATER. Pieces of fresh pine, bruised in a mortar, and distilled with twice their weight of water, till half the water has come over. After standing in a wide vessel, any floating oil is to be removed from the surface, and the water kept for use.

BURNETT'S (SIR WILLIAM) DISINFECTING FLUID. A neutral solution of zinc in commercial hydrochloric acid.

CACHOU AROMATISÉ. See **PERFUMERY**.

CAJEPUT LINIMENT. Soap liniment 7 oz., camphor $\frac{1}{2}$ oz., oil of cajeput 1 oz.

CAMPBOR LINIMENT, EXTEMPORANEOUS. Rectified spirit 17 fluid oz., strong water of ammonia $2\frac{1}{2}$ oz., camphor 2 oz., oil of lavender 50 minims.

CARRON OIL. Lime-water and linseed oil, equal quantities.

CASTILLON'S POWDERS. Sago meal, salep, tragacanth, each 1 dr.; prepared oyster-shells a scruple; coloured with cochineal. A drachm to be boiled with milk; in bowel complaints.

CEPHALIC SNUFF. Dried asarabacca leaves 3 parts, marjoram 1 part, lavender flowers 1 part; rub together to a powder. **BOELI'S** consisted of 2 drs. valerian, 2 drs. of snuff, 3 drops of oil of lavender, 3 drops of oil of marjoram; mix. This is said to relieve the eyes as well as the head,

CHAMBERLAINE'S PILLS. Milk of sulphur and vermilion.

Dr PARIS's statement that they contain sulphate of lime would probably surprise the proprietor, if not aware that a great part of the commercial milk of sulphur contains half its weight of that substance.

CHAMOMILE DROPS. Dr PARIS states that the nostrum sold under this name is merely spirit flavoured with essential oil of chamomile. A strong tincture of the flowers would probably be more efficacious.

CHAMOMILE PILLS. Equal parts of rhubarb and jalap made into a mass with extract of chamomile.—SCHÄDLER.

CHELSEA PENSIONER. Powdered rhubarb 2 drs., cream of tartar 1 oz., guaiacum 1 dr., sulphur 2 oz., 1 nutmeg grated fine, clarified honey 16 oz.; mix: take 2 spoonfuls night and morning: for chronic rheumatism.

CHELTENHAM SALTS (factitious). Sulphate of soda crystallised 34 grs., sulphate of magnesia 24 grs., chloride of sodium 50 grs., water 1 pint. Dissolve.

CHILBLAINS, Popular Remedies for. 1. Soap liniment 1 oz., cajeput oil $\frac{1}{2}$ oz., tincture or cantharides $\frac{1}{4}$ oz.; mix.

2. Sal ammoniac $\frac{1}{2}$ oz., vinegar 5 oz., spirit of rosemary 1 oz.; mix.

3. Oil of turpentine 1 oz., camphor $\frac{1}{4}$ oz., Goulard's extract $\frac{1}{4}$ oz.; mix.

4. Dr GRAVES'S *Preventive*. Sulphate of copper 10 grs., water 1 oz.; dissolve, brush over the parts with the lotion by means of a camel-hair pencil, and when dry apply a little simple ointment. Repeat this for some evenings in succession.

5. LEJEUNE'S BALSAM. See further on.

6. Sal enixum, alum, and sulphate of zinc, of each $\frac{1}{4}$ oz., water a pint; apply it frequently.

7. Muriatic acid $\frac{1}{2}$ oz., Friar's balsam 3 $\frac{1}{2}$ oz.; mix.

8. SWEDIAUR'S *Paste*. Bitter almonds 8 oz., honey 6 oz., powdered camphor $\frac{1}{2}$ oz., flower of mustard $\frac{1}{2}$ oz., burnt alum $\frac{1}{4}$ oz., olibanum $\frac{1}{4}$ oz., yolks of 3 eggs; beat together to form a paste. Rub a portion of it on the part affected, moistened with water, night and morning, then wash with warm water, and dry with a cloth.

9. WAHLER'S *Ointment for Broken Chilblains*. Black oxide of iron, bole, and oil of turpentine, of each 1 dr.; rub together, and add the mixture to 1 oz. of melted resin cerate.

10. *Another ointment for the same.* Locatelli balsam 1 oz.; citrine ointment $\frac{1}{4}$ oz., balsam of Peru 20 drops; mix.

11. *Russian remedy.* Dry the peelings of cucumbers, and when required for use soften the inner part with water, and apply it to the part affected.

12. Tincture of arnica, rose-water, glycerine, of each 3 parts, spirit of camphor 1 part.

13. Dr DEWAR. Sulphurous acid and glycerine, of each 1 part, distilled water 2 parts.

CHING'S WORM LOZENGES. The yellow lozenges contain 1 gr. of calomel in each, with sugar and sufficient mucilage (coloured with saffron) to form a paste. The brown contain $\frac{1}{2}$ gr. of calomel with $3\frac{1}{2}$ grs. of resinous extract of jalap, according to GRAY; or with 1 gr. of resin of jalap, according to Dr PARIS and others.

CHLORODYNE. Many formulæ have been given for this preparation. Mr GROVES gives the following as an improvement on the recipe of Dr OGDEN:

1. Take chloroform 4 drs., ether $1\frac{1}{2}$ drs., oil of peppermint 8 drops, resin of Indian hemp 16 grs., capsicum 2 grs.; macerate for 2 or 3 days, and filter. Then dissolve hydrochlorate of morphia 16 grs. in 1 fl. oz. of syrup, add perchloric acid and water $\frac{1}{2}$ dr. each, assisting the solution by a water-bath; then, when cold, add hydrocyanic acid (SCHEELE'S) 96 drops. Mix the solutions. Should be given with great caution.

2. OLDBERG'S formula is as follows:—Sulphate of morphia 1 part, menthol 1 part, ether 25 parts by measure, alcohol (820) 25 parts by measure, diluted hydrocyanic acid 50 parts by measure, chloroform 150 parts by measure, simple syrup q. s. to make 1000 parts by measure. See also, TINCT. CHLOROFORMI ET MORPHINÆ, B., *Pocket Formulary*.

CHOLERA MEDICINES. The following are some of the more popular remedies that have at various times been used during the visitations of this disease.

1. *Liverpool Preventive Powders.* Bicarbonate of soda 1 scruple, ginger 8 grs.; to be taken in a glass of water after breakfast and supper. These powders are said to have been administered with good effect among the workmen in the mining and manufacturing districts, during a former visitation of cholera.

2. Dr STEVENS' *Saline Powders:* Bicarbonate of soda

$\frac{1}{2}$ dr., chloride of sodium a scruple, chlorate of potash 7 grs.; mix, for 1 dose.

3. Mr HOPE'S *remedy*. Nitrous acid (red) 2 drs., peppermint water or camphor mixture 1 oz., tincture of opium 40 minims; dose, 1 to 2 teaspoonfuls in a cupful of gruel every 3 or 4 hours.

4. Spirit of wine 1 oz., spirit of lavender $\frac{1}{4}$ oz., oil of origanum $\frac{1}{4}$ oz., compound tincture of benzoin $\frac{1}{2}$ oz., spirit of camphor $\frac{1}{4}$ oz.; 20 drops on moist sugar. To be rubbed outwardly also.

5. *American remedy*. Equal parts of lard, maple-sugar, and charcoal; to be mixed, and the size of a nut swallowed.

6. Remedy recommended by the Board of Health, in premonitory diarrhoea:—Aromatic powder 3 drs., compound spirit of ammonia 3 drs., tincture of catechu 10 drs., compound tincture of cardamoms 6 drs., tincture of opium 1 dr., chalk mixture to make 20 oz. Dose, for an adult, 1 oz., for a child of 12 years, $\frac{1}{2}$ oz., for 7 years, $\frac{1}{4}$ oz., after each liquid stool.

7. Dr GRAVES' *Astringent Pills*. Acetate of lead 20 grs., opium 1 gr.; made into 12 pills. One every half hour till the watery discharges cease.

8. Mr BUXTON'S *remedy*. Twenty-five minims of diluted sulphuric acid in an ounce of water.

9. Dr BEAVEN'S *Preventive and Remedy*.

The Preventive. Sulphite of magnesia 2 drs.; sulphurous acid 2 oz., water 2 oz.; tincture of capsicum $\frac{1}{2}$ oz. Mix and dissolve. A teaspoonful night and morning.

The Remedy. Sulphite of magnesia 2 drs.; sulphurous acid 2 oz.; tincture capsicum $\frac{1}{2}$ oz., water 2 oz., sulphate of morphia 2 grs.; mix, and dissolve. A teaspoonful every half hour until relieved.

We have inserted the above, not to encourage quackery in reference to this terrible disease, but because the druggist may be called upon to supply these remedies, and expected to know their composition. For ELIXIR WORONEJE, see *Pocket Formulary*.

CHIRETTA PILLS AND MIXTURE. Dr REECE'S *Pills*. Extract of chiretta 2 drs., dried carbonate of soda 20 grs., ginger 15 grs.; mix, and divide into 36 pills. Two, twice

- a day. *Mixture*: Infusion of chiretta 8 oz., subcarbonate of soda 1 dr.; two tablespoonfuls 3 times a day.
- CITRATE OF MAGNESIA. See EFFERVESCING CITRATE OF MAGNESIA.
- CLARKE'S BLOOD MIXTURE. According to Dr FARQUARSON, this medicine contains 10 grs. of iodide of potassium in each dose.
- CLUTTON'S FEBRIFUGE SPIRIT AND TINCTURE. *Spirit*: The original formula is—oil of sulphur by the bell, oil of vitriol and sea salt, of each 1 oz.; rectified spirit 3 oz.; mix, digest for a month, and distil to dryness. *Tincture*: Febrifuge spirit 8 fluid oz.; angelica root, serpentary, cardamon seed, of each 1½ drs.; digest, and strain. Water acidulated with these and sweetened to the taste, forms a cooling diuretic and diaphoretic julep.
- COCHRANE'S COUGH MEDICINE. An acidulated syrup of poppies.
- COLLIER'S (Dr) WINE OF QUININE. Disulphate of quinine 18 grs., critic acid 15 grs., sound orange wine 1 bottle, or 24 fluid oz.
- COLLIER'S (Dr) CREAM OF TARAXACUM. See CREMOR TARAXACI, *Pocket Formulary*.
- COLLINS'S DISINFECTING POWDER. See DISINFECTING and DEODORIZING COMPOUNDS among the TRADE CHEMICALS.
- COLLODION. See TRADE CHEMICALS.
- COLLODION (STYPTIC). Dr RICHARDSON. A saturated solution of tannic acid and gun cotton in ether.
- Dr PAVESI. Collodion 100 parts, carbolic acid 10 parts, tannic acid and benzoic acid, of each 5 parts. Application to wounds.
- CONSUMPTION, *Popular Remedies for*. 1. Rum ½ pint, linseed oil, honey, garlic (beaten to a pulp), and loaf sugar, of each 4 oz., yolks of 5 eggs; mix; a teaspoonful night and morning.
2. *Breastplate*. Dissolve 1 oz. of aloes in 12 oz. of a strong decoction of fresh rue; fold a large piece of soft muslin in 8 folds, large enough to cover the chest and part of the stomach; steep in the solution and dry it in the shade; wear it on the chest constantly.
- COUGH LINCTUS. 1. *Rose Linctus*. Confection of roses 3 oz., paregoric elixir 1½ oz., diluted sulphuric acid 1 dr.

mix; a small teaspoonful now and then when the cough is troublesome.

2. Dr LATHAM'S *Cough Linctus*. DOVER'S powder $\frac{1}{2}$ dr., compound powder of tragacanth 2 drs., syrup of tolu $\frac{1}{2}$ oz., confection of hips and simple oxymel, of each 1 oz.; a small teaspoonful 3 or 4 times a day. For other formulæ, see LINCTUS, LINCTUS PAPAVERIS, LINCTUS SCILLÆ, &c., *Pocket Formulary*.

COUGH LOZENGES. See BATH LOZENGES, further back; also LOZENGES, further on.

CORN PLASTERS. See KENNEDY'S CORN PLASTER and EMPLASTRUM ÆRUGINIS, *Pocket Formulary*. Most of the advertised corn plasters contain verdigris. A few additional formulæ are subjoined,

1. Galbanum plaster 1 oz., prepared verdigris 1 scruple; melt, and mix.

2. Galbanum 1 oz., black pitch $\frac{1}{2}$ oz., simple diachylon $\frac{1}{4}$ oz., verdigris a scruple, sal ammoniac a scruple. Melt the first three together, and add the last two in fine powder.

3. Plaster of ammoniacum with quicksilver $1\frac{1}{2}$ oz., soap plaster $\frac{1}{2}$ oz., opium in fine powder $\frac{1}{2}$ dr.

MECHANICAL CORN PLASTERS. Any suitable adhesive plaster is spread on soft thick leather (buckskin), which is afterwards cut to a suitable size, and a hole punched in the centre. They are sometimes spread on amadou, or on vulcanized india rubber.

CORN SOLVENTS. One of the preparations sold under this name is probably a strong solution of carbonate of potash. A powder sold for the same purpose consists of carbonate of potash coloured with ochre or bole. A pinch is placed on the corn, and confined by means of a piece of adhesive plaster or rag. Sir Humphrey Davy's name was given to a remedy which consisted of carbonate of potash and salt of sorrel, similarly applied. The following is one of the advertised Corn and Bunion remedies:—Carbonate of soda 1 oz., finely powdered and mixed with $\frac{1}{2}$ oz. of lard. Applied on linen rag every night; the outer skin to be pared off every 2 or three days. It may be varied thus:—Dried carbonate of soda 4 drs., powder blue (smalts) a scruple, lard 4 drs.; mix. A very excellent and effective application, the invention of Mr GEZOW, a Russian apothecary, is the following:—Salicylic acid 1 part, extract of

Indian hemp, $\frac{1}{2}$ th part, collodion 8 parts. To be applied with a camel-hair brush, taking care to confine its application to the corn or bunion.

CAUSTIC FOR CORNS. 1. Tincture of iodine 4 drs., iodide of iron 12 grs., chloride of antimony 4 drs.; mix, and apply with a camel-hair brush, after paring the corn. It is said to cure in 3 times.

2. Strong acetic acid in glass tubes is used for this purpose.

COURT PLASTER. See *EMPLASTRUM ICHTHYOCOLLÆ*, *Pocket Formulary*.

CUSTARD POWDER. See *DIETETIC ARTICLES*.

DAFFY'S ELIXIR. This is similar to the compound tincture of senna; but different makers have their peculiar formulæ. The following are some of them. Avoirdupois weight seems to be intended:

1. Senna leaves 3 $\frac{1}{2}$ lbs., jalap, aniseed, caraway seed, of each 20 oz., rectified spirits 18 pints, sugar 5 lbs. Infuse the senna 2 or 3 times in sufficient boiling water to yield, when strained with pressure, 4 gallons of the whole. Add to this the tincture made with jalap and seeds, digested with the spirit for a week. Pour off the clear liquor and add the sugar, and brandy colouring if required.

2. **DICEY'S**, according to **GRAY**. Senna 5 oz., guaiacum shavings (some recipes add red sanders), dried elecampane root, seed of anise, coriander, and caraway, and root of liquorice, of each 2 $\frac{1}{2}$ oz., stoned raisins 8 oz., proof spirit 6 lbs.

3. **SWINTON'S**. Jalap 3 lbs., senna 2 oz., coriander seed, aniseed, liquorice root, and elecampane, of each 4 oz.; spirit of wine and water, of each a gallon.

4. Small senna 10 oz., bruised jalap, coriander seed, and aniseed, of each 2 $\frac{1}{2}$ oz., proof spirit a gallon. Digest 8 days, frequently shaking, and strain. Pour on the remaining ingredients 6 oz. of boiling water in which 2 drs. of salt of tartar have been dissolved; press strongly, and add the liquid to the tincture, with 3 oz. of treacle. Some recipes add rhubarb, in the proportion of about $\frac{1}{4}$ oz. to the gallon.

DALBY'S CARMINATIVE. 1. Carbonate of magnesia 1 oz., syrup of poppies 5 drs., tincture of wood-soot 1 oz., oil of caraway 25 drops, oil of peppermint 16 drops, water and spirit of wine, each $\frac{1}{2}$ oz. Mix.

2. Carbonate of magnesia 2 scruples, oil of peppermint 1 drop, oil of nutmeg 2 drops, oil of aniseed 3 drops, tincture of castor 30 drops, tincture of assafoetida 15 drops, tincture of opium 5 drops, spirit of pennyroyal 15 drops, compound tincture of cardamom 30 drops, peppermint water 2 oz. Mix.—Dr PARIS.

DANDELION COFFEE. The roots, collected at the end of the year, are dried at a gentle heat and reduced to powder. Some mix coffee with it. Others roast the root in the manner of coffee, but probably at the expense of its medicinal virtues. The better way is to dry and powder it, and direct it to be mixed with coffee when used. If considered necessary to give it more colour and flavour, it may be previously mixed with a sufficient quantity of roasted chicory, which should not exceed one eighth of the whole.

DARCET'S ALKALINE LOZENGES, or VICHY LOZENGES. Bicarbonate of soda 2 drs., refined sugar 14 oz., oil of peppermint 4 drops, mucilage of tragacanth q. s. Mix, and divide into 60 lozenges.

DAWSON'S LOZENGES. See BATH LOZENGES, above.

DELAMOTT'S GOLDEN DROPS. Muriate of iron 1 oz., spirit of sulphuric ether 7 oz.; dissolve and expose to sunshine in a closely stopped bottle till it becomes divested of colour. See BESTUCHEFF'S NERVOUS TINCTURE.

DERBYSHIRE'S PATENT EMBROCATION FOR PREVENTING SEA-SICKNESS. Boil 2 oz. of opium, 2 drs. of extract of henbane, 10 grs. of mace, and 2 oz. of mottled soap, in 3 pints of water for half an hour. When cold, add 1 quart of rectified spirit, and 3 drs. of spirit of ammonia.

DESLER'S SALVE. This is merely resin cerate.

DIGESTIVE PILLS. See BATH DIGESTIVE PILLS, DINNER PILLS, BAILLIE'S PILLS, WEBSTER'S, Lady, PILLS, Dr REECE'S CHIRAYTA PILLS.

DINNER PILLS. See BATH DIGESTIVE PILLS, WEBSTER'S, Lady, PILLS, &c. The following are a few additional formulæ:—

1. Rhubarb 30 grs., aloes 60 grs., ipecacuanha 12 grs., tincture of ginger q. s. to form a mass; to be divided into 24 pills.

2. Sir CHARLES BELL'S. Rhubarb 50 grs., mastic 6 grs., sulphate of quinine 4 grs.; in 12 pills.

DIXON'S ANTIBILIOUS PILLS. Equal parts of aloes, scam-

mony, and rhubarb, with the addition of a small quantity of tartar emetic, and made up with Castile soap.

DOVER'S POWDERS. The PULVIS IPECACUANHÆ COMPOSITUS of the Pharmacopœia. But the original powder consisted of nitre and sulphate of potash, each 4 oz., fused in a red-hot mortar, and afterwards reduced to powder, and mixed with 1 oz. each of ipecacuanha, opium, and liquorice.

DUPUYTREN'S POMMADE. See HAIR COSMETICS.

DUPUYTREN'S EYE SALVE. Nitric oxide of mercury 10 grs., sulphate of zinc 20 grs., lard 2 oz.; rub perfectly smooth.

DUNCAN'S FLUID EXTRACT OF SENNA. Senna 15 lbs. avoirdupois, boiling water 4 times its weight or q. s. Exhaust the senna by displacement, concentrate the liquor to 10 lbs. avoirdupois; dissolve in it 6 lbs. avoirdupois of treacle, previously concentrated over a water-bath, till it becomes nearly dry, on cooling: add 24 fluid oz. of rectified spirit, and water q. s. to make up 15 pints o. m. Dose, 2 drs. Each oz. corresponds with 1 oz. avoirdupois of senna.

DR DUNCAN'S LACTUCARIUM LOZENGES. As the TROCHISCI OPII (*Pocket Formulary*), substituting lactucarium for opium.

DUNCAN'S GOUT REMEDY. A preparation of colchicum with opium, &c.

DUTCH (OR HÆRLEM) DROPS. The basis of this popular remedy is said to be the residue which is left in redistilling oil of turpentine. The following is one of the imitations of it made in this country: Linseed oil 1 quart, resin 2 lbs., sulphur 1 lb.; boil together over a slow fire; when combined, remove from the fire, and add 1 pint of oil of turpentine and 50 drops of liquor of ammonia; stir well together and bottle.

EASTON'S TONIC SYRUP. See SYR. FERRI PROSPH. CUM QUIN. ET STRYCHNIA. See *Pocket Formulary*.

EATON'S STYPTIC. It is similar to that of HELVETIUS, which see below.

EAU DE FÉE. A hair wash. A strong solution of nitrate of lead.

EAU DE MAGNANIMITÉ. A tincture of ants, with aromatics.

EAU MÉDICINALE D'HUSSON. It is prepared, according to Dr WILSON, from the juice of colchicum flower with half the quantity of brandy; mix, and after standing a

few days, decant into small bottles. But it was more probably made from the root, as prescribed in the following formulæ:—(In one of the French Codices.) *Eau Colchique d'Husson*. Dry colchicum 60 parts in sherry 125 parts; 20 drops for a dose. (According to Mr WANT.)— $\frac{1}{4}$ oz. of the fresh root sliced, macerated in $\frac{1}{2}$ pint of proof spirit.

EAU DE COLOGNE (EAU DE MELISSE), &c. See PERFUMERY.

EAU DE JAVELLE. Dry chloride of lime 2 oz., carbonate of potash 4 oz., water 2 pints; mix the chloride with $1\frac{1}{2}$ pints of water, dissolve the potash in the remainder; mix the solution and filter.

EAU DE LUCE. See PERFUMERY.

EAU DE RABEL. See ACIDUM SULPHURICUM ALCOHOLISATUM, *Pocket Formulary*.

EAU SEDATIVE. RASPAIL. It may be imitated as follows:—Spirit of camphor 1 part, solution of ammonia ('923) 6 parts, salt 6 parts, distilled water to 100 parts. See *Pocket Formulary*.

EDINBURGH OINTMENT. White hellebore powder, sal ammoniac, and lard.

EFFERVESCING CITRATE OF MAGNESIA. Mr DYMOND. A granulated preparation, containing carbonate of magnesia and citric acid in equivalent proportions, and 15 per cent. of sugar. Thrown into water it effervesces, and citrate of magnesia is formed, analogous to Moxon's Magnesian Aperient. See further on. The preparation commonly sold as citrate of magnesia contains no magnesia at all. It is made with bicarbonate of soda and tartaric acid. See SODÆ CITRO-TARTRAS EFFERVESCENS, B. P., and MAGNESIÆ CITRAS, *Pocket Formulary*.

ELLERMAN'S DEODORIZING FLUID. It consists chiefly of persalts of iron. See DISINFECTING AND DEODORIZING COMPOUNDS.

ELIXIR DE GARUS. See *Pocket Formulary*.

ELIXIR LONGÆ VITÆ. Similar to BAUME DE VIE.

ELIXIR OF HALLER. See ELIXIR ACIDUM HALLERI, *Pocket Formulary*.

ELIXIR PAREGORIC. See TINCTURA CAMPHORÆ COMPOSITA, *Pocket Formulary*.

ELIXIR OF VITRIOL. (Mynsicht's Elixir.) See ACIDUM SULPHURICUM AROMATICUM, *Pocket Formulary*. For common sale, druggists used frequently to keep a more

ready and economical preparation, of which the following is one form : Compound tincture of cardamoms 1 lb., tincture of cinnamon 3 lbs. ; cinnamon water 2 lbs. ; mix, and add gradually 1½ lbs. of pure sulphuric acid.

ERVALENTA and REVALENTA. See DIETETIC COMPOUNDS.

ESSENCES. ESSENCES OF FLOWERS will be found under PERFUMERY. ESSENCE OF CELEBY, and other culinary essences, will be found, with allied compounds, in another place. A few concentrated infusions, and other strong preparations of drugs, not sanctioned by the Colleges, but very generally used, may be noticed here.

ESSENCE OF YELLOW BARK. Resinous extract of bark ½ oz., sulphate of quinine 60 grs., rectified spirit 6 oz., tincture of orange peel 2 oz. See EXT. CINCHON. FLAV. Liq. B.P. 1867.

ESSENCE OF CALUMBA, RHUBARB, SENNA. See LIQUOR CALUMBÆ, RHEI, SENNÆ, *Pocket Formulary*.

ESSENCE OF CAMPHOR. See LIQUOR CAMPHORÆ, *Pocket Formulary*.

ESSENCE OF CHAMOMILE. As a substitute for the infusion it may be made as LIQUOR CALUMBÆ, P. F. See CHAMOMILE DROPS for another preparation of this drug.

ESSENCE OF ERGOT. See ESSENTIA SECALIS CORNUTI, *Pocket Formulary*.

ESSENCE OF GINGER. Unbleached Jamaica ginger in coarse powder 5 oz., rectified spirit a pint; digest for 8 days and strain with pressure; or it may be made by percolation. As there is no established form, it varies in strength as prepared by different makers, and often contains Cayenne pepper.

ESSENCE OF GINGER (SOLUBLE) THRESH. Take of strong tincture (1 to 1) of finest Jamaica ginger 1 pint, add in small portions at a time finely powdered slaked lime, vigorously shaking after each addition, until the tincture ceases to lose colour, throw the whole upon a filter and pass through the residue, proof spirit until the product measures 2 pints. Now add drop by drop dilute sulphuric acid until the rich yellow colour of the tincture suddenly disappears, let stand for 24 hours, filter, dilute with water to 4 pints, shake with a little powdered pumice or silica (by no means lime or magnesia), and filter at 0° C. if possible.

ESSENCE OF CUBEBS. Mix powdered cubebs with ether, in

a well-stopped bottle; in twelve hours put the paste into a percolator, and add ether till the cubebs are nearly exhausted; distil off the ether in a water-bath, and preserve it for the same purpose. Dissolve the extract which remains in three times as much brandy. One dr. is equal to 2 drs. of the powder. A fluid extract is also made by concentrating the tincture.

ESSENCES OF MINT, PEPPERMINT, AND PENNYROYAL. The strength of these varies as prepared by different makers; some use 1 part of the essential oil to 3 of rectified spirit, but more usually, we believe, 1 part to 7. They are sometimes coloured with the leaves of the plant, or of spinach.

ESSENCE OF MUSTARD. Rectified spirit of turpentine 16 fluid oz., bruised black mustard seed 2 oz., camphor 4 oz., oil of rosemary $\frac{1}{2}$ oz., annatto to colour. Or, essential oil of mustard 1 part, rectified spirit 60 parts; to be sparingly sprinkled on piline, and applied as a mustard poultice.

ESSENCE OF SARSAPARILLA. Jamaica sarsaparilla 16 oz., lukewarm distilled water (100° to 112° F.) sufficient to cover it. Macerate for 6 hours, and strain. Bruise the root, macerate it again in sufficient warm water, and repeat the maceration with fresh water until it ceases to be much coloured. After straining, let the mixed liquids be immediately heated to 180° F., allowed to cool, and filtered. Evaporate the whole of the filtrate by a water or steam-bath, at a heat not above 160° , until reduced to 14 or 15 fl. oz; add 2 oz. of rectified spirit, and keep it in a close bottle in a cool place for a few days. Then carefully pour off the clear liquid from any sediment into a clean dry bottle. 1 fl. oz. represents 1 oz. of the root, or 8 oz. of the decoction. See also **EXTRACTUM SARZÆ LIQUIDUM**, **LIQUOR SARZÆ**, and **ESSENTIA SARSAPARILLÆ**, in *Pocket Formulary*. The latter is an elegant and efficacious preparation.

COMPOUND ESSENCE OF SARSAPARILLA. Jamaica sarsaparilla 16 oz.; proceed as above, but reserve the liquor of the last maceration for boiling the other ingredients; namely, guaiacum raspings, bruised liquorice root, sassafras, of each 2 oz., mezereon $\frac{3}{4}$ oz. Boil them in 4 or more pints of the weak infusion for half an hour, and strain; evaporate to 4 fl. oz.; let it cool, stirring it occasionally, and add 2 oz. of rectified spirit in which a few

drops of oil of sassafras have been dissolved. Evaporate the sarsaparilla liquid to 11 oz., and when cool add the other liquid. Proceed as for the former. One measure with 7 of water forms a near approximation to the Pharmacopœia Decoction.

ESSENCES (*Concentrated infusions*) of quassia, cascarilla, chiretta, gentian, &c., may be made as directed for LIQUOR CALUMBÆ, *Pocket Formulary*. Take 8 times the quantity of ingredients directed in the Pharmacopœia for one pint of infusion, and infuse them in one pint of boiling water for the time prescribed; strain with strong pressure, and again infuse the ingredients in nearly as much water as the liquor obtained is short of a pint. Strain again with pressure; mix the products, which will measure 18 or 19 oz.; add 2 oz. of rectified spirit, set aside for a few days in a well-closed bottle, and filter. Some substances, as chiretta, senna, calumba, &c., yield their active principles to cold water, which some prefer in these cases; but it is then necessary, before adding the spirit, to place the liquor (in a bottle) in a water-bath, and heat it to 180° F., in order to precipitate any albumen it may contain. When cold, filter and add the spirit. These Essences may also be made by percolation.

VINOUS ESSENCES (by fermentation). Dr B. LANE. See LIQUORES VINOSI, *Pocket Formulary*.

EXTRACTS, MEDICINAL. See *Pocket Formulary*,

EXTRACTS OF FLOWERS. See PERFUMERY, in this volume.

EXTRACT OF MALT. Evaporate sweet wort to the consistence of treacle. See *Pocket Formulary*.

FAIRTHORN'S (Dr) MILD PROVISIONAL PILLS. Sulphate of potash 1 scruple, extract of aloes 2 scruples, extract of senna 1 scruple, compound gamboge pill 50 grs., tartarised antimony 2 grs., compound powder of scammony 12 grs., Peruvian balsam 6 grs.; in 30 pills: one, two, or more occasionally, when required.

FORD'S BALSAM OF HOREHOUND. It contains the ingredients of paregoric elixir with squills, honey, and strong infusion of horehound and liquorice.

FORD'S LAUDANUM. A tincture of opium containing cinnamon and cloves.

FOTHERGILL'S (Dr) PILLS. Diaphoretic antimony, aloes, scammony, and extract of colocynth.

FRANK'S SOLUTION. See SOLUTION OF COPAIVA.

FREEMAN'S BATHING SPIRITS. Mix water and rectified spirit of each 3 gallons; dissolve in them soft soap 6 lbs., and camphor 8 oz.; add Daffy's elixir, 8 oz.

FRIAR'S BALSAM. COMPOUND TINCTURE OF BENZOIN, B. P.

GAZOGENE POWDERS. See PULVERES EFFERVESCENTES, *Pocket Formulary.*

GELÉE POUR LE GOÛTRE. See LINIMENT. IODURETUM GELATINOSUM, *Pocket Formulary.*

GINGERBREAD, PURGATIVE. Flour 14 oz., butter 4 oz., treacle 8 oz., p. ginger $1\frac{1}{4}$ oz., jalap 2 oz., caraway $\frac{1}{2}$ oz. Mix the powders, then add the butter, and lastly the treacle, previously warmed. Roll out, and divide into cakes of $\frac{1}{2}$ oz. each, containing each 6 or 7 grs. of jalap.

GLYCERINE JELLY. Used as an application to chaps and roughened parts of the skin. It may be made of pure glycerine thickened with tragacanth powder, and scented with otto of roses. An imitation may be prepared in the following manner: Mix good soft soap $\frac{1}{2}$ dr. intimately with purified honey 2 drs., gradually add pale olive oil 5 oz., stirring without intermission until all is taken up. Care must be taken not to mix in the oil too fast. Finally, perfume as desired.—*Pharm. Journal.*

GLYCERINE PASTE. A stiff glutinous compound, recommended by Dr TILT as a basis of plaster. It is made by boiling 100 or 150 grs. of common starch in 1 oz. of glycerine. See GLYCERINUM AMYLI, B. P.

GODBOLD'S VEGETABLE BALSAM. An acidulated syrup, or oxymel, of various herbs. The following is an imitation: Dissolve by heat 1 lb. of lump sugar in white wine vinegar 1 quart, in which 3 oz. of garlic have been steeped for 3 days; add tincture of tolu 2 drs.

GODFREY'S CORDIAL. The active ingredient is opium, and there is a great diversity in the strength of the compound as prepared by different makers. Many accidents have arisen from its too general use as a stupefactive for infants, but we believe its sale is now much less encouraged by druggists than formerly. The following are some of the more usual formulæ.

1. Heat together 7 lbs. (avoird.) of treacle, and 8 lbs. of water till united; when nearly cold add the following: rectified spirit 6 fl. oz., oil of sassafras 40 minims, oil of aniseed 10 drops, laudanum 4 oz. Mix and make up the weight if necessary to 15 lbs. It contains rather mor

than 9 minims (equal, according to some authorities, to 16 or 18 drops) of laudanum in each fluid ounce.

2. Treacle $3\frac{1}{2}$ lbs., water 6 lbs., spirit of wine 8 fl. oz., laudanum 4 fl. oz., oils of aniseed, sassafras, and caraway, of each $\frac{1}{2}$ dr. Mix. Contains 12 or 14 minims of laudanum in an ounce.

3. Sliced sassafras 2 oz., opium cut small 1 oz., bruised aniseed 8 oz., boiling water a gallon. Infuse, strain, and make the infusion into a syrup with 14 lbs. of treacle. If the whole of the active principles of the opium are extracted, this is much stronger than the preceding.

4. Make a syrup with 3 lbs. (avoird.) each of treacle and coarse sugar, and water sufficient to make up a gallon. Dissolve 24 drops of oil of sassafras, and 16 of oil of aniseed, in 3 fl. oz. of spirit of wine; add 10 fl. drs. of tincture of opium, and mix the whole with 8 pints, o. m., of the syrup. This is weaker than either of the preceding, containing only 5 minims of laudanum in a fluid ounce, or 1 drop in a drachm.

5. The *Philadelphia College of Pharmacy*, to prevent the mischief arising from the varying strength of this compound, directs it to be prepared as follows: Dissolve $2\frac{1}{2}$ oz. of carbonate of potash in 26 pints of water, add 16 pints of treacle; heat together over a gentle fire till they simmer, remove the scum, and, when sufficiently cool, add $\frac{1}{2}$ oz. of oil of sassafras dissolved in 2 pints of rectified spirit, and 24 fl. oz., of tincture of opium, previously mixed. The old wine measure is here intended. It contains about 16 minims of laudanum, or rather more than 1 grain of opium in each fluid ounce.

6. Sassafras 9 oz., seeds of coriander, caraway, and anise, of each 1 oz.; infuse in 6 pints of water, simmer the mixture till reduced to 4 pints; then add 6 lbs. of treacle, boil the whole for a few minutes, and when cold, add 3 fl. oz. of tincture of opium. Nearly the strength of No. 1.—Dr PARIS.

The above forms are inserted rather with a view to show the dangerous nature of this compound than to encourage its use. No terms are sufficiently strong to express the culpability of those who would place in the hands of ignorant persons, for administration to infants and children, compounds containing opium.

GODFREY'S SMELLING SALTS. Sesquicarbonate of ammonia resublimed with pearlsh, and a little spirit.—Dr PARIS.

- GOLDEN SPIRIT OF SCURVY GRASS.** It is said to be coloured with gamboge.
- GOLDEN OINTMENT.** Singleton's ointment, q. v. The ointment of nitric oxide of mercury is also called golden ointment. See SINGLETON'S GOLDEN OINTMENT, further on.
- GOUT PAPER.** See CHARTA ANTIRHEUMATICA, *Pocket Formulary*.
- GRANVILLE'S (Dr) COUNTER-IRRITANT LOTIONS.** See LINIMENT. AMMONIÆ COMPOSITUM, *Pocket Formulary*.
- GRAVES'S (Dr) GOUT PREVENTIVE.** Orange peel 2 oz., rhubarb 1 oz., hiera picra 2 oz., brandy a quart. Digest for a week.
- GREGORY'S POWDER.** Calcined magnesia 2½ oz., powdered Turkey rhubarb 1 oz., powdered ginger ½ oz. Mix. The above is Dr GREGORY'S formula. Some recipes add powdered chamomile. Rhubarb 1 oz., ginger ¼ oz., p. chamomile ½ oz., magnesia 2 oz. Mix. Some druggists prepare it with the heavy carbonates of magnesia instead of the calcined. See PULVIS RHEI COMPOSITUS, B. P.
- GREENOUGH'S TINCTURE.** See LIQUID PREPARATIONS FOR THE TEETH AND GUMS.
- GRIFFIN'S TINCTURE,** for coughs. Oil of caraway and anise, each 2 drs., saffron ½ oz., benzoic acid ½ oz., opium 5 drs., camphor ½ oz., spirit 6 oz., honey 6 oz. When mixed and dissolved, colour with burnt sugar.
- GRIFFITH'S MIXTURE.** This is MISTURA FERRI COMPOSITA of the B. P.
- GRINROD'S (Dr) REMEDY FOR SPASMS.** Sulphuric ether, aromatic spirit of ammonia, of each ½ oz.; acetate of morphia ¼ gr., camphor mixture 2 oz. Mix. A teaspoonful in a little water when required.
- GUESTONIAN EMBROCATION.** Oil of turpentine 1½ oz., olive oil 1½ oz., dilute sulphuric acid 3 fl. drs.—Dr PARIS.
- GUTHRIE'S BLACK OINTMENT.** 10 grs. of nitrate of silver, with 1 dr. of spermaceti ointment, and 10 drops of a solution of acetate of lead.
- GUTHRIE'S EYE OINTMENT.** A weaker preparation of the same kind; used in ophthalmia, but causes great pain. See UNG. HYD. NIT., *Pocket Formulary*.
- HALFORD'S (Sir H.) GOUT PILLS.** Acetic extract of colchicum 2½ grs., Dover's powder 1½ grs., compound extract of colocynth 1½ grs., in each pill. One for a dose.

- HARROGATE SALTS** (Dr DUFFIN'S). Sulphate of magnesia 2 drs., bitartrate of potash 10 grs., sal polychrest (potassæ sulphas cum sulphure, Ph. Ed.) $\frac{1}{2}$ dr.; in a pint of warm water. For another formula see **FACTITIOUS MINERAL WATERS** and salts for producing them.
- HELVETIUS' STYPTIC**. Melt together equal parts of alum and dragon's blood; when cold, powder the compound.
- HENRY'S MAGNESIA**. A solution of Epsom salts is precipitated by one of carbonate of potash in the cold; the precipitate is well washed, rose-water being used for the last washing; it is then made up while drying into large or small cubes.
- HILL'S BALSAM OF HONEY**. Balsam of tolu 2 oz., styrax 2 drs., opium $\frac{1}{2}$ dr., honey 8 oz., spirit of wine 32 fl. oz.
- HOFFMANN'S PILLS** contain corrosive sublimate, about $\frac{1}{8}$ th of a gr. in each. See *Pocket Formulary*.
- HOLLOWAY'S PILLS**. According to DORVAULT, these pills are composed of aloes 4 grms., rhubarb 1.7 grm., pepper .45 grm., saffron and Glauber salts of each .2 grm., made into 144 pills.
- HOLLOWAY'S OINTMENT**. White wax 12 $\frac{1}{2}$ parts, yellow wax 3 parts, turpentine 3 parts, resin 25 parts, spermaceti 3 parts, lard 50 parts, olive oil 62 $\frac{1}{2}$ parts.
- HOULTON'S LAUDANUM**. Opium 2 $\frac{1}{2}$ oz., distilled vinegar 32 fl. oz.; macerate 6 days with a gentle heat, and filter. Evaporate to an extract. Dissolve this in 5 fl. oz. of rectified spirit, and 35 fl. oz. of distilled water.
- HOOPER'S FEMALE PILLS**. These, according to Dr PARIS, consist of RUFUS' Pill, sulphate of iron, canella, and a portion of ivory-black. Mr GRAY gives two formulæ:
1. Sulphate of iron 8 oz., water 8 oz.; dissolve and add Barbadoes aloes 40 oz., canella 6 oz., myrrh 2 oz., opoponax $\frac{1}{2}$ oz. Makes a mass.
 2. Sulphate of iron 2 oz., powder of aloes with canella 16 oz., mucilage of tragacanth and tincture of aloes q. s. to form a mass. Divide 60 grains into 18 pills. According to a recent analysis, the iron is in a peroxidized state; probably the sulphate is partially calcined.
- The *Philadelphia College of Pharmacy* gives the following formula:
3. Barbadoes aloes 8 oz., dried sulphate of iron 2 oz. and 1 $\frac{1}{4}$ drs., extract of black hellebore 2 oz., myrrh and

soap of each 2 oz., canella 1 oz., ginger 1 oz., water q. s. to form a mass. Divide into pills of $2\frac{1}{2}$ grs. each.

HOOPING COUGH; popular remedies for. 1. Cochineal and salt of tartar mixture. This appears to have been first introduced by Dr LOBB, in 1765, and is still a favourite domestic remedy. 2. Salt of wormwood (subcarbonate (carbonate) of potash) 20 grs., powdered cochineal 10 grs., hot water $\frac{1}{4}$ of a pint; triturate together, strain and sweeten with white sugar (or sugar candy). Dose, a teaspoonful to a tablespoonful, according to the age.

3. *Fumigating powders*. Styra^x calamita and gum benzoin, of each a scruple, placed on hot cinders or a heated shovel, in the patient's room, every day.

HUXHAM'S TINCTURE OF BARK. THE COMPOUND TINCTURE OF BARK of the London Pharmacopœia is precisely that of HUXHAM, except that he used brandy instead of proof spirit.

ISSUE PEAS. Those in general use are unripe oranges (orange berries) turned in a lathe. The unturned berries are also used. Peas are also turned from orris root. NIEMANN gives the following composition for issue peas:—Yellow wax $1\frac{1}{2}$ oz., powdered turmeric 1 oz., powdered orris $\frac{1}{2}$ oz.; Venice turpentine q. s. These are more stimulating, and are used to increase the discharge. The following, according to Dr GRAY, will open an issue itself: yellow wax 6 oz., verdigis 2 oz., white hellebore 2 oz., cantharides 1 oz., orris $1\frac{1}{2}$ oz., Venice turpentine q. s.

JAMES'S POWDER. It is not known in what respect the mode of preparing this powder differs from the Pharmacopœia process for antimonial powder. Dr JAMES'S specification is vague and impracticable.

JAMES'S ANALEPTIC PILLS. Equal parts of JAMES'S powder, RUFUS' pill, and gum guaiacum, made into pills with tincture of castor. Dr PARIS has ammoniacum in the place of guaiacum. Another formula is: compound powder of aloes, aloes and myrrh pill, and JAMES'S powder, in equal quantities, formed into pills with tincture of castor and syrup.

JARAVE, SPANISH. See CERIVISIA SARZÆ, *Pocket Formulary*.

JESUITS' DROPS. WALKER'S. Balsam of capivi 6 oz., gum guaiacum 1 oz., Chio turpentine $\frac{1}{2}$ oz., subcarbonate of

- potash $\frac{1}{2}$ oz., cochineal 1 dr., rectified spirit 1 quart. See also ELIXIR ANTIVENEREUM, *Pocket Formulary*.
- KEATING'S LOZENGES are said to be imitated as follows:—
Lactucarium 2 drs., ipecacuanha 1 dr., squill $\frac{3}{4}$ dr., ext. liquorice 2 drs., sugar 2 oz. Make into a mass with tragacanth mucilage, and divide into lozenges of 20 grs. each.
- KENNEDY'S CORN PLASTER. Yellow wax 1 lb., Venice turpentine 2 oz., verdigris 1 oz., melted together, and spread on leather.
- KEYSER'S PILLS. Acetate of mercury, manna, gum arabic, of each 1 scruple, rose-water q. s. to form a mass, for 160 pills.
- KING'S CORDIAL. Dissolve in $\frac{1}{2}$ pint of proof spirit $1\frac{1}{2}$ drs. each of the oils of caraway and cinnamon; extract the stones from 3 lbs. of black cherries, and mash the fruit in a pan; grate one nutmeg; take 2 quarts of Madeira wine, 2 quarts of brandy, and 1 gallon of syrup; mix all together, and colour with red sanders wood.
- KIRKLAND'S NEUTRAL OINTMENT AND CERATE. See UNGUENTUM PLUMBI COMPOSITUM and CERATUM NEUTRALE, *Pocket Formulary*.
- KITCHINER'S (Dr) PERISTALTIC PERSUADERS. Turkey rhubarb in powder 2 drs., oil of caraway 10 drops, simple syrup 1 dr. by weight; mix, and divide into 40 pills. Dose, 2, 3, or more. "From 2 to 4 will generally produce one additional motion within 12 hours. The best time to take them is early in the morning."
- LAMPLOUGH'S PYRETIC SALINE. Mr BANNISTER, of the Somerset House Laboratory, analysed a sample of the preparation known under this name, and states that it contained 45·7 per cent. of tartaric acid, 52·4 per cent. of bicarbonate of soda, and 1·9 per cent. of chlorate of potash.
- LARTIGUE'S GOUT PILLS. Compound extract of colocynth 20 grs., extract of colchicum 60 grs., extract of opium 1 gr.; mix, and divide into 18 pills. Dose, one or more, according to their purgative effect.
- LEDOYEN'S DISINFECTING FLUID. It consists of about 20 oz. of nitrate of lead in a gallon of water. Its specific gravity should be 1·40.
- LEECH-BITES, ASTRINGENT FOR. Dissolve 1 part of crystallised perchloride of iron in 6 parts of collodion very gradually. A drop or two of the product forms an admirable

hæmostatic. [*French Journal.*] Dry maticho leaves, rubbed to powder between the fingers, will also be found an excellent styptic for this purpose.

LEE'S WYNDHAM'S PILLS. Gamboge 3 oz., aloes 2 oz., Castile soap 1 oz., nitre $\frac{1}{2}$ oz., extract of cow-parsnip 1 oz. In pills of 5 grs. each. *Amer. Journ. of Pharmacy.*

LEE'S ANTIBILIOUS PILLS. Aloes 12 oz., scammony 6 oz., gamboge 4 oz., jalap 3 oz., calomel 5 oz., soap 1 oz., syrup of buckthorn 1 oz., mucilage 7 oz.; mix, and divide into 5-grain pills.

LEROY'S PURGATIVE.	1	2	3
Scammony.....	12 drs.	16 drs.	24 drs.
Vegetable turbith...	6 drs.	8 drs.	12 drs.
Jalap.....	6 oz.	8 oz.	12 oz.
Brandy.....	10 pints imperial.		

Digest for 12 hours, strain, and add the following syrup:

Senna.....	6 oz.	8 oz.	12 oz.
Water.....	24 oz.	32 oz.	48 oz.

Infuse, strain with pressure, and add—

Brown sugar.....	32 oz.	36 oz.	48 oz.
------------------	--------	--------	--------

Make a syrup.

No. 4 is stronger than the above.

LEJEUNE'S BALSAM FOR CHILBLAINS. Camphor 1 dr., tincture of benzoin 5 drs.; dissolve, and add iodide of potassium 5 drs., extract of lead 10 grs., spirit of wine reduced to proof with rose-water $2\frac{1}{2}$ oz.; dissolve 10 drs. of white soap in $2\frac{1}{2}$ oz. of the same diluted spirit by a gentle heat, mix the solutions whilst still warm, and add any perfume. Let it cool in wide-mouthed bottles, and cork.

LIEBERT'S COSMETIC. For chapped nipples. Dissolve 10 grs. of nitrate of lead in 1 oz. of water. A pair of fine lead shields accompany the lotion, to be worn after applying it. The nipples must be carefully washed before the child is put to the breast.

LIGNUM'S ANTI-SCORBUTIC DROPS. These contain bichloride of mercury, and should not, therefore, be used without great caution.

LIQUEUR DORÉE. Peruvian bark, bitter orange peel, and cinnamon, of each 4 drs., saffron 2 drs., brandy 4 quarts, Malaga wine 2 quarts; digest for 4 days, strain, and add $2\frac{1}{2}$ lbs. of sugar.

LIQUID BLISTER. Powdered cantharides 5 oz., and sulphuric ether 15 ounces.—TOYNBEE.

LISTON'S ISINGLASS PLASTER. Soak 1 oz. of isinglass in 2 oz. of water, and dissolve it in 2 oz. of rectified spirit and $1\frac{1}{2}$ oz. of water, by the heat of a water-bath. Brush it over the surface of oiled silk, properly stretched. An improved kind is made by brushing one side of the peritoneal membrane of the cæcum of the ox (prepared in the same manner as gold-beater's skin) with the same solution, and the other side with drying oil.

LOCATELLI'S BALSAM. Melt together 4 oz. of yellow wax, 1 lb. of common oil, and 1 lb. of Venice turpentine, placing with them 4 oz. of alkanet root wrapped in a linen bag.

LOCOCK'S WAFERS. These owe most of their activity to morphia. They should therefore be used with caution.

LONG'S (ST JOHN) LINIMENT. See *LINIMENTUM TEREBINTHINÆ ACETICUM*, *Pocket Formulary*.

LOZENGES. See *CHING'S LOZENGES*, *DAWSON'S*, *DARCET'S*.

The medicated Lozenges which are sanctioned by different Pharmacopœias, and employed in practice, will be found under *TROCHISCI* and *PASTA*, in the *Pocket Formulary*. A few other formulæ are here added:

Absorbent Lozenges. Precipitated chalk 3 oz., heavy carbonate of magnesia 2 oz., nutmeg in fine powder 1 dr., sugar 12 oz., powdered gum 1 oz., water q. s. to form a stiff paste, which divide by a punch into lozenges of the usual size, and dry them gradually in a warm room.

Aperient Lozenges. Calomel 60 grs., pure scammony 80 grs., jalap 40 grs. (or jalapine 4 grs.), ginger 8 grs., cinnamon 4 grs., mucilage of tragacanth q. s. to form a stiff paste; mix the other powders accurately together, then with sugar q. s. to make lozenges of $\frac{1}{2}$ dr. each, lastly add the mucilage, beat the whole into a uniform mass, and divide it into 40 equal lozenges. Each contains $1\frac{1}{2}$ grs. calomel, 2 of scammony, and 1 of jalap.

Black Currant Paste. Soften 12 lbs. of picked black currants by heating them in a water-bath in a covered earthen vessel, pulp through a hair sieve, and evaporate to a paste, incorporating with it 1 lb. powdered sugar; roll it out into a sheet of proper thickness. Mr BARTLETT gives the following formula: 3 lbs. of powdered sugar, 3 lbs. of extract of black currants (the inspissated juice), 1 oz. of tartaric acid, 6 oz. of powdered gum; mixed,

rolled out, and cut, when dry, with a large pair of scissors into square pieces.

Black Currant and Ipecacuanha Lozenges. Black currant paste (as above) 8 oz., ipecacuanha 30 grs., tragacanth 90 grs.; in 240 lozenges.

Cough Lozenges (with Lactucarium). Powdered lactucarium 2 drs., extract of liquorice root 12 drs., ipecacuanha 30 grs., powdered squill 15 grs., refined sugar 6 oz., mucilage of tragacanth q. s., mix, and divide into 240 equal lozenges. Each contains $\frac{1}{2}$ gr. lactucarium, $\frac{1}{2}$ gr. of ipecacuanha, $\frac{1}{8}$ gr. of squill.

For other Cough Lozenges, see TROCHISCI ANTICATAR-
RHAELES, CUBEBAE, GLYCERHIZAE ET OPII, LACTUCE,
MORPHLE ET IPECAC., OPII, PAPAVERIS, SCILLÆ, TOLU-
TANI, &c., *Pocket Formulary*.

Digestive or Live-long Candy. 1. Powdered rhubarb 60 grs., heavy magnesia 1 oz., bicarbonate of soda 1 dr., finely-powdered ginger 20 grs., cinnamon powder 15 grs., powdered white sugar 2 oz., mucilage of tragacanth q. s.; beat together, and divide into parallelograms of 20 grs. each.

2. *Caraway Candy.* Rhubarb 60 grs., powdered cara-
ways 60 grs., oil of caraway 10 drops, ginger and cinna-
mon, each 15 grs., magnesia 6 drs., carbonate of soda 1 dr.,
sugar 2 oz., mucilage q. s.—as the last.

Edinburgh Lozenges. Extract of poppies 2 oz., powdered sugar 8 oz., powdered tragacanth 4 oz., water q. s.

Fruit Lozenges. Black currant paste 8 oz., red currant
paste (or the juice evaporated to a paste) 4 oz., syrup of
raspberries 4 oz., soften by a gentle heat, and beat in a
warm mortar with 2 lbs. of powdered sugar, and a drachm
of powdered citric acid, and, if required, a little mucilage
of gum tragacanth.

Marshmallow Lozenges. Marshmallow root powdered 2 oz.,
sugar 14 oz., mixed with some mucilage of tragacanth,
and orange-flower water.

Peppermint Lozenges. Rub together white sugar 6 oz., oil
of peppermint 36 drops, and the whites of two eggs.
Make into lozenges. See *Pocket Formulary*.

LYNCH'S EMBROCATION. Olive oil coloured with alkanet,
perfumed, and rendered stimulating by essential oils.

MADDEN'S ESSENCE. A strong acidulated infusion of roses.

MAHOMED'S ELECTUARY. Grocer's currants 1 oz., powdered
senna $\frac{1}{2}$ oz., powdered ginger 30 grs., oil of croton 1 drop,

- syrup of roses sufficient to make an electuary; two teaspoonfuls every morning.—BATEMAN.
- MAHY'S PLASTER** (American). Boil 12 oz. of white lead, 32 fluid oz. of olive oil, and a little water, stirring constantly until incorporated. Add yellow wax 4 oz., lead plaster 18 oz., and when these are melted stir in 9 oz. of powdered orris.
- MAREDANT'S NORTON'S DROPS.** Corrosive sublimate, gentian, ginger, and cochineal.
- MARSHALL'S CERATE.** Palm oil 5 oz., calomel 1 oz., acetate of lead $\frac{1}{2}$ oz., ointment of nitrate of mercury 2 oz.; mix.—Dr PARIS.
- MARSHALL'S EYE-DROPS.** These are said to consist of 2 grs. of nitrate of silver in 1 oz. of decoction of snails.
- MARSDEN'S ANTISCORBUTIC DROPS, MORTON'S, PERRY'S, LIGNUM'S,** and other antiscorbatic drops, contain corrosive sublimate.—Dr PARIS.
- MAGNESIA FLUID.** A solution of carbonate of magnesia in water by means of carbonic acid gas, forced into it by pressure. MURRAY'S and DINNEFORD'S should contain from 12 to 15 grs. of the carbonate in each fluid oz. See LIQUOR MAGNESIÆ CARBONATIS, B., *Pocket Formulary*.
- MATHIEU'S VERMIFUGE.** Tin filings 1 oz., fern root $\frac{3}{4}$ oz., worm seed $\frac{1}{2}$ oz., resinous extract of jalap 1 dr., sulphate of potash 1 dr., honey to form an electuary. A teaspoonful every 3 hours for 2 days; then substitute the following—jalap 2 scruples, sulphate of potash 2 scruples, scammony 1 scruple, gamboge 10 grs.; made into an electuary with honey, and given in the same dose.
- MEXICAN HAIR RENEWER.** Acetate of lead 1 part, precipitated sulphur 4 parts, glycerine 30 parts, water 160 parts.
- MINERAL WATERS (FACTITIOUS), AND SALTS FOR PRODUCING THEM.** See further on.
- MONTBAIN'S BARÈGE BALLS** (for Sulphur Baths). Sulphide of lime 8 oz., common salt 2 oz., Flanders glue 1 oz., extract of soapwort 1 oz.; make into 8 balls; to be kept from the air. M. MENIÈRE recommends, extract of soapwort $\frac{1}{2}$ oz., water 6 oz., lime in powder 4 oz., sulphur 3 oz., gelatine 1 oz.; dissolve the extract and gelatine in the water, add the lime and sulphur, heat gently, stirring it constantly, till the mass gets detached from the sides of the vessel; then form it into balls of $1\frac{1}{2}$ oz. each.

MORPHIA COLLODION. One part of hydrochlorate of morphia to 30 parts of flexible collodion. Applied with a camel's-hair brush, for neuralgic pains.

MORRISON'S PILLS. No. 1, consist of equal parts of aloes and cream of tartar; No. 2, consist of 2 parts of gamboge, 3 of aloes, 1 of colocynth, and 4 of cream of tartar, made into pills with syrup.

MORRISON'S ADHESIVE PASTE, for ringworm. See PASTA ADHESIVA, *Pocket Formulary*.

MOSELEY'S PILLS. Turkey rhubarb 60 grs., Jamaica ginger 24 grs., syrup and tincture of rhubarb q. s. to form a mass, to be divided into 24 pills.

MOXON'S EFFERVESCING MAGNESIAN APERIENT. The following have been proposed as imitations:

1. Heavy carbonate of magnesia 2 lbs., bicarbonate of soda 1 lb., tartaric acid 1½ lbs., refined sugar ½ lb., essence of lemons 40 minims; the powders to be all separately dried at a moderate temperature.

2. Sulphate of magnesia 1 lb., bicarbonate of soda 1 lb., tartaric acid ½ lb.; the ingredients to be well dried separately, at a moderate temperature. *Pharmaceutical Journal*.

3. Carbonate of magnesia 1 lb., sulphate of magnesia 2 lbs., bicarbonate of soda 2 lbs., potassio-tartrate of soda 2 lbs., tartaric acid 2 lbs.; to be separately dried, and mixed.—M. DURANDE.

MUNRO'S COUGH MEDICINE. 4 drs. of paregoric with 2 drs. of sulphuric ether, and 2 drs. of tincture of tolu. Dose, a teaspoonful in some warm water.

MURRAY'S (Sir J.) FLUID CAMPHOR. Each oz. contains 3 grs. of camphor and 6 grs. of carbonate of magnesia, dissolved by carbonic acid, and by pressure.

MURRAY'S GOUT SPECIFIC. It contains iodide of potassium, sulphate of magnesia, and an aromatic tincture. *Pharmaceutical Journal*.

MUSTARD LEAVES. See RIGOLLOT'S MUSTARD LEAVES.

MUSTARD TISSUE. See SINAPINE TISSUE.

NEPENTHE. Supposed to be a watery solution of opium, resembling BATTLE'S. See EXTRACTUM OPII LIQUIDUM, and LIQUOR OPII SEDATIVUS, *Pocket Formulary*.

NEURALINE. An anodyne application, for external use only. It is said to contain aconite and chloroform.

- NORRIS'S DROPS.** A solution of tartarized antimony, with a tincture of some vegetable substances, not ascertained.
- NOUFFLEUR'S (MADAME) WORM MEDICINE.** Powdered fern root 3 drs., to be given in the morning (the patient being prepared by an emollient clyster, and a supper of panada); followed in 2 hours by a bolus of calomel, scammony, and gamboge.
- OLLIVIER'S BISCUITS.** Beat up the whites of 2 eggs with 16 oz. of water, add a solution of 76 grs. of corrosive sublimate; collect, wash, and dry the precipitate, 1-7th of a gr. of which is contained in each biscuit of 2 drs.
- OPODELDOC.** *Lin. Saponis.*
- ORMSKIRK MEDICINE,** to prevent hydrophobia. Elecampane 1 dr., chalk 4 drs., Armenian bole 3 drs., alum 10 grs., oil of aniseed 5 drops.
- PARAMOUD.** See DIETETIC COMPOUNDS.
- PALMER'S ABBATED CHALYBEATE.** Mix 1½ parts of acetic acid with 40 of water, add 4 of proto-sulphate of iron, and 20 of syrup. Put into 4-oz. bottles, for No. 1 and No. 2 respectively, as much of the above as contains 2 and 4 grs. of sulphate of iron, and fill the bottles with a solution of carbonate of soda or of potash strongly charged with carbonic acid gas. Tartaric acid may be substituted for acetic.
- PAPIER EPISPASTIQUE D'ALBESPEYRES.** The *Pommade Epispastique* of the French Codex, spread on waxed paper. See UNGUENTUM EPISPASTICUM, *Pocket Formulary.*
- PAPIER EPISPASTIQUE DE VÉE.** This is of three strengths, distinguished by the colours white, green, and red. The composition is made by boiling cantharides for an hour with water, and lard, green ointment, or lard coloured with alkanet; adding white wax to the strained fats, and spreading on paper, silk, or linen. No. 1 is made with 10 oz. of cantharides to 4 lbs. of lard; No. 2, of 1 lb. of flies to 8 lbs. of green ointment; and No. 3, of 1½ lbs. to 8 lbs. of coloured lard; and to each are added 2 lbs. of white wax.—DORVAULT. See BLISTERING PAPER.
- PAPIER FAYARD.** *Gout Paper.* Euphorbium 3 drs., cantharides 6 drs., powdered and digested with 4 oz. alcohol; and 3 drs. Venice turpentine added to the strained tincture. Fine paper is dipped into it and dried in the air. MOHR directs 4 drs. of cantharides and 1 dr. euphorbium to be digested in 5 oz. of highly rectified spirit; filter and add

1½ oz. Venice turpentine previously liquefied with 2 oz. of resin. To be spread on the paper while warm.

PELLETIER'S ÆTHEREAL OPODELDOC. See BALSAMUM ACETICUM CAMPHORATUM, *Pocket Formulary*.

PERRY'S BALM OF LYNACUM. See BALM OF SYRIACUM.

PERSIAN INSECT POWDER. This is said to be the dried flowers of the *Pyrethrum roseum*, or red flowered Pyrethrum reduced to powder. The flowers of the *Pyrethrum caucasicum* are also stated to be used for the same purpose.

PETER'S PILLS. Aloes, jalap, gamboge, and scammony, of each 2 drs.; calomel 1 dr.

PILES, popular remedies for. Dr WARDLEWORTH'S PILLS contain 3½ grs. of pitch in each; 2 every night. For ELECTUARIES for the piles see CONF. SENNÆ, B. P., CONFECTIO RESINÆ, CONFECTIO SULPHURIS, ELECTUARIUM HÆMORRHOIDALE, all in *Pocket Formulary*. See also WARD'S PASTE, further on. For PILE OINTMENTS, see UNGUENTUM GALLÆ, UNGUENTUM GALLÆ CUM OPIÔ, UNGUENTUM HÆMORRHOIDALE, *Pocket Formulary*. Sir H. HALFORD'S Pile Ointment consisted of equal parts of citrine ointment and oil of almonds triturated in a glass mortar till perfectly smooth. Mr WARD'S is—Powdered nut-gall 2 drs., camphor 1 dr., melted wax 1 oz., tincture of opium 2 drs. Mix.

PILLS. See proprietors' names in alphabetical order. A great variety of formulæ for pills of every kind will be found in the *Pocket Formulary*.

PILLS, TO COAT WITH GELATINE . } See PILULÆ, *Pocket*
 —, TO SILVER } *Formulary*.

M. DURDEN recommends collodion as a covering for pills; others, a solution of gutta percha in chloroform; but the ready solubility of these materials in the stomach may be questioned. M. BLANCHARD uses balsam of tolu dissolved in ether. Mr BAILDON recommends chloroform instead of ether.

PLASMA. SCHACHT. See GLYCERINUM AMYLI, B. P.

PLUNKET'S OINTMENT FOR CANCER. See CAUSTICUM ANTICANCROSUM, *Pocket Formulary*.

POMADE DIVINE. Beef marrow 3 lbs.; put it into an earthen vessel, and cover it with cold water, and change the water daily for a few days, using rose-water the last day. Pour off and press out the water; add to the mar-

- row 4 oz. each of styrax, benzoin, and Chio turpentine, 1 oz. orris powder, $\frac{1}{2}$ oz. each of powdered cinnamon, cloves and nutmeg. Set the vessel in hot water, and keep the water boiling for 3 hours; then strain. For Pomades for the Hair, see HAIR COSMETICS.
- POND'S EXTRACT. An American proprietary preparation chiefly consisting of the witch hazel (*Hamamelis Virginica*).
- PORTLAND'S (DUKE OF) GOUT POWDERS. Equal quantities of the roots of gentian and birthwort, tops of germander, ground pine, and lesser centaury: all to be powdered and mixed together.
- POWELL'S BALSAM FOR COUGH. Mix together 2 drs. of syrup of tolu, 1 oz. of paregoric elixir, and 2 oz. of liquorice-juice.
- QUEEN OF HUNGARY'S WATER. Tops and flowers of rosemary 2 lbs., rectified spirit 3 lbs.; digest in a close vessel for 50 hours in a gentle heat, then distil by water-bath.
- RADCLIFFE'S ELIXIR. Aloes 6 drs., cinnamon, zedoary, and cochineal, each $\frac{1}{2}$ dr., rhubarb 1 dr., syrup of buckthorn 2 oz., proof spirit 16 fl. oz., water 5 fl. oz.—Dr PARIS. According to GRAY, it contains jalap, scammony, and senna.
- RASPAIL'S CAMPHOR CIGARETTES. These are merely camphor enclosed in a tube (a quill or paper tube may be used), confined by blotting paper, and used cold. Another kind of camphorated cigars is made by saturating dried colts-foot or other leaves with a strong solution of camphor, and rolling them in the form of cigars.
- REECE'S CHIRETTA PILLS. Extract of chiretta 2 drs., dried carbonate of soda 1 scruple, p. ginger 15 grs. Mix, and divide into 36 pills. Two twice a day.
- REGNAULD'S PECTORAL PASTE. Pectoral flowers (mullein, coltsfoot, catsfoot, and red poppies mixed) 16 oz., boiling water 3 lbs.; infuse, strain, and add to the clear liquid 6 lbs. of clean gum arabic; dissolve by a gentle heat, and having added 4 lbs. of white sugar, evaporate to a proper consistence, adding towards the end, 6 drs. of tincture of balsam of tolu.
- REYNOLDS' GOUT SPECIFIC. It is supposed to be a wine of colchicum.
- REVALENTA. It is said to be prepared from the seeds of the *Eryum lens*. See DIETETIC ARTICLES.

RIGA BALSAM FOR BRUISES. Mix 4 oz. of spirits of wine with 1 dr. of compound tincture of benzoin, and 2 drs. of tincture of saffron.

RIGOLLOT'S MUSTARD LEAVES. An admirable and cleanly substitute for mustard poultices. They appear to consist of flour of mustard attached to paper by some glutinous material. See *CHARTA SINAPIS*, B. P.

ROBINSON'S (Dr) STIMULATING PURGATIVE PILLS. Watery extract of aloes 1 dr., balsam of Peru 10 grs., oil of caraway 10 drops, scammony $\frac{1}{2}$ dr. Mix, and divide into 20 pills: 2 or 3 when required.

ROCHE'S EMBROCATION. Olive oil, with half its weight of oil of cloves and oil of amber.—Dr PARIS.

ROGÉ'S MAGNESIAN PURGATIVE.—Calcined magnesia 1 oz., carbonate of magnesia $\frac{1}{2}$ oz., citric acid $3\frac{1}{4}$ oz., sugar, rubbed with a few drops of essence of lemon, 6 $\frac{1}{4}$ oz. To form Aerated Magnesian Lemonade, put $\frac{1}{4}$ oz. of the powder into a soda-water bottle nearly filled with water, and immediately cork it securely.

ROSSITTER'S HAIR RESTORER. Rose-water 345 parts, glycerin 50 parts, milk of sulphur 2 parts, sugar of lead $1\frac{1}{2}$ parts.—HAGEE.

RUBINI'S CAMPHOR. A solution of camphor in its own weight of 60 o. p. spirit. *Pharmaceutical Journal*.

ROUSSEAU'S DROPS. See *VINUM OPII FERMENTATIONE PARATUM*, *Pocket Formulary*.

RUPINI'S STYPTIC. It contains (according to Dr A. T. Thomson) gallic acid, sulphate of zinc, spirit, and rose-water.

RYAN'S ESSENCE OF COLTSFOOT. Tincture of balsam of tolu 2 oz., compound tincture of benzoin 2 oz., spirit of wine 4 oz.—GRAY.

RYMER'S TINCTURE. A tincture of capsicum, camphor, cardamom, rhubarb, aloes, and castor, in proof spirit, with a small quantity of sulphuric acid.—Dr PARIS. The inventor states that it is impregnated with an aerial acid.

SALTS, MINERAL. See *WATERS, FACTITIOUS MINERAL*.

SCOTT'S PILLS. See *ANDERSON'S PILLS*.

SCOTT'S PLASTER. This appears to be a carefully prepared Emp. Plumbi, spread on calico. If it contain resin, the quantity is probably less than in Emp. Resinæ.

SEIDLITZ POWDERS (in separate powders). One contains

2 drs. of powdered Rochelle salts, and 40 grs. of bicarbonate of soda; the other powder is p. tartaric acid 35 grs.

SEIDLITZ POWDER, in one bottle. *Note*.—The powders are all to be thoroughly dried separately, at a gentle heat—the potassio-tartrate of soda (*Soda tartarata*) at a temperature not exceeding 110° F.; the others not higher than 120°. Take of potassio-tartrate of soda, dried, 15 oz., tartaric acid, dried, 5 oz. (or citric acid 4½ oz.), dry bicarbonate of soda 6 oz. Mix, and keep in a well-closed bottle. Dose, 3 drs. Keep dry. The above have no resemblance to the natural water of Seidlitz. See **MINERAL WATERS**.

SINAPINE TISSUE. A substitute for mustard poultices. Sheets of paper impregnated with essence of mustard and tincture of capsicum.

SINGLETON'S GOLDEN OINTMENT. Orpiment mixed with lard to the consistence of an ointment.

“There appears to be some mistake in this statement, as that sold us under this name had nearly the same composition as the ointment of nitric oxide of mercury of the *Pharmacopœia*. It did not contain a trace either of arsenic or sulphur.”—COOLEY.

SODA POWDERS. These usually contain in one paper 30 grs. of bicarbonate of soda, and in the other 25 grs. of tartaric acid (or 24 grs. of citric acid). For **SHERBET**, **LEMONADE**, and **GINGER-BEER POWDERS**, see **BEVERAGES**.

SMELLOME'S EYE-OINTMENT. Prepared verdigris 30 grs.; levigate with 30 drops of olive oil, and add 1 oz. of resinous cerate.

SPEEDIMAN'S PILLS. Rhubarb, aloes, myrrh, and extract of chamomile, of each 60 grs.; oil of chamomile 12 drops. Mix, and divide into 4-grain pills.

SOLOMON'S BALM OF GILEAD. An aromatic tincture, of which cardamoms form a leading ingredient, made with brandy.—Dr PARIS. It is thought to contain cantharides.

SOLOMON'S ANTI-IMPETIGINES is said to be a solution of corrosive sublimate.

SMITH'S (Dr HUGH) STOMACHIC PILLS. Aloes, rhubarb, aromatic powder, gum sagapenum, of each 1 dr.; oil of mint and oil of cloves, of each 10 drops; balsam of Peru q. s. In 5-grain pills; 2 to 4 every night.

SOLUTION OF COPAIVA. See SOL. COPAIBE ALKALIN.
Pocket Formulary.

SPILSBURY'S ANTI-SCORBUTIC DROPS. Corrosive sublimate 2 drs. (not 2 oz., as misprinted in the eighth edition of Dr PARIS's Pharmacologia), precipitated sulphuret of antimony 1 dr., gentian 2 drs., orange-peel 2 drs., red sanders 1 dr., proof spirit 16 fl. oz.; digest and strain.—Dr PARIS. We are informed that this is incorrect. Another formula is—Levigated crocus metallorum 18 drs., corrosive sublimate 135 grs., red sanders 1½ drs., gentian 6 drs., orange-peel 6 drs., brandy 48 fl. oz.; digest for 10 days, shaking frequently, and strain; dose, 5 to 20 drops.

SQUIRE'S ELIXIR. Opium 1 oz., camphor 1 oz., spirit of aniseed (compound) 4 pints, tincture of serpentaria 1 pint, water 4 pints, tincture of ginger ½ oz. Some recipes add a little aurum musivum.

STANDERT'S RED MIXTURE. Carbonate of magnesia 4 drs., powdered rhubarb 2 drs., tincture of rhubarb 1½ oz., tincture of opium 1 dr., oil of aniseed 24 drops, essence of peppermint 30 drops, water 1½ pints; mix. A popular remedy for bowel complaints in the West of England.

STANDERT'S STOMACHIC CANDY. Cardamom seed, ginger, rhubarb (all in fine powder), each 4 drs., lump sugar 4 oz., water 6 drs.; boil together, stirring constantly till the sugar is dissolved, then pour it into a proper mould.

STEEDMAN'S SOOTHING POWDERS. These appear, by analysis, to contain calomel, about 1 gr. in each, also a trace of morphia, with sugar.

STEER'S OPODELDOC. 1. Rectified spirit a quart, Castile soap 5 oz., camphor 2½ oz., oil of rosemary 2½ drs., oil of origanum 5 drs., weaker sol. ammonia 4 oz.; digest till dissolved, and pour while warm into wide-mouthed bottles.

2. Rectified spirit 8 pints o. m., white soap 20 oz., camphor 8 oz., water of ammonia 4 oz., oil of rosemary 1 oz., oil of horsemint 1 oz.; dissolve the soap in the spirit by a gentle heat, and add the other ingredients. Bottle whilst warm.—*Phil. Coll. of Pharmacy.*

STOUGHTON'S ELIXIR. 1. Gentian 36 oz., serpentaria 16 oz., dried orange-peel 24 oz., calamus aromaticus 4 oz., rectified spirit and water, of each 6 gallons, o. m.

2. Gentian 4 lbs., orange-peel 2 lbs., cochineal 2 drs., cardamom seed 1 oz., rectified spirit 8 gallons.
- STOREY'S WORM CAKES. Calomel 1 scruple, jalap 1 dr., ginger 2 scruples, sugar 1 oz., cinnabar to colour, syrup q. s. to form 10 cakes.
- STRUVE'S LOTION FOR HOOPING-COUGH. Emetic tartar 60 grs., water 2 oz., tincture of cantharides 1 oz.
- SWAIM'S VERMIFUGE. Worm seed 2 oz., valerian, rhubarb, pink root, white agaric, of each 1½ oz.; boil in sufficient water to yield 3 quarts of decoction, and add to it 30 drops of oil of tansy, and 45 drops of oil of cloves, dissolved in a quart of rectified spirit.—*American remedy.*
- SYDENHAM'S LIQUID LAUDANUM. See VINUM OPII, *Pocket Formulary.*
- TANJORE PILLS. See PILULE ARSENICI, *Pocket Formulary.*
- THIBAUT'S BALSAM for wounds. Digest flowers of St John's wort, one handful, in ½ pint of rectified spirit, then express the liquor, and dissolve in it myrrh, aloes, and dragon's blood, of each 1 dr., with Canada balsam ½ oz.
- TISSOT'S PUGRATIVE POWDERS. Jalap, rhubarb, senna, and soluble cream of tartar, equal quantities. Dose, 2 to 6 drs. Used in Normandy.
- TURLINGTON'S BALSAM. Rectified spirit 8 old wine pints, benzoin 12 oz., liquid styrax 4 oz., socotrine aloes 1 oz., balsam of Peru 2 oz., myrrh 1 oz., angelica-root ½ oz., balsam of tolu 4 oz., extract of liquorice 4 oz.; digest 10 days and strain.—*Phil. Coll. of Pharm.* The certified copy of the original recipe is more complex, containing three times as many ingredients.
- VALANGIN'S SOLUTION OF SOLVENT MINERAL. Arsenious acid (which has been mixed with chloride of sodium, and resublimed) 30 grs., hydrochloric acid 90 grs., distilled water 1 oz.; dissolve, and add distilled water to make up 30 fl. oz. Dose, from 1 drop, increased very gradually to 3. See LIQUOR ARSENICI HYDROCHLORICUS (L.), *Pocket Formulary.*
- VENLO'S VEGETABLE SYRUP. It is supposed to be a decoction of burdock, mint, dandelion, senna, &c., boiled with sugar, and a small portion of solution of sublimate added.
- WALKER'S JESUIT'S DROPS. See JESUIT'S DROPS.

WARBURG'S FEVER TINCTURE. See TINCT. WARBURGII, *Pocket Formulary*.

WARD'S PASTE. The same as CONFECTIO PIPERIS NIGRI of the London Pharmacopœia.

WARD'S WHITE DROPS. To 16 oz. of strong nitric acid add gradually 7 oz. of subcarbonate of ammonia; let it stand 2 or 3 hours; then put it into a bolt-head which it will only half fill, and to each 16 oz. put 4 oz. of pure quicksilver, and digest by a sand heat till the solution is complete; then gently increase the heat, and add a little more quicksilver at intervals till it will dissolve no more; then evaporate it in a glass or earthen dish placed in sand, till a pellicle appears, and set it aside to crystallize. Dissolve 1 lb. of the drained salt in 3 lbs. of rose-water by the heat of a sand-bath.

WARD'S ESSENCE FOR THE HEADACHE. Spirit of wine 2 lbs., roche alum in fine powder 2 oz., camphor 4 oz., essence of lemon $\frac{1}{2}$ oz., strong water of ammonia 4 oz.; stop the bottle close, and shake it daily for 3 or 4 days.

WARD'S RED PILL. Glass of antimony levigated with a fourth of its weight of dragon's-blood, made into a mass with wine, and divided into pills of a grain and a half each; one pill is a dose, on an empty stomach. In foulness of the stomach and bowels, and obstinate rheumatic disorders.

WARD'S DROPSY PURGING POWDER. Jalap 1 lb., cream of tartar 1 lb., red bole 1 oz.; mix; dose, from 30 to 40 grs., in broth or warm beer, repeated for 2 or 3 days, or oftener if necessary.

WARD'S SWEATING POWDER. Similar to DOVER'S POWDER.

WARD'S WASHING POWDER. See WASHING COMPOUNDS, TRADE CHEMICALS.

WARNER'S CORDIAL. Rhubarb 1 oz., senna $\frac{1}{2}$ oz., saffron 1 dr., liquorice $\frac{1}{2}$ oz., raisins 1 lb., brandy 3 pints; digest for a week, and strain.

WARTS, TO CURE. Strong acetic acid, or dichloroacetic acid, applied in tubes made for the purpose. Caution is required, that the sound flesh may not be involved. Equal parts of chromic acid and water. The skin around the wart should first be protected by painting it with oil.

WARWICK'S (COUNTESS OF) POWDERS. Scammony 2 oz., calx of antimony 1 oz., cream of tartar $\frac{1}{2}$ oz.; mix.

WEBSTER'S DIET DRINK. A decoction of syrup of sarsa-

- parilla, betony, dulcamara, guaiacum, liquorice, sassafras, turmeric, and thyme.
- WEBSTER'S (LADY) PILLS. See PILULA ALOES CUM MASTICHE, *Pocket Formulary*.
- WHITEHEAD'S ESSENCE OF MUSTARD. See ESSENCE OF MUSTARD for an imitation of it.
- WHITEHEAD'S MUSTARD PILLS. Dr PARIS says they consist of balsam of tolu and resin.
- WHITELAW'S ETHEREAL TINCTURE OF LOBELIA. See TINCTURA LOBELIÆ ETHEREA, *Pocket Formulary*.
- WILSON'S GOUT TINCTURE. A vinous infusion of colchicum.
- WINSLOW'S (MRS) SOOTHING SYRUP. Consists of sugar syrup strongly flavoured with an alcoholic tincture of fennel, anise, and a little caraway seed; or an alcoholic solution of their essential oils, and with or without an admixture of solution of sulphate of morphine in varying quantity.—*American Popular Health Almanac*.
- WISDOM'S (DR) EYE-WATER. Bole 2 oz., sulphate of zinc $\frac{1}{4}$ oz., camphor (dissolved in $1\frac{1}{4}$ oz. of rectified spirit) $\frac{1}{4}$ oz., water a gallon.
- WRIGHT'S PEARL OINTMENT. White precipitate 8 oz., extract of lead a pint; rub together and add 7 lbs. of white wax melted with 16 lbs. of olive oil.—*Pharm. Journal*.
- WORM LOZENGES. See CHING'S LOZENGES, and STOREY'S WORM CAKES; see also TROCHISCI ANTHELMINTICI, and TROCHISCI SANTONINI, *Pocket Formulary*.
- YOUNG'S PURGING DRINK. Carbonate of soda in crystals $2\frac{1}{2}$ drs., cream of tartar in crystals 3 drs., water 8 oz.; put it into a stone bottle, and secure the cork.

FACTITIOUS MINERAL WATERS,

AND

Salts for Producing them

AERATED OR CARBONATED WATERS

THESE require the aid of the powerful machine employed by soda-water manufacturers, to charge the waters strongly with carbonic acid gas. The gas is made from whiting and diluted sulphuric acid, and is forced by a pump into the watery solution. Sometimes the gas is produced by the mutual action of the ingredients introduced into the bottle of water, which must be instantly closed; but this method is found practically inconvenient, and is only adopted in the absence of proper apparatus. The quantity of gas introduced is directed, in the United States Pharmacopœias of 1870, to be 5 times the volume of liquid. The British and French Pharmacopœias direct the water to be charged with carbonic acid under a pressure of 4 atmospheres. For chalybeate and sulphuretted waters, the water should be previously deprived of the air it naturally contains, by boiling, and allowing it to cool in a closed vessel.

There are various manufacturers of aerated-water machines, and of syphon bottles for holding these waters when made. The names and addresses of these makers may be found in any trade directory.

SIMPLE AERATED WATER. *Carbonic Acid Gas Water.*
Water charged with five or more volumes of carbonic acid gas, as above.

ALKALINE AERATED WATERS. Aerated soda and potash waters should be made by dissolving a drachm of the carbonated alkali in each pint of water, and charging it strongly with carbonic acid gas. But the soda water of the shops generally contains little, or no soda.

AERATED MAGNESIA WATER. This is made of various strengths.

MURRAY'S AND DINNEFORD'S FLUID MAGNESIA may be thus made:—To a boiling solution of 16 oz. of sulphate of magnesia in 6 pints of water, add a solution of 19 oz. of crystallised carbonate of soda in the same quantity of water; boil the mixture till gas ceases to escape, stirring constantly; then set it aside to settle; pour off the liquid, and wash the precipitate on a cotton or linen cloth, with warm water, till the latter passes tasteless. Mix the precipitate, without drying it, with a gallon of water, and force carbonic acid gas into the mixture under strong pressure, till a complete solution is effected. See **LIQUOR MAGNESIÆ CARBONATIS, B. P.**

CARBONATED LIME WATER. *Carrara Water.* Lime-water (prepared from lime made by calcining Carrara marble) is supersaturated by strong pressure, with carbonic acid; so that the carbonate of lime at first thrown down is redissolved. It contains 8 grs. of carbonate of lime in 10 fl. oz. of water.

AERATED LITHIA WATER. This may be conveniently made from the freshly precipitated carbonate, dissolved in carbonated water, as directed for fluid magnesia. See **LIQUOR LITHIÆ EFFERVESCENS, B. P.**

SALINE CARBONATED WATERS

The following afford approximate imitations of these waters: The earthy salts, with the salts of iron, should be dissolved together in the smallest quantity of water. The other ingredients to be dissolved in the larger portion of the water, and the solution impregnated with the gas. The first solution may be then added or be previously introduced into the bottles. The salts, unless it is otherwise stated, are to be crystallised.

BADEN WATER. Chloride of magnesium 2 grs., chloride of calcium 40 grs., perchloride of iron $\frac{1}{4}$ gr. (or 3 minims of the tincture), chloride of sodium 30 grs., sulphate of soda 10 grs., carbonate of soda 1 gr., water 1 pint, carbonic acid gas 5 volumes.

CARLSBAD WATER. Chloride of calcium 8 grs., tincture of chloride of iron 1 drop, sulphate of soda 50 grs., carbonate of soda 60 grs., chloride of sodium 8 grs., carbonated water 1 pint.

EGER. Carbonate of soda 5 grs., sulphate of soda 4 scruples,

chloride of sodium 10 grs., sulphate of magnesia 3 grs., chloride of calcium 5 grs., carbonated water a pint. (Or it may be made without apparatus thus:—Bicarbonate of soda 30 grs., chloride of sodium 8 grs., sulphate of magnesia 3 grs., water a pint; dissolve and add a scruple of dry bisulphate of soda, and close the bottle immediately.)

EMS. Carbonate of soda 2 scruples, sulphate of potash 1 gr., sulphate of magnesia 5 grs., chloride of sodium 10 grs., chloride of calcium 3 grs., carbonated water a pint.

FRIEDRICHSHALL WATER. (LIEBIG.) Sulphate of soda 46½ grs., sulphate of magnesia 39½ grs., chloride of sodium 61 grs., chloride of magnesium 30 grs., bromide of magnesium 1-3rd of gr., sulphate of potash 1½ grs., sulphate of lime 10 grs., carbonate of lime 1-10th of gr., carbonate of magnesia 1 gr., carbonated water 16 oz.

HUNYADI JANOS WATER. (LIEBIG.) Sulphate of magnesia 123 grs., sulphate of soda 122 grs., chloride of sodium 10 grs., bicarbonate of soda 6 grs., carbonate of lime 7 grs., carbonated water 16 oz.

MARIENBAD. Carbonate of soda 2 scruples, sulphate of soda 96 grs., sulphate of magnesia 8 grs., chloride of sodium 15 grs., chloride of calcium 10 grs., carbonated water a pint. Or, bicarbonate of soda 50 grs., sulphate of soda 1 dr., chloride of sodium 15 grs., sulphate of magnesia 10 grs.; dissolve in a pint of water, add 25 grs. of dry bisulphate of soda, and cork immediately.

MARIENBAD PURGING SALTS. Bicarbonate of soda 5 oz., dried sulphate of soda 12 oz., dry chloride of sodium 1½ oz., sulphate of magnesia, dried, 2 oz., dried bisulphate of soda 2½ oz. Mix the salts, previously dried, separately, and keep them carefully from the air.

PULLNA WATER. Sulphate of soda 3717 grs., sulphate of magnesia 5168 grs., chloride of calcium 231 grs., chloride of magnesium 723 grs., chloride of sodium 231 grs., bicarbonate of soda 150 grs., carbonated water 2 gallons. One of the most active of the purgative saline waters.

PULLNA WATER, WITHOUT THE MACHINE. Bicarbonate of soda 50 grs., sulphate of magnesia, 4 drs., sulphate of soda 3 drs., chloride of sodium a scruple; dissolve in a pint of water; add, lastly, 2 scruples of bisulphate of soda, and close the bottle immediately.

SALTS FOR MAKING PULLNA WATER. Dry bicarbonate of

soda 1 oz., exsiccated sulphate of soda 2 oz., exsiccated sulphate of magnesia $1\frac{1}{2}$ oz., dry chloride of sodium 2 drs., dry tartaric acid $\frac{3}{4}$ oz. (or rather, dry bisulphate of soda 1 oz.).

PLOMBIÈRES WATER. Bicarbonate of sodium 0.210 gramme, chloride of calcium 0.028 gramme, sulphate of sodium 0.016 gramme, tartrate of potash and iron 0.011 gramme, carbonated water 1 litre.

SAINT-GALMIER WATER. A weak carbonated water containing minute quantities of the bicarbonates of soda, potash, lime, and magnesia, sulphates of soda and lime, silicate of alumina and chloride of magnesium.

SEIDLITZ WATER. This is usually imitated by strongly aerating a solution of 2 drs. of sulphate of magnesia in a pint of water. It is also made with 4, 6, and 8 drs. of the salts to a pint of water.

SEIDLITZ POWDERS. The common seidlitz powders do not resemble the water. A closer imitation would be made by using effloresced sulphate of magnesia instead of the potassio-tartrate of soda. A still more exact compound will be the following:—Effloresced sulphate of magnesia 2 oz., bicarbonate of soda $\frac{1}{2}$ oz., dry bisulphate of soda $\frac{1}{2}$ oz., mix and keep in a close bottle.

SEIDSCHUTZ WATER. Sulphate of magnesia 3 drs., chloride of calcium, nitrate of lime, bicarbonate of soda, of each 8 grs., sulphate of potash 5 grs., aerated water 1 pint.

SELTZER WATER. Chloride of calcium and chloride of magnesium, of each 4 grs.; dissolve these in a small quantity of water, and add it to a similar solution of 8 grs. of bicarbonate of soda, 20 grs. chloride of sodium, and 2 grs. of phosphate of soda; mix, and add a solution of $\frac{1}{4}$ of a gr. of sulphate of iron; put the mixed solution into a 20-oz. bottle, and fill up with aerated water. Much of the Seltzer sold is said to be nothing more than simple carbonated water, containing a little chloride of sodium. An imitation of Seltzer water is also made by putting into a stone Seltzer bottle, filled with water, 2 drs. bicarbonate of soda, and 2 drs. of citric acid in crystals, corking the bottle immediately. Sodaic powders are sometimes sold as Seltzer powders.

VICHY WATER. Bicarbonate of soda 1 dr., chloride of sodium 2 grs., sulphate of soda 8 grs., sulphate of magnesia 3 grs., tincture of chloride of iron 2 drops, aerated water a pint.

DORVAULT directs 75 grs. of bicarbonate of soda, 4 grs. of chloride of sodium, 1-5th gr. sulphate of iron, 10 grs. sulphate of soda, 3 grs. sulphate of magnesia, to a pint of water. By adding 45 grs. (or less) of citric acid an effervescing water is obtained.

M. SOUBEIRAN, relying on the analysis of LONGCHAMPS, imitates Vichy water by the following combination:—Bicarbonate of soda 135 grs., chloride of sodium $2\frac{1}{2}$ grs., cryst. chloride of calcium 12 grs., sulphate of soda $11\frac{1}{2}$ grs., sulphate of magnesia $3\frac{1}{2}$ grs., tartrate of iron and potash 1-8th gr., water 2 and 1-10th pints (1 litre), carbonic acid 305 cub. inches (5 litres). Dissolve the salts of soda and iron in part of the water, and add the sulph. magnes., and then the chlor. calc. in the remaining water. Charge now with the carbonic acid gas under pressure.

VICHY SALTS. Bicarbonate of soda $1\frac{1}{2}$ oz., chloride of sodium 15 grs., effloresced sulphate of soda 1 dr., effloresced sulphate of magnesia 1 scruple, dry tartarised potash and iron 1 gr., dry tartaric acid 1 oz. (or dry bisulphate of soda); mix the powders, previously dried, and keep them in a close bottle.

SALINE WATERS, &c., NOT CARBONATED.

SEA WATER. Chloride of sodium 4 oz., sulphate of sodium 2 oz., chloride of calcium $\frac{1}{4}$ oz., chloride of magnesium 1 oz., iodide of potassium 4 grs., bromide of potassium 2 grs., water a gallon.

A common substitute for sea water as a bath is made by dissolving 4 or 5 oz. of common salt in a gallon of water. The following mixture of dry salts may be kept for the immediate production of a good imitation of SEA WATER. Chloride of sodium (that obtained from evaporating sea water and not recrystallized, in preference) 85 oz., effloresced sulphate of soda 15 oz., dry chloride of calcium 4 oz., dry chloride of magnesium 16 oz., iodide of potassium 2 drs., bromide of potassium 1 gr. Mix, and keep dry. Put 4 or 5 oz. to a gallon of water.

BALARUC WATER. Chloride of sodium 1 oz., chloride of calcium 1 oz., chloride of magnesium $\frac{1}{2}$ oz., sulphate of sodium 3 drs., bicarbonate of sodium 2 drs., bromide of potassium 1 gr., water a gallon. Chiefly used for baths.

SULPHURETTED WATERS

SIMPLE SULPHURETTED WATERS. Pass sulphuretted hydrogen into cold water (previously deprived of air by boiling, and cooled in a closed vessel), till it ceases to be absorbed.

AIX-LA-CHAPELLE WATER. (GRAY.) 1. Pure water 2 gallons, carbonic acid 4 gallons, sulphuretted hydrogen 1-3rd of a gallon, chloride of sodium 423 grs., bicarbonate of sodium 177 grs., sulphate of sodium 90 grs., chloride of calcium 45 grs., chloride of magnesium 12 grs. Dissolve the sulphuretted hydrogen in its own volume of the water. Dissolve the salts in the remainder of the water, and then charge the solution with the carbonic acid. Divide the latter solution into 16 bottles; add an equal proportion of sulphuretted hydrogen to each bottle, and cork it immediately. 2. Bicarbonate of soda 12 grs., chloride of sodium 25 grs., chloride of calcium 3 grs., sulphate of soda 8 grs., simple sulphuretted water 2½ oz., water slightly carbonated 17½ oz.

BAREGES WATER. (Cauterets, Bagneres de Luchon, Eaux Bonnes, St Sauveur, may be made the same.) Crystallized hydrosulphate of soda (see Sodæ Hydrosulphas (Sodii Sulphidum Crystallizatum), *Pocket Formulary*), crystallized carbonate of soda, and chloride of sodium, of each 1½ grs., water (freed from air) a pint. A stronger solution for adding to baths is thus made:—Crystallized hydrosulphate of soda, crystallized carbonate of soda, and chloride of sodium, of each 2 oz., water 10 oz.; dissolve. To be added to a common bath at the time of using.

NAPLES WATER. Crystallized carbonate of soda 15 grs., fluid magnesia 1 oz., simple sulphuretted water 2 oz., aerated water 16 oz. Introduce the sulphuretted water into the bottle last.

HARROGATE WATER. Chloride of sodium 100 grs., chloride of calcium 10 grs., chloride of magnesium 6 grs., bicarbonate of sodium 2 grs., water 18½ oz. Dissolve, and add simple sulphuretted water 1½ oz.

HARROGATE SALTS. See Dr DUFFIN'S (further back).

CHALYBEATE WATERS

- SIMPLE CHALYBEATE WATER.** Water freed from air by boiling 1 pint, sulphate of iron $\frac{1}{2}$ gr.
- AERATED CHALYBEATE WATER.** Sulphate of iron 1 gr., carbonate of soda 4 grs., water deprived of air, and charged with carbonic acid gas, a pint. Dr PEREIRA recommends 10 grs. each of sulphate of iron and bicarbonate of soda to be taken in a bottle of ordinary soda-water. This is equivalent to 4 grs. of carbonate of iron.
- BRIGHTON CHALYBEATE.** Sulphate of iron, chloride of sodium, chloride of calcium, of each 2 grs., carbonate of soda 3 grs., carbonated water 1 pint.
- BUSSANG, FORGES, PROVINS,** and other similar waters, may be imitated by dissolving from $\frac{1}{2}$ to 2-3rds of a grain of sulphate of iron, 2 or 3 grs. of carbonate of soda, 1 gr. of sulphate of magnesia, and 1 of chloride of sodium, in a pint of aerated water.
- MONT D'OR WATER.** Bicarbonate of soda 70 grs., sulphate of iron 2-3rd gr., chloride of sodium 12 grs., sulphate of soda $\frac{1}{2}$ gr., chloride of calcium 4 grs., chloride of magnesium 2 grs., carbonated water a pint.
- PASSY WATER.** Sulphate of iron 8 grs., chloride of sodium $2\frac{1}{2}$ grs., sulphate of lime 17 grs., sulphate of magnesia 9 grs., alum 3 grs., water a pint.
- PYRMONT WATER.** Sulphate of magnesium 20 grs., chloride of magnesium 4 grs., chloride of sodium 2 grs., bicarbonate of sodium 16 grs., sulphate of iron 2 grs., Carrara water a pint.
- SPA WATER.** Chloride of calcium $26\frac{1}{2}$ grs., sulphate of magnesia 14 grs., protochloride of iron $15\frac{1}{2}$ grs., bicarbonate of soda 45 grs., water 2 galls., carbonic acid 5 vols. Decompose the chloride of calcium and sulphate of magnesia separately with carbonate of soda; collect and wash the precipitate, which add to the bicarbonate of soda and protochloride of iron dissolved in the water, and charge the solution with the carbonic acid.

VARIOUS AERATED MEDICINAL WATERS NOT
RESEMBLING ANY NATURAL SPRING

MIALHE'S AERATED CHALYBEATE WATER. Water a pint, citric acid 1 dr., citrate of iron 15 grs.; dissolve, and add 75 grs. of bicarbonate of soda.

TROUSSEAU'S MARTIAL AERATED WATER. Potassio-tartrate of iron 10 grs., artificial Seltzer water a pint.

BOUCHARDAT'S GASEOUS PURGATIVE. Phosphate of soda $1\frac{1}{2}$ oz., carbonated water a pint.

MIALHE'S IODURETTED GASEOUS WATER. Iodide of potassium 15 grs., bicarbonate of sodium 75 grs., water a pint; dissolve and add sulphuric acid, diluted with its weight of water, 75 grs. Cork immediately.

DUPASQUIER'S GASEOUS WATER OF IODIDE OF IRON. Solution of iodide of iron (containing 1-10th of dry iodide) 30 grs., syrup of gum $2\frac{1}{2}$ oz., aerated water $17\frac{1}{2}$ oz.

PERFUMERY

DISTILLED WATERS

THE simple distilled non-spirituous waters used in perfumery are chiefly those of rose, elder, orange flower, cinnamon, &c. The points requisite to be attended to in their preparation are, that the flowers be fresh, gathered after the sun has risen and the dew exhaled, and that sufficient water be used to prevent the flowers being burned in the retort, but not much more than is sufficient for this purpose. The quantities usually directed are—Roses 15 lbs., water 40 lbs.: distil 15 lbs. for *single*, and the same water with 15 lbs. of fresh roses, for *double* rose-water.

Orange-flowers 12 lbs., water 36 lbs.: distil 24 lbs. for *double* orange-flower water; this with an equal quantity of distilled water forms the *single*. The flowers should not be put into the still till the water nearly boils.

ELDER-FLOWER WATER, ACACIA-FLOWER WATER, AND BEAN-FLOWER WATER are prepared in the same manner as rose-water.

EAU DE NAPHE. This water is distilled in Languedoc from the leaves of the bigarade, or bitter-orange tree, but the preparation sold in England under this name is often prepared thus:—Orange flowers 7 lbs.; fresh yellow peel of the bigarade, or Seville orange $\frac{1}{2}$ lb., water 2 gallons; macerate for 24 hours, and distil 1 gallon. Ordinary orange-flower water is very generally sold for this water.

CINNAMON WATER A gallon should be distilled from 20 oz. of bruised cinnamon of best quality and 2 gallons of water.

STRAWBERRY WATER. Bruised strawberries 4 lbs., water a gallon; macerate for 12 hours, and distil 6 pints.

The waters prepared without distillation (by diffusing the essential oils through water, after mixing them with chalk, magnesia, china clay, or silica, or dissolving them in spirit) are seldom so well adapted for perfumery purposes as those distilled from the flowers, &c. Rose-water, made from the otto (8 drops of otto, previously mixed with a

drachm of precipitated chalk, diffused in a quart of distilled water, and afterwards distilled or simply filtered), is to most persons very agreeable; but that distilled from the flowers is preferred by others.

MUSK WATER, VIOLET WATER, JESSAMINE WATER, and some others, are made by mixing the spirituous essences with distilled or pure soft water. A usual proportion is 2 drs. to a pint.

SPIRITUOUS WATERS

The spirit employed in perfumery should be selected with great care; it should be perfectly free from grain-oil and other impurities. It should be 60 over-proof in strength, unless otherwise directed. The distillation should be effected by steam, or by the heat of a water-bath.

SIMPLE SPIRIT OF LAVENDER. Lavender-flowers (free from stalks) 2 lbs., rectified spirit 8 pints, water 16 pints; distil 8 pints.

SMYTH'S DISTILLED ESSENCE OF LAVENDER. Essential oil of English lavender 4 oz., rectified spirit (60° over-proof) 5 pints, rose-water 1 pint: mix, and distil 5 pints.

ESSENCE OF LAVENDER (by mixture). Essential oil of lavender 3½ oz., rectified spirit 2 quarts, rose-water ½ pint, tincture of orris ½ pint.

LAVENDER WATER. English oil of lavender 4 oz., spirit 3 quarts, rose-water 1 pint; mix and filter. (A commoner and cheaper preparation may be made with the French oil.)

ODORIFEROUS LAVENDER WATER. 1. Rectified spirit 5 gallons, essential oil of lavender 20 oz., oil of bergamot 5 oz., essence of ambergris ½ oz. Sometimes 4 oz. of orris-root are digested with the above.—Mr BRANDE.

2. Oil of lavender, oil of bergamot, of each 3 drs.; otto of roses and oil of cloves, of each 6 drops; musk 2 grs., true oil of rosemary 1 dr., honey 1 oz., benzoic acid 2 scruples; rectified spirit a pint, distilled water 3 oz.—Dr PEREIRA.

3. Oil of lavender 2 oz., essence of ambergris 1 oz., eau de Cologne a pint, rectified spirit a quart.

4. Oil of lavender 4 drs.; essence of bergamot, essence of lemon or cedrat, and otto of roses, of each 20 minims; essence of ambergris 1 dr., rectified spirit 3 pints, orange-

flower water 4 oz., rose (or distilled) water 12 oz., burnt alum 20 grs. Agitate frequently, then let it stand in a cool place for some days before filtering.

5. Oil of lavender 3 drs., oil of bergamot 20 drops, neroli 6 drops, otto 6 to 12 drops, essence of cedrat 8 or 10 drops, essence of musk 20 drops, rectified spirit 28 fluid oz., distilled (or orange-flower) water 4 oz.

6. *Eau de Lavande aux Millefleurs*. Oil of lavender 4 drs.; essence of bergamot, essence of lemon, otto of roses, of each 12 drops; essence of millefleurs 3 drs., essence of ambergris 1 dr., rectified spirit a pint and a half.

7. English oil of lavender 8 oz.; essence of musk 4 oz.; essence of ambergris and oil of bergamot of each 1½ oz., rect. spirit 2 gallons. Mix well. Very fine.—Mr COOLEY.

Note.—The oil of lavender in all the above should be the finest English oil;* that which first comes over is said to be the most fragrant. It should be kept for 12 months before using, either alone or mixed with an equal quantity of alcohol. Some makers prefer a mixture of old and new oil. The lavender water improves by age.

EAU DE COLOGNE—*Cologne Water*. 1. English oil of lavender, oil of bergamot, oil of lemon, oil of neroli, of each 1 oz.; oil of cinnamon ½ oz.; spirit of rosemary, and spirit of balm (*Eau des Carmes*), of each 15 oz.; highly rectified spirit 7½ pints. Let them stand together for 14 days, then distil in a water-bath.—Dr GRANVILLE.

2. Oil of bergamot, citron, and lemon, each 3 oz.; oils of rosemary, neroli, and lavender, of each 1½ oz.; oil of cinnamon 6 drs., rectified spirit 24 pints; compound spirit of balm (*Eau des Carmes*) 3 pints, spirit of rosemary 2 pints. Mix, and after standing a week, distil 24 pints.—*French Pharmacopœia*.

3. Essential oils of bergamot, of lemon, of neroli, of orange peel, and of rosemary, each 12 drops; cardamom seeds a drachm; rectified spirit a pint. It improves by age.—THROMMSDORF.

4. Essence of bergamot 40 minims, essence of lemon 45 minims, oil of rosemary 6 minims, oil of orange 22 minims, neroli 12 minims, highly rectified spirit 6 oz.

* Of late years, the high price of the English oil has precluded its use by the druggist. An excellent substitute will be found in the redistilled foreign oil, which is a great improvement on the ordinary kind. It may be obtained from most of the wholesale houses.

5. Alcohol a pint, oil of bergamot, oil of orange-peel, true oil of rosemary, cardamom seeds, of each a drachm; orange-flower water a pint. Mix, and distil a pint by water-bath.—Dr A. T. THOMSON.

EAU DES CARMES—*Eau de Melisse*. Fresh flowering balm 24 oz.; yellow rind of lemon, cut fine, 4 oz., cinnamon, cloves, and nutmeg (bruised), of each 2 oz.; coriander seed (bruised) 1 oz., dried angelica root 1 oz., rectified spirit a gallon. Macerate for 4 days, and distil in a water-bath.

AQUEBUSADE WATER. 1. Sage, angelica, wormwood, savory, sweet fennel, hyssop, balm, sweet basil, rue, thyme, marjoram, rosemary, angelica seed, origanum, red calamint, creeping thyme, lavender-flowers, of each 10 oz.; sweet flag root 5 oz., rectified spirit 2 gallons, water q. s. Distil 3 gallons.

2. (Simplified.) Balm, rosemary, thyme, calamus root, angelica seeds, lavender flowers, of each 4 oz.; rectified spirit 3 pints, water q. s. Macerate for a day, and distil 4 pints.

QUEEN OF HUNGARY'S WATER. *Spirit of Rosemary*. 1. Rosemary tops 2 lbs., rectified spirit, a gallon, water q. s. Distil carefully 1 gallon.

2. Spirit of rosemary (as No. 1) 4 pints, orange-flower water $\frac{1}{2}$ pint, essence of neroli 4 drops.

3. Simple spirit of rosemary 3 pints, simple spirit of lavender a pint, rose-water 8 oz.

EAU D'ANGE. Flowering tops of myrtle 16 oz., rectified spirit a gallon; digest, and distil to dryness in a water-bath. Or dissolve $\frac{1}{4}$ oz. essential oil of myrtle in 3 pints of rectified spirit. Mr GRAY gives under this name a water without spirit: Water 2 pints, benzoin 2 oz., storax 1 oz., cinnamon 1 dr., cloves 2 drs. Calamus a stick coriander seeds a pinch: distil.

HONEY WATER. *Eau de Miel*. 1. Rectified spirit 8 pints oil of cloves, oil of lavender, oil of bergamot, of each $\frac{1}{2}$ oz., musk 15 grs., yellow sanders shavings 4 oz., digest for 8 days, and add 2 pints each of orange-flower and rose-waters.

2. Oil of santal 20 drops, tincture of musk $2\frac{1}{2}$ oz., essence of bergamot $2\frac{1}{2}$ oz., oil of cloves 5 drs., oil of lavender 5 drs., rose-water 2 pints, orange-flower water 2 pints, spirits of wine a gallon; mix, and filter.

3. (With honey.) White honey 8 oz., coriander seed 8 oz., fresh lemon-peel 1 oz., cloves $\frac{1}{4}$ oz., nutmeg, benzoin, styrax calamita, of each 1 oz., rose and orange-flower water, of each $\frac{1}{4}$ oz. rectified spirit 3 pints; digest for a few days, and filter. Some receipts add 3 drs. of vanilla and direct only $\frac{1}{2}$ oz. of nutmeg, storax, and benzoin.

4. Coriander seeds 7 lbs., cloves 12 oz., storax 8 oz., nutmeg 8 oz., fresh lemon-peel 10 oz., calamus root 6 oz., rectified spirit 15 pints; macerate for a month, add water q. s. Distil 22 pints, and add to the distilled spirit 5 pints of orange-flower water, 24 drops otto of roses, a drachm of ambergris, and 2 oz. of fine vanilla; macerate for a week, and filter. The dry ingredients to be bruised or cut small.

LISBON WATER. To rectified spirit, 1 gallon, add the essential oils of orange-peel and lemon-peel, of each 3 oz., and of otto of rose $\frac{1}{4}$ oz.—PIESSE.

EAU DE PORTUGAL. To rectified spirit 1 gallon, add the following essential oils: of orange-peel 6 oz., of lemon-peel 1 oz., of lemon-grass $\frac{1}{4}$ oz., of bergamot 1 oz., and of otto of rose $\frac{1}{4}$ oz.—PIESSE.

EAU D'ÉLÉANCE. Spirit of jessamine 2 lbs., spirit of styrax 1 lb., spirit of hyacinth 1 lb., spirit of star aniseed $\frac{1}{4}$ oz., tincture of balsam of tolu 4 oz., tincture of vanilla 2 oz.

EAU DE MARESCHALE. Spirit of wine $1\frac{1}{2}$ pints, spirit of jessamine 1 oz., essence of bergamot $\frac{1}{4}$ oz., essence of violets 1 oz.

EAU ROMAINE. Spirit of jessamine 3 quarts, tincture of vanilla 1 quart, spirit of acacia flowers 1 quart, spirit of tuberose a pint, essence of ambergris 2 oz., tincture of benzoin 8 oz.

EAU DE MILLEFLEURS. Rectified spirit 2 pints, balsam of Peru $\frac{1}{4}$ oz., essence of bergamot $\frac{1}{2}$ oz., oil of cloves $\frac{1}{4}$ oz., essence of neroli $\frac{1}{2}$ dr., essence of musk 1 dr., orange-flower water 2 oz.

EAU SPIRITUEUSE D'HÉLIOTROPE. Vanilla 3 drs., double orange-flower water 6 oz., rectified spirit a quart; macerate for 3 days, and distil in a water-bath. It may be coloured with cochineal. The essence d'héliotrope of some perfumers appears, by the colour, not to have been distilled.

EAU D'ISAPHAN. Essential oil of bitter orange-peel 4 oz.,

oil of rosemary 3 drs., oil of mint 1 dr., oil of cloves 7 scruples, neroli 7 scruples, spirit of wine 14 pints. Used for the same purposes as eau de Cologne.

EAU SANS PAREILLE. Essential oil of lemon $\frac{1}{2}$ oz., of bergamot $2\frac{1}{2}$ drs., of cedrat $\frac{1}{4}$ oz., rectified spirit 6 pints, spirit of rosemary 8 oz.; mix. Some authorities state that it is improved by distillation.

EAU DE BOUQUET DE FLORE. 1. Honey water 2 oz., tincture of cloves 1 oz., tincture of calamus, of lavender, and of long cyperus, each $\frac{1}{2}$ oz.; eau sans pareille 4 oz., spirit of jessamine 9 drs., tincture of orris 1 oz., spirituous essence of neroli 20 drops.

2. Essence of violets $\frac{1}{2}$ oz., spirit of rosemary $\frac{1}{2}$ oz., essence of lemon 1 dr., rectified spirit 24 oz., rose-water 8 oz.

3. Spirit of rosemary 8 oz., rectified spirit 8 oz., lavender water 2 oz., oil of neroli 5 drops, cloves 1 dr., orris root 3 drs., rose-water 2 oz.; digest for a few days, and filter.

ESPRIT DE BOUQUET. English oil of lavender, oil of cloves and of bergamot, of each 2 drs.; otto of roses and oil of cinnamon, of each 20 drops; essence of musk 1 dr., rectified spirit a pint; mix.

EAU DE ROSIÈRES. Spirit of roses 4 pints, spirit of jessamine a pint, spirit of orange-flowers a pint, spirit of cucumber $2\frac{1}{2}$ pints, spirit of celery seed $2\frac{1}{4}$ pints, spirit of angelica root $2\frac{3}{4}$ pints, tincture of benzoin (simple) $\frac{3}{4}$ of a pint, balsam of Mecca a few drops.

EAU D'AMBRE ROYALE. Rectified spirit 2 lbs., tincture of musk seed 1 lb., essence of ambergris 1 oz., tincture of musk 1 oz., reduced with a proper proportion of orange-flower water.

ESPRIT DE SUAVE. Spirit of jessamine $1\frac{1}{2}$ pint, spirit of acacia flowers $1\frac{1}{2}$ pints, spirit of wine 12 oz., spirit of tuberose 8 oz., oil of cloves $1\frac{1}{2}$ drs., oil of neroli 30 drops, essence of bergamot $1\frac{1}{2}$ drs., tincture of musk 1 oz., rose-water 12 oz.

FLORIDA WATER. Oil of lavender, oil of bergamot, oil of lemon, of each 2 drs., tincture of turmeric, oil of neroli, of each 1 dr., oil of balm 30 drops, otto of roses 10 drops, rectified spirit 2 pints.

PARFUM DES ROIS. Spirit of wine 2 gallons, styrax 6 oz., benzoin 16 oz., aloes-wood 8 oz., spirit of rose 2 pints,

spirit of orange-flowers 2 pints, tincture of ambergris 8 oz., tincture of musk 8 oz., tincture of vanilla 16 oz.

ODOR DELECTABILIS. Rose-water, orange-flower water, each 4 oz., oil of lavender, oil of cloves, each 1 dr., oil of bergamot 2 drs., musk 2 grs., rectified spirit a pint.

NEW MOWN HAY. Ext. Tonquin Bean 2 pints; ext. geranium 1 pint; ext. orange-flower 1 pint; ext. rose 1 pint; ext. rose triple 1 pint; ext. jessamine 1 pint.

NEW VICTORIA PERFUME. 1. Cloves bruised 2 scruples; vanilla, cut small, 1 dr.; oil of cedrat 4 drops, oil of santal 1 dr., cinnamon 12 grs., oil of verbena 8 drops, otto of roses 8 drops, oil of neroli 20 drops, oil of lavender 1 dr., ambergris 16 grs., tincture of musk 1 dr., rectified spirit 16 fl. oz.; digest for a few days and filter. Or the whole except the musk and ambergris may be distilled in a water-bath, and these added to the distilled spirit.

2. Vanilla $\frac{1}{2}$ dr., yellow sanders 6 drs., cloves No. 16, neroli 3 drops, oil of lavender 6 drops, rectified spirit 4 oz.; digest for 3 days and add 4 oz. of orange-flower water, water q. s.; distil 6 oz., add essence of musk 1 dr.

JOCKEY CLUB BOUQUET. **PIESSE.** Extract of orris root 2 pints, esprit de rose triple 1 pint, esprit de pommade de rose 1 pint, extract of pommade of cassia, and tuberose, $\frac{1}{2}$ pint each, ext. of ambergris $\frac{1}{2}$ pint, oil of bergamot $\frac{1}{2}$ oz.

ESPRIT DE ROSE. 1. Macerate the fresh and picked flowers of the most fragrant varieties of the rose, with half their weight of rectified spirit, and distil in a water-bath to dryness.

2. Dissolve from 20 to 30 drops of otto in a pint of rectified spirit. A stronger solution, 6 or 8 drops of otto to an oz. of alcohol, forms essence of roses, or *esprit de rose triple*.

3. Also made by agitating and digesting the spirit with the perfumed oil or pommade of roses. See **EXTRACTS** (further on).

ESPRIT DE JASMIN. *Eau de Jasmin.* Is prepared by digesting and agitating pure spirit with oil or pommade of jessamine made with the flowers. (See **EXTRACTS** further on.)

Spirit of **JONQUIL**, **TUBEROSE**, **VIOLET**, &c., may be obtained by the same process.

ESPRIT DE VIOLETTE. *Eau de Violette.* Macerate 5 oz. of fine orris root in a quart of rectified spirit for some days,

and filter. It may also be obtained by the method just mentioned, or by mixing the product of both processes.

ESS BOUQUET. (KILNER.) Ext. musk 4 oz., ext. tuberose 2 oz., otto of roses 1 dr., bergamot 1½ drs., oil of neroli ½ dr., oil of verbena 8 drops, oil of pimento 10 drops, oil of patchouli 3 drops, oil of red cedar wood ½ dr.; oil of English lavender 12 drops, pure rectified spirit to make 4 pints. Mix and digest.

EAU ODORANTE DE JASMIN. *Compound spirit of jessamine*; for the handkerchief. Spirit of jessamine 1 pint, rectified spirit 1 pint, essence of ambergris 1 dr., simple tincture of benzoin 1 dr.

SPIRIT OF ORANGE-FLOWERS, SPIRIT OF ELDER-FLOWERS, AND SPIRIT OF ACACIA-FLOWERS. Fresh flowers 1 lb., rectified spirit 4 lbs., or pints, water 2 lbs.; distil 4 lbs., or pints.

SPIRIT OF ORANGE-PEEL, OF LEMON-PEEL, OF CITRON AND OF BERGAMOT. Fresh peel 1 lb., rectified spirit 6 lbs.; macerate for 2 days, and distil in a water-bath to dryness. Or, 1 oz. of the essential oil to 2 pints of spirit.

SPIRIT OF CINNAMON, OF CLOVES, OF NUTMEG, AND OF CALAMUS ROOT. Macerate 1 lb. of the bruised drug with 8 lbs., or a gallon, of rectified spirit, and distil as the last.

SPIRIT OF CUCUMBERS. Cucumbers grated 8 lbs., rectified spirit 1 lb.; distil 2 lbs.

SPIRIT OF ROSEWOOD. Rosewood shavings 1 lb., spirit 6 lbs., water 2 lbs.; distil 6 lbs. It is also made by adding the essential oil of rhodium to spirit.

SPIRIT OF ANGELICA. Dried angelica root 1 lb., rectified spirit a gallon. Macerate, and distil by water-bath to dryness.

SPIRIT OF BALSAM OF PERU. Balsam 3 parts, spirit 15 parts, carbonate of potash 1 part; macerate for 3 days, and distil by water-bath.

SPIRIT OF STRAWBERRIES, AND OF RASPBERRIES. Fresh fruit 3 lbs., rectified spirit 1 lb.; macerate 24 hours, and distil 2 lbs.

YLANG-YLANG. Ext. tonka 3 oz.,* ext. musk 4 oz., ext. tuberose 4 oz., ext. cassia 4 oz., ext. orris 8 oz., otto of

* EXTRACT OF TONKA is made as follows:—Tonka (or Tonquin) beans in coarse powder 1 lb., percolate with rectified spirit until 1 gallon is obtained.—KILNER.

orange 2 drs., otto of neroli $\frac{1}{2}$ dr., rectified spirit to make 4 pints; mix.

The following tinctures are chiefly used in the compound perfumes:

TINCTURE OF BALSAM OF PERU, AND OF TOLU. Digest 1 oz. of the balsam with 8 fl. oz. of rectified spirit for some days, shaking it occasionally, then filter. **TINCTURE OF BENZOIN** in the same manner.

TINCTURE OF ANGELICA. One part of the dried root to 8 parts of rectified spirit; as the last.

TINCTURE (common spirituous essence) OF LEMON, CITRON, ORANGE, AND BERGAMOT. An ounce of the fresh peel to $\frac{1}{2}$ pint of spirit, as above.

TINCTURE OF MUSK SEED. *Essence d'Ambrette.* Digest 16 oz. of bruised musk seed with 3 pints of rectified spirit for a month, and filter.

TINCTURE OF MUSK. China musk 2 drs., rectified spirit 16 oz. For more compound tinctures of musk, see **ESSENCE OF MUSK**, below.

TINCTURE OR ESSENCE OF AMBERGRIS. GUIBOURT directs 1 dr. of ambergris to be digested with a gentle heat in 3 oz. of rectified spirit. Another form is: Ambergris 1 dr., carbonate of potash 1 dr., spirit of roses 4 oz. (or rectified spirit 4 oz., otto 6 drops). Some recipes direct a weaker solution: 24 grs. of ambergris to 8 oz. of spirit. For other formulæ, see **ESSENCE**, further on.

TINCTURE OF CIVET. Bruise $\frac{1}{2}$ oz. of civet, $\frac{1}{2}$ oz. of ambergris, and the same of sugar candy, and macerate in a quart of rectified spirit for 6 weeks, in a warm place; then filter.

TINCTURE OR ESSENCE OF VANILLA. Vanilla cut very small 2 oz., rectified spirit a pint; infuse for 2 or 3 weeks. This is sometimes distilled, forming spirit of vanilla.

TINCTURE OF RHODIUM. Rosewood 1 lb., rectified spirit 3 or 4 pints; macerate for 3 or 4 weeks, and filter.

ESSENCE (OR TINCTURE) OF VETIVER. Take 2 lbs. of the root of vittie vayr cut small, and moisten it with a little water; let it macerate for 24 hours, then beat it in a marble mortar. Macerate it in sufficient spirit to cover it, for 8 or 10 days, and strain with pressure: filter through paper, and in a fortnight repeat the filtration. Sometimes the root is moistened with diluted sulphuric acid, which, after maceration, is neutralized by adding a suffi-

cient quantity of chalk, and the whole digested with spirit. The tincture when strained off, is distilled, and forms (with the addition of essence of balm and of roses) ESSENCE DE VETIVER DOUBLE.

ESSENCE OF PATCHOULI. Dried patchouli (*pucha pat*) 1 oz., rectified spirit a pint. It is generally combined with other perfumes.

ESSENCE OF TUBEROSE. Extract of tuberose 24 oz., extract of musk 4 oz., extract of jasmin 1 oz., otto of rose 1 dr., oil of neroli 10 drops, benzoic acid 2 drs., rectified spirits to make 4 pints. Mix.

ESSENCE OF WOOD VIOLET. (KILNER.) Extract of orris 12 oz., extract of tuberose 2 oz., extract of jasmin 1 oz., extract of musk 4 oz., bergamot 2 drs., English oil of lavender 1 dr., otto of verbena 10 drops, otto of bitter almonds 12 drops, otto of coriander 6 drops, otto of sweet flag 4 drops, otto of bay leaves 5 drops, benzoic acid 1½ drs., rectified spirit to make 4 pints. Mix.

EXTRACTS (*extraits*) are spirituous solutions of the odorous principle of flowers, obtained, indirectly, by agitating and digesting oils and pomatums which have been perfumed by the flowers (see HUILES ANTIQUES, under HAIR COSMETICS) with pure spirit. This is repeated with fresh oil until the spirit is sufficiently perfumed. When the same oil or pomade is treated with fresh spirit, inferior extracts, numbered 2, 3, &c., are obtained. These preparations are chiefly made in France.

EXTRACTS (EXTRAITS, or ESPRITS) OF JESSAMINE, VIOLETS, LILY OF THE VALLEY, are prepared by the process just mentioned.

EXTRAIT DE BOUQUET. Spirit (extrait) of jessamine 2 quarts, extract of violets 2 quarts, spirit of acacia-flowers, of rose, and of orange-flowers, each a quart, spirit of carnations a quart, flowers of benzoin ½ oz., essence of ambergris 1 oz.

EXTRAIT DE MARESCHALE. Essence of millefleurs 1½ oz., essence of jessamine 1 oz., essence of musk ½ oz., essence of ambergris ½ oz., essence of cedrat 20 drops, essence of violets 1 oz., sweet spirits of nitre 50 drops, true oil of rosemary 20 drops, rectified spirit 6 oz., oil of neroli 48 drops. Set aside for some time.

COMPOUND ESSENCES. Some of these contain a preparation of the substance whose name they bear, while others are

fictitious or imitative, being made up of a variety of other essences and volatile oils. Several of the formulæ are those of M. PIESSE.

ESSENCE OF AMBERGRIS. This name is applied both to the simple and compound tinctures of ambergris. See **TINCTURE OF AMBERGRIS** (further back.) Other formulæ may here be given.

1. Ambergris 4 oz., musk 2 oz., tincture of musk seed 7 pints. Digest with a gentle heat.

2. Ambergris 1 dr., musk $\frac{1}{2}$ dr., oil of cinnamon 18 drops, oil of rhodium 12 drops, rectified spirit 8 oz., spirit of roses 4 oz., carbonate of potash $1\frac{1}{2}$ drs., digest in a warm place for a few days, and strain. See also **ESSENCE ROYALE**.

ESSENCE OF CEDRAT. Dissolve $2\frac{1}{2}$ oz. of oil of cedrat in 1 gallon of spirit, and add bergamot $\frac{1}{2}$ oz.

ESSENCE OF CLOVE PINK. Esprit de rose $\frac{1}{2}$ pint, de fleur d'orange, and de fleur de cassie, each $\frac{1}{4}$ pint, esprit de vanille 2 oz., oil of cloves 10 drops.

COLOGNE ESSENCE. Oil of bergamot 2 drs., essence of lemon $\frac{1}{2}$ dr., essence of cedrat $\frac{1}{2}$ dr., true oil of rosemary 15 drops, rectified spirit (or spirit of balm) $1\frac{1}{2}$ oz.

ESSENCE OF HELIOTROPE. Spirituous extract of vanilla $\frac{1}{2}$ pint, of French rose pomatum $\frac{1}{4}$ pint, of orange-flower pomatum 2 oz., of ambergris 1 oz., add 5 drops of the essential oil of almonds.

ESSENCE OF HONEYSUCKLE. Spirituous extract of rose pomatum 1 pint, of violet 1 pint, of tuberose 1 pint; extracts of vanilla and tolu, of each 1 pint; oil of neroli 10 drops, essential oil of almonds 5 drops.

ESSENCE OF HOVENIA. Rectified spirit 1 quart, rose-water $\frac{1}{2}$ pint, essential oil of lemons $\frac{1}{2}$ oz., otto of roses 1 dr., oil of cloves $\frac{1}{2}$ dr., oil of neroli 10 drops.

ESSENCE OF JONQUIL. Spirituous extract of jasmine pomade 1 pint, of tuberose 1 pint, of orange-flower $\frac{1}{2}$ pint; add extract of vanilla 2 oz.

ESSENCE OF LILY OF THE VALLEY. Mix the following extracts: of tuberose $\frac{1}{2}$ pint, of jasmin 2 oz., of orange-flower 2 oz., of vanilla 3 oz., of cassia $\frac{1}{2}$ pint, of rose-water $\frac{1}{2}$ pint; add 3 drops of hydrocyanic acid. Keep together for a month, then bottle.

ESSENCE OF MAGNOLIA. Spirituous extract of orange-flower pomatum 1 pint, of rose pomatum 2 pints, of tube-

- rose pomatum $\frac{1}{2}$ pint, of violet pomatum $\frac{1}{2}$ pint; essential oil of citron 2 drs., and essential oil of almonds 10 drops.
- ESSENCE OF MIGNONETTE.** Digest 1 lb. of pomade de réséda in rectified spirit 1 pint for 14 days; filter off, and add 1 oz. of extrait d'ambre.
- ESSENCE OF MILLEFLEUR.** (KILNER.) Otto of roses 1 dr., otto of red cedar wood 1 dr., otto of orange 1 dr., otto of pimento 20 drops, extract of orris 6 oz., extract jasmine 2 oz., extract styrax 1 oz., extract tonka (tonquin) 4 oz., rectified spirits to make 4 pints. Mix.
- ESSENCE OF MOSS ROSE.** Spirituous extract of French rose pomatum 1 quart, esprit de rose triple 1 pint, extract of orange-flower pomatum 1 pint, of ambergris $\frac{1}{2}$ pint, and of musk 4 oz.
- ESSENCE OF MYRTLE.** Take the following extracts: of vanilla $\frac{1}{2}$ pint, of roses 1 pint, of orange-flower $\frac{1}{2}$ pint, of tuberose $\frac{1}{2}$ pint, of jasmin 2 oz. Mix, and allow to stand for a fortnight.
- ESSENCE OF MUSK.** Tincture of musk, of various strengths. The formula given previously (**TINCTURE OF MUSK**) is that of the Dublin Pharmacopœia, 1826. GUIBOURT directs 1 part of musk to 12 of proof spirit. Other authorities direct a smaller quantity of musk. A French work gives the following: Musk in the bag, cut small, 6 oz., civet 1 oz., tincture of musk seed 7 pints; digest in the sun, or in a warm place for 2 months.
- ESSENCE OF PATCHOULI.** Oil of patchouli 14 oz., otto of rose $\frac{1}{2}$ oz., rectified spirit 1 gallon.
- ESSENCE OF RONDELETIA.** Essence of bergamot, essence of lemon, oil of cloves, each 1 dr., otto of roses 6 drops, rectified spirit 1 pint.
- ESSENCE ROYALE.** Ambergris 1 dr., civet 15 grs., musk 30 grs., carbonate of potash 20 grs.; triturate together, and add oil of cinnamon 10 drops, oil of rhodium, and of neroli, 6 drops, otto of roses 6 drops, rectified spirit $\frac{1}{2}$ pint; digest, and filter.
- ESSENCE OF SWEET BRIAR.** Spirituous extracts of French rose pomatum 1 pint, of cassia and orange flowers, each $\frac{1}{4}$ pint, esprit de rose $\frac{1}{4}$ pint, with oils of neroli and lemon-grass, of each $\frac{1}{2}$ dr.
- ESSENCE OF SWEET PEA.** Essences of tuberose, orange-flower, and rose pomatum each $\frac{1}{2}$ pint, with essence of vanilla, 1 oz.

ESSENCE OF VERBENA. Essential oil of verbena 2 drs., rectified spirit 4 oz., essence of ambergris $\frac{1}{2}$ dr., orange-flower water $\frac{1}{2}$ oz.; mix. Another form is: Oil of verbena $\frac{1}{2}$ oz., essence of vanilla 40 drops, rectified spirit 4 oz.; mix and filter.

ESSENCE OF WHITE LILAC. Spirituous extract of tuberose pomade 1 pint, of orange-flower pomade $\frac{1}{2}$ pint; add essential oil of almonds 3 drops, and extract of civet $\frac{1}{2}$ oz.

FRANGIPANNI BOUQUET. Essence of vetiver 3 oz., ol. neroli 15 minims, oil of sandal wood $\frac{1}{2}$ dr., otto of rose 40 minims, essence of musk 3 drs., esprit de violette 3 oz., essence of ambergris 6 drs., rectified spirit to make up 20 oz.—PIESSE.

ESSENCES FOR SCENTING POMATUM.—*Millefleur*: Oil of lemon 3 oz., essence of ambergris 4 oz., oil of cloves 2 oz., oil of lavender 2 oz.—*Cowslip*: Essence of bergamot 16 oz., essence of lemon 8 oz., oil of cloves 4 oz., oil of orange-peel 2 oz., oil of jessamine 2 drs., eau de bouquet 2 oz., oil of bitter almonds 16 drops.—*For general use*: Essence of bergamot 16 oz., essence of lemon 8 oz., true oil of origanum and oil of cloves, each 2 oz., oil of orange-peel, $1\frac{1}{2}$ oz.

MISTURA ODORATA. Rectified spirit 48 oz., tincture of benzoin 4 oz., tincture of vanilla $\frac{1}{4}$ oz., tincture of musk $\frac{1}{2}$ oz., balsam of Peru $\frac{1}{2}$ oz., oil of cloves, of mace, and of cinnamon, each $\frac{1}{2}$ oz., oil of bergamot 1 oz., oil of cedrat 2 oz.—GIESKE.

SCENT FOR SNUFF. Oil of lavender 2 drs., essence of lemon 4 drs., essence of bergamot 1 oz.; mix. 1 dr. with 8 oz. of fine Scotch snuff, constitutes Queen's Snuff.

[The following Essences, Spirits, and Waters are given as specimens of some of the cheaper perfumes, as made in France.]

ESSENCES (SPIRITUOUS)

Essence (Spirituous) of Neroli. Spirit of wine $\frac{1}{2}$ pint, orange-peel, cut small, 3 oz., orris-root in powder, 1 dr., musk 2 grs.; let it stand in a warm place for 3 days, and filter.

Essence of Lemon. Spirit of wine $\frac{1}{2}$ pint, fresh lemon-peel 4 oz., as above.

Essence of Bergamot. Spirit of wine $\frac{1}{2}$ pint, bergamot-peel 4 oz.; as above.

- Essence of Violets.* Spirit of wine $\frac{1}{2}$ pint, orris root 1 oz.
Essence of Cedrat. Essence of bergamot (see *ante*) 1 oz.,
 essence of neroli 2 drs.
Essence of Jessamine. Essence of violets 1 oz., essence of
 cedrat 2 drs.
Essence of Musk. Spirit of wine $\frac{1}{2}$ pint, musk 16 grs.
Essence of Ambergris. Spirit of wine $\frac{1}{2}$ pint, ambergris
 24 grs.
Essence of Cloves. Spirit of wine $\frac{1}{2}$ pint, bruised cloves
 1 oz. Other essences in the same manner.

SPIRITS

- Spirit of Rose.* Spirit of wine $\frac{1}{2}$ pint, otto 6 drops.
Spirit of Jessamine. Spirit of wine $\frac{1}{2}$ pint, essence of
 jessamine (see further back) a drachm.
Spirit of Orange. Spirit of wine $\frac{1}{2}$ pint, essence of orange,
 or neroli, a drachm.
Spirit of Lavender. Spirit of wine $\frac{1}{2}$ pint, essential oil of
 lavender a drachm.
Spirit of Musk. Spirit of wine $\frac{1}{2}$ pint, essence of musk a
 drachm.
 Others in a similar manner.

SIMPLE WATERS

- Rose Water.* Distilled or rain water $\frac{1}{2}$ pint, spirit of roses
 a drachm.
*Jessamine Water, Musk Water, Violet, Orange-flower
 Water, &c.,* by adding a drachm of the above spirits to $\frac{1}{2}$
 pint of water.

AMMONIATED PERFUMES

- AMMONIATED COLOGNE WATER. A fragrant and reviving
 substitute for Spirit of Sal Volatile. Chloride of ammonium
 5 drs., carbonate of potash 8 drs., eau de Cologne 12 oz.,
 essential oil of cedrat and of bergamot, of each 15 drops
 (dissolved in an oz. of rectified spirit), orange-flower water
 8 oz.; mix, and carefully distil 15 or 16 oz.
 EAU DE LUCE. Mastic 2 drs., rectified spirit 9 drs.; dissolve
 and add to the clear tincture 30 drops of oil of lavender,
 10 drops of bergamot, and a pint of strong water of am-
 monia. This is more agreeable than the compound of the
 London Pharmacopœia, which, however, should always be
 used when prescribed medicinally.

ESSENCE FOR SMELLING SALTS. 1. English oil of lavender and essence of bergamot, of each a drachm, oil of orange-peel, or of cedrat, 8 drops, oil of cinnamon 4 drops, oil of neroli 2 drops, alcohol, and strongest water of ammonia, of each 2 oz.; or 4 oz. of strong ammoniated alcohol. See *SPIRITUS AMMONIÆ, L., Pocket Formulary.*

2. Ammoniated alcohol (see *SP. AMMONIÆ, L., Pocket Formulary*) 12 fl. oz. English oil of lavender, essence of bergamot and essence of lemon, of each a drachm cloves $\frac{1}{2}$ dr., camphor $\frac{1}{2}$ oz.; macerate for a week, and filter.—*Mr MAGGS.*

3. Essence of ambergris and musk 4 drs., otto of rose 20 drops, oil of lavender 1 dr., ammoniated alcohol 10 oz.; mix, and add strongest liquor ammonia 10 oz.—*Pharm. Journal.*

SMELLING SALTS. Sesquicarbonate of ammonia 40 oz., broken into small pieces not larger than a filbert, put into an air-tight $\frac{1}{2}$ -gallon jar; pour over it 20 oz. strong solution of ammonia (sp. gr. '880) previously perfumed according to taste, and immediately fix on the lid of the jar, taking care that it is properly secured. Keep in a cool place, opening and stirring it with a stiff spatula, every other day for a week; allow it now to remain for 2 or 3 weeks, at the end of which time it will have become hard.—*Mr ALLCHIN.*

GODFREY'S SMELLING SALTS. *Dr PARIS* says these are prepared by resubliming volatile salts with carbonate of potash and a little spirit of wine. They are usually scented with an alcoholic solution of essential oils.

ACETIC PERFUMES

AROMATIC SPIRIT OF VINEGAR. 1. Strong acetic acid 16 oz., camphor 1 oz.; when dissolved, add 1 oz. each of essential oils of cloves, lavender, and lemon. This is said to resemble *HENRY'S.*

2. Glacial acetic acid 8 oz., true oil of rosemary 20 grs., oils of bergamot 15 grs., of lavender 9 grs., of cloves 24 grs., neroli 4 grs., cinnamon 20 grs.; dissolve the oils in 2 drs. of rectified spirit. For another formula, see *Pocket Formulary.*

AROMATIC VINEGARS are made in France by infusing various flowers, &c., in distilled or finest wine vinegars, with or

without the addition of spirit. Others are made by distillation. As they are seldom required in this country, a few examples only will suffice.

ROSE VINEGAR. Red roses, picked and dried, $\frac{1}{2}$ lb., best vinegar 8 lbs.; macerate for a fortnight, with occasional stirring, and strain; then filter.

LAVENDER VINEGAR. Fresh lavender flowers 1 lb., vinegar 12 lbs.; macerate as above. It is sometimes distilled, drawing off 8 lbs.

DISTILLED ROSE VINEGAR. Pale roses, dried, 2 lbs., distilled vinegar 8 lbs.; distil 3-4ths by sand-bath, and add 2 lbs. of spirit of roses. It is occasionally coloured with cochineal and used as a cosmetic.

TOILET VINEGAR. Dried rose leaves 4 oz., otto of roses 90 drops, rectified spirit 10 oz., dilute acetic acid 40 oz.; macerate in a closed vessel for 14 days.—PIESSE.

VINAIGRE DE BULLY. Essence of bergamot 40 drops; essence of lemon 30 drops; oil of rosemary 20 drops; oil of balm 6 drops; oil of cloves, oil of lavender, oil of neroli, of each 4 drops; dissolve in rectified spirit 9 oz.; then mix with tincture of benzoin, tincture of styrax, tincture of tolu, of each 1 dr., distilled water 14 oz., diluted acetic acid 5 oz.; mix well, let stand, and filter.—LANGBECK.

ORANGE-FLOWER VINEGAR. Fresh orange-flowers $1\frac{1}{2}$ lbs., distilled vinegar 8 lbs., spirit of orange-flowers 1 lb.; macerate for 12 days, strain, and filter.

VINAIGRE VIRGINAL. Benzoin in powder 2 oz., rectified spirit 8 oz., white vinegar 2 lbs.; digest the benzoin in the spirit for 6 days, strain, and add the vinegar to the residue; macerate for 6 days, decant, and add to it the tincture; the next day filter. It is chiefly used as a cosmetic.

VINAIGRE DE COLOGNE. To each pint of eau de Cologne add an ounce of strong acetic acid.

VINAIGRE DE JOUVENCE. Spirit of cucumber 4 oz., spirit of storax 2 lbs., strong vinegar 8 lbs.

VINAIGRE DE FLORE. Equal parts of rose vinegar, vinaigre virginal, and orange-flower vinegar.

VINAIGRE DES QUATRE VOLEURS. *Thieves' Vinegar.* Dried tops of large and small (pontic) wormwood, rosemary, sage, mint, rue, lavender-flowers, of each 2 oz.; calamus root, cinnamon, cloves, nutmeg, garlic, of each $\frac{1}{4}$ oz.; camphor $\frac{1}{4}$ oz., concentrated acetic acid 2 oz., strong vine-

gar 8 lbs.; macerate the herbs, &c., in the vinegar for a fortnight, strain, press, and add the camphor dissolved in the acetic acid.

POT POURRI; SCENTED POWDERS; SACHETS OR SCENT BAGS; SCENT BALLS, PASTILS, &c.

POT POURRI. 1. Gather in the season the petals of the most fragrant kinds of roses (with which other flowers may be mixed, at pleasure, in small proportion); spread them out to dry in the sun, or in a warm room, sprinkle a little salt on them, and put them into a jar, in which they are to be kept covered up till wanted for use. Take of these rose leaves 4 oz., dried lavender-flowers 8 oz., vanilla, cloves, storax, and benzoin, all bruised, of each 1 dr., ambergris 20 grs., otto of roses 20 drops. Mix.

2. Calamus root, yellow sanders, of each 1 oz., vanilla 1 dr., musk 8 grs., ambergris 8 grs., cascarilla 1 oz., orris root 3 oz., cinnamon 1 oz., lavender flowers 1 oz., styrax 2 drs., benzoin 2 drs., cloves 2 drs., coriander seed 1 oz., nutmegs 2 drs., otto of roses 20 drops, oil of neroli 10 drops. The dry ingredients to be coarsely bruised. Mix.

3. *French*. Take the petals of the pale and red roses, pinks, violets, moss rose, orange-flower, lily of the valley, acacia-flowers, clove-gilliflowers, mignonette, heliotrope, jonquils; with a small proportion of the flowers of myrtle, balm, rosemary, and thyme; spread them out for some days, and as they become dry, put them into a jar with alternate layers of dry salt, mixed with orris powder, till the vessel is full. Close it for a month, then stir the whole up, and moisten it with rose-water.

4. Orris root 16 oz., dried acacia-flowers 8 oz., dried bergamot-peel 2 oz., musk seed $\frac{1}{2}$ oz., cloves $\frac{1}{2}$ oz.; pound them together.

5. Let rose leaves be quickly dried on a wicker tray, in a warm place. To a pint of the petals add powdered orris 2 oz., pimento $\frac{1}{2}$ oz., cascarilla $\frac{1}{2}$ oz., musk 2 grs., otto of roses 2 drops, bruised cloves $\frac{1}{2}$ oz.

SACHETS OR SCENT BAGS. The pot pourri No. 2 or 4 may be put into bags, alone, or with any perfume to increase the strength. Or coarsely powdered patchouli (a herb of the *Pogostemon* genus) may be used, with any other per-

fume. Or the bags may be filled with carded cotton mixed with any of the following scented powders.

SCENTED POWDERS, BALLS, &c.

Rose. Powdered starch 3 oz., carmine to colour, otto of roses 3 drops, orris powder 1 oz.

Violet. Orris powder 4 oz., essence of bergamot 20 drops, essence of ambergris 20 drops.

Poudre de Chypre. Oak moss is macerated in clean water for a day or two, and strongly pressed in a cloth; it is then moistened with rose-water mixed with a third of orange-flower water for two days, pressed and pulverised. It serves as a basis for other perfumes, the power of which it is said to increase.

Poudre à la Mousseline. Orris root 16 oz., coriander seed 8 oz., musk seed 2 oz., cinnamon, cloves, and sandal-wood, each 1 oz., star aniseed $\frac{1}{4}$ oz., mace, ginger, and violet ebony, of each 2 oz.; beat them to a powder, and pass through a sieve.

Poudre à l'Œillet. Red roses 48 oz., orris 48 oz., cloves 6 oz., bergamot-peel 20 oz., musk seed 24 oz., cinnamon 6 oz., long cyperus 6 oz., pale roses 26 oz., dried acacia-flowers, orange-flowers, and clove stalks, of each 8 oz.

Poudre à la Mareschale. Oak-moss in powder 2 lbs., plain starch powder 1 lb., cloves 1 oz., calamus 2 oz., cyperus 2 oz., rotten oak-wood powder 2 oz.; mix.—GRAY.

Jockey Club. Lavender flowers $\frac{1}{2}$ oz., rose leaves $1\frac{1}{2}$ oz., orris root 2 oz., vanilla beans $\frac{1}{2}$ dr., musk 4 grs., extract jasmin 2 drs., oil of sandal 20 drops, oil of neroli 5 drops, otto of rose 10 drops; mix.

Portugal. Dried orange-peel 1 oz., dried bergamot-peel $\frac{1}{2}$ oz., cloves 4 oz., storax 1 dr., ambergris 8 grs., benzoin a drachm, musk seed a scruple, musk 4 grs.

SCENTED BALLS, MEDALLIONS, &c. *Pastilles de Toilette odorantes.* These consists of perfumed powders made into a paste, and moulded to any desired form before drying. The above scent powders beaten up with mucilage of tragacanth will answer the purpose; or the following:

1. Beat the fresh petals of red roses in an iron mortar to a smooth paste, with a few drops of essence of ambergris, or other suitable perfume. It becomes sufficiently smooth to take a polish.

2. Powdered orris, oak moss, and poudre de mousseline, of each 1 oz.; lamp-black, or other colour, q. s. Form into a stiff paste, with a jelly made of 6 drs. of isinglass, 2 drs. of tragacanth, and boiling water q. s. Make it into beads by means of a pill-machine, or into any ornamental form by moulds.

3. Jessamine flowers 1 oz., powdered gum tragacanth $\frac{1}{2}$ oz., vermilion 2 oz.

4. Yellow sanders, cyperus, cloves, balsam of Peru, of each 2 drs., benzoin and styrax, of each $\frac{1}{2}$ oz., musk and civet, of each 10 grs., oil of cinnamon 5 drops, oil of rhodium 15 drops, essence of jessamine 1 dr., neroli 20 drops, ivory-black $1\frac{1}{2}$ oz., Paris plaster 2 oz., mucilage of tragacanth, made with rose-water. As the last.

PASTILS FOR BURNING. 1. Yellow sanders 3 oz., styrax 4 oz., benzoin 3 oz., olibanum 6 oz., cascarilla 6 oz., ambergris, 1 dr., Peruvian balsam 2 drs., myrrh $1\frac{1}{2}$ oz., nitre $1\frac{1}{2}$ oz., oil of cinnamon 20 drops, oil of cloves $\frac{1}{2}$ dr., otto 30 to 60 drops, oil of lavender $1\frac{1}{2}$ drs., balsam of tolu $1\frac{1}{2}$ oz., camphor $\frac{1}{2}$ oz., strong acetic acid 2 oz., charcoal 3 lbs.; mix s. a., and beat into a paste with mucilage of tragacanth, and form into conical pastils. A second and third quality may be made by using, respectively, 4 and 5 lbs. instead of 3 lbs. of charcoal. These are highly approved, but rather expensive.

2. *Clous fumans* (of the French Codex). Benzoin 2 oz., balsam of tolu $\frac{1}{2}$ oz., labdanum 1 dr., yellow sanders $\frac{1}{2}$ oz., light charcoal 6 oz., nitre $\frac{1}{4}$ oz., mucilage of tragacanth q. s. Reduce the substances to powder, and form into a paste with the mucilage, and divide into small cones with a tripod base.

3. Powdered cascarilla 8 oz., benzoin 4 oz., yellow sanders 2 oz., styrax calamita 2 oz., olibanum 2 oz., charcoal 3 lbs., nitre $1\frac{1}{2}$ oz., mucilage of tragacanth q. s.

4. Benzoin 1 oz., cascarilla 1 oz., myrrh 8 scruples, oil of nutmeg 4 scruples, oil of cloves 4 scruples, nitre $\frac{1}{2}$ oz., charcoal 6 oz., mucilage of tragacanth q. s.—**Dr PARIS.**

INCENSE. 1. Styrax $2\frac{1}{2}$ oz., benzoin 12 oz., musk 15 grs., burnt sugar $\frac{1}{2}$ oz., frankincense $2\frac{1}{2}$ oz., gum tragacanth $1\frac{1}{2}$ oz., rose-water sufficient to form a mass; to be divided into small tablets.—**Mr ASTLEY.**

2. Powdered cascarilla 2 oz., myrrh, styrax, benzoin,

thus, Burgundy pitch, each 1 oz. Mix.—Mr ATKINS (*Ph. Journal*).

MOUTH PASTILS. Dry compounds for perfuming or correcting bad odour in the breath.

Cachou Aromatisé. The basis of these compounds, as the name implies, was originally *catechu*, with which various odoriferous substances were combined. The catechu, however, is now often omitted. The following are some of the most approved forms :

1. Extract of liquorice 3 oz., oil of cloves $1\frac{1}{2}$ drs., oil of cinnamon 15 drops : mix, and divide into 1-grain pills, and silver them.

2. (M. CHEVALLIER'S.) Chocolate powder and ground coffee, of each $1\frac{1}{2}$ oz., prepared charcoal 1 oz., sugar 1 oz., vanilla (pulverised with the sugar) 1 oz., mucilage q. s. Make into lozenges of any form, of which 6 to 8 may be used daily to disinfect the breath.

3. *Cachou de Bologne.* *Bologna Catechu.* Extract of liquorice 3 oz., water 3 oz., dissolve by heat in a water-bath, and add catechu 1 oz., gum arabic $\frac{1}{2}$ oz.; evaporate to the consistence of an extract, and add (in powder) $\frac{1}{2}$ dr. each of mastic, cascarilla, charcoal, and orris : remove from the fire, and add oil of peppermint $\frac{1}{2}$ dr., essence of ambergris and essence of musk each 5 drops ; roll it flat on an oiled marble slab, and cut it into very small lozenges. Or it may be rolled into small pills, and silvered. They are chiefly used by smokers.

4. Catechu 7 drs., orris powder 40 grains, sugar 3 oz., oil of rosemary (or of peppermint, cloves, or cinnamon) 4 drops, or q. s. Proceed as for the last.

5. *Cachou Aromatisé.* Extract of liquorice and water, of each $3\frac{1}{2}$ oz.; dissolve in a water-bath, and add Bengal catechu in powder, 462 grains, and gum arabic in powder 231 grains ; evaporate to an extract, and then incorporate the following substances, first reduced to a fine powder :—Mastic, cascarilla, charcoal, and orris root, of each 30 grains ; melt the mass to a proper consistence, remove it from the fire, and then add English oil of peppermint 30 drops, tinctures of ambergris and musk, of each 5 drops ; pour it now on an oiled slab, and spread it out, by means of a roller, to the thickness of a sixpenny piece. When cool, apply some folds of blotting-paper to absorb any adherent oil, moisten the surface with water, and cover it

with sheets of silver leaf. Allow it to dry, and finally divide it into thin strips, and these again into small pieces, about the size of a fenugreek seed.—*Journal de Pharmacie*.

PASTILS OR LOZENGES, with chlorine, for disinfecting the breath. 1. Sugar flavoured with vanilla 1 oz., powdered tragacanth 20 grs., liquid chloride of soda q. s., any essential oil 2 drops. Form a paste, and divide into lozenges of 15 grs. each.

2. Dry chloride of lime 2 drs., sugar 8 oz., starch 1 oz., gum tragacanth 1 dr., carmine 2 grs. Form into small lozenges.

SKIN COSMETICS

WASHES FOR THE FACE, &c.

AQUA COSMETICA. *Cosmetic Lotion.* 1. Emulsion of bitter almonds 3 oz.; rose and orange-flower water, of each 4 oz.; borax 1 dr., tincture of benzoin 2 drs.; mix.—Dr COPLAND.

2. Elder-flower water a pint, borax $\frac{1}{2}$ oz., eau de Cologne 1 oz.; mix.

KALYDOR. The following is said to resemble KALYDOR* and GOWLAND'S lotion. Bitter almonds blanched 1 oz., corrosive sublimate 8 grs., rose-water 16 oz.

MILK OF ROSES. Sweet almonds 5 oz., bitter almonds 1 oz., rose-water $2\frac{1}{2}$ pints, white curd soap $\frac{1}{2}$ oz., oil of almonds $\frac{1}{2}$ oz., spermaceti 2 oz., white wax $\frac{1}{2}$ oz., English oil of lavender 20 drops, otto of roses 20 drops, rectified spirit a pint. Blanch the almonds, and beat them with the soap and a little of the rose-water. Melt together the oil of almonds, spermaceti, and white wax, and mix with the former into a cream, and strain it through fine muslin. Then add gradually the remaining rose-water, and lastly the spirit, with the essential oils dissolved therein.

2. A common kind is made by mixing 1 oz. of fine olive oil with ten drops of oil of tartar, and a pint of rose-water.

3. Bitter almonds 6 drs., sweet almonds 12 drs., blanch, dry, and beat up with 1 dr. of Castile soap; gradually adding 15 grs. of spermaceti, 30 grs. of white wax, and a drachm of almond oil, melted together. When thoroughly incorporated, add gradually six drops of otto of roses, dissolved in 6 oz. of rectified spirit, and 14 oz. of distilled water.

MILK OF CUCUMBERS. Made in the same manner as milk of roses, substituting juice of cucumbers for the rose-water.

* In a recent analysis of ROWLAND'S KALYDOR, the Editor failed to detect the presence of mercury.

MILK OF HOUSELEEK. Made as milk of roses, No. 1, substituting expressed juice of houseleek for a pint of the rose-water.

ALIBERT'S COSMETIC. Cucumber pomade (see further on) 3 oz., almond soap 1 oz., rose-water a quart. Mix the pomade and soap, and add the rose-water gradually.

SIEMMERLING'S COSMETIC. Make an emulsion with 1 oz. of sweet almonds, $\frac{1}{2}$ oz. bitter almonds, black cherry-water 10 oz.; and bichloride of mercury 5 grs., tincture of benzoin 5 drs., lemon juice $\frac{1}{2}$ oz.

WITHERING'S (Dr) COSMETIC. An infusion of horseradish in milk.

LAIT VIRGINAL. *Virgin's Milk.* Simple tincture of benzoin 2 drs., orange-flower water 8 oz. It may be varied by using rose or elder-flower water.

LAIT DE FRAICHEUR. Double rose-water 8 oz., tincture of benzoin 4 drs., balsam of Mecca $\frac{1}{2}$ oz.

SCHUBARTH'S COSMETIC EMULSION. Almond emulsion (made with rose-water) 8 oz., tincture of benzoin 3 drs.

ITALIAN COSMETIC WASH. Melilot water 12 oz., tincture of benzoin 2 drs.

AUGUSTIN'S. Rose-water 8 oz., salt of tartar 2 drs., tincture of benzoin 3 drs.

KITTOE'S LOTION FOR FRECKLES. 1. Chloride of ammonium 1 dr., spring-water a pint, lavender-water 2 drs. Apply with a sponge 2 or 3 times a day.

2. Sulpho-carbolate of zinc 2 parts, glycerine 25 parts, rose-water 25 parts, eau de Cologne 5 parts.

3. **WHITE'S LOTION FOR FRECKLES.** Perchloride of mercury 6 grs., diluted hydrochloric acid 1 dr., water 4 oz.; rectified spirit and rose-water, of each 2 oz., glycerine 1 oz. Apply sparingly at night, and wash off in the morning.

LEMON CREAM FOR SUNBURNS, FRECKLES, &c. Sweet cream 1 oz., new milk 8 oz., juice of lemon, brandy, or eau de Cologne 1 oz., alum 1 oz., sugar 1 dr. Boil and skim. Buttermilk is used for the same purpose.

LEMON EMBROCATION FOR FRECKLES. Borax 15 grs., lemon juice 1 oz., sugar candy $\frac{1}{2}$ dr.; mix the powders with the juice, and let them stand in the bottle, shaking occasionally, till they are dissolved.

PASTES, POMADES, COLD CREAMS, LIP-SALVE, &c.

POMMADE DE BEAUTÉ. Melt together in an earthen vessel placed in hot water, white wax $1\frac{1}{2}$ drs., spermaceti 2 drs., oil of sweet almonds $\frac{1}{2}$ oz., virgin olive oil $\frac{1}{2}$ oz., oil of poppies $\frac{1}{2}$ oz.; beat them with a few drops of balsam of Peru.

CUCUMBER POMATUM, for softening and cooling the skin. Clarified lard 4 lbs., veal suet 1 lb., juice of cucumbers 3 lbs.; melt the two former together, then beat them up assiduously with the juice. Next day, pour off the juice that has separated, and add the same quantity of fresh to the melted pomade. Repeat this six times, or until the pomade is sufficiently imbued with the odour of cucumbers. Then melt the pomade by a water-bath, and mix with it 3 drs. of powdered white starch; let it settle, and before it is too cold, pour it off into small pots, taking care not to disturb the dregs. See **UNGUENTUM CUCUMIS**, *Pocket Formulary*.

POMMADE D'HÈBÉ. Incorporate together juice of lily-bulbs 2 oz., Narbonne honey 2 oz., white wax 1 oz., rose-water 3 drs.; melt the wax with a gentle heat, and add the other ingredients. To be applied at night, and not wiped off till morning. To remove wrinkles. Probably cod-liver oil, used externally and internally, would be a more successful though less agreeable remedy.

PÂTE DIVINE DE VENUS. Mix equal parts of washed lard, fresh butter, and white honey; add balsam of Mecca and otto of roses, to perfume.

POMMADE DE NINON. Oil of sweet almonds 4 oz., washed lard 3 oz., juice of houseleek 3 oz.; mix. Softening and cooling.

POMMADE EN CRÈME. Melt together 1 dr. each of white wax and spermaceti, add oil of sweet almonds 2 oz.; pour it into a warm mortar, and gradually stir in $\frac{1}{2}$ oz. of rose or other perfumed water, and 1 dr. of tincture of tolu.

LEMON CREAM. Melt together 2 drs. of spermaceti and 1 oz. of oil of almonds; and as it cools, stir in 16 drops of essence of lemon.

COLD CREAM. 1. Oil of almonds 16 oz., white wax 4 oz.,

melt together in an earthen vessel, and when nearly cold, stir in, by little and little, 12 oz. of rose-water.

2. Melt together white wax 2 oz., oil of almonds 8 oz., and stir in 4 oz. of rose-water. Next day add 6 drops of otto of roses.

3. White wax and spermaceti, of each $\frac{1}{2}$ oz., oil of almonds 4 oz., orange-flower water 2 oz.; mix s. a.

4. As No. 3, but without the orange-flower water.

5. Lard 16 oz., white wax 2 oz., olive oil 1 oz., magistry of bismuth 1 oz.

6. White wax 1 oz., almond or olive oil 1 oz., rose-water 1 oz., glycerin 2 drs.

N.B. Those cold creams are generally preferred for present use which contain rose or other water, but they keep longer without them.

GRANULATED COLD CREAM. Melt together 1 oz. each of white wax and spermaceti, with 3 oz. of almond oil; when a little cooled, pour the mixture into a large Wedgewood mortar previously warmed, and containing about a pint of warm water. Stir briskly until the cream is well incorporated, add sufficient otto of roses to scent it, and pour the whole suddenly into a clean vessel containing 8 or 10 pints of cold water. Throw the whole on muslin, and shake out as much water as possible.—Mr OWEN

POMMADE DIVINE. Put 3 lbs. of beef marrow into an earthen vessel, and cover it with cold water, changing the water daily for a few days, and using rose-water the last day; press out the water and add to the marrow, styrax calamita, benzoin, Chio turpentine, each 4 oz.; orris powder 1 oz.; powdered cinnamon, cloves, nutmeg, of each $\frac{1}{2}$ oz. Place them in a well-tinned vessel in a water-bath, and keep the water boiling for three hours; then strain.

ALMOND PASTE FOR THE SKIN. 1. Powdered bitter almonds 4 oz., white of egg 1 oz., beat them well together to a smooth paste, with equal parts of spirits of wine and rose-water.

2. Sweet and bitter almonds, blanched, of each 2 oz.; spermaceti 2 drs., oil of almonds $\frac{1}{2}$ oz., Windsor soap $\frac{1}{2}$ oz., rose-water 1 oz., or q. s., otto of roses, and oil of bergamot, of each 12 drops; mix s. a.

3. (Camphorated.) To either of the above add 2 drs. of

powdered camphor. A few drops of oil of bitter almonds may be substituted for the otto and bergamot.

4. (*French.*) Blanch 12 oz. of bitter almonds and beat them in a mortar with a small quantity of rose or other water to a smooth paste; then add 7 oz. of rice flour, 3 oz., of bean flour, 1 oz. of orris powder, and when perfectly mixed, $\frac{1}{2}$ oz. of carbonate of potash dissolved in rose-water; again beat together, and add 3 oz. of spirituous essence of jessamine, 2 drops of oil of rhodium, and 1 of neroli.

ALMOND AND HONEY PASTE. Fine honey may be added to either of the preceding; or mix 16 oz. of clarified honey with 16 oz. of bitter almond powder; and add gradually, in alternate portions, 32 oz. of oil of almonds, and the yolks of 5 eggs.

HONEY PASTE, *Pâte au Miel.* It is sometimes made as the last; or by mixing clarified honey with cold cream, or some similar compound.

CAMPBOR BALLS, for rubbing on the hands, after washing them, to prevent chaps, &c. 1. Melt 3 drs. of spermaceti, and 4 drs. of white wax, with 1 oz. of almond oil, and stir in 3 drs. of powdered camphor. Pour the compound into small gallipots, so as to form hemispherical cakes. They may be coloured with alkanet, &c.

2. Lard 3 oz., white wax 2 oz., camphor $\frac{1}{2}$ oz.

3. Spermaceti 3 oz., white wax 1 oz., olive oil 4 oz.; melt together, and add $1\frac{1}{2}$ oz. of powdered camphor, and stir it well.

4. Melt 3 drs. of spermaceti, and 4 drs. of white wax, with 1 oz. of almond oil, and stir in 3 drs. of powdered camphor.

CAMPBOR ICE. Melt 1 dr. of spermaceti with 1 oz. of almond oil, and add 1 dr. of powdered camphor.

ALMOND POWDER. (*Cosmetic.*) This is prepared by grinding the marc or cake left after expressing the oil from sweet or bitter almonds. It is sometimes perfumed and mixed with other ingredients. It is used for cleansing the skin, and is less irritating than soap.

ALMOND WASH POWDER. 1. Almond powder (from expressed bitter almonds) 16 oz., rice flour 2 oz., powdered soap 1 oz., orris powder 1 oz., bergamot or other scent

q. p.

2. Almond powder (as above) 16 oz., powdered benzoin

$\frac{1}{4}$ oz., oil of bitter almonds 10 drops. For cleaning the hands and removing any unpleasant smell. To render it more detergent, 4 oz. of fine sand, or powdered pumice-stone, may be added.

3. GROUND MUSTARD, mixed with a little water and rubbed over the hands, removes strong odours from them. Linseed meal answers the same purpose.

ROSE LIP SALVE. 1. Oil of almonds 3 oz., alkanet $\frac{1}{2}$ oz.; digest with a gentle heat, and filter. Melt $1\frac{1}{2}$ oz. white wax and $\frac{1}{2}$ oz. spermaceti with the filtered oil, stir it until it begins to thicken, and add from 12 to 36 drops of otto of roses.

2. White wax 1 oz., oil of sweet almonds 2 oz., alkanet 1 dr.; digest till coloured, strain, and add 6 drops of otto of roses.

PERUVIAN LIP SALVE. As either of the above, substituting 20 or 30 drops of Peruvian balsam for the otto; 8 drops of oil of lavender may be added.

GRAPE LIP SALVE. *Pommade au raisin pour les lèvres.* Put into a glazed earthen pipkin $\frac{1}{2}$ lb. of fresh butter, $\frac{1}{4}$ lb. fine yellow wax, 1 oz. of alkanet, and 3 bunches of black grapes; boil together, and strain without pressure through linen.

FRENCH LIP SALVE. Lard 16 oz., white wax 2 oz.; nitre and alum in fine powder, of each $\frac{1}{2}$ oz.; alkanet to colour.

GERMAN LIP SALVE. Butter of cacao $\frac{1}{2}$ oz., oil of almonds $\frac{1}{4}$ oz.; melt together with a gentle heat, and add 6 drops of essence of lemon.

GANTS COSMETIQUES. These are white kid gloves, which have been turned inside out, and brushed over with a melted compound of wax, oil, lard, balsam, &c. The Peruvian lip salve, without any alkanet, may answer the purpose. For softening the hands.

FACE PAINTS. FARDS

FINE CARMINE (prepared from cochineal) is used alone, or reduced with starch, &c. And also the colouring matter of safflower and other vegetable colours, in the form of pink saucers, &c.

ROUGE is prepared from carmine, and the colouring matter of safflower, by mixing them with finely levigated French

chalk or talc, generally with the addition of a few drops of olive or almond oil. Sometimes fine white starch is used as the reducing ingredient. It is used in the form of powder, pomade, and *crépons*—the latter being pieces of crape imbued with the colouring matter. For common purposes, vermilion is used; and it is sometimes prepared for this purpose by mixing it with a few drops of almond oil and of mucilage of tragacanth, placing the mixture in rouge pots, and drying it by a very gentle heat.

ALMOND BLOOM. Boil 1 oz. of Brazil dust in 3 pints of distilled water, and strain; add 6 drs. of isinglass, 2 drs. of cochineal, 1 oz. of alum, and 3 drs. of borax; boil again, and strain through a fine cloth.—GRAY.

FACE WHITES. One of most innocent kind is prepared from Venetian talc, or French chalk, finely levigated. These are sometimes calcined, to increase their whiteness; but this diminishes their unctuousity and adhesiveness. Digestion with vinegar, and subsequent washing, are practised for the same purpose. Flake white (a fine variety of white lead) was formerly much used, but is now generally and justly condemned as unsafe; it is also liable to become brown under certain circumstances. Pearl or bismuth white, *blanc de perle* (magistery of bismuth*) is less injurious when pure, but is subject to the latter inconvenience. M. THENARD recommends oxide of zinc, with an equal weight of French chalk prepared by vinegar. Oxide of zinc, mixed with tallow, is almost invariably used by the members of the theatrical profession. Magnesia is said to be employed by the American ladies. White starch is used for the same purpose.

TOILET SOAPS, &c.

As the retail druggists and perfumers do not generally make their own soap in the first instance, it is only necessary to mention the means by which the soap, as it comes from the manufacturers, is prepared for the toilet.

SCENTED SOAPS, in general. Cut the best white curd soap, (or for some kinds, palm soap) into thin shavings, and

* Made by adding a little hydrochloric acid to the solution of the metal in nitric acid, and the magistery is precipitated by a small quantity of water; or the nitric solution is mixed with a weak solution of sea salt.

place it in a copper vessel, with sufficient distilled water, and heat it by a water-bath till the whole is uniformly liquefied. Let it cool to 135° F.; then add the colouring matter and perfumes. On the large scale these additions may be mixed with the liquid soap at the maker's, before it is poured into the frames. The quantity of perfume used will, of course, depend on the price at which it is to be sold.

ALMOND SOAP. To one hundredweight of the best hard white soap, melted as above, add 20 oz. of essential oil of bitter almonds. Soap really made from expressed almond oil is, we apprehend, rarely met with in commerce.

CARBOLIC SOAP. (KILNER). Cocoa-nut oil soap 75 oz., rect. spirit 10 oz., carbolic acid 6 oz., caustic potash 2 oz., oil of lemon 1 oz. Melt the soap and add to it the three last ingredients dissolved in the spirits. Mix well, and pour into moulds.

SAVON AU BOUQUET. Melt 60 lbs. of white curd soap as above, and 8 oz. of oil of bergamot, 1½ oz. each of oils of cloves, sassafras, and thyme, ¾ oz. of neroli, and 14 oz. or q. s. of brown ochre.

ROSE SOAP. Put into a copper vessel, placed in boiling water, 20 lbs. of white curd soap, and 30 lbs. of olive oil soap, both in thin shavings; add 5 lbs. of soft water, or rose-water; keep the heat below boiling till the soap is uniformly liquefied, then add 12 oz. of finely sifted vermilion, or enough to produce the required tint. Withdraw it from the fire, and when sufficiently cool, add ½ oz. of otto of roses, ½ oz. of oil of cloves, ½ oz. of oil of cinnamon, and 2½ oz. bergamot. For a cheaper article use less perfume.

WINDSOR SOAP. This is said to be made with lard. In France they use lard with a portion of olive or bleached palm oil. Dr PEREIRA states that it is made with one part of olive oil to nine of tallow. But a great part of what is sold is probably only curd (tallow) soap, scented with oil of caraway and bergamot. The brown is probably coloured with burnt sugar, or umber.

HONEY SOAP. White curd soap 1½ lbs., brown Windsor soap ½ lb.; cut them into thin shavings, and liquefy as directed above for scented soaps; then add 4 oz. of honey, and keep it melted till most of the water is evaporated; then

remove from the fire, and when cool enough, add any essential oil. According to PIESSE the *honey soap* usually sold consists of fine yellow soap perfumed with oil of citronella.

FLOATING SOAPS. These are made by liquefying, as described above, 30 lbs. of oil soap with about 5 lbs. of water, and agitating the mixture, by a suitable wooden apparatus turned by a handle, till the froth arises to the top of the vessel. It is then put into frames to cool. These soaps are variously perfumed and coloured.

TRANSPARENT SOAP. Cut fine white curd soap into thin shavings, and dry them with a gentle heat till they can be reduced to powder. Put 2 lbs. of this powder into a water-bath with 5 or 6 pints of rectified spirit of wine, and heat it gently (taking care that the water does not quite boil) till the solution is complete; add the perfume and pour into the frames. When cold, cut it into squares. They must be kept some time in a dry place before they can attain their full degree of transparency. By using a still, most of the spirit may be recovered for future use.

WASH BALLS. Savonettes. These are made from various kinds of soap, usually with the addition of powdered starch, or hair-powder, or of rice flour, together with perfuming and colouring ingredients. They are formed into spherical balls by taking a mass of the prepared soap in the left hand, and a conical drinking glass with rather thin edges in the right. By turning the glass and ball of soap in every direction, the rounded form is soon given; when dry, the surface is scraped to render it more smooth and even. One or two examples of this kind of soap will suffice.

COMMON OR LEMON WASH BALLS. Cut 6 lbs. of soap into very small pieces; melt it in a pint of water in which 6 lemons have been boiled. When melted, withdraw the soap from the fire, and add 3 lbs. of powdered starch, and a little essence of lemon; knead the whole into a paste and form into balls of the desired size.

CREAM WASH BALLS. White curd soap 7 lbs., powdered starch 1 lb., water or rose-water q. s. Beat the whole together, and form into balls.—GRAY.

CAMPHOR WASH BALLS. White soap 1 lb., spermaceti 1 oz., water q. s.; melt together and add 1 oz. of powdered camphor.

MRS SYMOND'S SOAP PASTE FOR THE HANDS. Best soft soap (from olive oil and potash) 16 oz., spermaceti 4 oz., best olive oil 1 oz., camphor $\frac{1}{4}$ oz., rectified spirit $\frac{1}{2}$ oz., soft water 1 pint, essence of lemon $\frac{1}{4}$ oz.; mix S.A. With 8 oz. of pumice-stone, powdered and sifted through fine book muslin, it forms sand soap paste.

POWDERED SOAP. Any of the hard soaps may be pulverised, if first cut into thin shavings, and kept at a gentle heat, till sufficiently dry. This process renders the soap more mild.

SHAVING POWDER. Melt together in a water-bath 1 lb. of white soap with 1 oz. of powdered spermaceti and $\frac{1}{4}$ oz. of chlorate of potash dissolved in a little water, or rose-water. Pour the liquefied soap into a shallow mould; when solidified, shave it fine and dry as above.

SHAVING PASTE. 1. Melt together 1 dr. each of spermaceti, white wax, and almond oil; beat it up with 2 oz. of the best white soap, and a little lavender or Cologne water.

2. Naples soap, beaten up with sufficient powdered soap to form a stiff paste.

3. White soft soap 4 oz., powdered Castile soap 1 oz., oil of olives or almonds $\frac{1}{4}$ oz.

SHAVING LIQUID. *Essence of Soap.* 1. White soap 3 oz., proof spirit 8 oz., distilled water 4 oz., carbonate of potash 1 dr., essence of lemon q. s. Dissolve the soap without heat, and add the potash and essence.

2. (*Italian Essence of Soap.*) White curd or Windsor soap 10 parts, rectified spirit 34 parts, rose or orange-flower water 34 parts. Digest with a gentle heat and filter.

3. Naples soap, or white soft soap, 16 oz., oil of olives $\frac{1}{2}$ oz., gum benzoin 1 dr., rectified spirit 24 oz. Digest. Rub a few drops on the beard, followed by warm water.

HAIR COSMETICS

HAIR POWDER. The basis of hair powder is finely powdered starch. It is variously scented, and was formerly tinted of various colours. The plain and violet hair powders are now principally used. The latter is perfumed with orris powder, or essence of violets, usually with the addition of bergamot, &c. GRAY gives the following species for scenting hair powder: powdered orris 1 lb., essence of bergamot 12 oz., oil of neroli 1 dr., musk 1 scruple. Hair powder is also perfumed with jessamine, roses, &c., by mixing the flowers with plain powder for 2 or 3 days, stirring the mixture twice or thrice a day, and then sifting out the powder from the flowers.

COMPOUNDS TO PROMOTE THE GROWTH OF THE HAIR

POMADES FOR THE CURE OF BALDNESS

1. **DUPUYTREN'S POMADE.** The recipe given by BATEMAN and RENNIE for this celebrated preparation, viz. almond oil, lard, suet, and essential oils, is remarkable as entirely omitting the active ingredient. It is probable that the preparation first employed by M. DUPUYTREN was more simple in its form than what he subsequently adopted, but cantharides was always the essential constituent. The first formula met with was: Tincture of cantharides (made according to the Paris Codex, 1 part flies to 8 of proof spirit) 1 part, lard 9 parts. The following are said more nearly to represent the compound in its improved and more elegant form. M. CAP prescribes: Beef marrow 2 oz., spirituous extract of cantharides (made by evaporating the above tincture) 8 grs., rose oil 1 dr., essence of lemon 50 drops. M. FONTAINE directs: Beef marrow 4 oz., calomel $2\frac{1}{2}$ drs., extract of cantharides 18 grs., attar of roses 2 drops. But the following by M. RECLUZ is said to have been acknowledged by M. DUPUYTREN as the true formula: Beef marrow 6

oz., nervine balsam* 2 oz., Peruvian balsam 2 oz., oil of almonds $1\frac{1}{2}$ oz., extract of cantharides 16 grs.; melt the marrow and nervine balsam with the oil, strain, add the balsam of Peru, and lastly the extract, dissolved in a drachm of rectified spirit. M. GUIBOUT says that no better than the following can be used: Beef marrow 1 oz., nervine balsam 1 oz., rose oil 1 dr., extract of cantharides (dissolved in spirit) 6 grs. These pomades should be rubbed on the scalp once or twice a day for some weeks. If any soreness is produced the pomade should be less frequently applied.

2. POMMADE CONTRE L'ALOPÉCIE. Fresh lemon-juice 1 dr., extract of bark (by cold water) 2 drs., marrow 2 oz., tincture of cantharides (as above) 1 dr., oil of lemon 20 drops, oil of bergamot 10 drops; mix. First wash the head with soap and water, containing a little eau de Cologne, then rub it dry. Next morning rub in a small lump of pomade, and repeat it daily. In 4 or 5 weeks the cure of baldness is effected.—Dr SCHNEIDER.

3. CAZENAVE'S REMEDY FOR BALDNESS. Beef marrow 1 oz., tincture of cantharides (as above) 1 dr., powdered cinnamon 1 dr. To be applied night and morning, the head being first washed with salt and water. Keep the hair short.

4. Dr CATTELL'S is the same, substituting 10 drops each of oils of origanum and bergamot for cinnamon.

5. Beef marrow 1 oz., castor oil $\frac{1}{2}$ oz., tincture of cantharides 1 dr., essential oil of bitter almonds and of lemon, each 12 drops.

6. Beef marrow 3 drs., almond oil 1 dr., sulphate of quinine 15 grs., otto 2 drops.—SOUBEIRAN.

7. Prepared lard 2 oz., white wax 2 drs.; melt together, remove from the fire, and add 2 drs. balsam of tolu, 20 drops of oil of rosemary, and in chronic cases 1 dr. of tincture of cantharides.—Dr NELIGAN.

8. Camphor 1 scruple, citrine ointment 2 drs., spermaceti cerate 6 drs.; mix. To be applied every night.

9. BATE'S *Unguentum Criniscum*. Labdanum 6 drs., bears'-grease 2 oz., honey $\frac{1}{2}$ oz., powdered southernwood 3 drs., ashes of red-root $1\frac{1}{2}$ drs., oil of nutmeg 1 dr., bal-

* This is made by melting together 4 oz. each of beef marrow and oil of mace, and adding 2 drs. of balsam of tolu, and 1 dr. each of oil of cloves and camphor, dissolved in $\frac{1}{2}$ oz. of rectified spirit.

sam of Peru 3 drs.; mix. Let the bald part be first rubbed with an onion till it is red, then apply the ointment. It should be used daily, or oftener, for 5 or 6 weeks.

10. Box leaves 2 oz., southernwood 2 oz., lard, marrow, or bears'-grease 8 oz.; digest together by the heat of a water-bath, and strain.

11. *Bears'-grease*. The most approved consists of 2 parts of prepared bears' fat, with 1 of beef marrow, scented at pleasure. We have placed this, on the ground of common report, among the preparations which may possess some efficacy, but reserve the compounds usually sold under this name for the POMATUMS. See further on.

12. *Pommade Philocome*. Powdered cinchona $\frac{1}{2}$ dr., oil of almonds 2 drs., beef marrow 6 drs., oil of bergamot 6 drops; balsam of Peru 20 drops; mix.—DOURVAULT.

13. Vaseline $\frac{1}{2}$ oz., castor oil $\frac{1}{2}$ oz., tincture cantharides 1 dr., water of ammonia $\frac{1}{2}$ dr., oil of rosemary 5 drops; mix. Apply to the roots of the hair two or three times a week, rubbing in well.

LIQUID COMPOUNDS FOR THE CURE AND PREVENTION OF BALDNESS

1. Dr LOCOCK'S *Lotions*. Oil of mace (expressed oil of nutmeg) $\frac{1}{2}$ oz., olive oil 2 drs., water of ammonia $\frac{1}{2}$ dr., spirit of rosemary 1 oz., rose-water $2\frac{1}{2}$ oz.; mix. Mr ASTLEY recommends the following modification: Oil of mace $\frac{1}{4}$ oz., olive oil 2 drs., oil of rosemary 4 drops; incorporate them carefully, then add gradually $3\frac{1}{2}$ oz. of rose-water, 2 drs. of solution of carbonate of ammonia, and 2 drs. of rectified spirit.

2. SIR ERASMUS WILSON'S. Eau de Cologne 2 oz., tincture of cantharides 2 drs.; oil of rosemary and oil of lavender, of each 10 drops.

3. Mr ACTON'S. Equal parts of rectified spirit, castor oil, and eau de Cologne.

4. Mr ACTON'S, *stronger*. Equal parts of honey-water and tincture of cantharides.

5. Tincture of cantharides 3 drs., acetate of copper 3 grs., oil of almonds and castor oil, of each a fl. oz., with any essential oil to scent it. A small quantity to be applied to the roots of the hair every morning.

6. Vinegar of cantharides (Lond. Pharm.) $\frac{1}{2}$ oz., eau de Cologne 1 oz., rose-water 1 oz.; mix.

7. Castor oil, lavender-water, and tincture of cantharides, in equal quantities.

8. Glycerine 2 oz., tincture of myrrh 1 oz., eau de Cologne 1 oz., tincture of cantharides $\frac{1}{2}$ oz., distilled water 24 oz.—*Pharm. Journ.*

9. *American Shampoo Liquor.* Rum 3 quarts, spirit of wine 1 pint, water 1 pint, tincture of cantharides $\frac{1}{2}$ oz., carbonate of ammonia $\frac{1}{2}$ oz., salt of tartar 1 oz. Rub it on, and afterwards wash with water. By omitting the salt of tartar it nearly resembles balm of Columbia.

10. **FRENCH RECEIPT.** Rum 500 parts, alcohol 75 parts, distilled water 75 parts, tincture of cantharides 3 parts,* carbonate of potash 3 parts, carbonate of ammonia 5 parts. Dissolve the salts, mix the liquids, and filter. Wet the skin of the head with this lotion for several minutes, then wash it with water.

11. **Dr LANDERER'S.** Bay leaves 2 oz., cloves $\frac{1}{2}$ oz., spirit of lavender 4 oz., spirit of thyme 2 oz.; digest for 6 days, filter, and add $\frac{1}{2}$ oz. of ether. To be rubbed on every morning.

12. Put into a still 4 lbs. of honey, 12 handfuls of the tendrils of vine, and the same of rosemary tops; distil very slowly till the liquor begins to taste sour.

13. **Dr CATTELL'S.** See **WASHES FOR THE HAIR**, further on.

14. A few drops of the best paraffin oil well rubbed into the roots of the hair or the bald part twice or thrice a week. Ought not to be used by candle-light.

Note.—The above ointments and liquids require to be used for some weeks, in order to produce a decided effect, either in curing or preventing baldness. Those which contain cantharides in any form are the most active, and must be used with caution. They should be applied once or twice a day, according to the effect produced; but if the scalp becomes sore, their use must be omitted for a time, or longer intervals allowed, as the case may require. When employed to prevent the hair falling off, or from becoming grey, they need not be applied so frequently as for baldness.

* Made according to the French Codex.

The following require no particular caution, being less active than the preceding.

POMATUMS, LOTIONS, &c.

FOR EMBELLISHING, STRENGTHENING AND CLEANSING
THE HAIR

Pomatums, or Pomades

These are composed usually of animal fats, variously perfumed. The lard, veal fat, beef and mutton suet, bears' fat, and beef marrow, employed for this purpose, require to be prepared with great care. The following is, perhaps, the best mode: Cut the raw fat into pieces, carefully removing the fleshy and bloody portions of membrane, &c., and beat it in a marble mortar; melt it in a well-tinned vessel placed in boiling water, and strain the melted fat through a hair-sieve without pressure (reserving the residue to be heated again and pressed for more fat, to be used for commoner purposes). Keep the melted fat for some time gently warm, without disturbing it; remove any scum which may have arisen, and pour off the clear fat, taking care that none of the dregs or watery liquid which have subsided, pass with it. A mixture of these fats forms the basis of many varieties of pomades. Sometimes a little white wax is added. A greater degree of whiteness is said to be given, by adding to the liquefied fat a few grains of citric acid. The same end is promoted by assiduously beating the pomade, while cooling, with a wooden spatula.

To perfume pomatums, various essential oils, &c., are added (see COMMON POMATUM); but the finer sorts are perfumed by infusing fresh flowers in the melted fats for some hours, and straining; or, in other cases, the simple pomade is thinly spread on plates of glass set in frames, and the fresh flowers stuck in the scored surface of the fat, changing the flowers daily till the pomatum is sufficiently perfumed. As these compounds can seldom be prepared to advantage by the retailer, a few varieties only require to be noticed here.

COMMON POMATUM. Mutton suet (prepared as above) 1 lb., common lard 3 lbs.; melt together in a water-bath, pour it into an earthen basin, and beat it assiduously with a

wooden spatula. When sufficiently cool, add 2 oz. or q. s. of essence of bergamot or of lemon, and continue the stirring till nearly cold.

ROSE POMATUM. Prepared lard 16 oz., prepared suet 2 oz.; melt with a gentle heat, and add 2 oz. of rose-water, and 6 drops of otto of roses. Beat them well together, and pour into pots before it is too cold. For making jessamine, violet, and orange pomade put the same quantity of water, and 1 dr. of the essence.

MARROW POMATUM. Beef marrow and beef suet, coloured with a little annotto, may be employed for this and other yellow pomatums. For the perfumes employed for these and other pomatums, see **ESSENCES FOR SCENTING POMATUMS**, under **PERFUMERY**.

POMADE FOR BEAUTIFYING THE HAIR. Oil of sweet almonds a pint, spermaceti 1½ oz., purified lard 2 oz.; melt with a gentle heat; when nearly cold, add any agreeable scent, and pour it into pots or wide-mouthed bottles.

BEARS' GREASE (ARTIFICIAL). Bears' grease is imitated by a mixture of prepared veal suet and beef marrow. It may be scented at pleasure; oil of lavender, with a very little oil of thyme, is sometimes used. The following are some of the compounds sold under this name:

1. Prepared suet 3 oz., lard 1 oz., olive oil 1 oz., oil of cloves 10 drops, compound tincture of benzoin 1 dr.; mix.
2. Lard 1 lb., solution of carbonate of potash 2 oz.
3. Olive oil 4 flasks, white wax 4 oz., spermaceti 2 oz.; scented with otto of roses and oil of bitter almonds.

GREEN BEARS' GREASE. Bears' grease digested with fresh walnut leaves, and strained. This is repeated with more leaves till the pomade is sufficiently coloured; it is then scented with oils of rosemary, thyme, and bergamot.

GERMAN POMADE, FOR STRENGTHENING THE HAIR. Take 8 oz. of purified marrow, melt it in a glass or stoneware vessel, and add 1½ oz. of fresh bay leaves, 1 oz. of orange leaves, 1 oz. of bitter almonds, ½ oz. of nutmegs, ½ oz. of cloves, and 1 dr. of vanilla, all bruised; cover the vessel, and let the whole digest for 24 hours, with a gentle heat; strain while warm through linen, and stir it as it cools.

HARD OR ROLL POMATUM. 1. Suet 5 lbs., white wax 8 oz., spermaceti 2 oz., oil of lavender and essence of ambergris, of each ½ oz.

2. Beef suet 16 oz., white or yellow wax 1 oz., with 1 dr. of oil of lavender or of bergamot.

3. Lard melted with one third or half its weight of white wax, and poured into semi-cylindrical paper moulds when nearly set. This is sold under the name of *cosmetique*. It is sometimes coloured to match the hair. See further on.

COLOURED POMATUMS. The colouring matters employed are annatto, alkanet, marigold, carmine, gamboge, indigo, cobalt blue, umber, ivory black, &c.

CIRCISSIAN CREAM. Two flasks of oil, 3 oz. of white wax, 2 oz. of spermaceti, $\frac{1}{2}$ oz. of alkanet root. Digest the oil with the alkanet till coloured, strain, melt the wax and spermaceti with the oil, and when sufficiently cool add $2\frac{1}{2}$ drs. of English oil of lavender, and $\frac{1}{2}$ dr. of essence of ambergris.

COSMOLIN (VASELIN) CREAM. Cosmolin 24 oz. (troy), white wax 12 oz. (troy), spermaceti 12 oz. (troy), glycerine 3 fl. oz., oil of geranium 1 dr. Melt the wax and spermaceti, add the cosmolin, then stir until nearly cold, add the glycerine and oil, and stir until cold.

CRYSTALLINE CREAM. Oil of almonds 8 oz., spermaceti 1 oz.; melt together; when a little cooled add $\frac{1}{2}$ oz. or less of essence of bergamot, or other perfume; put it into wide-mouthed bottles, and let it stand till cold.

Camphorated crystalline cream may be made by using camphorated oil (*Lin. Camphoræ*) instead of oil of almonds.

CASTOR OIL POMADE. Castor oil 4 oz., prepared lard 2 oz., white wax 6 drs., essence of bergamot 2 drs., oil of lavender 20 drops, eau de Cologne $\frac{1}{2}$ oz.; stir till cold.

CRYSTALLINE CASTOR OIL POMADE. Castor oil 16 oz., spermaceti $1\frac{1}{4}$ oz.; melt together, and when a little cool, add 1 oz. of essence of bergamot, $\frac{1}{2}$ dr. oil of verbena, $\frac{1}{2}$ dr. oil of lavender; pour it into wide-mouthed bottles, and let it stand till cold.

CASTOR OIL AND GLYCERINE POMADE. Dissolve white wax $1\frac{1}{2}$ oz. with a moderate heat in 3 oz. of castor oil, triturate with 9 oz. of castor oil and 2 oz. glycerine until cool. Then add essence of lemon 5 drs., ess. of bergamot 2 drs., oil of lavender 1 dr., oil of cloves 10 drops. Rub annatto gr. x with 1 dr. of water till smoothly suspended, add 1

dr. of alcohol, and stir the colouring into the pomade until thoroughly mixed.—AMERICAN RECIPE.

FOX'S CREAM. Marrow pomatum 2 oz., oil of almonds 2 oz.; melt, and add while cooling, with constant stirring, essence of jessamine or of bergamot 2 drs.—BATEMAN.

GLYCERINE AND LIME CREAM. Glycerine, oil of sweet almonds, and lime water, of each 8 oz., tincture of cantharides 1 oz. Perfume as desired.

LIME JUICE AND GLYCERINE. Lime juice $\frac{1}{2}$ pint, rose-water $\frac{1}{4}$ pint, glycerine 2 oz., rect. spirit 2 oz., oil of lemon 30 drops.

QUILLAI BARK. Stir a piece of the bark in some water till a strong lather is produced, and rub it on the hair with a rough towel. Thus used it is an excellent remedy for dandruff.

HUILES ANTIQUES

PERFUMED OILS FOR THE HAIR

The basis of these oils is either almond oil, olive oil, or oil of ben; whichever is used should be perfectly fresh, and of the finest quality. The perfume is communicated in three ways: by infusing the flowers in the oil at a gentle heat; by placing layers of flowers alternately with folded cotton soaked in the oil, in proper frames, and pressing out the oil when sufficiently imbued with the odour of the flower; or simply by adding essential oils, &c., to the fixed oil. An example or two of each method will be sufficient.

OIL OF ROSES, BY INFUSION. Heat in a water-bath 1 lb. of virgin olive oil, and 1 lb. of picked fresh petals of Provence roses. Let these remain together in a water-bath for half an hour, then remove from the bath, and leave them together for 24 hours, stirring them twice during the time. Strain through a cloth, and express all the oil. To this oil add fresh roses, and proceed as before; repeating this for 5, 6, or 7 times, till the oil is sufficiently perfumed.

OIL OF JESSAMINE, PERFUMED WITH THE FLOWERS. Fold pieces of white cotton cloth twice or four times; moisten them with fine olive oil, slightly pressing them, and place them in proper frames. Then place on the cloths a rather thick layer of freshly-gathered and dry jessamine flowers, carefully deprived of all green parts. In 24 hours care-

fully remove the flowers, and replace them by fresh ones, till the oil is sufficiently perfumed. The oil is then expressed. The same method is employed in preparing oils from other delicate flowers; such as violet, lily of the valley, &c.

OIL OF ROSES, COMMON. Fine olive or almond oil a pint, otto of roses 16 drops. If required red, colour the oil with alkanet root, and strain before adding the otto. For common sale, essence of bergamot or of lemon is often substituted, wholly or in part, for the more expensive otto.

PERFUMED OIL OF BERGAMOT, LEMON, ORANGE, &c. To oil of ben, or finest almond or olive oil, add essential oil of bergamot, lemon, &c., q. s. For common purposes a drachm of the essential oil may be added to 16 oz. of oil. Some recipes, however, direct as much as $1\frac{1}{2}$ drms. or 2 drms.

OIL OF AMBERGRIS AND MUSK. Ambergris 2 drs., musk $\frac{1}{2}$ dr.; grind them together in a mortar, then with a small quantity of oil; add more oil to make up a pint, and let them stand together for 12 days, stirring them occasionally. Then decant or filter. Add half a pint of oil to the residue for an oil of second quality.

COMMON OIL OF MUSK, OIL OF BENZOIN, OIL OF STYRAX, &c., may be obtained by mixing a strong tincture of these drugs with fine oil, agitating them frequently together, and after remaining some hours at rest, decanting the clear oil.

HUILE COMOGÈNE. Mix equal parts of oil and spirit of rosemary with a few drops of oil of nutmeg. To be used daily.

HUILE DE PHÉNIX. Clarified beef marrow 4 oz., lard 2 oz., oil of mace 4 oz.; melt together, and strain through linen into a warm mortar; stir, and when it begins to cool add the following solution, and stir constantly till it is quite cold: oil of cloves, lavender, mint, rosemary, sage, and thyme, of each $\frac{1}{2}$ dr.; balsam of tolu 4 drs., camphor 1 dr., rectified spirit 1 oz. Put the spirit and balsam into a phial, and place it in warm water till the solution is complete, then add the camphor and essential oils.

HUILE PHILOCOME D'AUBRIL. Triturate together, without heat, equal parts of cold-drawn nut oil, almond oil, and prepared beef marrow, adding any essential oil as a perfume.

HUILE VERTE. Macerate 1 dr. of guaiacum with 1 lb. of olive oil; strain, and add any essential oil to perfume it.
—GRAY.

MARROW OIL. Clarified beef marrow, or marrow pomatum, with enough almond or olive oil to bring it to the desired consistence.

FLUIDE DE JAVA. This consists of beef marrow, white wax, fine olive oil, and essential oils at pleasure.

MACASSAR OIL. The oil made by the natives in the island is obtained by boiling the kernel of the fruit of a tree resembling the walnut, called in Malay *badeau*. The oil is mixed with other ingredients, and has a smell approaching to that of creasote. But the Macassar oil sold in this country has probably no relation to the above, except in name. The following is given by GRAY: Olive oil 1 lb., oil of origanum 1 dr.; others add $1\frac{1}{2}$ drs. of oil of rosemary. The following French compound is probably named Macassar oil rather to denote its properties than from any resemblance either to the product of Macassar, or to the oil sold under this name in England.

HUILE DE MACASSAR DE NAQUET. Oil of ben 14 pints, nut oil 7 pints, spirits of wine 1 quart, essence of bergamot 3 oz., tincture of musk 3 oz., spirit of orange (*esprit de Portugal*) 2 oz., otto of roses 2 drs., alkanet to colour it. Digest them together with a gentle heat for an hour, and shake frequently for a week.

BRILLIANTINE. A solution of castor oil in eau de Cologne, 1 part in 4. Another formula is as follows:—Glycerine and eau de Cologne, of each 1 part; honey, 2 parts; rectified spirit 4 parts.

WASHES FOR THE HAIR

VEGETABLE EXTRACT FOR CLEANSING AND STRENGTHENING THE HAIR. 1. Southernwood 2 oz., box leaves 6 oz., water $\frac{1}{2}$ pints. Boil gently in a saucepan for $\frac{1}{4}$ of an hour, strain, and to each pint of the liquid add 2 oz. of spirit of rosemary and $\frac{1}{2}$ dr. of salt of tartar or 1 dr. of Naples soap.

2. Boil 1 lb. of rosemary in 2 quarts of water, and add to the filtered liquor 1 oz. of spirit of lavender, and $\frac{1}{4}$ oz. of Naples soap, or salt of tartar.

3. Incinerate 2 oz. each of rosemary, maidenhair, southern-

wood, myrtle berries, and hazel bark; make a strong solution of the ashes, with which wash the hair at the roots every day. Keep the hair short.—Dr CATTELL.

4. Borax 1 oz.; powdered camphor $\frac{1}{2}$ oz., boiling water a quart. When cold, filter for use. Damp the hair with it frequently.

WASH FOR REMOVING SCURF AND PROMOTING THE CURLING OF THE HAIR. 1. Beat up the yolk of an egg with a pint of clean rain-water. Apply it warm; and afterwards wash the head with warm water.

2. Lime-water a pint, distilled vinegar $\frac{1}{4}$ of a pint; mix.

WASH FOR IRRITABLE ERUPTIONS OF THE SCALP OR FACE.
Dr HEADLAND. Rimmel's toilet vinegar 6 oz., glycerine 2 oz., carbolic acid $\frac{1}{4}$ dr.

COMPOUNDS FOR STIFFENING THE HAIR

EAU COLLANTE. Dissolve without heat 8 oz. of clear gum in 2 lbs. of distilled or rose water, and filter through coarse filtering paper.

BANDOLINE, OR FIXATEUR. Vegetable mucilage, with sufficient spirit to preserve it. Mucilage of quince seed is used; mucilage of picked Irish moss, carefully strained, is said to answer still better. The following is employed by some London perfumers: Finest picked gum tragacanth, reduced to a coarse powder, 1 oz., rose-water a pint; put them into a wide-mouthed vessel, and shake them together daily for 2 or 3 days; then strain with gentle pressure through fine linen or cambric. If required to be coloured, infuse cochineal in the water employed, before making the mucilage. Another form is—linseed (not bruised) a tablespoonful, water $\frac{1}{2}$ pint; boil for 5 minutes and strain.

POMMADE COLLANTE, FOR FALSE CURLS. Melt together in an earthen pipkin 24 oz. of fine Burgundy pitch and 8 oz. of white wax, and add 1 oz. of pomatum; remove from the fire and add 4 oz. of brandy or other spirit, replace it on the fire till it boils slightly, then strain through linen, adding bergamot or other perfume, and cast it into moulds.

HAIR DYES

*** It may be well to remind our readers that all medical authorities strongly condemn the use of lead in hair dyes.*

1. **ORFILA'S HAIR DYE.** Take 3 parts of litharge and 2 of quicklime, both in an impalpable powder, and mix them carefully. When used, a portion of the powder is mixed with hot water or milk, and applied to the hair, the part being afterwards enveloped in oil-skin or a cabbage-leaf for 4 or 5 hours.

2. Litharge 2 parts, slaked lime 1 part, chalk 2 parts, all finely powdered, and accurately mixed. When required for use, mix the powder with warm water, and dip a brush into the mixture, and rub the hair well with it. After 2 hours, let the hair be washed.

3. Litharge 4½ oz., quicklime ¾ oz.; reduce to an impalpable powder, and pass it through a sieve. Keep it in a dry, well-closed bottle. Wash the hair first with soap and water, then with tepid water; wipe it dry, and comb with a clean comb. Mix the dye in a saucer with hot water to the consistence of cream, and apply it to the hair, beginning at the roots. Place over it four folds of brown paper, saturated with hot water, and drained till cool; and over this an oilskin cap and a nightcap. Let it remain from 4 to 8 hours, according to the shade required. When removed, oil the hair, but do not wet it for 3 or 4 days.

4. **CHEVALLIER'S.** Mix 5 drs. of freshly slaked lime with 1½ oz. of water, and strain through silk; put the milk of lime into a 4-oz. bottle. Dissolve 5 grs. of acetate of lead in sufficient water, and add enough slaked lime to saturate the acetic acid (a drachm or rather more), let it settle, pour off the supernatant liquor, wash the precipitate with water, and add it to the milk of lime.

5. **CHESTNUT HAIR DYE.** *a.* We have met with the following, but do not guarantee it: Permanganate of potash gives the hair a beautiful chestnut-brown colour, varying according to the strength of the solution of the salt. A good formula is: Permanganate of potash 1 dr., powdered gum arabic 2 drs., rose-water 3 oz.; mix. Apply carefully with a tooth brush, so as to avoid staining the skin.—*Chemist and Druggist.*

b. Citrate of bismuth 1 oz., rose-water 2 oz., distilled

water 2 oz., rectified spirit 5 drs., water of ammonia q. s. Apply as usual, and follow with hyposulphite of soda 12 drs., distilled water 4 oz.; mix.

6. HAIR RESTORER. Sulphur 45 grs., acetate of lead 20 grs., glycerine $\frac{1}{2}$ oz., water to 10 oz.; mix.

7. GOLDEN HAIR DYE. *Aureoline*. A solution of peroxide of hydrogen in water, containing from 3 to 6 per cent., by weight, of the peroxide.

8. DARK BROWN HAIR DYE. *a*. Pyrogallic acid 4 grs., distilled water 2 oz.; mix. Apply this with a tooth-brush for 16 or 18 minutes, then brush the hair over with *b*.

b. Crystallised nitrate of silver 1 dr., gum arabic 1 dr., distilled water 2 oz.; mix.

9. Dr HANMANN'S. Levigated litharge 11 oz., powdered quicklime 75 oz., hair powder 37 oz.; mix. When used, a portion of the powder is mixed with warm water in a saucer, and applied to the hair with the fingers, taking care to cover the hair to the roots. Cover the whole with a sheet of cotton wadding moistened with water, and this with a folded cloth. Let it remain on for 3 hours; or better, for the night.

10. WARREN'S. Sifted lime 16 oz., white lead 2 oz., litharge in fine powder 1 oz.; mix well together, and keep dry. To dye *black*, mix a little powder with water to the consistence of cream. To dye *brown*, use milk instead of water. Apply with a small sponge.

ESSENCE OF TYRE. GRECIAN WATER. EAU D'EGYPT, EAU DE CHINE. These are solutions of nitrate of silver; in applying them it must be remembered that they stain the *skin* as well as the hair. Hence there is more difficulty in applying than with the preceding; but they are considered to impart a finer colour to the hair, with the disadvantage, however, of rendering it dry and crisp. The following are some of the most approved formulæ:

1. Dr CATTELL'S. Nitrate of silver 11 drs., nitric acid 1 dr., distilled water 1 pint, sap green 3 drs., gum arabic 1 dr.; mix.

2. Nitric acid 1 dr., nitrate of silver 10 drs., sap green 9 drs., mucilage 5 drs., distilled water 37 $\frac{1}{2}$ fl. oz.

3. Silver 2 drs., iron filings 4 drs., nitric acid 1 oz., distilled water 8 oz. Digest, and decant the clear solution. To be carefully applied with a close brush.

4. Hydrosulphate of ammonia 1 oz., solution of potash 3 drs., distilled water 1 oz.; mix. Apply this with a tooth-brush for 16 or 20 minutes, then brush the hair over with the following: Nitrate of silver 1 dr., distilled water 2 oz., using a clean comb to separate the hair.

MERCURIAL BLACK DYE. A weak solution of perchloride of mercury, used for some days, followed by a wash containing hyposulphite of soda, *not without danger*.

BROWN HAIR DYE. Acetate of lead 2 drs., hyposulphite of soda 1 dr., rose-water 14 oz., glycerine 2 oz. Dissolve the acetate of lead and hyposulphite in separate portions of the rose-water; filter separately, mix the solutions and add the glycerine.

PYROGALLIC STAIN. Distil coarsely powdered nutgalls in a retort, dissolve the solid acid which sublimes in a little hot water, add the solution to the acid liquid which afterwards passes over, separate the floating oil, shake the liquid with charcoal, filter, and add a little spirit.

TINCTURE OF WALNUT. A strong tincture of the shells of green walnuts, scented.

LA FOREST'S COSMETIC WASH FOR THE HAIR. Red wine 1 lb., salt 1 dr., sulphate of iron 2 drs. Boil for a few minutes, and add common verdigris 1 dr.; leave it on the fire for 2 minutes, withdraw it, and add 2 drs. of powdered nutgall. Rub the hair with the liquid; in a few minutes dry it with a warm cloth, and afterwards wash with water.

POMATUMS, OR COSMETIQUES, in sticks for the hair.

BLACK POMATUM, in sticks, for the eyebrows, whiskers, &c. Prepared lard melted with a third of its weight of wax in winter, or half in summer, is coloured with levigated ivory black, and strained through tammy, or any material which will permit the fine particles of ivory black to pass through. Stir it constantly, and when it begins to thicken pour it into paper moulds.

BROWN AND CHESTNUT POMATUMS are prepared in the same way, but coloured with umber, &c. The white variety is made as Hard Pomatum.

EBONY POMATUM, in pots. Melt 4 oz. of white wax with 12 oz. of any kind of pomatum, and add 2 oz. of levigated ivory-black. Proceed as above, and pour into pots.

POMMADE DE JEUNESSE. Pomatum mixed with magistery of bismuth. It is said to turn the hair black.—GRAY.

DEPILATORIES

FOR REMOVING SUPERFLUOUS HAIRS

These require caution, as they are apt to injure the skin. We have omitted those which contain yellow sulphide of arsenic (orpiment), as there is danger of its being absorbed, and the object can be accomplished without its use. The powders require to be kept in well-closed bottles or boxes, and no more should be mixed with liquid than is required to be used at once.

1. Mix lime and water to a thick cream, and pass through the mixture 25 or 30 times its volume of sulphuretted hydrogen gas. When the gas escapes stop the process. The pulpy mass is spread on paper, and applied for 12 or 15 minutes, and then washed off with a sponge and water. The only objection to this is its disgusting smell.

2. CHINESE. Quicklime 16 oz., pearlash 2 oz., liver of sulphur 2 oz. Reduce to a fine powder, and keep it in a well-closed bottle. Use it as No. 4.

3. Mr REDWOOD recommends a strong solution of sulphuret (sulphide) of barium, with sufficient powdered starch to form a paste; to be left on for a few minutes, then scraped off with the back of a knife.

4. BOUDET'S DEPILATORY. Crystallized hydrosulphate of soda 3 parts, quicklime in powder 10 parts, starch 10 parts; mix. To be mixed with water, and applied to the skin, and scraped off in 2 or 3 minutes with a wooden knife.

TEETH AND MOUTH COSMETICS

TOOTH POWDER

General Directions.—The dry ingredients should be finely pulverized, and the whole well mixed, which is best effected by triturating the powders together, or agitating them in a bottle, and afterwards passing the whole through a sieve. Some ingredients are usually levigated, or ground with water, as prepared chalk, coral, &c. The tooth powders which contain acids, and acid salts, should not be frequently used. For children those only which contain very soft powders should be permitted; the heavy carbonate of magnesia is very suitable for them.

AMERICAN TOOTH POWDER. Coral, cuttle-fish bone, dragon's blood, of each 8 oz., burnt alum and red sanders, of each 4 oz., orris 8 oz., cloves and cinnamon of each $\frac{1}{2}$ oz., vanilla 2 drs., rosewood $\frac{1}{2}$ oz., rose pink 8 oz.

ANTISEPTIC TOOTH POWDER. Prepared or precipitated chalk 2 oz., dry chloride of lime 10 grs., oil of cassia or of cloves 5 drops; mix. It may be coloured, if preferred, by a little levigated bole.

ANTISCORBUTIC TOOTH POWDER. Extract of rhatany $\frac{1}{2}$ oz., prepared charcoal 2 oz., cinnamon $\frac{1}{4}$ oz., cloves $\frac{1}{4}$ oz.

AROMATIC TOOTH POWDER. Calamus aromaticus 4 drs., charcoal 1 dr., soap 1 dr., oil of cloves 12 drops.—**PITTSCHAFT.**

ASIATIC TOOTH POWDER. Prepared coral 4 oz., Venetian red 3 drs., ochre 5 drs., pumice 5 drs., musk 1 gr.; mix. Or, bole 3 parts, chalk 2 parts, ochre 1 part, pumice 1 part, musk to scent.

CADET'S OR DR COOMBE'S. Sugar 1 oz., charcoal 1 oz., Peruvian bark $\frac{1}{2}$ oz., cream of tartar $1\frac{1}{2}$ drs., cinnamon 24 grs.

CAMPHORATED CHALK. Camphor (pulverized by the aid of a few drops of spirit) 1 oz., prepared or precipitated chalk

3 oz. Some makers put only 1 part of camphor to 7 of chalk.

COMPOUND CAMPHORATED TOOTH POWDER. Camphor 1 oz., precipitated chalk 2 oz., cuttle-fish bone $\frac{1}{2}$ oz., myrrh 2 drs., borax 2 drs., lake or rose pink 1 dr., or q. s.

CARTWRIGHT'S DENTIFRICE. Prepared chalk 7 oz., orris 1 oz., Castile soap $\frac{1}{2}$ dr.

CARABELLI'S. Cuttle-fish bone $1\frac{1}{2}$ oz., prepared oyster shells $1\frac{1}{2}$ oz., cinnamon, orris, and lime-tree charcoal, of each 3 drs., vanilla 10 grs.

CHARCOAL, PREPARED. The charcoal, made in iron cylinders, from willow, is to be preferred. It should be reduced to an impalpable powder, and kept from the air. Charcoal of areca nut is highly commended. That of the shells of cocoa-nuts is said to be used for the same purpose. Dr HEIDER prefers the charcoal of the lime tree.

CHARCOAL TOOTH POWDER. Prepared charcoal 1 oz., powdered chalk 3 oz.—GRAY.

CHARCOAL TOOTH POWDER (*French*). Prepared charcoal 1 oz., sugar 1 oz., oil of cloves 3 drops; mix.

CHARCOAL WITH BARK. Charcoal 1 oz., red cinchona 1 oz., powdered sugar $\frac{1}{2}$ oz., with a few drops of some essential oil. See also RHIGHINI'S, further on.

CHARCOAL WITH BARK (*French recipe*). Charcoal 1 oz., Peruvian bark $\frac{1}{2}$ oz., oil of cinnamon, mint, or other oil, 2 drops, essence of ambergris 30 drops.

CHARCOAL WITH QUININE. Charcoal 1 oz., sulphate of quinine 2 to 4 grs., magnesia 4 to 8 grs., otto of rose (or other perfume) 2 drops.

CARBONIC DENTIFRICE (DESFORGES'). Willow charcoal 4 oz., cinchona bark 4 oz., cloves $\frac{1}{2}$ dr.

CIRCISSIAN DENTIFRICE (Dr HALIFAX'S). Prepared hartshorn 2 oz., sulphate of potash 2 oz., cuttle-fish bone 8 oz., orris 4 oz., yellow sandal-wood 1 oz., rose pink 3 oz., oil of rhodium 30 drops. Mix the dry ingredients, previously reduced to a fine powder, and add the oil of rhodium.

CORAL DENTIFRICE (*Poudre Dentifrice* of the French Pharmacopœia). Red coral, bole, cuttle-fish bone, of each 3 oz., dragon's blood $1\frac{1}{2}$ oz., cochineal 3 drs., cream of tartar $4\frac{1}{2}$ oz., cinnamon 6 drs., cloves 1 dr.; reduce separately to powder, mix and grind on porphyry.

DESCHAMP'S ALKALINE DENTIFRICE. Venetian talc 4 oz.,

bicarbonate of soda 1 oz., carmine 4 or 5 grs., oil of mint or other perfume 15 drops.

DESCHAMP'S ACID DENTIFRICE. Venetian talc 4 oz., cream of tartar 1 oz., carmine 4 or 5 grs., oil as the last.

DESFORGES'. See **CARBONIC DENTIFRICE.**

DETERGENT TOOTH POWDER. Bicarbonate of soda 1 oz., powdered Castile soap $\frac{1}{2}$ oz., sulphate of potash $\frac{1}{2}$ oz., sugar of milk $\frac{1}{2}$ oz., orris root 4 oz., oil of bitter almonds 4 drops. Coloured at pleasure.

ELEPHANT'S (MRS) TOOTH POWDER. Bole 1 oz., myrrh, bark, and orris, each $\frac{1}{4}$ oz. All to be finely powdered, and mixed.

FLORENTINE DENTIFRICE. Prepared oyster shells 14 drs., orris 6 drs., cream of tartar 3 drs., lake to colour.

FRENCH TOOTH POWDER. (See **CORAL TOOTH POWDER**, further back; also **GALVANIC**, **DESCHAMP'S**, &c.) Peruvian bark, burnt crust of bread, and sugar, in equal proportions.

GALVANIC DENTIFRICE. Triturate 2 leaves of gold-leaf and 3 of silver-leaf with 2 drs. of sulphate of potash and 1 dr. of alum; then add white sugar 2 drs., common salt 1 dr., pellitory of Spain $\frac{1}{2}$ dr., prepared hartshorn 1 oz., sulphate of quinine 10 grs.; mix. Colour with finest smalts (powder blue), rose, pink, or lake. **FOZEMBAS'** recipe is: 2 leaves of gold, 2 leaves of silver, alum 3 drs., salt $1\frac{1}{2}$ drs., white sugar $1\frac{1}{2}$ drs., pepper 15 grs., opium 5 grs., coral 3 drs., Peruvian bark 3 drs. Grind the gold and silver with the salt and alum, and add the latter ingredients. For the *double* galvanic tooth powder, put twice the above quantities of gold, silver, alum, salt, pepper, and opium. The galvanic action of the metals is thought to stimulate the gums.

GERMAN TOOTH POWDER. Peruvian bark 6 drs., red sanders 2 drs., oil of cloves and of bergamot 3 drops.

GROSVENOR'S TOOTH POWDER. Prepared oyster shells and coral, of each 12 oz., orris root 2 oz., oil of rhodium 6 drops.

HEMET'S DENTIFRICE. It is said to consist of cuttle-fish bone 6 oz., cream of tartar 1 oz., orris $\frac{1}{2}$ oz.; mix.

JAMET'S. Orris 16 oz., magnesia 4 oz., pumice-stone 8 oz., cuttle-fish bone 8 oz., sulphate of quinine 4 oz., cascarilla 1 oz., sugar of milk 16 oz., oil of mint 1 oz., oil of cinnamon 2 drs., oil of neroli 1 dr., essence of ambergris 1 dr.

- KEMMEBER'S.** Wood-soot $1\frac{1}{2}$ oz., strawberry-root $\frac{1}{2}$ oz., and a few drops of eau de Cologne.
- LAVENDER TOOTH POWDER.** Crimson lake 1 dr., Chinese blue (or Turnbull's blue) a scruple; mix and add bicarbonate of soda $\frac{1}{2}$ oz., cuttle-fish bone 2 oz., precipitated chalk 6 oz., oil of lavender 8 drops.
- LARDNER'S TOOTH POWDER.** See CHARCOAL TOOTH POWDER.
- LEFOULON'S TOOTH POWDER.** Scurvy-grass, horse-radish, guaiacum, cinchona, mint, pellitory root, calamus, rhatany, of each equal quantities. Reduce to an impalpable powder. A little calcined magnesia is sometimes added.
- MAURY'S CARBONIC TOOTH POWDER.** Charcoal 8 oz., cinchona 4 oz., sugar 8 oz., oil of mint $\frac{1}{2}$ oz., oil of cinnamon $\frac{1}{4}$ oz., tincture of ambergris $\frac{1}{2}$ dr.
- METGES' TOOTH POWDER.** Prepared chalk $3\frac{1}{2}$ lbs., lake or rose pink 1 lb., orris 2 lbs., cream of tartar 12 oz., levigated pumice 1 oz., sugar 9 oz., oil of cloves 1 dr.
- MIALHE'S RATIONAL DENTIFRICE.** Sugar of milk 3 oz., pure tannin 3 drs., lake 1 dr., oil of mint 8 drops, oil of aniseed 8 drops, neroli 4 drops.
- MYRRH DENTIFRICE.** Myrrh 1 oz., cuttle-fish bone 4 oz., orris 3 oz.; mix.
- NICHOL'S TOOTH POWDER.** Cuttle-fish bone, prepared chalk, orris, of each 1 oz.; cassia $\frac{1}{2}$ oz., myrrh $\frac{1}{2}$ oz.; mix.
- PALMER'S TOOTH POWDER.** Prepared chalk 1 lb., camphor 1 oz., orris 1 lb., cuttle-fish bone 4 oz., rose pink 1 oz.
- PEARL DENTIFRICE.** Precipitated chalk 16 oz., talc 8 oz., finest smalts $\frac{1}{2}$ oz., or q. s. to give it a slight tint.
- PELLETIER'S QUININE DENTIFRICE.** Sulphate of quinine 4 grs., prepared red coral 1 oz., myrrh a scruple. For the coral may be substituted levigated bole 2 drs., precipitated chalk 6 drs.
- REGNAUD'S DENTIFRICE.** Calcined magnesia $\frac{1}{2}$ oz., sulphate of quinine 8 grs., carmine (or cochineal) $\frac{1}{2}$ dr., oil of peppermint 3 drops.
- RHATANY TOOTH POWDER.** Rhatany root 2 oz., cuttle-fish bone 4 oz., prepared chalk 8 oz., borax 1 dr.
- RIGHINI'S CHARCOAL AND BARK.** Charcoal 4 parts, yellow bark 1 part.
- ROSE DENTIFRICE.** Lake $\frac{1}{2}$ dr., myrrh 2 drs., bicarbonate of soda 2 drs., orris 2 oz., cuttle-fish bone 2 oz., precipi-

tated chalk 6 oz., otto of roses 16 drops; or it may be coloured with rose pink to any desired shade.

RUSPINI'S DENTIFRICE. Cuttle-fish bone 8 oz., prepared hartshorn 2 oz., alum 1 oz., cream of tartar 2 oz., orris 1 oz., oil of rhodium 6 drops.

RUSSIAN TOOTH POWDER. Peruvian bark 2 oz., orris root 1 oz., sal ammoniac $\frac{1}{2}$ oz., catechu 6 drs., myrrh 6 drs., oil of cloves 6 or 8 drops.

SAUNDERS' DENTIFRICE. Prepared chalk 2 oz., cuttle-fish bone 1 oz., orris 1 oz., myrrh $\frac{1}{2}$ oz., sulphate of quinine 10 grs.

DR SCHOEPP'S TOOTH POWDER, against mercurial salivation. Alum 2 scruples, cinchona bark 1 oz.

VIOLET TOOTH POWDER. Orris root 2 oz., cuttle-fish bone 4 oz., precipitated chalk 12 oz., bicarbonate of soda $\frac{1}{2}$ oz., essence of violets 1 dr., pure percyanide of iron and crimson lake or rose pink, enough to give it a pale violet colour. See *PULVIS DENTIFRICIUS*, *Pocket Formulary*.

TOOTH PASTES

Any of the above tooth powders may be made into a paste with honey, clarified honey, or honey of roses. A little perfumed spirit may be added. A common objection to these pastes or electuaries is their liability to fermentation, or effervescence. Some makers keep the paste in the bulk for a considerable time, till the effervescence has completely subsided, and then put it up in pots for sale. Others heat the honey, stir in the powders, and keep the mixture warm till any effervescence produced by the action of the acidity of the honey on the cretaceous powder has subsided. It would perhaps be preferable in all cases to use the *prepared* honey (see *MEL DEPURATUM*, *Pocket Formulary*) for these purposes. Electuaries of this kind are termed by the French *opiat*s, although they may contain no opium in any form.

CORAL TOOTH PASTE. *Opiat Dentifrice.* 1. Prepared coral 5 oz., cream of tartar 3 oz., cuttle-fish bone 3 oz., cochineal $\frac{1}{2}$ dr., Narbonne honey 16 oz.—*DESFORGES*.

2. *Opiat Dentifrice Rouge.* Prepared coral 8 oz., cochineal 1 oz., cinnamon 2 oz., alum 3 drs., honey 20 oz., water 1 oz.; triturate the cochineal with the alum and water, add the honey, then the coral and cinnamon; leave

the whole for 24 hours, or till the effervescence has subsided; then rub it with a few drops of oil of cloves, or other aromatic oil, and put it into covered pots for sale.

DYON'S CHARCOAL PASTE. Triturate $\frac{1}{2}$ dr. of chlorate of potash with $\frac{1}{2}$ oz. of mint water, and add gradually 1 oz. of powdered charcoal.

METGES' TOOTH PASTE. METGES' tooth powder (see further back) 48 oz., Narbonne honey 32 oz., syrup 64 oz., cochineal 1 oz., alum 1 oz., water 4 oz.; triturate the cochineal and alum with the water, and strain, add the honey and syrup, and lastly the powder.

PELLETIER'S ODONTINE. This is said to consist of magnesia and butter of cacao, aromatised with some essential oil.

ROSE TOOTH PASTE. Cuttle-fish bone 3 oz., prepared or precipitated chalk 2 oz., orris 1 oz., lake or rose pink to give it a pale rose colour, otto of roses 16 drops, honey of roses q. s.

RED OR CHERRY PASTE. See CORAL PASTE, No. 2.

ROSEMARY PASTE. Levigated bole 4 oz., myrrh 1 oz., oil of rosemary 2 drs. (dissolved in 1 oz. rectified spirit), clarified honey q. s.

SALINE TOOTH PASTE. Sulphate of potash 1 oz., bay salt $\frac{1}{2}$ oz., clarified honey q. s., eau de Cologne 2 drs. or essence of ambergris 30 drops.

VANILLA TOOTH PASTE (French). Charcoal 1 oz., white honey 1 oz., vanilla sugar 1 oz., Peruvian bark $\frac{1}{2}$ oz., and a few drops of any essential oil. The vanilla sugar may be made by triturating a drachm of saturated tincture of vanilla with 1 oz. of pure sugar, and drying the mixture with a gentle heat.

WHITE TOOTH PASTE. 1 (French). Orris, sal ammoniac, cream of tartar, of each 2 oz., tincture of cinnamon and tincture of vanilla, of each $\frac{1}{2}$ oz., oil of cloves 60 drops, clarified honey and syrup to form a paste.

2. Precipitated chalk 4 oz., sulphate of potash $\frac{1}{2}$ oz., prepared honey sufficient to form a paste; to be flavoured with a few drops of otto of roses or oil of cinnamon, &c.

LIQUID PREPARATIONS FOR THE TEETH AND GUMS

ASTRINGENT TINCTURE FOR THE TEETH AND GUMS. 1.

Borax, alum, bay salt, of each a drachm, spirit of camphor, tincture of myrrh, of each 1 oz., spirit of scurvy-grass (or of horseradish) 4 oz., tincture of rhatany 2 oz.; mix, and shake occasionally for a day or two, then filter. A teaspoonful in a wineglassful of water, to rinse the mouth after cleaning the teeth, or at any other time.

2. Tannin 1 dr., rose-water 4 oz., spirit of wine 2 oz., spirit of scurvy-grass (or of horseradish) 2 oz., essence of bitter almonds a few drops.

ODORIFEROUS TINCTURE OF MYRRH. 1. Choice Turkey myrrh 3 oz., eau de Cologne a quart; digest for 7 days, and filter.

2. To 18 fluid oz. of tincture of myrrh add 2 oz. of essence of Cologne. (See PERFUMERY, further back.) If the tincture should not be quite clear, add a few grs. of burnt alum, shake frequently, and filter in a day or two.

BORATED TINCTURE OF MYRRH. 1. Myrrh 1 lb., eau de Cologne 16 lbs., borax 1 lb., distilled water 3 lbs., syrup 3 lbs., essence (or tincture) of roses 6 drs., rhatany root 4 oz.; digest for 10 or 12 days and filter.—Mr COCKLE.

2. Borax 1 oz., shell-lac $\frac{1}{2}$ oz., myrrh 2 oz., spirit of camphor 2 oz., honey of roses 2 oz., rectified spirit a pint, Cologne essence 2 drs., orange-flower or rose-water 4 oz.; digest for a few days in a warm place, shaking occasionally, and filter.

3. Borax 1 oz., shell-lac $\frac{1}{2}$ oz., water 8 oz.; boil together to $\frac{1}{4}$ oz., and add spirit of scurvy-grass a pint, camphor $\frac{1}{2}$ oz., myrrh 2 oz.; digest and filter.

. Borax is very readily soluble in glycerin.

ANTISCORBUTIC ELIXIR. Cinchona 3 oz., guaiacum 5 oz., pellitory 3 oz., orange-peel 2 drs., cloves 5 drs., saffron $\frac{1}{2}$ dr., benzoin, 2 drs., spirit of wine or brandy 32 oz.; digest and filter.—DESFORGES.DESFORGES' EXTRACT OF PELLITORY. Pellitory root 5 oz., cinchona 1 oz., benzoin $1\frac{1}{2}$ drs., essence of peppermint 3 drs., brandy a quart.ELIXIR OF ROSES. Cloves 1 dr., cinnamon 3 oz., ginger 2 oz., spirit of wine $2\frac{1}{2}$ pints, oil of orange-peel 1 dr., otto

of roses 15 drops, essence of peppermint 1 oz. ; mix, digest for 15 days and filter.

LEFANDINIÈRE'S ELIXIR. Rased guaiacum wood $\frac{1}{2}$ oz., pellitory 1 dr., nutmegs 1 dr., cloves $\frac{1}{2}$ oz., oil of rosemary 10 drops, oil of bergamot 4 drops, brandy a pint ; macerate for a fortnight, and filter.

EAU DE BOTTOT. Aniseed 4 oz., cinnamon 1 oz., cloves 1 oz., cochineal 2 drs., oil of mint 2 drs., spirit of wine or brandy 8 lbs. ; macerate 8 days, and filter.

2. Tincture of cedar wood 1 pint, tincture of myrrh 1 oz., mixed with the following essential oils : of peppermint $\frac{1}{2}$ dr., of spearmint $\frac{1}{4}$ dr., of cloves 10 drops, of roses 10 drops.—PIESSE.

EAU DENTIFRICE DE STAHL. Spirit of wine or brandy 2 gallons, rose-water 3 quarts, pellitory 5 oz., cypress root 3 oz., tormentil 3 oz., balsam of Peru 3 oz., cinnamon 5 drs., goats' rue 1 oz., rhatany 1 oz. : macerate for 6 days, shaking it occasionally ; let it rest for 24 hours, and pour off the clear. Add to the clear liquor, oil of mint $1\frac{1}{2}$ drs., cochineal 4 drs. ; in 3 or 4 days, filter.

EAU DU DE O'MEARA. A tincture made of pellitory, vetiver, cloves, orris, and coriander, with creasote, &c.

BOBIES' ODONTALGIC ELIXIR. Pellitory root 2 oz., simple spirit of lavender 16 oz., muriate of ammonia $\frac{1}{2}$ dr. ; digest 24 hours and filter.

GREENOUGH'S TINCTURE. Bitter almonds 2 oz., Brazil wood $\frac{1}{2}$ oz., cinnamon $\frac{1}{2}$ oz., orris root $\frac{1}{4}$ oz., cochineal, alum, salt of sorrel, each 1 dr., spirit of wine 32 fluid ounces, spirit of scurvy grass 1 oz.

HUDSON'S PRESERVATIVE. Tincture of myrrh, tincture of bark, cinnamon water, of each 3 oz., arquebusade water 1 oz., powdered gum $\frac{1}{2}$ oz.

CHELLENHAM DENTAL TINCTURE. Camphor $4\frac{1}{2}$ oz., myrrh 2 oz., bark 5 oz., rectified spirit 36 fluid oz., distilled water 8 oz.

LEFOULON'S ELIXIR FOR THE TEETH. Fresh roots of horse-radish, fresh leaves of scurvy grass and of mint, of each 6 drs., guaiacum, cinchona, pellitory, calamus, and rhatany, each 5 drs., proof spirit a quart ; macerate for 16 days, and strain.

EAU DE MADAME DE LA VEILLIÈRE POUR LES DENTS. Cinnamon 2 oz., cloves 6 drs., fresh lemon-peel $1\frac{1}{2}$ oz., dried

rose petals 1 oz., scurvy-grass 8 oz., spirit 3 lbs.; macerate 24 hours, and distil in a water-bath.

RUSPINI'S TINCTURE. Orris 8 oz., cloves 1 oz., spirit 32 fluid oz., essence of ambergris 1 oz. (or ambergris a scruple); macerate 14 days, and filter.

FRENCH ELIXIR FOR THE TEETH. Rose-water 16 oz., spirit of scurvy-grass 2 oz., tincture of galbanum 1 oz.; colour with cochineal.

ALKALINE LOTION, for preventing injury to the teeth from acid medicine. Bicarbonate of soda 4 drs., distilled water 8 oz., eau de Cologne 2 drs., aromatic spirit of ammonia 1 dr. The mouth to be rinsed with the lotion immediately after swallowing any medicine containing an acid.

QUILLAI TOOTH-WASH. Quillai bark 3 oz., star anise 1 oz., cloves 2 drs., cinnamon 2 drs., oil of peppermint 12 drops, carmine 5 grs., proof spirit 28 oz. Let the drugs be coarsely powdered, and exhaust them by maceration or displacement with the proof spirit, add the oil and carmine, and filter.

LOTION OF CHLORINATED SODA, for purifying the breath, cleansing the mouth, removing unpleasant odours, &c. Liquid chlorinated soda 1 oz., distilled water 19 oz.; mix. A teaspoonful in a glass of water. The same direction applies to most of the above.

STRONGER TINCTURES, SOLUTIONS, OR ESSENCES FOR TOOTHACHE

THESE are applied by moistening a little cotton wool or lint with the liquid, and introducing it into the cavity of the decayed and aching tooth. Where there is no cavity, they are sometimes applied to the gums surrounding the affected tooth. Most of them are stated by their several inventors to give "immediate relief." The cavity should be dried with lint before applying the remedy.

1. **M. PIESTER'S.** Water of ammonia, with half the quantity of tincture of opium; applied as above.

2. Creasote 1 dr., spirit of camphor 2 drs., laudanum 1 dr. Creasote is also used alone: so is *carvacrol*, a liquid of similar properties, so also is carbolic acid. **LAENNEC** prescribes 1 part of creasote and 10 of alcohol.

3. M. COTTEREAU'S. Ether saturated in the cold with camphor, and then a few drops of ammonia added.

4. Mr BLAKE'S. Finely powdered alum 1 dr., spirit of nitric ether 7 drs.

5. *Paraguay-roux, or Compound Tincture of Para Cress.* Flowers of Para cress 4 parts, Italian elecampane (*Inula bifrons*) 1 part, pellitory root 1 part, rectified spirit 8 parts; macerate 14 days, and strain.

6. Mr BLANDE'S *Tincture.* Bruised pellitory $\frac{1}{2}$ oz. camphor 3 drs., opium 1 dr., oil of cloves $\frac{1}{2}$ dr., rectified spirit 6 oz.; digest for ten days, and strain.

7. Pellitory, ginger, cloves, camphor, of each 1 oz., tincture of opium 4 oz., spirits of wine 16 oz.; macerate for 8 days, and strain.

8. Camphor 1 dr., ether 4 drs.; dissolve.

9. Camphor 2 drs., chloroform 1 dr., spirit of sal volatile $\frac{1}{2}$ dr.

10. Opium 2 oz., mastic 1 oz., balsam of tolu 1 dr., camphor 1 oz., oil of cloves 1 dr., rectified spirit 16 fluid oz., oil of bitter almonds 8 drops.

11. Creasote and chloroform, of each 2 drs., Sydenham's laudanum 4 drs., tinct. benzoin 1 oz.

12. Chloral and camphor, of each 1 dr., morphia 2 grs., oil of peppermint 2 drs.

13. BOERHAAVE'S *Odontalgic.* Rectified spirit 1 oz., camphor $\frac{1}{2}$ oz., opium 1 scruple, oil of cloves 80 drops.

14. LEMAZURIER'S *Odontalgic.* Cherry-laurel water 2 oz., acetate of morphia 1 gr. Wash the mouth with warm water, to a glass of which a few drops of this mixture have been added.

15. Oil of rosemary 2 oz., tincture of galbanum 1 oz.; mix. Cotton wetted with this, is to be introduced into the ear.

16. RHIGINI'S. Alcohol 4 drs., creasote 6 drs., tincture of cochineal 2 drs., oil of peppermint 3 drops.

17. Mr DRUITT'S. Tannin 20 grs., mastic 5 grs., ether 2 drs. Wash the mouth with warm water containing a little carbonate of soda; lance the gums, and apply the tincture to the cavity of the tooth on cotton.

18. Mr TOMES recommends a solution of mastic in chloroform. The mastic serves to retain the chloroform, but the latter may be used alone on cotton or lint. Mr BEATSON uses a solution of copal in chloroform.

19. Chloroform and white of egg, equal parts, digested for 4 hours, then applied on lint.

20. A drop on cotton, of a strong solution of thymol in spirits, applied to the hollow tooth.

21. WALKER'S TOOTHACHE TINCTURE (DUTCH REMEDY). Pellitory root finely cut 1 part, guaiacum resin in powder 4 parts, rectified spirit 24 parts; macerate during 7 days, express and filter.

22. HENBANE FUMIGATION FOR TOOTHACHE. A popular remedy is to throw henbane seed on hot cinders, inverting a cup over them to receive the smoke and epyreumatic oil produced. The cup is then filled with hot water, and the steam conveyed to the affected side of the mouth. Dr DOWNING'S ANEURALGICON would probably prove a more effective means of applying remedies of this kind. See GUTTE ODONTALGICÆ, *Pocket Formulary*.

PILLS, OR PASTES, &c., FOR TOOTHACHE

MASSES ODONTALGIQUES

1. DE HANDEL'S. Opium 12 grs., camphor 24 grs., cajeput oil 4 drops, tincture of cantharides 4 drops, extract of henbane and of belladonna, of each 24 grs., distilled water of opium q. s.

2. VOGLER'S. Powdered opium 1 oz., mastic 2 drs., sandarach 2 drs., dragon's blood $\frac{1}{2}$ dr., oil of rosemary 8 drops, spirit to form a paste; to be applied near the affected tooth.

3. Powdered alum 1 dr., powdered mastic $\frac{1}{2}$ dr., spirit of nitric ether q. s. to form a paste.

4. RUST'S. Opium 5 grs., oil of cloves 3 drops, extract of henbane 5 grs., extract of belladonna 10 grs., powdered pellitory sufficient to form a paste.

CEMENTS, &c., FOR STOPPING THE CAVITIES OF TEETH

THESE are harder than the preceding, and are intended to remain in the tooth for an indefinite time. In all cases the cavity should be previously cleansed from all extraneous matters, and wiped perfectly dry with a piece of lint or blotting paper.

1. SOUBEIRAN'S. Powdered mastic and sandarach, of

each 4 drs., dragon's blood 2 drs., opium 15 grs.; mix with sufficient rectified spirit to form a stiff paste. A solution of mastic, or of mastic and sandarach, in half the quantity of alcohol, is also used, applied with a little cotton or lint.

2. Sandarach 12 parts, mastic 6 parts, amber in powder 1 part, ether 6 parts. Applied with cotton. Or simply a paste of powdered mastic and ether. Or a saturated ethereal solution of mastic, applied with cotton.

3. TAVEARE'S CEMENT is made with mastic and burnt alum. BERNOTH directs 90 parts of powdered mastic to be digested with 40 of ether, and enough powdered alum added to form a stiff paste.

4. Gutta percha, softened by heat, is recommended. Dr ROLLE advises melting a piece of caoutchouc at the end of a wire, and introducing it while warm.

5. GAUGER'S CEMENT. Put into a quart bottle 2 oz. of mastic and 3 oz. of absolute alcohol; apply a gentle heat by a water-bath. When dissolved, add 9 oz. of dry balsam of tolu, and again heat gently. A piece of cotton dipped in this viscid solution becomes hard when introduced into the tooth, previously cleansed and dried as above.

6. Mr ROBINSON'S. After washing out the mouth with warm water, containing a few grains of bicarbonate of soda, and cleaning the cavity as above directed, he drops into it a drop of collodion, to which a little morphia has been added, fills the cavity with asbestos and saturates with collodion, placing over all a pledget of blotting paper.

7. OSTERMAIER'S CEMENT. Mix 12 parts of dry phosphoric acid with 13 of pure and pulverised quicklime. It becomes moist in mixing, in which state it is introduced into the cavity of the tooth, where it quickly becomes hard. The acid should be prepared as directed under TRADE CHEMICALS (ACID PHOSPHORIC).

8. Oxide of zinc made into a paste with pyrophosphoric acid.—*Lancet*.

9. Oxide of zinc (recently made by burning) 200 parts, powdered silica 8 parts, powdered borate of soda 4 parts, powdered glass 5 parts. Mix and pass through a very fine sieve, to be kept in a well-stoppered bottle. When required for use a little of the powder is mixed quickly

with a concentrated solution of chloride of zinc, so as to make a thick paste, which is pressed into the cavity of the tooth and will harden in less than 10 minutes. — *Pharmaceutical Journal*.

10. SILICA. This name has been given to a mixture of Paris plaster, levigated porcelain, iron filings, and dregs of tincture of mastic, ground together.

11. WIRIH'S CEMENT. It is said to consist of a viscid alcoholic solution of resins, with powdered asbestos.

12. METALLIC CEMENT. Amalgams for the teeth are made with gold or silver, and quicksilver, the excess of the latter being squeezed out, and the stiff amalgam used warm. Inferior kinds are made with quicksilver and tin, or zinc. A popular nostrum of this kind is said to consist of 40 grs. of quicksilver and 20 of fine zinc filings, mixed at the time of using. Mr EVANS states that pure tin, with a small portion of cadmium, and sufficient quicksilver, forms the most lasting and least objectionable amalgam. The following is the formula: Melt 2 parts of tin with 1 part of cadmium, run it into ingots, and reduce it to filings. Form these into a fluid amalgam with mercury, and squeeze out the excess of mercury through leather. Work up the solid residue in the hand, and press it into the tooth. Or, melt some bees'-wax in a pipkin over the fire, throw in 5 parts of cadmium, and, when melted, add 7 or 8 parts of tin in small pieces; pour the melted metals into an iron or wooden box, and shake them till cold, so as to obtain the alloy in a powder. This is mixed with $2\frac{1}{2}$ or 3 times its weight of quicksilver in the palm of the hand, and used as above.

Another cement consists of about 73 parts of silver, 21 of tin, and 6 of zinc, amalgamated with quicksilver. An amalgam of copper is said to be sometimes used. But this class of stoppings is altogether disapproved of by other authorities. Pure leaf-gold seems the least objectionable.

13. MARMORATUM. Finely levigated glass, mixed with tin amalgam.

14. POUDDRE METALLIQUE. The article sold under this name in Paris appears to be an amalgam of silver, mercury, and ammonium, with an excess of mercury, which is pressed out before using it.

15. FUSIBLE METAL. Melt together 8 parts of bismuth,

5 of lead, 3 of tin, and $1\frac{1}{2}$ or 1.6 of quicksilver, with as little heat as possible.—CHAUDET.

16. NON-EXPENSIVE METALLIC TOOTH-STOPPING. Take 1 part of sulphate of mercury, 1 part of copper in fine powder; rub them well together with a little warm water; when the amalgam is formed, wash well, and remove the surplus mercury by pressing it through chamois leather.—*Pharm. Journ.*

17. EXPENSIVE METALLIC TOOTH-STOPPING, AND MUCH PREFERABLE. Take pure gold, pure gelatine, 1 part of each, pure silver 2 parts, melt, and when refrigerated, reduce to a powder by means of a file; wash well and dry. In the moment of using it, add sufficient mercury to form a plastic paste.—*Pharm. Journ.*

18. PASTE FOR DESTROYING THE SENSIBILITY OF THE DENTAL PULP PREVIOUS TO STOPPING. Arsenious acid 30 grs., sulphate of morphia 20 grs., creasote q. s. Unsafe; it is only inserted by way of warning, against what may prove an unsuspected cause of mischief. Pepsine in the form of paste, is also used for the same purpose.

PIVOTS FOR ARTIFICIAL TEETH. An alloy of platinum and silver.

SPRINGS FOR ARTIFICIAL TEETH. Equal parts of copper, silver, and palladium.—CHAUDET.

EARACHE, SIMPLE CURE FOR. Take a common tobacco-pipe, place a wad of cotton in the bowl, drop upon it 8 or 10 drops of chloroform, and cover with another wad of cotton; place the stem to the affected ear, then blow into the bowl, and in many cases the pain will cease almost immediately.—*American Journal.*

BEVERAGES, DIETETIC ARTICLES, AND CONDIMENTS

BEVERAGES: AND POWDERS FOR PREPARING THEM

WE have placed here such beverages as are rather employed as a refreshing luxury than either medicinally or as regular articles of diet. Wines, spirits, &c., are necessarily excluded. The medicinal mineral waters will be found elsewhere.

GINGER BEER. 1. Infuse 3 oz. of bruised ginger in 4 gallons of boiling water till cold. Strain through tammy or flannel. Dissolve in the liquor 5 lbs. of loaf sugar, and add half a pint of solid yeast, and $2\frac{1}{2}$ oz. of cream of tartar. In cold weather it will be necessary to set the cask near the fire, so as to excite brisk fermentation. As soon as this subsides, rack off the clear liquor, return it into the cask previously washed out, and allow it to work for a day or two longer. Then draw it off and bottle it.—
Mr DONOVAN.

2. Ginger sliced 1 oz., dried orange peel $\frac{1}{2}$ oz.; tie them in a bag, and boil with 16 lbs. of water, and strain; add $\frac{3}{4}$ of an oz. of tartaric or citric acid, 25 drops of essence of lemon, and 24 oz. of loaf sugar. When sufficiently cool, add 2 tablespoonfuls of fresh yeast; let it work for 12 hours and bottle it.

3. Ginger sliced $\frac{3}{4}$ oz., essence of lemon (rubbed with sugar) $\frac{1}{2}$ dr., lump sugar 12 oz., boiling water 8 lbs.; infuse till cold and strain. Ferment as above, with 3 or 4 spoonfuls of yeast, and bottle.

4. Boil $2\frac{1}{2}$ oz. of bruised ginger and 3 lbs. of sugar in $3\frac{1}{2}$ gallons of water for 20 minutes; put into a large pan 1 oz. cream of tartar, and the juice and rind of 2 lemons; pour the boiling liquor over them, and stir the whole well together; when milk-warm add $\frac{1}{4}$ pint of good ale yeast,

cover it, and let it work for 2 or 3 days, skimming it frequently; then strain it through a jelly-bag into a cask, add $\frac{1}{2}$ pint of brandy, bung down close, and in 2 or 3 weeks bottle it in the usual way.

5. Boil 22 oz. bruised ginger in 3 gallons of water for $\frac{1}{2}$ an hour; add 20 lbs. of white sugar, 18 oz. of lemon-juice, 1 lb. of honey, and 15 gallons of water, and strain through a cloth. When cold add the white of an egg and $\frac{1}{2}$ oz. of essence of lemon; after standing 4 days bottle, and lay the bottles in a cellar for three weeks.

6. Prepare a clear decoction or infusion of ginger with sugar and lemon as above; but instead of fermenting it with yeast, charge it strongly with carbonic acid gas by means of a machine.

7. *Imperial Pop.* Cream of tartar 3 oz., ginger 1 oz., white sugar 24 oz., lemon-juice 1 oz., boiling water a gallon and a half; when cool, strain, and ferment with 1 oz. of yeast, and bottle.

GERAMBING, OR LIMONIATED GINGER BEER. 1. Boil $4\frac{1}{2}$ oz. of ginger with 11 quarts of water; beat up 4 eggs to a froth, and add them with 9 lbs. of sugar to the preceding. Take 9 lemons, peel them carefully, and add the rind and juice to the foregoing. Put the whole into a barrel, add 3 spoonfuls of yeast, bung down the barrel, and in about 12 days bottle it off. In 15 days it will be fit for drinking; but it improves by keeping.

2. To 10 gallons of water add $11\frac{1}{2}$ lbs. of loaf sugar and the whites of 10 eggs well beaten; boil till the scum rises, and add 6 oz. of bruised ginger; boil for 20 minutes, then pour the hot liquor on the rinds of 12 lemons thinly peeled; when cold, put into a barrel the juice of 12 lemons, 1 oz. of isinglass, a gill of brandy, and a spoonful of yeast, and fill the barrel with the liquor. In a fortnight it will be ready to bottle.

GINGER BEER POWDERS. Fine powder of Jamaica ginger 4 or 5 drs., bicarbonate of soda $3\frac{1}{2}$ oz., refined sugar in powder 14 oz., essence of lemon 30 drops: mix, and divide into 5 dozen powders. Or 4 to 5 grs. of ginger, 28 of bicarbonate of soda, 112 of sugar, and $\frac{1}{2}$ drop of essence of lemon, in each powder. In the other powder, put 32 grs. of tartaric acid; or 35 grs. if a more decidedly acidulated beverage is required. Or from 30 to 33 grs. of citric acid.

Other formulæ are also in use. Dr PEREIRA gives the following:—Bicarbonate of soda 30 grs., white sugar 1 dr., powdered ginger 5 grs., in each blue paper; and 25 grs. of tartaric acid in each white paper. The following is from the *Pharmaceutical Journal*:—Sugar 2 drs., sesquicarbonate of soda 2 scruples, ginger 4 or 5 grs., essence of lemon $1\frac{1}{2}$ or 2 drops, in each blue paper; with 35 grs. of tartaric acid.

GINGER BEER POWDER IN ONE BOTTLE. (The soda, acid, and sugar must be very carefully dried, separately, and at a temperature not exceeding 120° F.) Fine powder of Jamaica ginger 4 or 5 drs., bicarbonate of soda $3\frac{1}{2}$ oz., double-refined sugar 14 oz., essence of lemon 30 drops, tartaric acid $4\frac{1}{2}$ oz. The acid and soda should not be too finely powdered. Mix the recently dried powders in a warm mortar, and immediately put the mixture into dry bottles, and cork securely. A measure holding 3 drs. should accompany each bottle.

LEMON JUICE (factitious). Dissolve 4 oz. of citric acid in 3 pints of water, with 8 drops of essence of lemon (rubbed with the acid, or dissolved in a little spirit or tincture of fresh lemon-peel). After standing a few days filter it, and keep it in well-closed bottles.

ORANGE JUICE (factitious). Citric acid 1 oz., water 2 pints, oil of orange-peel 4 drops, tincture of orange-peel $\frac{1}{2}$ oz. As the last.

KING CUP; OR LEMON DRINK WITHOUT ACID. 1. Pour a quart of cold water on the thin peel of 1 or 2 lemons: let them infuse 6 or 8 hours; then strain.—Mr BRANDE.

2. Pour a pint of boiling water on the outer rind of one lemon, a small piece of dried orange-peel, and a moderate-sized lump of sugar.

LEMONADE, ACIDULATED (NON-AERATED). 1. Fresh lemon-juice 4 oz., fresh lemon-peel (thinly peeled) $\frac{1}{2}$ oz., white sugar 4 oz., boiling water 3 pints. Strain when cold.—Mr BRANDE.

2. *Imperial.* Cream of tartar $1\frac{1}{2}$ drs., a slice of thin lemon-peel, a lump of sugar; pour on them a quart of boiling water. Strain when cold. To be taken as a cooling drink.

3. *Common.* Cut 2 lemons into slices, add 2 oz. of sugar, and pour on them a quart of boiling water. It is sometimes made with cold water.

4. *French*. Syrup of citric acid 2 oz., water a quart, spirit of lemon-peel a teaspoonful.

5. Juice and thin peel of 1 lemon, citric acid 1 dr., sugar 3 oz., boiling water a quart. It may be varied by substituting for the sugar, syrup of raspberries, or of other fruits.

AERATED OR EFFERVESCING LEMONADE. This may be made by putting into each bottle (soda-water bottle) 1 oz. or 1½ oz. of syrup of lemons, and filling it up with simple aerated water from the machine. The syrup is made by dissolving 30 oz. of lump sugar in 16 oz. of fresh lemon-juice, by a gentle heat. It may be aromatized by adding 30 or 40 drops of essence of lemon to the sugar; or by rubbing part of the sugar on the peel of 2 lemons; or by adding to the syrup an ounce of a strong tincture of fresh lemon-peel, or of the distilled spirit of the same.

EFFERVESCING LEMONADE, WITHOUT A MACHINE. Put into each bottle 2 drs. of sugar, 2 drops of essence of lemon, ½ dr. bicarbonate of potash, and water to fill the bottle; then drop in 35 or 40 grs. of citric or tartaric acid in crystals, and cork immediately, placing the bottles in a cool place, or preferably, in iced water. Mr BARTLETT recommends 2 scruples of sesquicarbonate of soda, 2 drs. of sugar, 4 drops of essence of lemon, and half a pint of water, lastly, a drachm of tartaric acid in crystals. Care must be taken to avoid accidents from the bursting of the bottles. Another form is:—Into a soda-water bottle nearly filled with water, put 1 oz. of sugar, 2 drops of essence of lemon (dropped on the sugar), 20 grs. of bicarbonate of potash in crystals; and, lastly, 30 to 40 grs. of citric acid, also in crystals. Cork immediately.

MILK LEMONADE. Dissolve 1½ lbs. of sugar in a quart of boiling water, add ½ pint of fresh lemon-juice, and the same of sherry; and, lastly, two thirds of a pint of cold milk. Stir together, and strain.

DRY LEMONADE, OR ACIDULATED LEMONADE POWDER. Citric acid ¼ oz., refined sugar 8 oz., essence of lemon 36 drops. Some recipes direct a larger quantity of acid, others a much larger proportion of sugar.

EFFERVESCING LEMONADE POWDERS. Bicarbonate of soda 3½ oz., refined sugar 14 oz., essence of lemon 60 drops. [Sometimes 12 or more grains of the powdered yellow

rind of lemon-peel are added to colour it.] Mix, and divide into 60 powders, of 140 grs. in each blue paper. In the white papers put from 30 to 32 grs. of citric acid, or from 32 to 35 grs. of tartaric acid. Or the mixed alkaline powder and the acid may be put into separate bottles, furnished with measures holding the proper quantity of each.

EFFERVESCING LEMONADE POWDERS IN ONE BOTTLE. *Note.*

—The powders must all be separately and carefully dried, at a moderate temperature, before mixing, and when mixed must be carefully secured from the air.

1. Bicarbonate of soda 1 oz., refined sugar $3\frac{1}{2}$ oz., tartaric acid $1\frac{1}{4}$ oz., essence of lemon 30 drops; mix, and put into well-corked bottles.

2. Mix $3\frac{1}{2}$ oz. of bicarbonate of soda, 14 oz. of double refined sugar, 60 drops of essence of lemon, and $\frac{1}{2}$ oz. to $4\frac{1}{2}$ oz. of tartaric acid.

3. *Lemon-Kali. Acidulated Kali.* Sesquicarbonate of soda 8 oz., tartaric acid 8 oz., refined sugar 16 oz., essence of lemon 100 drops: mix.—*Pharmaceutical Journal.*

ORANGEADE, OR SHERBET. 1. Juice of 4 oranges, thin peel of 1 orange, lump sugar $\frac{1}{2}$ oz., boiling water 3 pints.

2. Juice and peel of 1 large orange, citric acid $\frac{1}{4}$ dr., sugar 3 oz., boiling water a quart.

EFFERVESCING OR AERATED ORANGEADE, OR SHERBET.

1. Mix 1 lb. of syrup of orange-peel, a gallon of water, and 1 oz. of citric acid, and charge it strongly with carbonic acid gas with a machine.

2. Syrup of orange juice $\frac{3}{4}$ oz., aerated water $\frac{1}{2}$ pint.

3. Simple syrup $\frac{1}{2}$ fl. oz., tincture of orange-peel $\frac{1}{2}$ dr., citric acid 1 scruple; fill the bottle with aerated water.

4. Put into a soda-water bottle $\frac{1}{2}$ oz. to 1 oz. of syrup of orange-peel, 30 grs. of bicarbonate of potash, 8 oz. of water, and, lastly, 40 grs. of citric acid in crystals, and cork immediately.

5. Put into each bottle 2 or 3 drs. of sugar, 2 drops of oil of orange-peel, 30 grs. of bicarbonate of potash, or 25 grs. of bicarbonate of soda; water to fill the bottle, and 40 grs. of citric acid, as before.

AERATED SHERBET OR ORANGEADE POWDERS. Powdered sugar $14\frac{1}{2}$ oz., powdered orange-peel 12 grs., oil of orange-peel 60 drops, essence of cedrat 12 drops, bicarbonate of soda $3\frac{1}{2}$ oz.; mix, and put 145 grs. in each blue paper.

In the white paper put 32 grs. of tartaric (or 30 grs. of citric) acid. Or the alkaline and acid powders may be put into separate bottles, with a measure holding the proper proportion of each. The orange-peel may be omitted.

AERATED SHERBET POWDERS IN ONE BOTTLE. Double-refined sugar 14½ oz., powdered orange-peel 12 grs., bicarbonate of soda 3½ oz., essence of cedrat 12 drops, oil of orange-peel 60 drops, tartaric acid 4 oz. The powders must be carefully dried, mixed quickly, and afterwards kept dry, and securely corked. A measure holding nearly 3 drs. of the powder should accompany each bottle.

ORANGEADE POWDER, NOT AERATED. Citric acid ½ oz., sugar 8 oz., oil of orange-peel 20 drops.

SODA POWDERS. The usual proportions are—30 or 32 grs. of bicarbonate of soda in each blue paper, and 25 or 26 grs. of tartaric acid in each white paper.

ACIDULATED EFFERVESCING POWDERS; for making effervescing drinks with concentrated syrups of lemon, ginger, &c. Put into separate papers, distinguished by their different colours, 20 grs. of bicarbonate of soda, and 28 grs. of citric or tartaric acid. One of each powder to be dissolved separately in 1-3rd of a tumbler of water, and a teaspoonful of the syrup added to the acid solution, and the liquids mixed.

[The **CONCENTRATED SYRUPS** are thus made—

Concentrated Syrup of Ginger. Simple syrup 7½ fl. oz., essence of ginger (1 part ginger to 4 of spirit) ½ oz.

Concentrated Syrup of Lemon-peel. Strong tincture of lemon-peel* 1 oz., simple syrup 15 fl. oz.

Concentrated Syrup of Orange-peel. Strong tincture of fresh orange-peel* ½ oz., simple syrup 7½ fl. oz.

[Syrup of raspberries, pine-apples, and other fruit, may be used with the above powders in the same way.]

For Seidlitz and other Medicated Powders, see **MINERAL WATERS and POWDERS**, at the end of **PATENT MEDICINES**, &c.

SPRUCE BEER. Water 10 gallons, treacle or lump sugar (according to the colour required) 6 lbs., essence of spruce 4 oz.; add yeast, and ferment as for ginger beer.

* These tinctures are thus made: fresh lemon-peel thin and cut small, 1 oz.; rectified spirit 8 oz.; digest for some days, and strain. Fresh peel of Seville oranges 4 oz.; spirit 16 oz.

SPRUCE BEER POWDERS. In each blue paper put 5 scruples of powdered sugar, 28 grs. of bicarbonate of soda, and 10 grs. essence of spruce. In each white paper 30 grs. of tartaric acid.

TREACLE BEER. 1. Brown sugar 1 lb., treacle 1 lb., bruised ginger 1 oz., hops $\frac{1}{4}$ oz.; boil for a few minutes in 3 quarts of water, strain, and add 5 quarts of cold water; add a spoonful of fresh yeast, let it work all night, and bottle it in the morning.

2. Treacle 14 lbs., hops $1\frac{1}{2}$ lbs., water 36 gallons, yeast 1 lb.; boil the hops with the water, add the treacle, and strain; cool to 80° F., and ferment with the yeast. In winter $\frac{1}{2}$ oz. of cayenne pods with the hops is an improvement.—*Family Friend*.

CAPILLAIRE. To a pint of boiling water add 3 oz. of fine maidenhair; remove from the fire, cover, and set near the fire for 3 hours; strain, and add $\frac{1}{4}$ pint of orange-flower water. Boil a gallon of fine syrup till reduced to 7 pints, then add the infusion, and boil for 10 minutes; strain through a jelly-bag, and when quite cold, bottle the syrup. It is used to give a fine flavour to water.

LIMONIATED CAPILLAIRE. Refined sugar 24 oz., water 12 oz.; dissolve by a gentle heat; and add essence of lemon 30 drops, neroli 3 drops, citric acid 2 oz., orange-flower water 4 oz.

SYRUP OF PINE APPLE. Expressed juice of pine apple a pint; loaf sugar 2 lbs. Boil gently, and when cold filter.

SIROP D'ORGEAT. See SYRUPUS AMYGDALÆ, *Pocket Formulary*. Another formula for this excellent syrup is the following: Take 20 oz. of sweet and 8 oz. of bitter almonds, 9 lbs. of white sugar, and 4 pints of water. Blanch the almonds, dry them well, beat them with a portion of the sugar, and add gradually two thirds of the water; strain through linen, wash the almonds on the strainer with the rest of the water, and dissolve the sugar in the strained liquor by a gentle heat. Pour the syrup into an earthen vessel, remove the scum, and when nearly cold, add 2 oz. of orange-flower water.

ACIDULATED RASPBERRY SYRUP. Put 6 lbs. of raspberries into a china or glass bowl, or an earthen pan not glazed with lead, with a quart of water in which has been dissolved $2\frac{1}{2}$ oz. of tartaric (or preferably citric) acid, and let it remain 24 hours; then strain it, taking care not to

bruise the fruit. To each pint of clear liquor add $1\frac{1}{2}$ lbs. of pounded loaf sugar, and stir it with a silver spoon till dissolved; leave it for a few days, then bottle it close. A little of this syrup, or of either of the two following, with water, forms a refreshing drink in warm weather, and in some febrile disorders.

ACIDULATED STRAWBERRY SYRUP. As **RASPBERRY SYRUP**, using 2 oz. of citric acid, instead of $2\frac{1}{2}$ oz. of tartaric acid.

RASPBERRY VINEGAR. Put a pint and a half of best wine vinegar to 3 lbs. of fruit in a glass or porcelain vessel; leave them together for a fortnight, then strain without pressure. Or put an equivalent quantity of strong acetic acid to the fruit, in the same way. Or it may be made as directed above for **ACIDULATED RASPBERRY SYRUP**.

WHEY POWDER. Sugar of milk in fine powder 2 oz., powdered white sugar 7 oz., gum arabic $\frac{1}{2}$ oz.; mix. An ounce dissolved in a quart of water is used as a substitute for whey.

WHEY may be made by adding a little infusion of rennet (prepared calf's stomach) to milk, and gently heating it till curdled. It is also made by heating a quart of milk nearly to boiling, and adding either a little lemon-juice, orange-juice, solution of citric acid, vinegar, or white wine, or cream of tartar, sufficient to coagulate it. It is then strained. If required bright, beat up the white of an egg with a portion of the whey, mix with the rest, boil for a moment, and run it through a jelly-bag. See **SERUM LACTIS**, *Pocket Formulary*.

One or two recipes for Confectionery may be introduced here.

ORANGE MARMALADE. 1. Procure some large Seville oranges with clear skins, peel them, squeeze out the pulp and juice, taking care to remove all the pips. Boil the peels, divided into quarters, till they are sufficiently tender; scrape clean all the inside from them, lay them in folds, and cut them into very thin slices about an inch long. Weigh the juice, pulp, and boiled peel; then add broken lump sugar equal in weight to the whole, and boil for half an hour, carefully removing the scum. Then put it into pots, and when quite cold, cover them over.

2. Instead of using all Seville oranges, let only half or a third of them be bitter, and the rest, common sweet

oranges. Proceed in the same way as above. Some add honey.

CURRANT JELLY. Pick the currants, put them in an earthen jar, and place it in boiling water till the juice is extracted. Strain through a sieve without pressing them, and boil the juice in an enamelled saucepan with its weight of loaf sugar, removing the scum as it rises. When it will jelly on the back of a cold spoon, it is sufficiently done. A little of the jelly dissolved in warm water forms an agreeable beverage.

DIETETIC ARTICLES

As the ingredients of some of the following compounds are usually sold by druggists, who may be expected to furnish information as to the manner of using them, and as they may all be regarded as auxiliaries to medical treatment, some notice of them here seems desirable, though it must necessarily be brief and incomplete.

ARROW-ROOT. [West Indian arrow-root is the fecula of the tubers of the *Maranta arundinacea*; East Indian arrow-root is obtained from the *Curcuma augustifolia*; South Sea or Tahiti arrow-root from the *Tacca pinnatifida*. They have all the same properties, and are used in the same manner.] Mix a dessert-spoonful of arrowroot with sufficient cold water to form a soft paste; rub it till quite smooth, and add half a pint of boiling water, stirring it briskly. Boil it for a minute or two, and when removed from the fire add a teaspoonful of sherry or other white wine (where wine is admissible), with a little grated nutmeg or lemon-peel and sugar to the taste. For young children, milk should be used instead of water, and the wine omitted; it is also more nourishing in this form for those invalids with whom milk agrees.

Tous les Mois. [The fecula of a species of *Canna*.] It is used in the same way as arrow-root; but rather less is required. It forms a more tenacious but less transparent jelly.

SAGO. [The granulated fecula of the pith of one or more species of the Sago Palm.] Wash an ounce of pearl sago in cold water; then boil it very gently in a pint of fresh water, stirring it frequently till dissolved. It may be

flavoured with wine, spices, and sugar, as directed for arrow-root. For children, and for consumptive and debilitated patients, it may be made with milk instead of water. The common sago, being in larger grains, requires more time to dissolve, and is usually steeped for some hours before boiling it.

TAPIOCA. [Obtained from the tuberous roots of the Cassava (*Jatropha manihot*). It is usually sold in small lumps formed by drying the fecula on hot plates.] It is used in the same way as sago, but requires to be previously steeped for some hours, or to be simmered for a longer time. It forms a clear jelly, which may be flavoured with wine, spices, and sugar, as directed for arrow-root; but it is more nourishing when made with milk.

SAGO POSSET, for invalids. Macerate a tablespoonful of sago in a pint of water for 2 hours on the hob of a stove, then boil for 15 minutes, assiduously stirring. Add sugar with an aromatic, such as ginger or nutmeg, and a tablespoonful or more of white wine. If the wine be not permitted, flavour with lemon-juice.

SAGO OR TAPIOCA MILK, for invalids. Take an ounce of either of these feculæ, and soak it in a pint of cold water for an hour; then pour off this water, and, adding 1½ pints of good milk, boil slowly until well incorporated.—**Dr A. T. THOMSON.**

TAPIOCA PUDDING, for invalids. Beat up ½ oz. of sugar with the yolks of 2 eggs, and stir the mixture into a pint of tapioca milk.—**Dr THOMSON.**

Arrow-root milk and pudding may be made like the corresponding preparations of tapioca.

PANADA, for invalids. Place in a saucepan some very thin slices of bread crumb, and add rather more water than will cover them. Boil now until the bread becomes pulpy, strain off the superfluous water, and beat up the remainder into the consistence of gruel. Sweeten with white sugar, and add, if permitted a little sherry wine.

BARLEY WATER. See **DECOCTUM HORDEI**, and **DECOCTUM HORDEI COMPOSITUM**, *Pocket Formulary*. **ROBINSON'S PATENT BARLEY** is a convenient preparation; printed directions accompany it.

ASSES' MILK, ARTIFICIAL. Eringo root, pearl barley, sago, rice, of each 1 oz. Wash them with cold water, then boil them with 3 pints of water to 1½ pints, and strain. Put a

teaspoonful to a cup of boiling water, and sweeten to the taste. [*Bonbons de lait d'ânesse* are made with sugar of milk, white sugar, gum, and starch or arrow-root.]

LINSEED TEA. Take $1\frac{1}{2}$ oz. of clean linseed, and $\frac{1}{2}$ an oz. of bruised liquorice root; put them into a warm teapot or jug, and pour on them 2 pints of boiling water; let them stand, covered, near the fire, for 3 or 4 hours, stirring them occasionally; then strain. To save time, the ingredients may be boiled for 15 or 20 minutes, instead of infusing them; but the tea so made is less agreeable.

ICELAND MOSS. Infuse an ounce of picked Iceland moss for 15 minutes in half a pint of hot water; strain off the water and boil the moss in a quart of fresh water till reduced to a pint and a half. Half an ounce of liquorice root may be added, towards the end of the boiling, if agreeable: or milk may be used instead of water.

JELLY OF ICELAND MOSS. See *GELATINA LICHENIS*, *Pocket Formulary*. Another form is the following:—Infuse 2 lbs. of the moss for half an hour in sufficient boiling water to cover it; drain the moss, and boil it in $2\frac{1}{2}$ gallons of water for an hour, and strain. Boil the moss with fresh water, adding an oz. of isinglass; strain; mix the product of the two boilings, and let it stand till clear. Evaporate the clear liquid to the consistence of a stiff jelly, adding, towards the end, 6 lbs. of fine lump sugar, 2 oz. of French brandy, and half an ounce of orange-flower water. It may be taken, almost at pleasure, dissolved in water or milk.

ICELAND MOSS CHOCOLATE. See *CHOCOLATA LICHENIS*, *Pocket Formulary*.

IRISH MOSS, OR CARRAGEEN. Steep a $\frac{1}{4}$ of an ounce of the moss in cold water for a few minutes: then withdraw it, shaking the water from each sprig, and boil it in a quart of milk till it attains the consistence of jelly, and sweeten to the taste. A decoction of the same quantity of moss in a quart of water is also used as a demulcent in coughs, &c. **BLANC-MANGE** may be made by washing $\frac{1}{2}$ oz. of the moss as above, and boiling it in $1\frac{1}{2}$ pints of new milk to such a consistence that it will retain its form when cold, sweetening and flavouring it to the taste. An agreeable jelly may be made by boiling it with water instead of milk, and adding lemon- or orange-juice or peel, wine, &c.

CEYLON MOSS. Boil $\frac{1}{2}$ oz. of the prepared moss in a quart

of water for 25 minutes; or till a spoonful taken out forms a firm jelly in 2 or 3 minutes; then flavour with wine, cinnamon, or with lemon- or orange-juice or peel; and sweeten to the taste. Boil for five minutes longer, and press through a jelly-bag, or doubled muslin. Pour it into earthen moulds, and leave it undisturbed till it has set. If the jelly is required bright it must be clarified with white of egg, as directed for gelatine jelly. For **BLANC-MANGE** add 1 oz. of prepared moss to a quart of boiling water, and boil gently till reduced to a third; add 2 pints milk and the flavouring ingredients, and pour into earthen moulds.

AUSTRALIAN MOSS. This has been introduced for the same use as Irish and Ceylon mosses, but has not been very generally adopted. Soak $\frac{1}{2}$ oz. of the moss in water for an hour or two, pour away the water, and boil the moss in a quart of fresh water till dissolved. Strain through a hair sieve, and sweeten and flavour to the taste.

SALEP. [The dried root of some species of orchis.] Boil $\frac{1}{2}$ oz. of salep powder in a pint of water till of the proper consistence: strain, and sweeten and flavour to the taste.

HARTSHORN JELLY. Boil 4 oz. of true hartshorn shavings (previously washed in warm water) in a quart of water till reduced to a pint; strain, and sweeten and flavour to the taste. For children and consumptive patients, the simple jelly may be mixed with milk and a little sugar. To make a bright jelly for the table, boil 4 oz. of washed hartshorn shavings in $1\frac{1}{2}$ pints of water, till reduced to $\frac{3}{4}$ of a pint, and add 2 oz. of sugar, and a tablespoonful of lemon, or orange-juice. Strain with pressure; beat up the white of an egg with a little cold water, mix this thoroughly with the jelly, and evaporate the liquid till a little taken out solidifies on cooling. Add a little fresh lemon-peel, and strain through a jelly-bag.

GELATINE JELLY. Steep 1 oz. of **NELSON'S** or other purified gelatine in half a pint of cold water for ten minutes; then add the same quantity of boiling water, and stir till it is dissolved, applying heat if required: add the juice and peel of two lemons, sugar, and wine sufficient to make up the whole to a pint and a half. If required bright, have ready the white and shell of an egg well beaten together, stir them briskly into the jelly, boil for 2 or 3 minutes without stirring, and pass through a jelly-bag. As a nourishing

diet for children and invalids, a little of the gelatine simply dissolved in water may be mixed with milk, or the dry gelatine dissolved in milk by heat.

ISINGLASS JELLY. Isinglass is used in the same way as gelatine, but as it is not wholly soluble in water, it requires straining. To make a bright jelly, it requires more eggs for its clarification than gelatine. A very pleasant jelly is made with the Acidulated Raspberry or Strawberry Syrup (see further back) thus: Dissolve $1\frac{1}{2}$ oz. of isinglass in a very little water, put this to a quart of the syrup, warm it and stir it well; then strain it into a mould. In warm weather put 2 oz. of isinglass.

ARROW-ROOT BLANC-MANGE. Beat up 2 oz. of genuine arrow-root with a little cold milk to the consistence of cream; pour on it $1\frac{1}{2}$ pints of boiling milk, stirring it all the time. Flavour and sweeten to the taste, boil for 10 minutes, stirring it constantly, pour into moulds, and leave it until next day.

BLANC-MANGE. This may be made with either isinglass or gelatine. Boil $\frac{1}{2}$ oz. in 16 fluid oz. (the old wine pint) of new milk; stir it constantly till it boils, let it simmer for a few minutes till the isinglass is dissolved; strain, add sugar to the taste, and a few drops of almond flavour, or other flavouring ingredients, and pour into moulds.

CHOCOLATE. This is prepared from the finest cocoa-nuts (seeds of *Theobroma cacao*) after roasting, winnowing, &c., by grinding them on a hot stone or plate, or beating them in a hot mortar to a smooth paste. Sugar is generally added, and vanilla or other flavouring ingredients.

CHOCOLATES, MEDICATED. See CHOCOLATA, *Pocket Formulary*.

WHITE CHOCOLATE. White sugar 3 lbs., rice flour $27\frac{1}{2}$ oz., English or Indian arrow-root 8 oz., tincture of vanilla $\frac{1}{2}$ oz., butter of cacao 8 oz., powdered gum arabic 4 oz.; form a paste with boiling water, and put it into moulds.

COCOA. This should also be prepared from the seeds of *Theobroma cacao*; and the rock, roll, and flake cocoas often consist of this alone. But most of the paste cocoa, and soluble cocoa powder, is mixed with saccharine and farinaceous matters. This is the case with much of the "*Homœopathic*" Cocoa, which professes to be unadulterated, but generally contains potato starch. A common proportion for soluble cocoa appears to be two thirds of

pure cocoa, and one third of sugar and farina; the latter being one or more of the following:—Wheat flour, sago meal, potato flour, arrow-root, &c. The *Paste Cocoa* often contains only about half its weight of cocoa, the rest being sugar and water, with sometimes the addition of sago meal or other farina.

GUARANA. An alimentary and medicinal substance, imported in the form of cakes from Brazil, where they are used as we use chocolate, mixed with water and sugar, and taken as a beverage. Guarana is very rich in caffeine. See *Pocket Formulary*.

BROMA. This consists of about 8 oz. of pure cocoa, $3\frac{1}{2}$ of sugar, and $4\frac{1}{2}$ of sago-meal, arrow-root, &c.

WACAHA DES INDES. Roasted cacao beans (chocolate) in powder 2 oz., sugar 6 oz., cinnamon $\frac{1}{4}$ oz., vanilla (powdered with part of the sugar) $\frac{1}{2}$ dr., ambergris 3 grs., musk $1\frac{1}{2}$ grs. Sometimes a drachm of prepared annotto is added, and the ambergris and musk omitted.

RACAHOUT DES ARABES. This is professedly a preparation of acorns (perhaps those of the *Quercus ballotta*, which are naturally sweet, or of other kinds deprived of their bitterness by being buried in the earth); but it is imitated by the following:—1. Chocolate in powder 1 oz., rice flour 3 oz., sugar 9 oz., potato arrow-root 3 oz., vanilla (pulverized with part of the sugar) 1 dr.; mix.

2. Chocolate in powder 4 oz., salep 1 oz. (or powdered tragacanth 1 oz.), potato arrow-root 5 oz., sugar (flavoured with vanilla) 8 oz.—CADET.

DICTAMIA. Sugar 7 oz., potato arrow-root 4 oz., flour of brent barley (*Triticum monococcum*) 3 oz., Trinidad and Granada chocolate, each 1 oz., vanilla 15 grs.

PALAMOUD. Chocolate 1 oz., rice flour 4 oz., potato arrow-root 4 oz., red sanders, in fine powder, 1 dr.; mix. [In the above, by chocolate is meant the cacao beans roasted and pulverized without addition. Indian arrow-root or *tous les mois* may be substituted for the potato arrow-root.]

FECULUM SAXONIA. Barley flour 21 oz., sugar 7 oz., cinnamon 1 dr. Mix, and bake them in an oven, enveloped in a paste of wheat flour, and placed in an earthen vessel. When sufficiently baked, remove the crust, and when the contents are cool, reduce them to powder. About $\frac{1}{2}$ oz. to 1 oz. is boiled with broth, &c., as a nourishing diet.

It is often medicated with the addition of sarsaparilla, bark, &c.

FABINACEOUS FOODS, &c. The following compounds are accompanied with full directions for use :

BASTER'S *Soojee and Compound Farina.* Wheat flour, with sugar.

BRIGHT'S *Nutritious Farina.* The basis is said to be potato starch.

BRIGHT'S *Breakfast Powder.* A combination of chocolate with his nutritious farina.

BRADEN'S *Farinaceous Food.* Chiefly wheat flour carefully baked.

BULLOCK'S *Semola.* Wheat flour, from which a portion of the starch has been removed, so as to leave a definite proportion of gluten.

CHAPMAN'S *Entire Wheat Flour.* Is what its name states.

DENHAM'S *Farinaceous Food.* As BRADEN'S, with perhaps a mixture of barley flour.

GARDINER'S *Alimentary Preparation.* Very finely ground rice flour.

HARD'S *Farinaceous Food.* Carefully baked wheat flour.

HUNT'S *Breakfast Powder.* Rye, carefully roasted as coffee. [For DANDELION COFFEE, see further back, under "DRUGGISTS' NOSTRUMS."]

LEATH'S *Alimentary Farina.* Baked wheat flour, with sugar, potato flour, and a small quantity of Indian corn meal, and tapioca.

MAIDMAN'S *Nutritious Farina.* Potato flour, tinged with some pink colouring matter.

PALMER'S *Vitaroborant.* See ERVALENTA, below.

PLUMBE'S *Farinaceous Food.* South Sea arrow-root combined with pea flour.

Prince of Wales's Food. Potato flour.

Semolina. A hard kind of wheat containing much gluten, ground into coarse grains. But some articles sold under this name appear to be compounds of gluten, artificially granulated, resembling BULLOCK'S *Semola*.

ERVALENTA, REVALENTA ARABICA, LENTIL POWDER, &c. These consist chiefly either of the European or Egyptian lentil.

Ervalenta. WARTON'S consists of the French or Ger-

man lentil, with either Indian corn, or, more probably, a species of corn called *Durra* or *Doura*, used by the Arabs. But Dr SCHENK states that what is sold in Paris consists of the flour of French beans and Indian corn.

Revalenta Arabica. A mixture of the red (Egyptian or Arabian) lentil with barley flour. Some samples contain sugar, others salt and a flavouring ingredient.

Lentil Powders. Some consist entirely of lentil flour (French, German, or Egyptian, or both kinds mixed). Others contain barley flour in addition. NEVILL'S consists of 1 oz. of curry powder to 4 lbs. of lentil flour. The *Lancet* gives the following recipes for lentil powder: 1. Arabian lentil flour 2 lbs., barley flour 1 lb., salt 3 oz.

2. Pea flour 2 lbs., Indian corn flour 1 lb., salt 3 oz.

GRUEL is made either from oatmeal, or from groats or grits (oats deprived of their cuticle), either whole or crushed (Emden groats). Dr THOMPSON directs 3 oz. of groats, previously washed, to be boiled slowly in 4 pints of water, till reduced to 2 pints, then strained through a sieve. The Emden groats require less boiling. Dr KITCHENER directs one or two tablespoonfuls of oatmeal (according as the gruel is preferred thin or thick) to be well mixed with 3 spoonfuls of cold water, gradually added, a pint of boiling water poured on it, and the whole boiled for 5 minutes, constantly stirring it; it is then skimmed and strained through a hair sieve; a little butter is usually added, and sometimes milk, with salt, or otherwise sugar and spices to the taste. Thorough trituration of the oatmeal and cold water, and constant stirring of the gruel while on the fire, render long boiling unnecessary.

BOILED WHEAT. Steep the wheat in water for 10 or 12 hours, then boil it for half an hour. [As a substitute for vegetables, and to obviate constipation.—Mr BULLOCK.]

BEEF TEA. Professor LIEBIG directs 1 lb. of beef, free from fat, to be minced very small, mixed with an equal weight of cold water, and heated slowly to boiling; when it has boiled for a minute or two, strain through a cloth. It may be coloured with roasted onion or burnt sugar, and salted to the taste. Dr SEYMOUR directs 2½ lbs. of lean beef cut small to be put into 3 pints of cold water, and simmered slowly, without boiling, till reduced to a pint and a half; then carefully strained. In another formula we read—

Macerate raw beef (recently killed), chopped very fine, $\frac{1}{2}$ lb., in distilled water $22\frac{1}{2}$ oz., with common salt 50 grs., dilute hydrochloric acid 16 drops, for an hour and a half; strain through a fine hair sieve. Give two tumblers daily.

EXTRACT OF MEAT. Cut the lean of freshly killed meat very small, put it into eight times its weight of cold water, and heat it gradually to the boiling point. When it has boiled for a few minutes, strain it through a cloth, and evaporate the liquor gently by a water-bath to a soft mass. 2 lbs. of meat yield 1 oz. of extract. Fat must be carefully excluded, or it will not keep.—LIEBIG.

LIEBIG'S FOOD FOR INFANTS. Malt flour, to which is added a small quantity of bicarbonate of potash or soda.

LIEBIG'S SOUP FOR CHILDREN. Mix $\frac{1}{2}$ oz. wheat flour, $\frac{1}{2}$ oz. malt flour, $7\frac{1}{2}$ grs. bicarbonate of potash; add 1 oz. of water and 5 oz. of milk. Heat with constant stirring over a gentle fire till it begins to thicken. Remove from the fire and stir for 5 minutes. Heat once more, and again remove and stir. Heat again to ebullition. Separate the bran from the soup by a fine sieve.

TROPHAZOME. Mince 16 oz. of meat, free from fat, very fine, pour on it 8 oz. of cold or lukewarm water (not exceeding 100° F.); mix well, and let it stand for an hour, stirring it 3 or 4 times. Press out the liquid (about 6 oz.); mix 8 oz. more of water with the meat, stir it occasionally, and in half an hour strain with pressure. Repeat this with 8 oz. more water. Break up the pressed meat, and put it into a small tin vessel; place this in a water-bath of cold water, heat gradually to the boiling point, and keep it boiling for 20 minutes. Mix the fluid which exudes with the others, add salt, spices, and other flavouring ingredients, and boil for 20 minutes in a covered vessel. It may be thickened with 1 oz. of semola.—Mr BULLOCK.

MEAT BISCUITS. A thick extract of meat (made by boiling freshly killed beef or other meat, and evaporating the strained liquid) is kneaded with wheaten flour, and the dough rolled out and divided into biscuits, which are dried or baked in an oven. They are kept in the form of biscuits or coarsely ground: 1 ounce makes a pint of rich soup, which may be salted or flavoured to the taste.

BREAD, AERATED. DAUGLISH'S patent. This is prepared without yeast, to the saving, it is said, of 10 per cent. of the weight of the flour. Aerated water is forced by

machinery into the mass of flour enclosed in an iron vessel. The dough is then kneaded by machinery inside the vessel. The pressure being now removed, the dough instantly rises. The whole process occupies no more than half an hour, instead of eight or ten hours, as when yeast is used.

BREAD, UNFERMENTED. Mix carefully $\frac{1}{2}$ oz. of bicarbonate of soda and $\frac{1}{4}$ oz. of salt with 4 lbs. of flour; mix this with a quart (or rather 41 or 42 fluid oz.) of very cold water, previously mixed with $\frac{1}{2}$ a fluid oz., and 20 minims of hydrochloric acid of 1.16 specific gravity, into a thin dough, with as little kneading as possible, and let it be immediately placed in the oven; it requires rather more time than fermented bread. By mixing 26 measures of the acid with 46 of water, a diluted acid is obtained, of which a fluid ounce and a half may be taken for every $\frac{1}{2}$ oz. packet of soda.

A pamphlet on the subject, directs for *Brown Bread*, 3 lbs. of wheat-meal and 10 drs. (Apoth. weight) of bicarbonate of soda to be well mixed, and made into dough, with 25 oz. of cold water, previously mixed with $12\frac{1}{2}$ fluid drs. of hydrochloric acid.

The proportions used first by Dr WHITING, and subsequently by Mr DODSON, in the preparation of unfermented bread and biscuits, were: 7 lbs. of wheaten flour, 350 to 500 grs. of bicarbonate of soda, $2\frac{1}{4}$ pints of water, with hydrochloric acid 420 to 500 grs., or as much as may be sufficient. The soda, dissolved in a small quantity of water, is first carefully kneaded with the dough, and the acid being afterwards rapidly mixed in, the bread is baked without delay.

A third formula is that of Mr J. SAVOY, and is recommended as excellent by Dr PEREIRA. Intimately mix with 1 lb. of flour, sesquicarbonate of soda 40 grs., and powdered white sugar a teaspoonful, in a large basin with a wooden spoon. Then gradually add cold water about $\frac{1}{2}$ a pint, previously mixed with 50 drops of pure hydrochloric acid, and stir constantly, so as to form very speedily an intimate mixture. Divide into two loaves, and put into a quick oven immediately.

Another form of unfermented bread is as follows:—Mix 1 oz. of carbonate of soda, $\frac{3}{4}$ oz. of tartaric acid, and $\frac{1}{2}$ oz. of salt, with 7 lbs. (half a peck) of flour; mix the whole thoroughly, taking care that all the ingredients are per-

fectly dry; add, in 2 or 3 portions, 4 pints of cold water, and incorporate quickly; place it in tins, and send it to the oven immediately. If not baked in tins, less water must be used.

JONES'S PATENT FLOUR contains all the ingredients ready mixed, and may therefore be kneaded with water, and baked at once, without further additions. To 1 cwt. of perfectly dry wheat flour, $10\frac{1}{2}$ oz. (avoird.) of dry, finely powdered tartaric acid are added. Mix well, pass through a flour-dressing machine, and allow it to remain for 2 or 3 days. Then add, all in fine powder and dry, 12 oz. of bicarbonate of soda, 24 oz. of common salt, and 8 oz. of loaf sugar. Mix all thoroughly together, and pass through a flour-dressing machine, when it will be ready for use. 1 lb. made into bread will require 10 oz. of water, or for biscuits 6 oz., and the dough must be baked at once in a well-heated oven.

(Biscuits and cakes made without yeast, and containing no butter, are prescribed in some dyspeptic cases; of such the following are examples:—)

ABERNETHY BISCUITS. Make into a stiff biscuit paste, 1 quart of milk, 6 eggs, 8 oz. of loaf sugar, and $\frac{1}{2}$ oz. of caraway seeds, with flour sufficient to bring the whole to the required consistence. Make the biscuits thin, dock them with holes to prevent them from swelling up, and bake in an oven at a moderate heat.

SPONGE BISCUITS. Beat the yolk of 12 eggs for $\frac{1}{2}$ an hour, then put in $1\frac{1}{2}$ lbs. of finely powdered sugar, and whisk it briskly until it rises in bubbles; beat the whites to a strong froth, and whisk them well with the sugar and yolks; then work in 14 oz. of flour, with the rinds of 2 lemons grated. Bake in tin moulds buttered, for 1 hour in a quick oven, sifting over them a little fine sugar.

RICE CAKE. Beat the yolks of 15 eggs for nearly half an hour with a whisk, mix well with them 10 oz. of finely sifted white sugar; put in $\frac{1}{2}$ lb. ground rice, a little orange-flower water or brandy, and the rinds of 2 lemons grated; then add the whites of 7 eggs, well beaten up, and stir the whole for $\frac{1}{4}$ of an hour; put into a hoop, and bake for $\frac{1}{2}$ an hour in a quick oven.

(Bread and biscuits made with ginger as an ingredient may also be useful in assisting a weak digestion. The two following recipes are by M. SOYER:—)

COMMON GINGERBREAD. Put on a slab or table 1 lb. of flour; make a ring of it; put $\frac{1}{2}$ pint of treacle in, mix well together, working it so as to form a stiff paste. Put some flour into a basin, to which add this dough, which will keep thus for 7 or 8 weeks. When wanted, put in any quantity of ground ginger, according to taste or desire. Mix well, roll thin, and cut into pieces about the size of a crown; put them on a baking sheet, and bake for a few minutes, till crisp. To every pound of paste, an ounce of butter may be used, if preferred. These cakes will keep a long while in an air-tight case.

GINGER CAKE. Take $\frac{1}{2}$ lb. of sugar, $\frac{1}{2}$ lb. of butter, 1 $\frac{1}{2}$ oz. of ground ginger, 6 eggs, beat well, stir in 1 $\frac{1}{2}$ lbs. of flour, and add as much milk, a little warm, as will make a stiff dough for bread. Bake in a pan for 2 hours.

GLUTEN BREAD (for diabetic patients). It is made with the gluten of flour, a small portion only of the starch being retained.

Dr ALDRIDGE recommends, for preparing the gluten, second best flour. This must be made into a stiff paste with cold water, and then kneaded with the hands under a current of water, on a slanting board placed in a two-gallon jar, until starch can no longer be detected by tincture of iodine in small portions taken from the mass. A stone of flour yields from 3 $\frac{1}{2}$ to 4 $\frac{1}{2}$ lbs. of gluten. The gluten may also be obtained from starch manufactories.

To prepare the bread, the following materials should be mixed. Fresh moist gluten 24 oz., bicarbonate of ammonia 3 $\frac{1}{4}$ drs., common salt 1 $\frac{1}{2}$ drs., powdered caraway 48 grs., wheaten flour 4 $\frac{1}{2}$ oz., powdered bran 1 $\frac{1}{2}$ oz., salt butter 4 oz. These quantities yield 24 oz. of bread, when baked. It may be baked in small and flat circular tin pans, placed on a moderately heated hot hearth. It is difficult to bake this bread properly with yeast, as prescribed by BOUCHARDAT; and if made without butter, or with the total exclusion of starch, it becomes extremely unpalatable.

Dr PERCY proposes the following method:—Take the matter left after removing the starch from 16 lbs. of rasped potatoes, $\frac{3}{4}$ lb. of mutton suet, 12 eggs, $\frac{1}{2}$ lb. of butter, and $\frac{1}{2}$ oz. of carbonate of soda; mix, and add 2 oz. of diluted hydrochloric acid; divide into 8 cakes, and bake immediately in a quick oven.

[Various alimentary preparations have from time to time been introduced, the basis of which is the gluten which remains in extracting the starch from wheat flour by the mechanical process. Mr GENTIL's *gluten flour* is a mixture of this with wheat flour. It contains 42 per cent. of gluten, and yields a nutritious and digestible *gluten biscuit*, *gluten bread*, and, with cocoa, *gluten chocolate*. Mr BULLOCK's *semola* and Mr VERON's *granulated gluten* were of the same nature. 30 parts of white flour, 10 of fresh gluten, and 7 of water, form a paste resembling *Italian macaroni*, *vermicelli*, &c.]

BAKING POWDER. Refer to DRUGGISTS' NOSTRUMS, further back.

CUSTARD POWDER. Rub up together gum tragacanth 2 oz., potato starch 1 lb., powdered turmeric 2½ drs., with oil of bitter almonds (deprived of its prussic acid) ½ dr., and essence of lemons 1 dr. Put up into 1-ounce packets. (From 1 pint of new milk take 2 tablespoonfuls to work up with the powder; boil the remaining milk with 2 ounces of lump sugar, and pour it, while boiling, into the basin, stirring quickly until thoroughly mixed. Bake as a custard.)—Mr SCHOLEFIELD.

Without the colouring, this forms *Blanc-mange Powder*.

RENNET WINE (*for dyspeptic persons*). Take the fresh rennet bag of a calf, cut off and throw away 3 inches of the upper or cardiac extremity, slit the rest longitudinally, wiping gently with a dry napkin, but not removing the clean mucus, cut it into small pieces, and put into a common wine-bottle filled up with good sherry. Let it remain corked for three week. One teaspoonful in a wine-glassful of water, taken after meals, assists digestion.—Mr GEO. ELLIS.

PREPARATIONS OF COD-LIVER*

THESE are designed for such invalids as are likely to derive benefit from cod-liver oil, but are unable to overcome their disgust for that useful medicine. In such cases the fresh liver itself may be found of equal service. The fol-

* Dr Squibb says, that a 5 per cent. solution of gum arabic tends greatly to cover the taste of cod-liver oil; and that some salt herring eaten just before taking it, would render the taste imperceptible.

lowing methods of cooking it are adapted from those of M. SOYER.

POTATOES AND COD-LIVER. Take 1 lb. of fresh cod-liver; peel and steam 2 lbs. of floury potatoes. Cut the liver in 4 pieces, place it over the potatoes, then again steam them, letting the oil from the liver fall on them. When done, make some incisions in the liver with a knife, to extract the oil remaining. Afterwards, the liver may be eaten with some anchovy sauce,—and the potatoes, mixed with the oil, with a little salt and pepper. (Simple cod-oil, as tasteless as can be procured, may be made in the same manner to yield a palatable dish with potatoes.)

RICE AND COD-LIVER. Boil $\frac{1}{2}$ lb. of rice in 2 quarts of water. When nearly done, remove 3 parts of the water, then put over the rice 1 lb. of cod-liver, cut in large dice. Put the saucepan in a slow oven for about 30 minutes, by which time it will be well cooked. Take the liver out, which serve as above. Stir up the rice with a fork, and serve it with salt and pepper. If no oven, care must be taken to cook it over a slow fire, or else it will burn.

TAPIOCA AND COD-LIVER. Boil $\frac{1}{2}$ lb. of tapioca till soft, in 2 quarts of water, drain in a cullender, then put it back in the pan; season with salt and pepper, add $\frac{1}{2}$ pint of milk, put over 1 lb. of fresh cod-liver, cut in 8 pieces. Set the pan near the fire to simmer slowly for half an hour, or a little more, till the liver is quite cooked. Press on it with a spoon to squeeze out the oil. Take away the liver, and mix the tapioca. If too thick, add a little milk, boil for a few minutes, add a little more salt and pepper, and serve. Here, again, a slow oven is better than a fire.

COD-ROE AND COD-LIVER. Take a cod's roe, cut open the skin which surrounds it; put the eggs in a basin, pour water over them, mashing with the hand, to separate them; throw away the water. Add $\frac{1}{2}$ lb. of salt and a teaspoonful of pepper. Let them soak all night. Wash well 2 or 3 times, leaving about a gill at the bottom. Put over it now 2 lbs. of the liver, in 6 or 8 pieces; put the stew-pan on a very slow fire or in an oven for an hour; then take out the liver, which serve as usual. Add about a gill of melted butter to the roe, when it will be ready.

COD-LIVER SAUCE. $\frac{1}{2}$ lb. of cod-liver, previously boiled, and cut into large dice, may be added to a little anchovy sauce, and $\frac{1}{2}$ pint of melted butter. (This may be eaten with

potatoes or fish. Cod-liver oil may be used as a sauce in the same way, disguising its taste in the butter by pepper, vinegar, or anchovies, and gradually increasing its amount as the patient becomes habituated to it.)

COD-LIVER OIL AND KETCHUP (Mr FAIRTHORNE). Add 2 drs. of tomato or walnut ketchup, to each ounce of the oil and shake well before taking.

The taste of cod-liver oil may be very successfully covered by shaking it up with an equal quantity of fresh milk, and adding to every two ounces of such mixture 5 drops of true oil of cinnamon. The mixture will remain fresh for a long time.

CONDIMENTS, AND VARIOUS CULINARY COMPOUNDS

CURRY POWDER. The recipes for "true Indian Curry Powder" are numerous, and vary much in the number and proportion of the ingredients. The total quantity of powder in each of the following recipes being nearly equal, the relative proportion of the different colouring, pungent, and flavouring ingredients will at once be seen. Dr KITCHENER complains that the proportion of cayenne is generally so large that a proper quantity of the powder cannot be used to obtain the benefit of the other ingredients; and the Editor of the *Pharmaceutical Journal* justly observes that many recipes contain too large a proportion of turmeric. All the ingredients should be of fine quality, and recently ground.

	1	2	3	4	5	6	7	8	9	10	11	12
Turmeric . . .	9	6	6	8	9	9	9	4	6	6	7	8
Coriander seed .	9	16	12	22	10	9	16	12	11	16	13	12
Mustard, <i>scored</i>	3	—	—	—	3	3	—	4	—	—	—	—
Cayenne . . .	2½	4	1½	1	1¼	1½	1½	2	1	2	1	1
Pepper, black or } long . . . }	6	3	8	2	3	3	2½	4	5	1½	5	4
Pimento . . .	—	—	—	—	—	—	—	—	2	—	—	—
Cloves . . .	1	—	—	—	½	—	—	—	½	—	—	—
Cinnamon . . .	2	—	—	—	—	—	—	—	3	—	—	—
Cardamom . . .	2	3	—	—	1½	1	—	—	—	—	—	2
Ginger . . .	1	—	—	—	3	3	—	—	2	1½	—	4
Mace . . .	1	—	—	—	—	—	¼	—	—	—	—	—
Fenugreek . . .	—	—	3	—	—	—	—	¼	2	3½	3	—
Cumin . . .	—	—	2	—	1	—	3	2	—	—	3	1

The addition of 1 oz. of garlic, or 2 oz. of shallots, to 2 lbs. of either of the above, will be approved by some palates.

The true Indian curry is said to be thus made: Coriander seed 6 drs., turmeric 5 scruples, fresh ginger $4\frac{1}{2}$ drs., cumin seed 18 grs., black pepper 54 grs., poppy seed 94 grs., garlic 2 heads, cinnamon a scruple, cardamom 5 seeds, 8 cloves, 1 or 2 chillies, half a cocoa-nut grated; all but the last to be ground on a stone.

BENGAL CHITNI. Chillies $1\frac{1}{2}$ lbs., unripe mangoes (or apples) 1 lb., red tamarinds 2 lbs., sugar candy 1 lb., fresh ginger root $1\frac{1}{2}$ lbs., garlic $\frac{1}{2}$ to $1\frac{1}{2}$ lbs., sultana raisins $1\frac{1}{2}$ lbs., fine salt 1 lb., and 5 bottles of the best vinegar; soak the chillies for an hour in the vinegar, then grind all with a stone and muller to a paste.

ITALIAN TAMARA. Coriander seed, cloves, and cinnamon, of each 8 oz.; anise and fennel seeds, of each 4 oz.; mix.

MIXED SPICES AND SAVOURY HERBS. 1. *Kidder's Sweet Spice.* Equal weights of cloves, mace, nutmegs, cinnamon, and sugar.

2. *Kidder's Savoury Spice.* Equal weights of salt, pepper, cloves, nutmegs, and mace.

3. *Ragout.* Salt 16 oz., pepper 8 oz., nutmeg, ginger, and allspice, each $\frac{1}{4}$ oz.; lemon-peel 8 oz., mustard flour 8 oz.; cayenne 2 oz.; mix.

4. *Sausage.* Pepper 5 lbs., cloves $1\frac{1}{2}$ lbs., nutmegs $1\frac{1}{2}$ lbs., ginger $2\frac{1}{2}$ lbs., aniseed $\frac{1}{2}$ lb., coriander seed $\frac{1}{2}$ lb.; mix.

5. *Dr KITCHENER'S Savoury Ragout.* Salt 2 oz., mustard, black pepper, and grated lemon-peel, of each 1 oz., allspice, ginger, and nutmeg, of each $\frac{1}{2}$ oz., cayenne $\frac{1}{4}$ oz.

6. *Soup Herb and Savoury Powder.* Mix 3 parts of No. 7 with 1 part of No. 5.

7. *Dr KITCHENER'S Soup Herb Powder, or Vegetable Relish.* Dried parsley, winter savory, sweet marjoram, lemon thyme, of each 2 oz., dried lemon-peel and sweet basil, of each 1 oz.; mix. They should be carefully dried in a Dutch oven, powdered, passed through a hair sieve, and kept in closely covered bottles. For sauces, soups, &c.

8. *Pease Powder.* Pound together in a marble mortar 2 oz. each of dried mint and sage, $\frac{1}{4}$ oz. each of celery

seed and black pepper, and rub them through a hair sieve.

HORSE-RADISH POWDER. Take up the roots in November or December, dry them carefully with a gentle heat, and reduce to powder.

SOLUBLE CAYENNE. To 1 lb. of the best cayenne pepper add as much rectified spirit as will form it into a paste, cover this up for 2 hours; then place it in a percolator, and gradually pour on it more spirit till a pint of liquid is procured. A little water cautiously poured on the pepper will displace most of the remaining spirit. Distil off most of the spirit for future use, and add to the residue 3 lbs. of fine salt and evaporate the mixture to dryness by the heat of a water-bath. It is usually coloured, but is better without being so.

CULINARY ESSENCES, TINCTURES, &c.

ALMOND FLAVOUR. Essential oil of bitter almonds 1 part, rectified spirit 7 parts. Some put 1 part of oil to 15 of spirit; others, 1 part to 3. It should not be used without a caution as to the quantity to be used; or rather, the oil should be first deprived of its hydrocyanic acid, by mixing it with a solution of chloride of iron and cream of lime, with a little peroxide of mercury, and, after a few days' contact, carefully re-distilling the oil.

FLAVOURING ESSENCE. Oil of bitter almonds (freed from hydrocyanic acid) 8 drops, essence of lemon 12 drops, oil of cinnamon 8 drops, oil of nutmeg 4 drops, highly rectified spirit 1 oz. A few drops to be added to puddings, custards, &c.

LEMON FLAVOUR. Fresh lemon-peel, cut thin, 3 drs., essence of lemon 1 dr., alcohol 3 oz. Another method is to rub a lump of sugar on clean, dry lemons, till the yellow rind is taken up by the sugar; then scrape off the saturated part of the sugar, and keep it in a closely covered pot for use.

TINCTURE OF CINNAMON (KITCHENER'S). Bruised cinnamon 3 oz., a bottle of Cognac brandy; digest for a fortnight, and strain. [TINCTURES OF ALLSPICE, NUTMEG, CLOVES, in the same manner.]

ESSENCE OF CINNAMON. Bruised cinnamon 2 drs., oil of cinnamon 1 dr., highly rectified spirit 3 oz.; digest, and strain.

ESSENCES OF NUTMEG, MACE, CLOVES, [ALLSPICE, &c.

These are made from the spices and their essential oils, as ESSENCE OF CINNAMON.

ESSENCE OF CELERY. Celery seed $\frac{1}{2}$ oz. to 1 oz., brandy 4 oz.; digest for 8 or 10 days, and filter.

ESSENCE OF CARAWAY. Bruised caraway seed 1 oz., rectified spirit 8 oz., oil of caraway $\frac{1}{4}$ oz., brown sugar $\frac{1}{4}$ oz.; digest for 8 or 10 days, and filter.

AROMATIC ESSENCE OF GINGER. Freshly grated ginger 3 oz., fresh thin lemon-peel 2 oz., brandy $1\frac{1}{2}$ pints; macerate for 10 days.—Dr KITCHENER.

ESSENCE OF PEAR, AND ESSENCE OF PINE APPLE. See TRADE CHEMICALS.

ESSENCE OF CAYENNE (KITCHENER'S). Put $\frac{1}{2}$ oz. of cayenne pepper into $\frac{1}{2}$ pint of brandy; let it steep for a fortnight, then pour off the clear liquor. A much stronger essence is sometimes kept, prepared by percolation, as directed for SOLUBLE CAYENNE.

SPIRIT OF SAVOURY SPICES (KITCHENER'S). Black pepper 1 oz., allspice $\frac{1}{2}$ oz., nutmeg $\frac{1}{4}$ oz. (all pounded); infuse in 16 fl. oz. of brandy for 10 days.

SPIRIT OF SOUP HERBS (KITCHENER'S). Lemon thyme, winter savory, sweet marjoram, sweet basil, of each 1 oz., grated lemon-peel and shallots, each $\frac{1}{2}$ oz., celery seed 1 dr.; infuse in a pint of brandy for 10 days.

SPIRIT OF SAVOURY SPICES. Infuse half the Savoury Ragout Powder (see MIXED SPICES, &c., No. 5, further back) in a quart of brandy for 10 days.

KITCHENER'S SOUP HERB AND SAVOURY SPICE SPIRIT. A mixture of equal measures of the last two.

CULINARY VINEGARS, SAUCES, &c.

TARRAGON VINEGAR. Put fresh tarragon leaves into a stone jar, and pour on them a sufficient quantity of the best wine vinegar to cover them. Set the jar in a warm place for 14 days; then strain through a jelly-bag. [In the same way may be made, *elder flower, basil, green mint, and burnet vinegars*. *Cress and celery vinegars* are made with $\frac{1}{2}$ oz. of the bruised seed to a quart of vinegar. *Horse-radish vinegar*, with 3 oz. of the scraped root, 1 oz. of minced shallots, 1 dr. of cayenne to a quart of vinegar. *Garlic vinegar* is made with 2 oz. of minced garlic to a

quart of wine vinegar. *Shallot vinegar* in the same proportion. *Chilli vinegar*, with 50 English chillies, cut or bruised (or $\frac{1}{4}$ oz. cayenne pepper), to a pint of the best vinegar; digest for 14 days.]

CAMP VINEGAR. Take 12 chopped anchovies, 2 cloves of garlic minced, 1 dr. of cayenne, 2 oz. of soy, 4 oz. of walnut catsup, and a pint of the best vinegar; digest for a month, and strain.

2. Vinegar a quart, walnut catsup a pint, mushroom catsup 4 tablespoonfuls, garlic 4 heads, cayenne $\frac{1}{4}$ oz., soy 2 tablespoonfuls, port wine 2 glasses, 3 anchovies, and a tablespoonful of salt; put them into a bottle, shake daily for a month, and decant.

CURRY VINEGAR. Infuse 3 oz. of curry powder in a quart of vinegar, near the fire, for 3 days.

RASPBERRY VINEGAR. Macerate 2 lbs., of fresh raspberries with a pint of the best vinegar for 14 days and strain. Or to a quart of the juice add 2 oz. of strong acetic acid, or enough to render it sufficiently acid.

ESCHALOT WINE. Bruised shallots 3 oz., sherry wine a pint; infuse for 10 days. An ounce of scraped horse-radish and a drachm of thin lemon-peel may be added. [“The most elegant preparation of the onion tribe.”—Dr KITCHENER.] Wines of several herbs may be made in the same proportion as the vinegars.

FRENCH MUSTARD. This is sold with a great variety of flavours. A good substitute may be made by mixing good flour of mustard with the liquor of walnut and other pickles; or with the flavoured vinegars above. The following is one of the published recipes:—Salt 12 oz., scraped horse-radish 8 oz., a clove of garlic, $\frac{1}{2}$ oz. of sugar, a gallon of French vinegar (hot but not boiling). Macerate for 24 hours, and strain. Mix with flour of mustard q. s.

MUSTARD FOR THE TABLE. Mix 8 spoonfuls of flour of mustard with 2 of salt, and 9 of water. Mix to a smooth paste, add 6 spoonfuls more of water, and mix.

ESSENCE OF ANCHOVIES. Beat 1 lb. of anchovies in a Wedgewood mortar, and put them into a pipkin with 4 oz. of vinegar; boil for a few minutes, and rub the pulp through a hair sieve. Boil the bones in $1\frac{1}{2}$ lbs. of water, strain, and add 2 oz. of salt, and 2 oz. of flour of starch, and the pulped anchovies; let it boil, and pass it through a hair sieve. It is usually coloured with powdered bole,

or with annotto. Gum tragacanth is sometimes used to stiffen it instead of flour. Another method is to simmer anchovies in their own weight of water for 2 or 3 hours, removing any scum that may rise, strain with pressure through a strong canvas bag, and filter through flannel. This has the pure flavour of the fish, but a little cayenne and salt may be added, to preserve it.

ANCHOVY PASTE. Pound the fish in a mortar, and rub the pulp through a fine sieve. Put it into pots, and cover with clarified butter.

MUSHROOM CATSUP. Press the mushrooms in a tincture press and boil the juice with $\frac{1}{2}$ oz. black pepper-corns, 1 oz. pimento, $\frac{1}{2}$ oz. of ginger, $\frac{1}{4}$ oz. cloves, $1\frac{1}{2}$ oz. shallots, and 8 oz. of salt, to each gallon. Some add 4 oz. of brandy. Or sprinkle the mushrooms with salt (a pound to 2 pecks), stir occasionally for 2 days, then squeeze them gently in a hair-sieve, and boil the liquor with pepper and other spices.

WALNUT CATSUP. 1. Mix 6 half sieves of green walnut-shells with 2 or 3 lbs. of salt in a wooden vessel; let them stand 6 days, beating them frequently till they become pulpy; then drain off the juice, and boil it up with 4 oz. of ginger and allspice, and 2 oz. of long pepper and cloves.

2. Juice of walnuts 1 gallon, anchovies 2 lbs., shallots 1 lb., cloves, mace, and black pepper, of each 1 oz., and a clove of garlic. Boil for a short time, and bottle it.

LEMON PICKLE. Slice 6 lemons, rub them with salt, lay them in a stone jar, with 2 oz. each of allspice and white pepper; add $\frac{1}{4}$ oz. each of mace, cloves, and cayenne, and 2 oz. each of horse-radish and mustard seed; pour over them 2 quarts of hot distilled vinegar; and, after standing for a few days, strain. Some add garlic or shallots.

QUIN SAUCE. Mushroom catsup $\frac{1}{2}$ pint, walnut pickle $\frac{1}{4}$ pint, port wine $\frac{1}{2}$ pint, 6 anchovies, and 6 shallots (both pounded); soy a tablespoonful; cayenne $\frac{1}{2}$ dr.; simmer together gently for 10 minutes, strain, and bottle.

WATERLOO SAUCE. Vinegar 4 pints, port wine 1 pint, cayenne 1 oz., walnut catsup $\frac{1}{2}$ pint, mushroom catsup $\frac{1}{2}$ pint, essence of anchovies 4 oz., powdered cochineal 1 oz., garlic 12 cloves.

EPICUREAN SAUCE. Indian soy 2 oz., walnut catsup, mushroom catsup, of each 8 oz., port wine 2 oz., bruised white

pepper $\frac{1}{2}$ oz., shallots 3 oz., cayenne $\frac{1}{2}$ oz., cloves $\frac{1}{2}$ oz. Macerate for 14 days in a warm place, strain, and add white wine vinegar to make up a pint.

SAUCE, SUPERLATIVE (Dr KITCHENER'S). Port wine and mushroom catsup, of each a pint; walnut or other pickle liquor, $\frac{1}{2}$ pint, powdered anchovies 4 oz.; fresh lemon-peel cut thin, sliced shallots, and scraped horse-radish, of each 1 oz., allspice and black pepper, of each $\frac{1}{2}$ oz.; cayenne 1 dr.; curry powder 3 drs.; celery seed 1 dr.; put them into a wide-mouthed bottle, stop it close, shake daily for a fortnight, and strain: a $\frac{1}{4}$ pint of soy may be added.

[A variety of sauces may be made by mixing, in different proportions, the ingredients of the last 4 sauces.]

CASSAREEP. The expressed juice of the roots of the bitter cassava; used as a condiment in the West Indies.

SOY. Boil a gallon of the seeds of *Dilochos soja* till soft, add a gallon of bruised wheat, keep them in a warm place for 24 hours; add a gallon of salt, and 2 gallons of water, and after keeping them bunged up in a stone jar for 2 or 3 months, press out the liquor.

PICKLES

A few recipes are here given as illustrations of the methods employed in preparing these condiments. For full particulars the reader is referred to the popular treatises on Cookery. The best vinegar (pickling, or No. 24 vinegar) should be employed. Some prefer the crystal or white vinegar (distilled vinegar, or rather pure diluted wood-vinegar), especially for white pickles; but the best wine vinegar is more agreeable. Stoneware jars, not glazed with lead, should be used to keep the pickles in; or otherwise green glass jars.

SPICED VINEGAR, FOR PICKLES GENERALLY. Bruise in a mortar 2 oz. of black pepper, 1 oz. of ginger, $\frac{1}{2}$ oz. of allspice, and 1 oz. of salt. If a hotter pickle is desired, add $\frac{1}{2}$ dr. of cayenne, or a few capsicums. For walnuts, add also 1 oz. of shallots. Put these into a stone jar, with a quart of vinegar, and cover them with a bladder wetted with the pickle, and over this place a piece of leather. Set the jar on a trivet near the fire for 3 days, shaking it 3 times a day, then pour it on the walnuts or other vegetables. For walnuts it is used hot, but for cabbage,

&c., cold. To save time, it is usual to simmer the vinegar gently with the spices, which is best done in an enamelled saucepan.

BEEF ROOT. Boil the roots till 3 parts done (from 1½ to 2½ hours); then take them out, peel them, and cut them in thin slices. Put them into a jar, and pour on them sufficient cold spiced vinegar (as above) to cover them.

CABBAGE, WHITE. Cut it into thin slices, put them into an earthen pan, sprinkle them with salt, and let them lie for 2 days; then drain them and spread them out before the fire for some hours; put them into a stone jar, and add sufficient white vinegar, or pale malt vinegar, to cover them, and add a little mace and white pepper-corns.

CABBAGE, RED. Remove the outer leaves and stalks, and cut the cabbage in quarters, then shred them into a cullender, and sprinkle them with salt; next day drain them, put them into a jar, and pour on them sufficient cold spiced vinegar to cover them. Others hang up the cabbage for a few days to dry, then shred the leaves, and put them in layers in a jar with a little salt, pepper, and ginger, and fill up with cold vinegar. Others use vinegar without spice.

CAULIFLOWER AND BROCCOLI. These should be sliced, and salted for 2 or 3 days, then drained, and spread upon a dry cloth before the fire for 24 hours; then put into a jar, and covered with spiced vinegar. Dr KITCHENER says that, if vegetables are put into cold salt and water (a ¼ lb. of salt to a quart of water) and gradually heated to boiling, it answers the same purpose as letting them lie some days in salt.

CUCUMBERS. *Gherkins.* Small cucumbers, but not too young, are wiped clean with a dry cloth, put into a jar, and boiling vinegar, containing a handful of salt, poured on them. Boil up the vinegar every 3 days, and pour it on them till they become green; then add ginger and pepper, and tie them up close for use. Or cover them with salt and water (as above) in a stone jar, cover this, and set them on the hearth before the fire for 2 or 3 days, till they turn yellow; then put away the water, and cover them with hot vinegar, set them near the fire, and keep them hot for 8 or 10 days, till they become green; then pour off the vinegar, cover them with hot spiced vinegar, and keep them close.

MANGOES. Large cucumbers, or small melons, are split so that a marrow-spoon may be introduced, and the seeds scooped out; they are then parboiled in brine strong enough to float an egg, dried on a cloth before the fire, filled with mustard seed and a clove of garlic, and then covered with spiced vinegar. True mangoes the same.

MUSHROOMS. Clean them with water and flannel, throw them into boiling salt and water in a stewpan, and let them boil for a few minutes. Drain them in a cullender, and lay them on a linen cloth, covering them with another. Put them into bottles with a blade or two of mace, and fill up with white vinegar, pouring some melted mutton fat on the top, if they are intended to be kept long.

NASTURTIUMS, FRENCH BEANS, and other small green vegetables, are pickled in the same manner as gherkins.

ONIONS. 1. Let them lie in strong salt and water for a fortnight; then take them out and peel them; put them in fresh salt and water for another fortnight; take them out, wash them clean, and let them lie in fresh water all night. Next day put them on a cloth to drain; then put them in a jar, and pour over them hot spiced vinegar. If you wish them of a nice colour use white vinegar.

2. Peel small silver button onions, and throw them into a stewpan of boiling water; as soon as they look clear, take them out with a perforated spoon, and lay them on a folded cloth, covered with another, and when quite dry, put them into a jar, and cover them with hot spiced vinegar. When quite cold, bung them down, and cover with bladder wetted with the pickle, and leather.

WALNUTS. Take 100 young walnuts, lay them in salt and water for 2 or 3 days, changing the water every day. (If required to be soon ready for use, pierce each walnut with a larding-pin, that the pickle may penetrate.) Wipe them with a soft cloth, and lay them on a folded cloth for some hours. Then put them in a jar, and pour on them sufficient of the above spiced vinegar, hot, to cover them. Or they may be allowed to simmer gently in strong vinegar, then put into a jar with a handful of mustard seed, 1 oz. ginger, $\frac{1}{4}$ oz. mace, 1 oz. allspice, 2 heads of garlic, and 2 split nutmegs, and pour on them sufficient boiling vinegar to cover them. Dr KITCHENER recommends the walnuts to be gently simmered with the brine, then laid on a cloth

for a day or two, till they turn black, put into a jar, and hot spiced vinegar poured on them.

TOMATOES. AS GHERKINS. See CUCUMBERS.

PICCALILLI, INDIAN, OR MIXED PICKLES. 1. To each gallon of strong vinegar put 4 oz. of curry powder, 4 oz. of good flour of mustard, 3 oz. of bruised ginger, 2 oz. of turmeric, 8 oz. of skinned shallots, and 2 oz. of garlic (the last two slightly baked in a Dutch oven), $\frac{1}{4}$ lb. of salt, and 2 drs. of cayenne pepper. Digest these near the fire, as directed above for spiced vinegar. Put into a jar, gherkins, sliced cucumbers, sliced onions, button onions, cauliflower, celery, broccoli, French beans, nasturtiums, capsicums, large cucumbers, and small melons. All, except the capsicums, to be parboiled in salt and water, drained, and dried on a cloth before the fire. The melons and large cucumbers to be prepared as directed above for mangoes. Pour on them the above pickle.

2. Take 1 lb. of ginger-root, and $\frac{1}{2}$ lb. of garlic (both previously salted and dried), 2 gallons of vinegar, $\frac{1}{2}$ oz. of turmeric, $\frac{1}{4}$ lb. of long pepper. Digest together for 2 or 3 days near the fire in a stone jar; or gently simmer them in a pipkin or enamelled saucepan. Then put in the above vegetables, or almost any except red cabbage and walnuts, all previously salted and dried.

BRINE, OR PICKLE, FOR PORK, &c. Brown sugar, bay salt, common salt, of each 2 lbs., saltpetre $\frac{1}{2}$ lb., water a gallon; boil gently, and remove the scum. Another meat pickle is made with 12 lbs. of salt, 2 lbs. of sugar or treacle, $\frac{1}{2}$ lb. of nitre, and sufficient water to dissolve it. To cure HAMS, mix 5 oz. of nitre with 8 oz. of coarse sugar; rub it on the ham, and in 24 hours rub in 2 lbs. of salt, and in a fortnight 2 lbs. more. The above is for a ham of 20 lbs.; it should lie in the salt a month or 5 weeks.

WESTPHALIAN ESSENCE—CAMBRIAN ESSENCE OF WOOD SMOKE. These appear to be crude pyroligneous acid, or wood vinegar, and are used to give to hams, &c., the smoked flavour. The following has been published as the recipe for Essence of Smoke, but we apprehend it is far from being correct:—Macerate for several weeks $\frac{1}{2}$ dr. of Barbadoes tar, 1 dr. of liquid burnt sugar, 5 drs. each of port wine and vinegar, 2 drs. of salt, and 7 oz. of water.

TRADE CHEMICALS

MISCELLANEOUS PREPARATIONS, & COMPOUNDS
EMPLOYED IN THE ARTS, IN DOMESTIC ECO-
NOMY, CHEMICAL RESEARCH AND AMUSE-
MENT, &c.

THIS division of the work comprises those chemical compounds which are employed for other purposes than those of medicine, and which have not been noticed in the former parts of this volume. It includes a variety of miscellaneous articles which are sometimes sold by the retail druggist, or the materials of which he is expected to furnish, or with the composition of which it is desirable he should be acquainted. The limits of the work do not admit of a minute description of the processes and manipulations employed in the manufacture of such chemicals as are only made on the large scale, and never by the retailer; nor of those chemical arts which have no immediate connection with the trade.

ACETATES. Such as are employed in medicine will be found in the *Pocket Formulary*. The only acetates requiring notice here are the following :

ACETATE OF ALUMINA. This is made, for the use of dyers and calico printers, by decomposing acetate of lime with alum. It may be conveniently made by adding to a boiling solution of 5 parts of alum, a solution of 6 parts of sugar of lead. When the mixture is cold, the clear liquid is poured off; from which the dry salt may be obtained by careful evaporation. It contains, besides acetate of alumina, some sulphate of potash.

ACETATE OF IRON, OR IRON LIQUOR. Usually made, for the use of dyers, by digesting scraps of iron in redistilled wood-vinegar. (See IRON LIQUOR, further on.)

ACETATE OF LIME. Impure acetate (or pyrolignite) of lime, is made by neutralizing pyroligneous acid with cream of lime or chalk, and evaporating to dryness. By using pure acetic acid a purer acetate is obtained.

ACETATE OF SODA. By mixing the above impure acetate of lime, in solution, with a solution of sulphate of soda, filtering and evaporating the clear liquid, an impure acetate of soda is obtained; which by repeated crystallization is rendered colourless, and fit for yielding pure concentrated acetic acid by distillation with sulphuric acid.

ACETIMETRY. The strength of vinegar is estimated for the duty by an instrument named an acetimeter, which determines the quantity of acetic acid present by the specific gravity of the vinegar after neutralization by slaked lime. Dr URE's plan is to add to a given weight of vinegar, bicarbonate of potash till exactly neutralized; every 2 grs. of the bicarbonate indicate 1 gr. of real acetic acid. In this and the following operations it is convenient to use a tube graduated into 100 equal divisions, numbered from the top downwards (see ALKALIMETRY, further on). The quantity of test solution used is then seen at once. In the present case the 200 grs. of the alkaline bicarbonate being dissolved in sufficient water to fill the graduated portion of the measure, each of the divisions used in neutralizing 100 grs. of vinegar is equivalent to 1 per cent. of absolute acetic acid. Consult the larger manuals.

ACID, ACETIC. See VINEGAR. For the methods of procuring this concentrated acetic acid, see ACIDUM ACETICUM, *Pocket Formulary*. The acetic acid of the British Pharmacopœia contains 33 per cent. of real acetic acid. A strong acid, very suitable for making aromatic spirit of vinegar, is procured by distilling crystallized verdigris in an earthen retort coated with clay, into a series of 3 globes, connected by opposite tubulures, and kept constantly cool, the last being furnished with a WELTER's safety tube. The acid which comes over is usually coloured, and requires to be rectified by a slow and careful redistillation in a glass retort. Acetic acid of moderate strength may be rendered stronger by redistilling it over acetate of potash, rejecting the first portions that come over, and taking care that the temperature does not rise above 572° F. By redistilling it, and rejecting the

first and last portions, glacial acetic acid is procured. The same acetate of potash may be used repeatedly. The process of the Dublin Pharmacopœia yields a good product of glacial acetic acid.

ACID, BORACIC. *Boric Acid.* The commercial acid is derived from the *suffioni* and *fumarole* of Tuscany. These are jets of steam issuing from the interior of the earth, and holding a very minute quantity of the acid in solution. The acid is obtained by condensation of the vapour, and subsequent evaporation and crystallization. For the laboratory, boric acid may be made by adding hydrochloric acid to a hot saturated solution of borax. As the liquid cools, crystals of the acid separate and fall to the bottom. The crystals may be purified by recrystallization from solution in hot water. Largely used in the arts, and as an antiseptic for the preservation of food. See **ACIDUM BORICUM**, *Pocket Formulary*.

ACID, CARBOLIC. See *Pocket Formulary*.

ACID, CHROMIC. *Chromic Anhydride.* It may be obtained by pouring 1 vol. of a saturated solution of bichromate of potash in a thin stream into $1\frac{1}{2}$ vols. of strong sulphuric acid, stirring the whole time. As the liquid cools, the anhydride crystallizes from it in crimson needles. The crystals are drained on a porous tile till nearly dry, and then recrystallized. See **ACIDUM CHROMICUM**, *Pocket Formulary*.

ACID, CINNAMIC. It is most readily procured by distilling genuine balsam of tolu by a gentle heat. The white crystalline mass which condenses on the neck of the retort is purified by pressing it between blotting paper, dissolving in boiling water, and recrystallizing.

ACID, FLUORIC. *Acid, Hydrofluoric.* The anhydrous acid is made by distilling powdered fluor spar with twice its weight of oil of vitriol in a leaden, or better, a silver alembic, the pipe of which fits into a bottle of the same material, surrounded with ice. But as it is usually required in a diluted state, water equal in weight to the spar may be put into the receiver. Great care must be taken, as the acid, both in its gaseous and liquid form, is very destructive.

ACID, HIPPURIC. Mix the urine of the horse with milk of lime, boil for some minutes, and strain. Boil down the clear liquid to $\frac{1}{8}$ th of its bulk, avoiding burning; add

hydrochloric acid, press the impure acid, boil it with fresh milk of lime, and again precipitate with hydrochloric acid.

ACID, HYDROBROMIC. See **ACIDUM HYDROBROMICUM DILUTUM**, *Pocket Formulary*.

ACID, IODIC. See **ACIDUM IODICUM**, *Pocket Formulary*.

ACID, LACTIC. See **ACIDUM LACTICUM**, *Pocket Formulary*.

ACID, MURIATIC, OR HYDROCHLORIC. Commercial hydrochloric acid is largely produced by the action of sulphuric acid on common salt, in the manufacture of sulphate of soda, which salt is used in making soda ash and washing soda. From the impurity of the ingredients used in its manufacture, the acid is apt to be contaminated with arsenic and sulphurous acid, as well as with sulphuric acid, and iron. It may be purified from arsenic by redistilling it over strips of bright copper. See **ACIDUM HYDROCHLORICUM**, *Pocket Formulary*.

ACID, NITRIC, AND FUMING NITROUS ACID. See **ACIDUM NITRICUM**, and **ACID. NITRIC. FUMANS**, *Pocket Formulary*.

ACID, NITRO-HYDROCHLORIC. *Aqua Regia*. This is used in the arts, chiefly as a solvent for gold. By the mutual action of nitric and hydrochloric acids a compound of chlorine, nitrogen, and oxygen is formed. The best proportions and strength of the acid are variously stated. Colourless nitric acid must be used. Mr **ELKINGTON** employs 21 parts of nitric acid, sp. gr. 1.45; 17 parts of hydrochloric acid, 1.15 sp. gr.; and 14 parts of water. This dissolves 5 parts of gold. For the nitro-hydrochloric acid employed by dyers as a solvent for tin, see **DYES, &c.**, further on.

ACID, OLEIC. See *Pocket Formulary*.

ACID, OXALIC. Digest by the aid of heat 1 part of treacle, or of potato starch, in 5 parts of nitric acid, sp. gr. 1.42, diluted with 10 parts of water, so long as gaseous products are evolved. By evaporation the acid is obtained in crystals, and must be recrystallized until sufficiently pure. Mr **LEWIS THOMPSON** directs 28 oz. of sugar, 184 oz. of nitric acid of 1.245 sp. gr., to be digested at 125° F. It yields 30 or 31 oz. of acid. M. **SCHLESINGER** directs 4 parts of dry sugar and 33 of nitric acid of 1.38 sp. gr. to be boiled to one sixth of the original volume, and allowed to crystallize. This is the best method of operating on a

small scale, when the amount of product is not the principal object. Considerable quantities of oxalic acid are made, on the large scale, by heating sawdust with a mixture of caustic potash and soda, and by subsequently decomposing the oxalate of soda formed.

ACID, PHOSPHORIC. *Phosphoric Anhydride.* See *Pocket Formulary*. Dry phosphoric acid is thus obtained: On a flat plate place a large bell glass, and under it a small porcelain cup or crucible. Introduce into the latter a piece of phosphorus, dried with blotting paper, and set it on fire by a heated wire. Let the bell glass be raised on one side to admit sufficient air to maintain combustion; and as the phosphorus is consumed, introduce successive pieces, taking care that the glass does not become too hot. When the quantity of acid is considerable, knock it from the plate with an iron spoon, and put it into stoppered bottles. Several glasses may be used at once. Dry phosphoric acid is used as a desiccating body, having the strongest attraction for water of any known substance. It is used also in making a stopping for teeth—see **TEETH CEMENTS**.

ACID, HYPOPHOSPHOROUS. See *Pocket Formulary*.

ACID, PICRIC. *Trinitrophenic Acid. Carbazotic Acid.*
Produced by the action of nitric acid on carbolic acid.

ACID, PYROGALLIC. Heat powdered nutgalls in a dish covered with thin filtering paper pasted to its edges, and surmounted with a bell-formed receiver, kept cool. A solution of the condensed acid, decolorized by animal charcoal, and mixed with spirit, is used to stain the hair (and skin) brown.

ACID, SULPHURIC. This is only made on a large scale; but the commercial acid requires purification for many chemical as well as pharmaceutical purposes. See **ACIDUM SULPHURICUM PURUM**, *Pocket Formulary*.

DRY OR ANHYDROUS SULPHURIC ACID. *Sulphuric Anhydride.* Into a retort, placed in a freezing mixture, and having a receiver attached, put some dry phosphoric acid and add 3-4ths of its weight of strong sulphuric acid. Remove the retort from the freezing mixture, and place the receiver attached to it there; a gentle heat being now applied to the retort, the anhydrous acid is obtained in silky crystals.

ACID, SALICYLIC. May be procured from the volatile oil of

Gaultheria procumbens by heating it with strong potash ley till no more methylic alcohol is given off, the residue being precipitated with strong hydrochloric acid. The precipitate is washed with cold water and crystallized from alcohol. It is also obtained artificially from carbolate of soda by the action of carbonic acid, at an elevated temperature. (See *Pocket Formulary*.) KOLBE, KIERSCH, and GODEFROY have made experiments with this acid, from which it would appear that it possesses disinfecting and considerable antiseptic properties.

ACID, SULPHUROUS. For the mode of obtaining an aqueous solution of this acid, see **ACIDUM SULPHUROSUM**, *Pocket Formulary*. The following are cheaper methods of obtaining it for bleaching purposes, &c. BERTHIER directs a mixture of 100 parts black oxide of manganese, and 12 or 14 of sulphur, to be heated in a glass retort, and the gas received into water kept very cold. Mr REDWOOD directs $\frac{1}{2}$ oz. of powdered charcoal to be acted on by 4 fl. oz. of oil of vitriol. Treacle is sometimes used instead of charcoal; as well as linseed oil.

ACID, TANNIC. *Tannin*. See **ACIDUM TANNICUM**, *Pocket Formulary*.

ACIDS, MIXED, FOR GALVANIC BATTERIES. 1. For troughs for general purposes, medical galvanism, &c.: Nitric acid 1 fl. oz., sulphuric acid $1\frac{1}{2}$ fl. oz., water 4 pints.

2. For BUNSEN'S Battery. For the glass vessel: nitric acid at 36° or 40° Baumé. For the coke cylinder: sulphuric acid 1 part, water 9 parts.

3. FARADAY. Oil of vitriol 2 fl. oz., nitric acid 1 fl. oz., water 5 pints. It should be tried by dipping into it a piece of sheet zinc. A continuous succession of bubbles should be produced.

4. For GRENET'S Battery. For the porous cell: bichromate of potash 10 parts, sulphuric acid 10 parts, water 100 parts.

5. For Mr SMEE'S Battery. *a*. One measure of sulphuric acid to 7 of water. The intensity of its action is increased by the addition of a few drops of nitric acid, but this tends to destroy the plates. *b*. Water saturated with common salt. In electro-metallurgy the water should only contain a sixteenth of sulphuric acid.

6. For Mr GROVE'S Battery. For the outer vessel:

1 pint of sulphuric acid to 7 of water; for the inner: concentrated nitric acid.

7. For DANIELL'S Battery. For the porous tube containing the zinc: 1 part of sulphuric acid with 10 of water. For the outer cylinder: a saturated solution of sulphate of copper, with a tenth part of sulphuric acid.

8. Nitro-sulphuric acid, for Dr T. WRIGHT'S Batteries. Nitric acid 1 part, sulphuric acid 5 parts. The zinc plate is immersed in a solution of chloride of ammonium or of salt, the platinized zinc in the above acid. The platinizing requires to be repeated every time the plate is washed.

9. For WATTS' Battery. Water 4 galls., sulphuric acid 2 lbs., nitric acid 1 oz. For the porous cell: a concentrated solution of common salt.

ACIDIMETRY. Acids generally are estimated by the quantity of alkalis* or carbonated alkalis required to neutralize them. Weigh 100 grs. of the acid and dilute it with a few times its weight in water. Then add sufficient dry or crystallized carbonate of soda, or carbonate of potash, to exactly neutralize the acid. The alkalimeter tube may be used for the solution of the alkali. By a reference to the table of chemical equivalents, the quantity of real acid of any kind represented by the quantity of alkali required to neutralize it may be estimated. Consult the larger manuals.

ACIDULATED KALI. See BEVERAGES.

ALBUMINOUS SIZE. Beat up the white of an egg with twice its bulk of cold water, until well incorporated. Used as a varnish for leather binding and kid gloves; also to size drawing paper.

ALCOHOL. May be obtained by the dehydration of rectified spirit of wine. See ALCOHOL, *Pocket Formulary*.

ALKALIMETRY. The quantity of real alkali contained in the commercial alkalis (common soda, soda ash, potashes, pearlash, salt of tartar, &c.) is ascertained by the quantity of an acid solution of known strength required to neutralize it. For this purpose a tube termed an ALKALIMETER is used, which will hold 1000 grains of distilled water. The tube may be three fourths of an inch internal diameter, and 9½ inches in length; or five eighths of an inch diameter, and 14 inches in height; and should be graduated

* The caustic alkalis are preferable and more convenient for this purpose than the carbonated ones.

into 100 equal divisions numbered from the top downwards. The quantity of test acid used is then at once seen: This consists of sulphuric acid diluted with water; so that each measure (10 grs.) is equivalent to 1 gr. of pure soda. To use it, dissolve 100 grains of the impure soda in 3 oz. of hot water, filter, and wash the filter. Then add to the solution the test acid until the litmus or cabbage paper ceases to show an alkaline reaction. The same acid will serve for potash, if, roughly speaking, the number be multiplied by 4 and divided by 3.—PARNELL. There are several other methods of performing the process given in the larger manuals.

ALLOYS AND AMALGAMS. A few only of these metallic compounds require notice here:—

Fusible Metal. 1. Tin 8 parts, lead 4, bismuth 3; melt together, removing the scum. Used as a metal-bath.

2. DARCET'S for the same purpose: Bismuth 8 parts, lead 5, tin 3.

3. Lead 3 parts, tin 2, bismuth 5. This melts at 197° Fahrenheit.

4. *For anatomical injections:* Melt together with a gentle heat 174 parts of tin, 312 of lead, 514 of bismuth, with a little charcoal: remove from the fire, and add 100 parts of mercury previously heated. It is fluid at 173°; solid at 140° Fahrenheit.

Brass. Mostly made of copper and zinc, the proportions varying according to the required colour and the purpose for which it is intended.

Britannia Metal. Tin 150, copper 3, antimony 10.

Bronze. 1. *For medals and small castings:* Copper 95 tin 4.

2. Copper 89, tin 8, zinc 3.

3. *Ancient.* Copper 100, tin 7, lead 7.

4. *KELLY'S.* Copper 91, zinc 6, tin 2, lead 1.

5. *For gilding.* Copper 14, zinc 6, tin 4.

6. *For cutting instruments.* Copper 100 parts, tin 14 parts.

7. *Bell-metal.* Copper 78, tin 22.

Cock metal. Copper 20 lbs., lead 8 lbs., litharge 1 oz., antimony 3 oz.

German Silver. 1. Copper 40½, nickel 31½, zinc 25½, iron 2½.

2. Pure copper 55, nickel 23, zinc 17, iron 3, tin 2.

Gold, Factitious. 1. Platinum 7, copper 16, zinc 1; fuse together.

2. *Aluminium Gold (pale).* Copper 100, aluminium 10. See AURUM MUSIVUM.

Common Gold. Copper 16, silver 1, gold 2.

Or-molu. Contains a larger proportion of copper and less zinc than ordinary brass.

White Brass. POREL. Melt zinc with 10 per cent. each of copper and iron. This alloy has the fracture and appearance of zinc, but is tougher than cast iron. It does not rust, nor adhere to metal moulds.

Solders. 1. For Gold: Pure gold 12 parts, silver 2, copper 4.

2. *Soft Solder.* Tin 2 parts, lead 1.

3. For brass: Brass 2 parts, zinc 1.

4. For aluminium: Tin 45, aluminium 10.

Type Metal (ordinary). Lead 9, antimony 1.

Alloys for Electrotype. Cliché Moulds. Bismuth 8 parts, tin 4, lead 5, regulus of antimony 1; melt repeatedly together, and pour out in drops, till perfectly mixed.

Amalgam for Electrical Machines. 1. Fuse 1 oz. of zinc with $\frac{1}{2}$ oz. of tin, at as low a temperature as possible; then add $1\frac{1}{2}$ oz. of quicksilver, previously made hot; mix, pour out, and when cold, reduce it to a powder, and triturate it with sufficient quicksilver to bring it to a proper consistence.

2. Zinc 1 part, tin 1, quicksilver 2; melt together.

3. Zinc 2, tin 1, mercury 5.

4. LA BEAUME'S. Pour into a chalked wooden box 6 oz. of quicksilver; pour into an iron ladle $\frac{1}{2}$ oz. beeswax, with 2 oz. of purified zinc, and 1 oz. of grain tin; set it over a brisk fire, and when the metals are melted, pour them into the box, avoiding the dross. When cold, reduce it to a powder, and mix it with lard. Keep it in a box, covered with tallow, and spread it on leather for use.

Liquid Amalgam for Silvering Globes, &c. Pure lead 1 oz., grain tin 1 oz.; melt in a clean ladle, and immediately add 1 oz. of bismuth. Skim off the dross, remove the ladle from the fire, and, before the metal sets, add 10 oz. of quicksilver. Stir together, avoiding the fumes.

Amalgam for Varnishing Figures. Melt 2 oz. of tin with $\frac{1}{2}$ oz. of bismuth, and add $\frac{1}{2}$ oz. of quicksilver. When

cold, grind it with white of egg, and apply to the figure.

ALUM is prepared by lixiviating calcined aluminous schist, and concentrating the solution to 1.4 or 1.5 density, and then adding the requisite quantity of chloride of potassium, soap-boilers' ash, or kelp, to supply the alkali. By recrystallization it is obtained colourless. In some manufactories sulphate of ammonia, from gas-liquor is added to the lixivum, instead of chloride of potassium. Alum is also manufactured from clay, cryolite or Greenland spa, Bauxite (a mineral containing a large proportion of hydrated alumina), and blast-furnace slag; also by DUNCAN and NEWLAND'S process, from beetroot sugar solution. Roman or cubic alum is crystallized from a solution, the temperature, of which is not allowed to exceed 104° F. It differs from common alum in containing a large quantity of base, a portion of which separates, if the solution be heated to 120° F. Another kind of alum, sometimes used as a mordant, consists almost entirely of sulphate of alumina, and is made by boiling fine clay, free from iron, with sulphuric acid, and cooling the solution so as to obtain a solid mass. See DYES, &c.

ALUMINIUM, TO FROST. The metal is plunged into a solution of caustic potash. The surface, becoming frosted, does not tarnish on exposure to the air.—MACADAM.

ALUMINIUM BRONZE. 10 parts of aluminium are melted with 90 parts of copper. It is said to be as tenacious as steel.

AMADOU. Prepared from *Boletus igniarius*, *B. fomentarius*, and some other allied species of fungi. The fungus is cut into thin slices, the hard external parts removed, and the rest beaten with a mallet till soft. This forms *surgeon's agaric*. If intended for *German tinder* it is soaked in a solution of nitre and sometimes sprinkled with gunpowder and carefully dried.

AMALGAMS. See ALLOYS, further back.

AMMONIA, SULPHATE AND CARBONATE OF. An impure sulphate of ammonia suitable for agricultural purposes is obtained by neutralizing the ammoniacal liquor of gas-works with sulphuric acid. By recrystallization and filtration through animal charcoal it may be obtained in a state of greater purity. The carbonate (hydrated sesquicarbonate) is obtained by mixing either this sulphate or

sal-ammoniac with chalk, and subliming it in iron retorts into leaden receivers. It is further purified by resubliming it with a gentle heat. See AMMONIÆ CARBONAS, *Pocket Formulary*.

ANATOMICAL INJECTIONS. *For coarse preparations.* 1.

Tallow, resin, and wax, equal parts; melt over a slow fire; and add red lead or vermilion sufficient to colour.

2. *For delicate preparations.* A strong solution of isinglass, coloured as required.

3. AMALGAM INJECTION. *Also for delicate parts.* Melt together 1 oz. each of bismuth, lead, and zinc, and, when melted, add 2 oz. of quicksilver.

ANATOMICAL SUBJECTS, AND ANIMAL SUBSTANCES TO PRESERVE.

1. M. GANNAL'S *Solution*. Common salt 2 lbs., alum 2 lbs., nitre 1 lb., water 4 gallons. M. GANNAL injects into the carotid artery a solution of sulphate of alumina, of density 1.286. From 5 to 7 pints are required in summer, but less will suffice in winter.

2. (WICKERSHEIMER.) In 3000 parts of boiling water, dissolve 100 of alum, 25 of sodium chloride, 12 of potassium nitrate, 60 of potassa, and 10 of arsenious acid; let cool and filter. To every ten litres of the filtrate add 4 litres of glycerin, and one litre of methylic alcohol.

3. Dr BABINGTON injects pyroxylic spirit into the aorta, and a little into the cavity of the perineum and the rectum.

4. CHLORAL HYDRATE is a powerful antiseptic; made into a weak solution, it may be successfully employed for the preservation of anatomical preparations.

5. Mr GOADBY, *for insects, and for preparations of their organs*. Bay salt 4 oz., alum 2 oz., corrosive sublimate from 2 to 4 grains, water 1 to 2 quarts. The weaker proportions should always be employed in the first instance. Let the insect or its organs be covered with the fluid, which should be changed frequently.

6. *For Mollusca*. Bay salt $\frac{1}{2}$ oz., arsenic $\frac{1}{2}$ dr., sublimate 2 grs., water a quart; dissolve.

7. Mr PIGNE, *for preserving pathological specimens*. Creasote 3 to 6 drops, water a pint.

8. CARBOLIC ACID, often sold for creasote, may be used in the same manner.

9. PURE GLYCERIN. Found especially useful in preserv-

ing the fresh tints of fishes, &c., intended for exhibition in museums.

10. Dr STAPYLTON, for *pathological specimens*. In a quart of saturated solution of alum dissolve $\frac{1}{2}$ dr. of nitre. A recent preparation immersed in this liquid becomes discoloured; but within a few days the colour returns. It is then put into a saturated and filtered solution of alum. M. REBOULET proposes: Water 16 parts, chloride of lime 4, alum 2, nitre 1.

11. Chloride of tin 4 (or corrosive sublimate 5) parts, in 100 of water, with 2 of hydrochloric acid.—Mr COOLEY.

12. For *preserving animals*. Alum 32 oz., nux vomica 3 oz., water 5 pints; boil to $4\frac{1}{2}$ pints. When cold, filter, or decant. This serves for injection. The residue, mixed with yolk of egg, is used for anointing the interior of the skin and fleshy parts left in skinning animals.

13. For *preserving feathers*. Strychnine 16 grs., rectified spirit a pint. (*Dangerous*.) See STUFFING BIRDS, &c. ANNOTTO, PURIFIED. To a boiling solution of pearlash add as much annatto as it will dissolve. When cold, decant the clear solution, and neutralize with diluted sulphuric acid, avoiding any excess. Wash the precipitate with a little cold water, and dry it.

ANNOTTO, SOLUTION OF. Boil equal weights of annatto and pearlash with water, and dilute to the required colour.

ANTI-ATTRITION, AND AXLE GREASE. 1. One part of fine black-lead, ground perfectly smooth, with $\frac{1}{4}$ parts of lard. Some recipes add a little camphor.

2. BOOTH'S AXLE GREASE. (Expired Patent.) Dissolve $\frac{1}{2}$ lb. common soda in 1 gallon of water, add 3 lbs. of tallow, and 6 lbs. palm oil (or 10 lbs. palm oil only); heat them together to 200° or 210° F.; mix, and keep the mixture constantly stirred till the composition is cooled down to 60° or 70° F. A thinner composition is made with $\frac{1}{2}$ lb. of soda, a gallon of water, a gallon of rape oil, and $\frac{1}{4}$ lb. of tallow or palm oil. See also LUBRICATING COMPOUNDS.

ANTI-FERMENT. Sulphite of lime; or equal parts of sulphite of lime and ground black mustard seed. Used to check the fermentation of cider.

AQUA FORTIS. *Double aqua fortis* is nitric acid of 1.36 specific gravity; *single aqua fortis* about 1.22.—DR

PERRIRA. A compound acid was formerly used under this name by dyers, and for cleaning brass, consisting of strong spirit of nitre 20 lbs., oil of vitriol 7 lbs., water 30 lbs.

AQUA REGIA. See ACID, NITRO-HYDROCHLORIC, further back.

AQUARIUM, MARINE, SALTS FOR, &c. Artificial sea-water may be used instead of the natural. A rough imitation is formed by mixing 100 oz. of fresh water with 3 oz. common salt, 1 oz. of Epsom salts, 200 grs. of chloride of magnesium, and 40 grs. of chloride of potassium.

Or, more precisely, the real constitution of sea-water may be imitated in the following manner:—Mix with 970,000 grains of rain-water, 27,000 of chloride of sodium, 3600 of chloride of magnesium, 750 of chloride of potassium, 29 of bromide of magnesium, 2300 of sulphate of magnesia, 1400 of sulphate of lime, 35 of carbonate of lime, 5 of iodide of sodium. These all being finely powdered and mixed first, are to be stirred into the water, through which a stream of air may be caused to pass from the bottom, until the whole is dissolved. On no account is the water to be boiled, or even heated.

Into this water, when clear, the rocks and seaweed may be introduced. As soon as the latter are in a flourishing state, the animals may follow. Care must be taken not to have too many of these, and to remove immediately any that die. The loss by evaporation is to be made up by adding clean rain-water. The aquarium, whether of fresh or of salt water, will require occasionally artificial aeration. This may be done by simply blowing through a glass tube which reaches to near the bottom; or, better still, in the following way:—Take a glass syringe which can be easily worked. Having filled it with water, hold it with the nozzle about 2 inches from the surface of the water in the aquarium, into which the contents are to be discharged quickly, and with a sort of jerk. By this means a multitude of small bubbles are forced down into the fluid. This operation should be repeated for a considerable number of times.

ARABINE. Gum arabic dissolved in water, and precipitated by alcohol.

ARGENTUM MUSIVUM. Fuse $\frac{1}{2}$ oz. each of grain tin and bismuth in a crucible, and add $\frac{1}{2}$ oz. of mercury.

ARBORE DIANÆ. See TREES, METALLIC.

AROMATIC PASTILS. See PERFUMERY.

AURUM MUSIVUM. *Mosaic gold. Bisulphide or bisulphuret of tin.* See STANNI SULPHURETUM, *Pocket Formulary*. 1.

Dr URE directs 12 oz. of tin to be melted, and 3 oz. of mercury added. This amalgam is triturated with 7 oz. of sulphur and 3 of sal ammoniac, and the powder put into a matrass, which is bedded deep in sand, and kept for several hours at a gentle heat. The heat is then raised, and continued for several hours, taking care not to raise it so high as to blacken the mass.

2. Melt together in a crucible, over a clear fire, equal parts of sulphur and the white oxide of tin; keep it continually stirred with a glass rod, until the compound appears as a yellow flaky powder. This may be used as a cheap bronze powder.

BALDWIN'S PHOSPHORUS. Heat nitrate of lime till it melts; keep it fused for 10 minutes, and pour it into a heated iron ladle. When cool, break it into pieces, and keep it in a closely stoppered bottle. After exposure to the sun's ray's it emits a white light in the dark.

BALLS FOR HORSES. See VETERINARY FORMULARY.

BALLS, ASH. The ashes of fern, or other kinds of wood ashes made into balls.

BALLS, HEEL. 1. Melt together 4 oz. of mutton suet, 1 oz. of beeswax, 1 oz. of sweet oil, $\frac{1}{2}$ oz. oil of turpentine, and stir in 1 oz of powdered gum arabic, and $\frac{1}{2}$ oz. of fine lamp-black.

2. Beeswax 8 oz., tallow 1 oz., powdered gum 1 oz., lamp-black q. s. These are used not merely by the shoemaker, but to copy inscriptions, raised patterns, &c., by rubbing the ball on paper laid over the article to be copied. ULLATHORNE'S Balls answer the purpose very well. For copying ancient monumental brasses a similar compound, coloured with bronze-powder, instead of lamp-black, is sometimes employed.

BALLS FOR SCOURING—BREECHES BALLS, CLOTHES BALLS.

1. Bath-brick 4 parts, pipe-clay 8 parts, pumice 1, soft-soap 1; ochre, umber, or other colour to bring it to the desired shade q. s.; ox-gall to form a paste. Make it into balls and dry them.

2. Pipe-clay 4 oz., fuller's-earth $\frac{1}{2}$ oz., whiting $\frac{1}{2}$ oz.,

white pepper $\frac{1}{2}$ oz., ox-gall sufficient to form it into a paste.

3. Pipe-clay 3 oz., white pepper 1 dr., starch 1 dr., orris powder $1\frac{1}{2}$ drs. It may be kept in powder, or formed into balls, as above.

BALLS, BLACKING. See **BLACKING.**

BALLS, FURNITURE. See **FURNITURE PASTE.**

BARIUM, PEROXIDE OF. Heat pure baryta to low redness in a platinum crucible; then gradually add chlorate of potash in the proportion of 1 part of the latter to 4 of the former. Cold water removes the chloride of potassium, and the peroxide remains as a hydrate.

BARIUM CHLORATE. See **CHLORATE OF BARIUM.**

BEEBLE POISON. Put a drachm of phosphorus in a flask with 2 oz. of water; plunge the flask into hot water, and when the phosphorus is melted, pour the contents into a mortar with 2 or 3 oz. of lard. Triturate briskly, adding water, and $\frac{1}{2}$ lb. of flour, with 1 or 2 oz. of brown sugar. — *Pharmaceutical Journal.* Plaster of Paris, with oatmeal, is said to destroy cockroaches.

BEEBLE WAFERS. Red lead 4 parts by weight, flour and brown sugar, of each 1 part. Cats and dogs will not touch it.

BENZINE COLLAS. Benzine is sold under this name.

BENZOL. *Benzine.* A volatile liquid, procured by distilling light coal naphtha at a temperature not exceeding 200° F., by the method patented by Mr MANSFIELD. It is a solvent for fats and oils, and hence is used for cleansing silks, and other stuffs. It likewise dissolves gutta percha and many other substances; and also, with heat and long digestion, India rubber.

BENZOLINE. A product of the fractional distillation of American rock oil, sold for sponge lamps. At a temperature less than 100° F. it will ignite if brought near a light, so that great care must be taken with it, and it should not be kept in quantity.

BEVERAGES, AND POWDERS FOR PREPARING THEM. See further back.

BIRD LIME. Boil the middle bark of the holly 7 or 8 hours in water; drain it, and lay it in heaps in the ground, covered with stones, for 2 or 3 weeks, till reduced to a mucilage. Beat this in a mortar, wash it in rain-water, and knead it till free from extraneous matters. Put it

into earthen pots, and in 4 or 5 days it will be fit for use. An inferior kind is made by boiling linseed oil for some hours, until it becomes a viscid paste.

BISULPHURET OF CARBON. *Bisulphide of Carbon. Disulphide of Carbon.* This is used in the arts, as a solvent for India rubber, gutta percha, &c. To procure it, MULDER recommends the following process as the most convenient. Provide an iron bottle (a quicksilver bottle answers very well), and make a second opening into it. To one opening adapt a copper tube bent twice at right angles; and to the other a straight tube dipping into the bottle. Having nearly filled the bottle with pieces of charcoal (recently heated to redness), and having screwed on the bent and straight tubes, place the bottle in a furnace, closing the mouth of the latter tube with a stone or clay cover in two pieces, hollowed in the centre so as to fit the upper part of the bottle, and defend it from the action of the fire. Connect the curved tube with a WOOLFE'S bottle half filled with water, and placed in a freezing mixture; and when the iron bottle is sufficiently heated, introduce by the straight tube, fragments of sulphur, and immediately close the mouth of the tube with a plug. The bisulphuret, as it comes over, falls to the bottom of the water. Separate it from the water and distil over dry chloride of calcium, or it may be purified by shaking up with mercury. It is said the offensive odour of the bisulphide may be removed by agitation with copper filings. See also WAGNER'S *Chemical Technology*.

BLACKING, LIQUID, FOR SHOES, &c. [*Note.*—By ivory-black, bone-black, which is usually sold under this name, is intended. True ivory-black has a more intense colour, but is too dear for general use.] 1. Ivory-black, 3 oz., treacle 2 oz., sweet oil $\frac{1}{2}$ oz.: mix to form a paste; add gradually $\frac{1}{4}$ oz. of oil of vitriol, and then half a pint of vinegar, and $1\frac{1}{2}$ pints of water, or sour beer. Some prefer mixing the oil of vitriol with sweet oil.

2. Ivory black 2 lbs., treacle 2 lbs., sweet oil $\frac{1}{2}$ lb.; mix and add $\frac{1}{2}$ lb. oil of vitriol, and beer or vinegar to make up a gallon.

3. Ivory-black 3 lbs., treacle 4 lbs., vinegar, a pint, oil of vitriol 8 oz., water a gallon.

4. Ivory-black 2 lbs., neat's-foot oil 4 oz.; mix, and add

3 quarts of sour beer, or vinegar, and a spoonful of any kind of spirits; stir till smooth, and add 2 oz. of oil of vitriol, and sprinkle on it $\frac{1}{2}$ dr. of powdered resin. Then boil together 3 pints of sour ale with a little log-wood, and $\frac{1}{4}$ oz. of Prussian blue, 3 oz. of honey, and 8 oz. of treacle. Mix, but do not bottle it for 2 or 3 days.

5. Ivory-black 8 oz., brown sugar or treacle 8 oz., sweet oil 1 oz., oil of vitriol $\frac{1}{2}$ oz., vinegar 2 quarts. Mix the oil with the treacle, then add the oil of vitriol and vinegar, and lastly, the ivory-black.

BLACKING FOR DRESS BOOTS. 1. Gum 8 oz., treacle 2 oz., ink a pint, vinegar 2 oz., spirit of wine 2 oz. Dissolve the gum and treacle in the ink and vinegar, strain, and add the spirit.

2. To the above add 1 oz. of sweet oil, and $\frac{1}{4}$ oz. lamp-black. [These are applied with a sponge, and allowed to dry out of the dust. They will not bear the wet.]

3. Beat together the whites of two eggs, a tablespoonful of spirit of wine, a lump of sugar, and a little finely powdered ivory-black to thicken.

BLACKING WITHOUT POLISHING. Treacle 4 oz., lamp-black $\frac{1}{2}$ oz., yeast a tablespoonful, 2 eggs, olive oil a teaspoonful, oil of turpentine a teaspoonful. Mix well. To be applied with a sponge, without brushing.

BLACKING, INDIA RUBBER (PATENT). Ivory-black 60 lbs., treacle 45 lbs., vinegar (No. 24) 20 gallons, powdered gum 1 lb., India-rubber oil 9 lbs. (The latter is made by dissolving by heat 18 oz. of India rubber in 9 lbs. of rape oil.) Grind the whole smooth in a paint mill, then add, by small quantities at a time, 12 lbs. of oil of vitriol, stirring it strongly for half an hour a day for a fortnight.

BLACKING, PASTE. 1. These pastes may be made with the ingredients of liquid blacking, using sufficient vinegar, in which a little gum has been dissolved, to form a paste. Make it into cakes, and dry it.

2. *German Blacking.* Powdered bone-black is mixed with half its weight of molasses and one eighth of its weight of olive oil; and to this is added afterwards one eighth of its weight of hydrochloric acid, and one fourth of its weight of strong sulphuric acid. The whole is to be then mixed up with water into a sort of unctuous paste.—LIEBIG.

3. BAILEY'S *Blacking Balls*. Bruised gum tragacanth 1 oz., water 4 oz.; mix, and add 2 oz. of neat's-foot oil, 2 oz. of fine ivory-black, 2 oz. of Prussian blue, 4 oz. of sugar-candy; mix, and evaporate to a proper consistence.

For HEEL BALL, see BALLS, further back.

BLACKING FOR HARNESS. 1. Isinglass or gelatine $\frac{1}{4}$ oz., powdered indigo $\frac{1}{4}$ oz., soft soap 4 oz., logwood 4 oz., glue 5 oz. Boil together in 2 pints of vinegar till the glue is dissolved; then strain through a cloth, and bottle for use. This appears an unchemical composition, but is inserted like many similar ones, because it is in actual use. The next is of a different character.

2. Melt 8 oz. of beeswax in an earthen pipkin, and stir into it 2 oz. of ivory-black, 1 oz. of Prussian blue ground in oil, 1 oz. of oil of turpentine, and $\frac{1}{4}$ oz. of copal varnish. Make it into balls. To be applied with a brush and polished with an old handkerchief.

3. Treacle $\frac{1}{2}$ lb., lamp-black 1 oz., yeast a spoonful, sugar-candy, olive oil, gum tragacanth, isinglass, each 1 oz., an ox gall. Mix all together with 2 pints of stale beer, and let it stand before the fire for an hour.

BLACK REVIVER. 1. Bruised nutgalls 1 lb., logwood 1 lb., water 5 quarts; boil to 4 quarts, and add sulphate of iron 4 oz., dissolve, and strain. When cold, add 8 oz. of ox-gall.

2. Galls 3 oz., logwood 1 oz., copperas, iron filings, and sumach, of each 1 oz., vinegar 2 pints.—GRAY'S SUPPLEMENT.

BLACK JAPAN. True asphaltum $1\frac{1}{2}$ lbs., boiled linseed oil 4 pints, burnt umber 4 oz. Heat together till the whole is incorporated, remove from the fire, and when sufficiently cool, add as much oil of turpentine as will bring it to a proper consistence.

BLEACHING LIQUID. Solutions of chloride of lime, and chloride of soda, are sold for this purpose, with directions for use. The following is also used: Mix 3 lbs. of common salt and 1 lb. of black oxide of manganese with as much water as will form a paste. Put the mixture into a retort, and add 2 lbs. of oil of vitriol previously diluted with 4 lbs. of water. Pass the gas into a solution of 1 lb. of common pearlash, or 11 oz. of caustic potash, in 6 lbs. of water. The retort may be placed, after a short period, in hot water, to extricate the remaining gas. In bleaching

cotton by chloride of lime, 1 lb. is dissolved in 3 gallons of water for each pound of cloth; it is afterwards passed through diluted hydrochloric or sulphuric acid (1 part of acid to 30 of water), and then washed.

BLIGHTS, REMEDIES FOR.

Apple-tree canker. Having brushed off the white down and the red stain underneath it, anoint the places with a mixture of train oil and Scotch snuff.

White Blight of Apple-tree. Apply a decoction of foxglove mixed with fresh cow-dung into a paste.

Fly in Turnips. Steep the seed before sowing in train oil; or distribute slaked lime over the field as soon as the plants have appeared; or contrive, if possible, to fumigate the field with brimstone.

Mildew of Wheat. 1. *To prevent it.* Dissolve in $3\frac{1}{2}$ gallons of cold water $3\frac{1}{2}$ oz. of sulphate of copper for every 3 bushels of sowing grain. Throw this quantity of wheat into another vessel, and pour over it the prepared liquid, until it rises 5 or 6 inches above the corn. Stir thoroughly, and remove all the grains that swim. Throw the mixture into a basket, so as to drain off the liquid; wash it well in soft water, and dry before sowing.

2. *To remove it.* A solution of 1 lb. of salt to the gallon of water, sprinkled with a flat brush over the growing corn.

Smut in Wheat. *To prevent it.* Boil 3 gallons of water, and slake in it about 36 lbs. of quicklime; add 3 gallons more of cold water, and pour the hot mixture on $4\frac{1}{2}$ bushels of the grain placed in a tub, stirring incessantly. Turn over the mixture now and then for 24 hours. Allow the liquid to drain off, and sow the limed wheat as soon as it is sufficiently dry. See WHEAT, STEEP FOR.

Blight in Vines. A solution of pentasulphuret of calcium may be painted over the branches and twigs until they acquire a continuous coating of sulphur mixed with carbonate of lime. *M. Merceuil* raises the earth round the stems of the vine to about a foot deep, then places in each excavation a good handful of flowers of sulphur, heaping it round the stems. The earth is then replaced. This is done at the beginning of August.

For Lice, Aphides, and Red Spiders. See WASHES FOR VERMIN IN PLANTS.

BLUE FOR LINEN. The ordinary kinds of cake blue consist

of indigo and starch. LIEFCHILD'S patent blue is thus made: Mix 4 parts of Chinese blue, 1 of TURNBULL'S, and 1 of oxalic acid; gradually add boiling water until the whole is dissolved, and lastly 4 parts of sulphate of indigo. The latter is made with 1 part of indigo and 4 of sulphuric acid, neutralised with carbonate of ammonia.

BOILERS, TO PREVENT INCRUSTATIONS IN. M. DUCLOS proposes to mix the water with which the boiler is to be supplied with the following solution, in the proportion of $1\frac{1}{4}$ per cent. Chloride of barium 125 parts, hydrochloric acid (sp. gr. 1.20) 25 parts, water 450. The water must then be allowed to clear in large reservoirs, where both the sulphate and carbonate of lime become decomposed, with formation of chloride of calcium. Any excess of acid must be neutralized before using, which may be done by placing pieces of limestone in the pipes.

BONES, SULPHATED. To a bushel of ground bones, add from 10 to 14 lbs. of oil of vitriol, previously mixed with half its bulk of water. This is sometimes mixed with an equal weight of salt and a sufficient quantity of bran. Turnip seed may be mixed with this compound and the two sown together.

BOROGLYCERIDE, an antiseptic preparation patented by PROFESSOR BARFF, more especially designed for the preservation of food, but recommended also as a dressing after surgical operations.

Boroglyceride is prepared by heating together 62 parts of boracic acid and 92 of glycerin, at a tolerably high temperature until combination is complete.

BOOT-TOP LIQUID. 1. Solution of chloride of tin 3 drs.; French chalk, or Venetian talc, in powder, 1 oz.; salt of sorrel $\frac{1}{2}$ oz., flake white 1 oz., burnt alum $\frac{1}{2}$ oz., powdered cuttle-fish bone 1 oz., white arsenic 1 oz., boiling water a quart. Probably sulphate of baryta might be substituted for arsenic, the use of which it is desirable to discourage.

2. Sour milk 3 pints, cream of tartar 2 oz., oxalic acid 1 oz., alum 1 oz.—REDWOOD.

3. Wash the tops with soap and water, and scrape them with the back of a knife. Then apply the following with a harefoot brush.—Oxalic acid 1 oz., water a pint. Use the back of a knife as before; then polish with the following: Powdered gum arabic $\frac{1}{2}$ oz., red spirits of lavender 2 oz., powdered turmeric $\frac{1}{2}$ oz.; pencil this over

the top, let it half dry, then polish by rubbing it, one way only, with a flannel till it shines.

4. Sour milk 3 pints, butter of antimony 2 oz., cream of tartar 2 oz., citric acid, alum, burnt alum, of each 1 oz.—GRAY'S SUPPLEMENT.

5. *White Top*. One ounce each of magnesia, alum, cream of tartar, and oxalic acid; $\frac{1}{2}$ oz. of salt of sorrel, and $\frac{1}{2}$ oz. of sugar of lead; dissolve in a quart of water, and apply with a sponge.

6. *Brown Top*. Oxalic acid, alum, annotto, of each 1 oz.; isinglass $\frac{1}{2}$ oz., sugar of lead $\frac{1}{2}$ oz., salt of sorrel $\frac{1}{2}$ oz.; boil together in a quart of water for 10 minutes. Apply with a sponge.

BOOKBINDERS' STAINS, FOR LEATHER.—*Black*. A solution of 1 part of sulphate of iron in 6 of water. *Blue*. A solution of indigo. (See CHEMIC BLUE.) *Brown*. A solution of pearlash, or of common soda.

BOOT POWDER. Finely powdered French chalk, or Venetian talc.

BREAD, UNFERMENTED. See DIETETIC ARTICLES, further back.

BRONZE POWDER. The best methods of preparing these powders are probably kept secret. The following are some of the published recipes;

1. Gold leaf, or alloys of gold, reduced to powder by grinding them with sulphate of potash, or with honey, and washing away the extraneous matter with hot water, and drying the metallic powder.

2. Dutch metal, and other similar alloys, treated in the same way.

3. Verdigris 4 oz., tutty 2 oz., sublimate 1 dr., borax 1 dr., nitre 1 dr., mix them into a paste with oil and fuse the mixture in a crucible. This has failed in some hands—perhaps from the tutty being factitious.

4. Mix together 100 parts of sulphate of copper, and 50 of crystallized carbonate of soda; apply heat till they unite. Powder the mass when cold, and add 15 parts of copper filings; mix well and keep it at a white heat for 20 minutes. Wash and dry the product.

See also AURUM MUSIVUM, and ARGENTUM MUSIVUM, further back.

BRONZING LIQUIDS, FOR BRONZING COPPER MEDALS, FIGURES, INSTRUMENTS, &c. For copper medals, electro-

type casts, &c. 1. Sal ammoniac 1 dr., oxalic acid 15 grs., vinegar a pint; after well cleaning the article to be bronzed, warm it gently, and brush it over with the liquid, using only a small quantity at a time. When rubbed dry, repeat the application till the desired tint is obtained.

2. *Bronze for Plaster Figures.* Dissolve palm soap in water, and add a mixed solution of sulphate of copper and sulphate of iron until no further precipitate occurs. Dry the precipitate, and mix it with oil of turpentine, or linseed oil. Sulphate of copper alone produces too bright a green. Palmitate of iron is yellow. These may be precipitated separately, and mixed to the desired shade after being triturated with the oil. Ten ounces of soap will require 3 ounces of sulphate of copper.

3. Sal ammoniac 1 oz., cream of tartar 3 oz., salt 6 oz.; dissolve in a pint of hot water, add 2 oz. of nitre and 2 oz. of nitrate of copper dissolved in $\frac{1}{2}$ pint of water.

4. *To give an antique appearance to bronze figures, &c.* Salt of sorrel 1 oz., sal ammoniac 2 oz., white vinegar 14 oz.

5. *For copper binding screws and other small articles.* A diluted solution of perchloride of platinum.

6. *For electrotype medals.* A weak solution of hydro-sulphuret (hydro-sulphide) of ammonia, or of sulphuret (sulphide) of potassium.

7. Immediately on removing the electrotype cast from the solution, brush it over with good black lead; then heat it moderately, and brush it over with a painting brush, the slightest moisture being used.

8. *For electrotype copper medals.* Boil 2 oz. of carbonate of ammonia and 1 oz. of acetate of copper in $\frac{1}{2}$ pint of vinegar, until nearly all the vinegar is evaporated. Pour into this a solution of 62 grains of sal ammoniac, and $15\frac{1}{2}$ grains of oxalic acid, dissolved in $\frac{1}{2}$ pint of vinegar; boil the whole and filter. Apply it to the medal (which should be perfectly bright, and previously warmed) with a camel-hair pencil for half a minute; then pour boiling water on it; wipe it with soft cotton very slightly moistened with linseed oil, and rub it with clean cotton. The medals may also be bronzed by applying oxide of iron (jeweller's rouge, or crocus) in the same manner as directed above, for plumbago; or a mixture of these may be used.

9. *Tin Castings.* Wash them over, after being well

cleaned and wiped, with a solution of 1 part of sulphate of iron and 1 of sulphate of copper, in 20 parts of water : afterwards with a solution of $\frac{1}{2}$ parts of verdigris in 11 of distilled vinegar ; leave for an hour to dry, then polish with a soft brush and colcothar.

BRONZING BALL. See BALLS (HEEL).

BRONZING PASTES, PARISIAN. 1. Plumbago 1 oz., sienna 2 oz., rouge $\frac{1}{2}$ oz. Add a few drops of hydrosulphate of ammonia and water.

2. Chrome lead 2 oz., Prussian blue 2 oz., plumbago $\frac{1}{2}$ lb., sienna powder and lake carmine, each $\frac{1}{4}$ lb. Add sufficient water to make a paste.

BROSSE DE CORAIL. The roots of lucerne (*Medicago sativa*) cleaned, dried, and hammered at the end. Used as a tooth-brush.

BROWNING, OR BRONZING LIQUIDS, FOR GUN BARRELS. 1. Aquafortis $\frac{1}{2}$ oz., sweet spirit of nitre $\frac{1}{2}$ oz., spirit of wine 1 oz., sulphate of copper 2 oz., water 30 oz., tincture of perchloride of iron 1 oz. ; mix.

2. Sulphate of copper 1 oz., sweet spirit of nitre 1 oz., water a pint ; mix. In a few days it will be fit for use.

3. Sweet spirit of nitre 3 oz., gum benzoin $1\frac{1}{2}$ oz., tincture of perchloride of iron $\frac{1}{2}$ oz., sulphate of copper 2 drs., spirit of wine $\frac{1}{2}$ oz., mix, and add 2 lbs. of soft water.

4. Tincture of perchloride of iron $\frac{1}{2}$ oz., spirit of nitric ether $\frac{1}{2}$ oz., sulphate of copper 2 scruples, rain-water $\frac{1}{2}$ pint.

The above are applied with a sponge, after cleaning the barrel with lime and water. When dry, they are polished with a stiff brush or iron scratch brush.

BRUNSWICK BLACK. Melt asphaltum, and add to it half its weight of boiled linseed oil ; mix, and when sufficiently cool, add enough oil of turpentine to bring it to the proper consistence.

BUG POISON. 1. Spirit of wine 1 oz., spirit of turpentine 8 oz., camphor $\frac{1}{2}$ oz. : mix.

2. Distilled vinegar, or diluted wood vinegar, a pint ; camphor $\frac{1}{2}$ oz. ; dissolve.

3. Corrosive sublimate 3 oz., hydrochloric acid 3 oz., oil of turpentine 12 oz., water 6 pints. Or, 1 oz. of sublimate, 2 oz. of hydrochloric acid, a pint of oil of turpentine, and a pint of decoction of tobacco.

4. Strong mercurial ointment 1 oz., soft soap 1 oz., oil of turpentine a pint.

5. White arsenic 2 oz., lard 13 oz., corrosive sublimate $\frac{1}{4}$ oz., Venetian red $\frac{1}{4}$ oz.

6. Scotch snuff mixed with soft soap.

7. *For Floors.* Corrosive sublimate 1 lb., sal ammoniac 1 lb., hot water 8 gallons.

It is said that if a branch of narrow-leaved dittany or pepperwort (*lepidium ruderale*) be suspended in a room, all the bugs will settle in it, and may be taken. Fumigating the rooms with sulphur is a troublesome and disagreeable process, and not always successful.

BURNETT'S (Sir W.) DISINFECTING FLUID. See DISINFECTING AND DEODORIZING COMPOUNDS.

BUTTER, TO PRESERVE. Powder finely, and mix together, 2 parts of the best salt, 1 of loaf sugar, and 1 of nitre. To each pound of butter, well cleansed from the milk, add 1 oz. of this compound. It should not be used under a month. [Butter that has an unpleasant flavour is said to be improved by the addition of $2\frac{1}{2}$ drs. of bicarbonate of soda to 3 lbs. of butter. A turnipy flavour may be prevented by only feeding the cows with turnips immediately after milking them.]

BUTTER OF ANTIMONY. The liquid chloride of antimony, commercially known by this name, is usually made by dissolving crude or roasted black antimony in hydrochloric acid with the addition of a little nitric acid. It usually contains pernitrate of iron.

BUTYRIC ETHER. Saponify butter with a strong solution of potash, dissolve the soap in the smallest quantity of alcohol by the aid of heat, add a mixture of alcohol and sulphuric acid till the solution is acid, and distil as long as the product has a fruity odour. Redistil the product from chloride of calcium. It is sold as essence of pineapple.

CAMPHINE. *Highly rectified oil of turpentine.* ENGLISH'S patent camphine is made by passing the vapour of oil of turpentine through caustic solutions of potash, soda, or lime; or through sulphuric acid.

CAMPHOR, ARTIFICIAL. This is formed by passing hydrochloric acid gas into oil of turpentine.

CAMPHOR BALLS. See SKIN COSMETICS, after PERFUMERY.

CANDIES. These belong rather to the confectioner than to

the druggist. The green stalks of angelica, the peels of orange, lemon, and citron, green roots of ginger, &c., are first boiled in water till soft, then in syrup till they are transparent, and dried in a stove, at a heat not exceeding 104° F. Candied horehound is made by boiling lump sugar with a little strong decoction or infusion of dried horehound, till a portion taken out and cooled becomes solid. It is then poured on to a slab, or into paper or tin moulds dusted with powdered sugar.

CANTHARIDINE BLISTER. LISSONDE. Melt white wax 45 parts in olive oil 28 parts with a gentle heat, add turpentine 24 parts, camphor 1 part, cantharidin 2 parts; stir, let the mass cool an instant, then pour out or spread on cloths. It raises the epidermis in a few hours.

CANTON'S PHOSPHORUS. Place calcined oyster shells in layers alternately with sulphur, and heat strongly in a covered crucible for an hour.

CAOUTCHOUC, SOLVENTS FOR. See SOLVENTS.

CAPSULES, GELATINOUS. These are used to contain copaiva and other nauseous liquids which do not dissolve gelatin, so that the liquids may be swallowed without inconvenience. They are made by "dipping the bulbous extremity of an iron rod into a concentrated solution of gelatine. Just before the rod is withdrawn, it is to be rotated, in order to diffuse the gelatine equally over the bulb." When sufficiently hardened, they are removed, placed on pins to dry, and when dry, filled with the balsam or oil, and the orifice closed with liquid gelatin. They are usually of an olive form, and contain 10 grs. of balsam in each. See Dr PEREIRA'S 'Elements,' article COPAIVA. M. GIRAUD recommends the following composition for capsules: Transparent gelatin 12 parts, syrup of gum 2 parts, syrup 2 parts, water 10 parts. Melt in a warm bath, remove the scum, and dip the mould, previously oiled, into the compound.

CARBON. See CHARCOAL.

CARBONIC ACID. See GASES.

CARMINE. See PIGMENTS.

CASE-HARDENING POWDER. This is ferroproussiate of potash (ferrocyanide of potassium) dried, and finely powdered. By sprinkling it on iron heated to bright redness, the metal becomes case-hardened, or superficially converted into steel. The iron should be plunged into cold water as

soon as the powder has acted on it. The following compound is used for the same purpose : Sal ammoniac 2 oz., burnt bone dust 2 oz., HENWOOD'S composition $\frac{1}{2}$ oz. Used as the former.

CAYENNE, SOLUBLE. See CONDIMENTS, further back.

CEMENTS AND LUTES, VARIOUS.

Shell-lac Cement. Fine orange shell-lac, bruised, 4 oz., highly rectified spirit 3 oz. Digest in a warm place, frequently shaking, till the shell-lac is dissolved. Methylated spirit may be substituted for spirit of wine, where the smell is not objectionable. A most useful cement for securely joining almost any material. See GLUE, LIQUID.

Shell-lac Cement without Spirit. Boil 1 oz. of borax in 16 oz. of water, add 2 oz. powdered shell-lac, and boil in a covered vessel till the lac is dissolved. Cheaper than the above, and for many purposes answers very well. Both are useful in fixing paper labels to tin, and to glass when exposed to damp.

Armenian Cement for Glass, China, &c. 1. KELLER'S. Soak 2 drs. of cut isinglass in 2 oz. of water for 24 hours; boil to 1 oz., add 1 oz. methylated spirit of wine, and strain through linen. Mix this, while hot, with a solution of 1 dr. of mastic in 1 oz. of methylated spirit, and triturate with $\frac{1}{2}$ dr. powdered gum ammoniac, till perfectly homogeneous.

2. Dr URE'S *Diamond Cement.* Isinglass 1 oz., distilled water 6 oz., boil to 3 oz., and add $1\frac{1}{2}$ oz. of methylated spirit. Boil for a minute or two, strain, and add while hot, first $\frac{1}{2}$ oz. of a milky emulsion of ammoniac, and then 5 drs. of tincture of mastic.

Cement used in the East for joining jewels, glass, and metals.—Mr ETON. Dissolve 5 or 6 pieces of gum mastic, each about the size of a large pea, in just as much methylated spirit as will render it liquid. Soften some isinglass by steeping it in water; having dried it, dissolve as much of it in good brandy as will make a two-ounce phial of strong glue, to which must be added two small bits of gum ammoniacum, rubbing until they are dissolved. Mix the two solutions; keep in a close phial; and when it is to be used, set the phial in boiling water.

Liquid Cement, for glass, porcelain, wood, &c. Macerate for several hours 6 parts of glue in small pieces, in 16 parts of water, add 1 part hydrochloric acid and $1\frac{1}{2}$ parts of sul-

- phate of zinc, and expose the mixture for several hours to a temperature of 150° F.
- HENLE's Cement, for Glass or Earthenware.** Shell-lac 2 parts, Venice turpentine 1 part, fuse together, and form into sticks.
- Cheese Cement, for Earthenware, &c.** Mix together:—white of egg beaten to a froth, quicklime, and grated cheese, and beat them to a paste. See **GLUE, CASEIN.**
- Curd Cement.** Add $\frac{1}{2}$ pint of vinegar to $\frac{1}{2}$ pint of skimmed milk; separate the curd and mix it with the whites of 5 eggs well beaten, and sufficient powdered quicklime to form a paste. It resists water and a moderate degree of heat.
- Glass Flux, for mending broken China, &c.** Mix 3 parts of red lead, 2 of fine white sand, and 3 of crystallized boracic acid, fuse the mixture, levigate it, and apply it with thin mucilage of tragacanth. Heat the repaired article gently, so as partly to fuse the cement.
- Cement for joining Spar and Marble Ornaments, &c.** Melt together 8 parts of resin, 1 of wax, and stir in 4 parts, or as much as may be required, of Paris plaster. The pieces to be made hot.
- HENSLER'S Cement.** Grind 8 parts of litharge, 2 of recently burnt lime, and 1 of white bole, with linseed-oil varnish. Very tenacious, but takes a long time to dry.
- SINGER'S Cement for Electrical Machines and Galvanic Troughs.** Melt together 5 lbs. of resin, and 1 lb. of bees-wax, and stir in 1 lb. of red-ochre (highly dried, and still warm), and 4 oz. of Paris plaster, continuing the heat a little above 212° F., and stirring constantly till all frothing ceases. Or (*for troughs*), resin 6 lbs., dried red-ochre 1 lb., calcined plaster of Paris $\frac{1}{2}$ lb., linseed oil $\frac{1}{2}$ lb.
- Mucilage for Minerals.** The following is recommended for mending fossils and minerals. Starch 2 drs., white sugar 1 oz., gum arabic 2 drs., water q. s. Dissolve the gum, add the sugar, and boil until the starch is cooked.—*Druggists' Circular.*
- Cement for Wood, Porcelain, and Glass.** Dissolve 30 grs. of sulphate of aluminum in two thirds of an ounce of water, and add it to $8\frac{1}{2}$ fl. oz. of a strong solution of gum arabic.
- Botany Bay Cement.** Botany Bay gum, melted and mixed with an equal quantity of brickdust.
- Cap Cement.** As **SINGER'S**; but 1 lb. of dried Venetian

red may be substituted for the red ochre and Paris plaster.

Bottle Cement. Resin 15 parts, tallow 4 (or wax 3) parts, highly dried red ochre 6 parts, or ivory black q. s. The common kinds of sealing wax are also used.

Turner's Cement. Beeswax 1 oz., resin $\frac{1}{2}$ oz., pitch $\frac{1}{2}$ oz. Melt, and stir in fine brickdust q. s.

Coppersmiths' Cement. Powdered quicklime, mixed with bullocks' blood, and applied immediately.

Engineers' Cement. Equal weights of red and white lead, with drying oil, spread on tow, or canvas. For metallic joints, or to unite large stones, in cisterns, &c.

Cement for Steam Pipes. Good linseed oil varnish, ground with equal weights of white lead, oxide of manganese, and pipe-clay.

Iron Cement, for closing the joints of iron pipes. Iron borings, coarsely powdered, 5 lbs., powdered sal ammoniac 2 oz., sulphur 1 oz., water sufficient to moisten it. It quickly hardens; but if time can be allowed, it sets more firmly without the sulphur. It must be used as soon as mixed, and rammed tightly into the joints.

GAD'S Hydraulic Cement. Powdered clay 3 lbs., oxide of iron 1 lb., boiled oil to form a stiff paste.

Cement for Masonry of Chambers of Chlorine, &c. Equal parts of pitch, resin, and plaster of Paris.

Roman Cement. A mixture of clay, lime, and oxide of iron, separately calcined, and finely powdered. It must be kept in closed vessels, and mixed with water when used.

Oxychloride of Zinc Cement. M. SOREL. In liquid chloride of zinc marking from 50° to 60° of Baumé's areometer, dissolve 3 per cent. of borax or sal ammoniac: add oxide of zinc which has been heated to redness, until the mass is of a proper consistence. The cement, when hard, becomes as firm as marble. It may be cast in moulds, like plaster of Paris, or used in the construction of mosaic works, &c.

Marine Cement. See GLUE, MARINE.

MAISSIATT'S Cement, as an airtight covering for bottles, &c.

Melt India-rubber (to which 15 per cent. of wax or tallow may be added), and gradually add finely powdered quicklime, till a change of odour shows that a combination has taken place, and a proper consistence is obtained.

Cement for attaching Metal Letters to Plate Glass. Copal

- varnish 15 parts, drying oil 5 parts, turpentine 3 parts, oil of turpentine 2 parts, liquefied glue 5 parts; melted in a water-bath, and 10 parts of slaked lime added.
- Cement to fasten Leather to Metal.* A hot solution of gelatin. With this wash the metal; and the leather, previously stepped in a hot infusion of gall-nuts, is to be pressed tightly on the surface and allowed to cool, when it will adhere very firmly.
- Cement for India-rubber.* Bisulphide of carbon 4 oz., best India-rubber 1 oz., isinglass 2 drs., gutta percha 1½ oz.—*Pharmaceutical Journal.*
- Cement to fasten India-rubber to Wood or Metal.* Dissolve 1 part powdered gum shellac in 10 parts of strong solution of ammonia. It is at first slimy, in 3 to 4 weeks becomes liquid; then hard and impermeable.—*Druggists' Circular.*
- Cement for fixing labels to Tin, or other Metallic Substances.* Mucilage of tragacanth 10 parts, honey 10 parts; dry wheaten flour 1 part.
- Japanese Cement.* Mix rice flour intimately with cold water, and boil gently.
- French Cement.* Mix thick mucilage of gum arabic with powdered starch.
- Common Paste.* To a dessert-spoonful of flour add gradually half a pint of cold water, and mix till quite smooth; add a pinch of powdered alum (some add also as much powdered resin as will lie on a sixpence), and boil for a few moments, stirring constantly. The addition of a little brown sugar, and a few grains of corrosive sublimate, is said to preserve it for years.
- Transparent Cement (LENHER).* Dissolve 5 parts of caoutchouc in 3 parts of chloroform, and add to the solution 1 part of mastic.
- Soft Cement.* Melt yellow wax with half its weight of common turpentine, and stir in a little Venetian red, previously well dried, and finely powdered. As a temporary stopping for joints and openings in glass and other apparatus, where the heat and pressure are not great. See the next.
- Lutes or Cements for closing the joints of Apparatus.* 1. Mix Paris plaster with water to a soft paste, and apply it immediately. It bears nearly a red heat. It may be rendered impervious by rubbing it over with wax and oil.

2. Slaked lime, made into a paste with white of egg, or a solution of gelatin.

3. *Fat Lute*. Finely powdered clay moistened with water, and beaten up with boiled linseed oil. Roll it into cylinders, and press it on the joints of the vessels, which must be perfectly dry. It is rendered more secure by binding it with strips of linen moistened with white of egg.

4. Linseed meal beaten to a paste with water.

5. Strips of moistened bladder, smeared with white of egg.

Luting for Acids. 1. Dissolve 1 part of India-rubber in 2 parts of linseed oil, by heat, and work into a stiff paste with 3 parts, or as much as sufficient, of white clay.

2. A concentrated solution of silicate of soda, made into a paste with powdered glass.

Lutes for Coating Retorts. 1. Dissolve 1 oz. of borax in $\frac{1}{2}$ pint of water, and add slaked lime to form a thin paste. Brush this over the retort, and let it dry gradually. Then apply a coating of slaked lime and linseed oil beaten together. Let it dry a day or two before use, and fill up any cracks which may appear with lime and linseed oil.

2. *For bearing a stronger heat*: Stourbridge clay, mixed with a little sand to prevent it splitting off, a little cut tow, or horse-dung, or asbestos, is usually added; to increase its coherence. It should be beaten to a stiff paste, and rolled out before application. The glass should be first rubbed over with a little of the lute mixed with water, then placed in the centre of the paste, rolled out to about $\frac{1}{4}$ or 1-3rd of an inch in thickness, and the edges of the latter raised and moulded to the glass, taking care to press out all the air.

MOHR'S Lute. Mix equal parts of brickdust and litharge, and beat them into a paste with linseed oil. Apply this with a stiff brush, and dust it over with coarse sand: dry it in a warm place.

For Cements for plugging teeth, see **TEETH AND MOUTH COSMETICS**, further back.

CHARCOAL. Wood charcoal is made by burning wood with only a partial access of air. For chemical purposes, that made in iron cylinders, in the manufacture of wood vinegar, is preferred.

CHARCOAL, ANIMAL. The most common form is that of

bone black (commonly called *ivory black*), made by distilling bones (from which the grease has been removed by boiling) in iron or earthen retorts, the ammoniacal liquor, &c., being collected in proper receivers. The residue is bone black. When used for the purpose of decolorization, further treatment is required: either the bone earth may be entirely removed by hydrochloric acid, as directed in the *Pharmacopœia*; or more commonly, in the refining of sugar, and for other manufacturing purposes, this is only partially effected in some such way as the following: Mix 8 lbs. of the bone black, coarsely powdered, with sufficient water to form a paste, and add 1 lb. of hydrochloric acid. In an hour pour boiling water on the mass, let it settle, pour off the liquid, add more water, and repeat this till the water comes off free from taste. Drain and press the black in a cloth, and dry it. Its power is increased by mixing it with a little potash, heating it to dull redness in a covered crucible, and again washing it.

CHARCOAL, ALUMINIZED. This was recommended by Dr STENHOUSE as a cheap and very efficient decolorizing agent. Dissolve in water 54 parts of the sulphate of alumina of commerce, and mix with 92½ parts of finely powdered wood charcoal. When the charcoal is saturated, evaporate to dryness, and heat to redness in covered Hessian crucibles till the water and acid are dissipated. The charcoal contains 7½ per cent. of anhydrous alumina.

CHARCOAL FROM COAL-TAR. Heat gently in an iron pot 1 lb. by weight of coal-tar pitch till it melts. Add 2 lbs. of fluid coal-tar, and mix. Stir in 7 lbs. of hydrate of lime in very fine powder. The thick mass is now roasted, stirring all the time, till it is reduced to a fine powder. It is then ignited in a covered crucible till all the vegetable matter is carbonized. The charcoal when cold is digested with dilute hydrochloric acid, and finally washed with water in a filter, and dried. Dr STENHOUSE recommended this as an admirable form for decolorization. For such liquids as decoction of logwood it is four times as efficient as animal charcoal.

CHAMELEON, MINERAL. Mix equal weights of black oxide of manganese and pure potash, and heat them in a crucible. Keep the compound in closely stopped bottles. A solution of it in water passes through various shades of colour from green to red.

- CHEMIQUE, OR CHEMIC BLUE—SULPHATE OF INDIGO.** To 7 or 8 parts of oil of vitriol, in a glass or earthen vessel, placed in cold water, add gradually 1 part of fine indigo in powder, stirring the mixture at each addition with a glass rod, or piece of tobacco-pipe. Cover the vessel for 24 hours, then dilute with an equal weight of water. Sometimes it is sold without diluting. The German (Nordhausen) fuming acid answers best, 4 or 5 parts of it being sufficient for 1 of indigo. For dyeing silk, &c., carbonate of potash, soda, or ammonia, is added, to neutralize the acid, taking care not to add it in excess.
- CHLORALUM.** Chloride of aluminum is made by acting on alumina, the basis of clay, with hydrochloric acid, and evaporating to form crystals. Under the above name it is extensively used as a disinfectant and deodorizer, decomposing sulphuretted hydrogen and coagulating albumen.
- CHLORATE OF BARIUM.** Saturate solution of chloric acid with freshly precipitated carbonate of barium, filter, and crystallize.
- CHLORATE OF POTASH. LIEBIG.** Dissolve chloride of lime in water, add solution of chloride of potassium, and boil to dryness. Dissolve the mass in hot water, and filter if necessary: on cooling, a large quantity of chlorate of potash is deposited. For another process, see **POTASSÆ CHLORAS**, *Pocket Formulary*.
- CHLORIDE OF LIME, OR CHLORINATED LIME. Bleaching Powder.** Chlorine gas (slowly evolved from a mixture of 10 parts of common salt and 10 to 14 parts of binoxide of manganese, placed in an alembic of lead, and heated by steam, and with 12 to 14 parts of oil of vitriol previously diluted with a fourth of its weight of water, added) is conveyed into a chamber where sifted slaked lime is thinly spread on shelves. It is so cheaply made by the large manufacturers for bleaching purposes that it is seldom prepared by druggists. The liquid chloride of lime may be made either by triturating the dry chloride with a little cold water till perfectly smooth, then adding more water, and filtering the solution, or by passing chlorine gas into a mixture of lime and water. The *Brit. Pharmacopœia* directs a solution of 1.055 sp. gr. The *Paris Codex* directs 1 part of dry bleaching powder to be diffused in 45 of water. **SOUBEIRAN** directs 1 part to 50; but prefers passing the gas from 1 part of binoxide of manganese and 4 of hydro-

chloric acid into a mixture of 1 part of lime and 50 of water. M. OPYL states that for the production of a good chloride of lime, the temperature of the chambers must be as low as possible. Winter is always found the most favorable season for its manufacture.

CHLORIDE OR HYPOCHLORITE OF POTASH. *Eau de Javelle*. See LIQ. POTASSÆ CHLORINATÆ, *Pocket Formulary*.

CHLORIDE OF SODA, OR CHLORINATED SODA. See SODA CHLORINATA, and LIQUOR SODÆ CHLORATÆ, in the *Pocket Formulary*. A more ready way of preparing it, for other purposes than for dispensing prescriptions, is the following: Diffuse 1 lb. of chloride of lime in 30 lbs. of water. Dissolve 2 lbs. of crystallized carbonate of soda in 15 lbs. of water. Mix the solutions, let the mixture settle, pour off the clear liquid, and filter it.

CHLORINE. See GASES.

CHLOROFORM. See CHLOROFORMUM, *Pocket Formulary*.

CHOCOLATE. See CHOCOLATA, *Pocket Formulary*.

CHROMATE OF POTASH. Mix 4 parts of chrome iron ore (chromate of iron) with 2 of pearlash and 1 of nitre, and heat the mixture in a reverberatory furnace for several hours. Lixivate and crystallize. The chromate is converted into bichromate by adding sulphuric acid, or rather acetic acid, to the solution.

CHROMATE OF LEAD. See PIGMENTS.

CHROMIC OXIDE. Prepared by igniting potassium bichromate at a red heat, and well washing and drying the residue.

CHROMIC ACID. See ACID, CHROMIC.

CLOTHES, POWDER TO KEEP AWAY MOTHS FROM. Mix powdered pipe-clay $1\frac{1}{2}$ lbs., white pepper and starch, of each 1 oz., root of Florentine iris $1\frac{1}{2}$ oz., with spirits of wine 2 oz. To be dusted over the clothes when laid by.

COCHINEAL COLOURING.—Mr PALMER. Macerate best carmine 1 oz. in strong solution of ammonia 6 oz. until it is dissolved. Heat gently to drive off excess of ammonia. Put into a quart wine bottle, and add rectified spirit 4 oz., and white sugar 3 lbs. Fill up with warm water, and shake until the sugar is dissolved.

COLLODION. See *Pocket Formulary* for its medical preparations, and PHOTOGRAPHY in this volume.

COLLODION, ELASTIC (*for surgical purposes*). See COLLODIUM FLEXILE, *Pocket Formulary*.

COLOURS, VARIOUS. The principal dry colours will be found under **PIGMENTS**. Other colours are noticed below.

COLOURS FOR DRUGGISTS' SHOW BOTTLES. In making these, distilled water should be used, and rather more of the colour than will fill the carboys made, to avoid the necessity of adding water to fill up after filtration, as this sometimes renders them turbid. The carboys should be perfectly clean and also dry, or otherwise rinsed out with a portion of the filtered liquid.

Amber. 1. Digest dragon's blood 1 part, with sulphuric acid 4 parts, and dilute the solution obtained to the required shade with distilled water.—*Pharmaceutical Journal.*

2. Deep chrome yellow 3 oz., pearlash 9 oz., water 2 gallons; boil gently for half an hour, take it off to cool, and add 6 oz. hydrochloric acid, and water to the desired colour.

Amethyst. Dissolve 5 grs. of salicylic acid in a little solution of ammonia, mixing this with 2 gallons or enough water to fill the show bottle. To this is to be added a few drops of solution of perchloride of iron, and afterwards a few drops of hydrochloric acid.

1. *Blues.* Sulphate of copper 4 to 8 oz., water a gallon, oil of vitriol 1 oz.

2. *Royal Blue.* Sulphate of copper 8 oz., water a gallon; dissolve, and add water of ammonia till the full colour is developed: then water to make up 2 gallons.

3. *Finest Royal Blue.* Nitrate of copper 3 oz., water sufficient to dissolve it; add water of ammonia as long as the colour becomes deeper, then water to make up 2 gallons.

4. *Paler.* Crystallized acetate of copper $\frac{1}{2}$ oz., hydrochloric acid $1\frac{1}{2}$ oz., water of ammonia q. s., water to make up 2 gallons.

5. *Light Blue.* Crystallized acetate of copper a scruple, water of ammonia 2 oz., water 2 gallons.

Green. 1. An infusion of saffron added to the above blues.

2. Sulphate of copper 4 oz., bichromate of potash $\frac{1}{2}$ dr., water q. s.

3. By adding to the above deep or light blues a small quantity of chromate or bichromate of potash till the

desired tint is produced, various shades of green may be obtained.

4. Sulphate of copper, with chloride of sodium or of iron.

5. *Emerald Green*. Nitrate of copper 3 oz., hydrochloric acid 4 oz., nitric acid 4 oz., water 2 gallons.

6. *Very Fine Emerald*. Nickel 3 oz., hydrochloric acid 4 oz., nitrous acid 2 oz.; digest for 24 hours, add 2 gallons of water and filter.

Red. 1. Dried rose petals 8 oz., boiling water a gallon; digest for 12 hours, strain; digest the roses with more water, and strain; mix the infusions, add a pint of diluted sulphuric acid, and filter.

2. Cochineal $\frac{1}{2}$ oz., boiling water a gallon; digest, strain, add $\frac{1}{2}$ oz. of sulphuric acid, and water to make up 2 gallons.

3. *Crimson*. Iodine, and iodide of potassium, of each 2 drs.; triturate with a drachm of water, and add 3 gallons of water, and 4 oz. of hydrochloric acid.

4. *For External Lamps*. Camphine, strongly coloured with alkanet root.

Pink. 1. Infuse $\frac{1}{2}$ oz. of good madder in a quart of boiling distilled water; when cold, add 1 oz. of strong ammonia, and filter into 2 gallons of distilled water.

2. Dissolve chloride or nitrate of cobalt in water, and add sufficient carbonate of ammonia to redissolve the precipitate first thrown down; then water q. s.

Purple. To the last, add sufficient of the blue No. 3 to give the desired shade.

Lilac. Smalts 4 oz., nitric acid 4 oz.; let it stand 24 hours, add 2 gallons of water, 1 oz. of alum, and 4 oz. of water of ammonia. Or rather as the purple.

Magenta. Dissolve acetate of rosaniline in a sufficient quantity of water to the required shade.

Yellow. 1. Chromate or bichromate of potash, with water q. s.

2. Bichromate of potash 2 drs., pure water 4 oz.; dissolve, and add 4 oz. sulphuric acid, and 2 gallons of water.

3. Bichromate of potash, nitrate of potash, and water.

Olive. Sulphate of iron 3 oz., sulphuric acid 3 oz., water 2 gallons; dissolve, and add the green No. 5, q. s. to brighten the colour,

LIQUID COLOURS FOR MAPS, &c. See **INKS, DYES, CHEMIQUE BLUE, LAKE LIQUOR, &c.** Gamboge and some of the cake colours, mixed with water, are also used.

COLOURS FOR CONFECTIONERS. Many fatal accidents occur from confectionery being coloured with poisonous pigments. The following may be safely used: Cochineal and its preparations, sap green, vegetable lakes, Prussian blue; a mixture of a yellow lake and Prussian blue for green.

COLOURING FOR BRANDY, &c. Sugar melted in a ladle till it is brown, and then dissolved in water or lime-water.

COLOURS FOR LIQUEURS. Pink is given by cochineal; yellow by saffron or safflower; violet by litmus; blue, by sulphate of indigo, saturated with chalk; green, by the last with tincture of saffron, or by sap green.

COLOURS FOR LEATHER. See **BOOKBINDERS' STAINS**, further back.

COLOURS, IMPROVED VEHICLES FOR. 1. One measure of saturated solution of borax, with 4 of linseed oil. The pigment may be ground with the oil, or the mixture.

2. A solution of shell-lac with borax, as in making **COATHURP'S WRITING FLUID**. See **INK**.

3. Water colours, mixed with gelatin, and afterwards fixed by washing with a solution of alum.

4. Curd of milk, washed and pressed, then dried on fine net, and when required for use, mixed with water and the colouring matter.

CONDY'S FLUID. (Patent.) The green fluid appears to contain the manganates, the red fluid the permanganates, of soda and potash. The latter is said to be double the strength of the *Liquor Potassæ Permanganatis*, B. P. Condy's "Ozonized Water" is a weaker solution of the permanganates, "for toilet purposes."

COPPER, OXIDE OF. The black oxide is made by calcining the nitrate; or by adding caustic potash to sulphate of copper, in solution, and washing and drying the precipitate. The red oxide may be made as directed for **Bronze Powder, No. 4**, or in the moist way, thus: Pour a solution of 27 parts of sugar in 60 of water, over 9 parts of hydrated oxide of copper, weighed in its compressed but still moist state. A solution of 18 parts of caustic potash in 60 of water is added, and the whole agitated

together without heat, and filtered. The clear liquid heated in a warm bath, and continually stirred, deposits the red oxide, and the liquid becomes colourless.

NITRATE OF COPPER. Dissolve copper in nitric acid to saturation, evaporate to dryness, redissolve, filter, and evaporate, so that the salt may crystallize. Or add a solution of sulphate of copper to a solution of nitrate of lead, so long as sulphate of lead is precipitated; filter, evaporate, and crystallize. For the other salts of copper, see *Pocket Formulary*.

COSMOLIN. See **VASELIN**, further on.

COTTON POWDER. See **GUN COTTON**.

CRAYONS FOR WRITING ON GLASS. Fuse in a cup 4 parts of spermaceti, 3 of tallow, and 2 of wax; stir in 6 parts of minium, and 1 of potash; keep warm for half an hour, and then pour into glass tubes of the thickness of a lead pencil. If cooled rapidly, the mass may be screwed up and down in the tube, and cut at the end to a fine point. The glass to be written on must be clean and dry,

CYANIDE OF POTASSIUM. See *Pocket Formulary*.

DEPILATORIES. See **HAIR COSMETICS**, further back.

DEXTRIN, OR STARCH GUM. Heat 4 gallons of water in a water-bath to between 77° and 86° Fahrenheit; stir in 1½ or 2 lbs. finely ground malt; raise the temperature to 140°, add 10 lbs. of potato or other starch: mix all thoroughly, raise the heat to 158°, and keep it between that and 167°, for 20 or 30 minutes. When the liquor becomes thin, instantly raise the heat to the boiling point, to prevent the formation of sugar. Strain the liquor, and evaporate it to dryness, as the dextrin will not keep long in a liquid form. Another method is to boil solution of starch with a few drops of sulphuric acid, to filter the solution, and to add alcohol to throw down the dextrin. See **GUM [BRITISH]** for another form of dextrin.

DEXTRIN VARNISH. BARON DE SYLVESTRE. Dextrin 2 parts, water 6 parts, rectified spirit 1 part.

DIASTASE. Macerate ground malt in cold water; strain with pressure; heat the clear solution in a water-bath to 158° Fahrenheit to coagulate the albumen; filter again, and add rectified spirit as long as diastase fails. If required very pure, redissolve it in water, and again precipitate with spirit. Dry it at a low temperature. Well-

malted barley contains about '002 per cent. of pure diastase; one part of which is capable of converting 2000 parts of starch into dextrin or sugar.

DIETETIC ARTICLES. For these see another division of the work.

DISINFECTING AND DEODORISING COMPOUNDS. 1. **SIR WM. BURNETT'S Patent Solution.** It is made by dissolving zinc in commercial hydrochloric acid to saturation.

2. **ELLERMAN'S Deodorizing Fluid.** This consists chiefly of perchlorides and chlorides of iron and manganese. In a report addressed to the Metropolitan Board of Works in 1859, Drs HOFFMAN and FRANKLAND stated that the perchloride of iron was the cheapest and most efficient deodorizer that could be applied to sewage: $\frac{1}{2}$ gallon deodorized 7500 gallons of sewage. 1 bushel of lime, or 3 lbs. of chloride of lime, would do the same.

3. **LEDOYEN'S Solution.** This is a solution of nitrate of lead, and contains about 20 oz. of the salt in a gallon. The specific gravity should be 1.40. A similar compound may be made by mixing $13\frac{1}{2}$ oz. of litharge with 6 pints of water and adding 12 oz. of nitric acid at 1.38 specific gravity (or 8 oz. at 1.50); and digesting at a gentle heat till the solution is complete.

4. **SIRET'S COMPOUND.** Sulphate of iron 20 lbs., sulphate of zinc $3\frac{1}{2}$ lbs., wood or peat charcoal 1 lb., sulphate of lime $26\frac{1}{2}$ lbs., mix and form into balls. To be placed in cess-pools, &c., to deodorise them. M. SIRET has subsequently modified this compound, thus: Sulphate of iron 100 parts, sulphate of zinc 50, tan or oak-bark powder 40, tar 5, and oil 5 parts.

5. **COLLINS' Disinfecting Powder.** Mix 2 parts of dry chloride of lime with 1 of burnt alum. To be set in shallow dishes in rooms, &c., with or without the addition of water.

6. **CARBOLIC ACID.** After infectious diseases, apartments may be purified by washing the woodwork floors, &c., with soft soap and water, to which carbolic acid has been added in the proportion of 1 pint of acid to 4 gallons of water. Rooms and houses may also be disinfected by sprinkling some of the acid on a hot iron plate, in the latter case so placed that the fumes may be made to ascend through the dwelling. See **CARBOLIC ACID, Pocket Formulary.**

7. CALVERT'S Powder. Carbolic acid (20 to 30 per cent.), alumina, and silica.

8. M'DOUGAL'S Powder. Carbolate of lime 33 parts, sulphate of magnesia 59 parts, water 8 parts. These powders can be sprinkled about a room, thrown on sewage or dissolved in water and so applied.

8. SANITAS. We believe the efficacy of this disinfectant depends upon its property of evolving peroxide of hydrogen and camphoric acid by the atmospheric oxidation of turpentine.

9. CONDY'S DISINFECTING FLUIDS. See back.

10. THYMOL. An excellent, but expensive disinfectant, non-poisonous, and non-irritant. See THYMOL, *Pocket Formulary*.

11. See also CHLORIDE OF LIME, CHLORIDE OF SODA, ACID SALICYLIC, CHLORALUM, further back. Peat charcoal also possesses powerfully deodorising properties.

FOOT POWDER. M. PAULCKE prepares a mixture of salicylic acid, soap, talc, and starch in the form of powder, to be applied to the feet, which, whilst rendering them firm, is said to induce an agreeable softness, and to remove all unpleasant smell arising from perspiration.

DUBBING, CURRIERS'. Made by boiling cuttings of sheepskins in common cod-oil.

DRYERS FOR PAINTERS. White copperas 1 lb., sugar of lead 1 lb., white lead 2 lbs.; ground with boiled oil.

DRYING AND BOILED OIL. Linseed oil is mixed with powdered litharge, and heated till it becomes thick. A pale drying oil is obtained by mixing with linseed oil sufficient dry sulphate of lead to form a milky liquid, and shaking it repeatedly for some days, letting it stand exposed to the light. When it has become quite clear, it may be poured off from the dregs. The sulphate of lead, when washed from the mucilage, may be again used for the same purpose. LIEBIG directs 1 lb. of acetate of lead to be dissolved in half a gallon of rain-water, and 1 lb. of finely powdered litharge added: the mixture is either boiled, or exposed for a longer time to a moderate heat, and frequently stirred, till no more particles of litharge can be seen. A white deposit is formed, which may be left in the liquid or separated by filtration; 20 lbs. of linseed oil, in which 1 lb. of levigated litharge has been diffused, are gradually added to the lead solution, pre-

viously diluted with an equal bulk of water, and the mixture frequently stirred. It is then left to clear itself in a warm place; but to obtain it bright it must be filtered through coarse paper or cotton. It may be bleached by exposure to the sun. The lead solution which subsides from the mixture may be filtered and used again, after dissolving it in 1 lb. of litharge as before. The oxide of lead contained in the oil may be removed from it by agitating it with diluted sulphuric acid, and letting it stand to settle. See **LINSEED OIL, REFINED.**

DUPUYTREN'S POMADE. See **HAIR COSMETICS.**

DYES AND COMPOUNDS USED IN DYEING. A few of the principal colouring matters and mordants may here be noticed; for further information, the reader is referred to Dr URE'S 'Dictionary of the Arts,' CALVERT'S 'Dyeing and Calico Printing,' CROOKE'S 'Handbook of Dyeing and Calico Printing,' and "Bleaching, Dyeing, and Calico Printing," 'Churchill's Technological Handbooks.'

Blue Dyes. The most important of these is indigo. Being insoluble in water, it is prepared for use by sulphate of iron, and alkalies, or lime, by fermentation and alkalies, and by solution in sulphuric acid.

1. Triturate 1 lb. of indigo with water and a little caustic potash; then add 3 lbs. of lime, and afterwards $2\frac{1}{2}$ lbs. of sulphate of iron in solution, stirring them well together. The solution contains refined indigo, which is soluble in lime and alkalies. The cotton, linen, &c., to be dyed is repeatedly dipped in the solution, and afterwards rinsed in water soured with hydrochloric acid.

2. To 45 or 50 gallons of water, heated to 122° F., add 12 oz. of indigo, 8 oz. of madder, 8 oz. of bran, and $2\frac{1}{4}$ oz. of potash. In 36 hours introduce 12 oz. more of potash, and the same in 12 hours after. In 72 hours, add a little lime to check the fermentation. Wool, silk, linen and cotton may be dyed in this bath. Another form of this dye is—Indigo 2 parts, common soda 5 parts, lime 2 parts, clarified honey 1 part, water as much as may be sufficient. Keep it warm in an earthen jar till the indigo is dissolved.

3. A solution of indigo in sulphuric acid (see **CHEMIC BLUE**) is used as a dye, but a purer tone of colour is obtained by the following method: The sulphate of indigo, mixed with water, is heated in a copper kettle; wool is

immersed in it, and the whole is allowed to cool for 24 hours. The wool is then taken out, washed till the water comes off colourless and free from acid; it is then boiled in water containing about 2 per cent. of pearlash, or other alkaline carbonate, for a quarter of an hour. The quantity of pearlash should be equal to one third the weight of the indigo.

Logwood, with verdigris, or sulphate of copper, gives a blue dye, bordering on violet: with alum and tartar, a violet.

Prussian Blue is sometimes used in dyeing, after being triturated and digested for 24 hours with its weight of hydrochloric acid. A blue is also given by immersing silk, &c., in a solution of peracetate of iron, then in a solution of prussiate of potash, and afterwards rinsing it in acidulated water. Boiling water is sufficient to discolour articles thus dyed.

Red Dyes. The various shades of red are given by madder, cochineal, lac dye, safflower, &c.; fixed by albuminous or tin mordants. Less permanent dyes are produced by Brazil wood, peach wood, and archil. Some of these require peculiar treatment. Safflower contains a yellow as well as a red colouring matter. The first, being soluble in cold water, is extracted by putting the safflower in a bag and kneading it under water. The safflower, thus deprived of the yellow matter, yields its red colour to alkaline liquids: at the time of using which, lemon juice or some other acid is added sufficient to saturate the alkali. Pink saucers are made by adding lemon juice to an alkaline infusion of washed safflower and allowing the colouring to deposit. Madder also contains a dun colouring matter which deteriorates the red unless previously removed. This may be partially effected by washing it in cold water: another mode is to treat the madder with its own weight of sulphuric acid, which carbonizes the other matters, but leaves the red colour uninjured. As madder gives out but little of its red colouring matter to water, the decoction is not strained off, but the madder left in the bath. With acetate of iron, madder yields a purple tint. Lac dye, as imported from India, requires acids for its solution. See LAC SPIRIT, further on.

Yellow Dyes. These are given by French berries, quercitron bark, turmeric, weld, yellow wood, &c. Also by

some mineral colours, as the following: The material to be dyed is first padded in a solution of bichromate of potash (8 oz. to a gallon of water), then in a solution of acetate or nitrate of lead. Cotton is dyed yellow by alternate dippings in iron liquor and lime water, or solution of pearlash. A yellow colour is given to silk by passing it through a mixture of equal measures of nitric acid (sp. gr. 1.288) and water, heated to 95° or 100° F., and from thence into a stream of water, or a mixture of chalk and water. This is termed *mandarining*.

Nankeen Dye is made by boiling annotto with an equal weight of pearlash in sufficient water. Orange is given by annotto; or by a mixture of red and yellow dyes; or by the successive application of acetate of alumina, a bath of quercitron, and the madder-bath.

Greens are given to woollens by first dyeing them blue, immersing the article in acetate of alumina, drying it, and finally immersing it in a quercitron-bath. For silks, the order is reversed.

Browns are given by catechu, by walnut-peels with alum, by red-wood and copperas, by madder and black dye, &c.

Drabs are given by fustic with iron liquor.

Blacks. These are given by salts of iron, with galls, sumach, and logwood. The best black cloths are previously dyed blue with indigo.

By the mixture of various dyes, every variety of shade is produced: and often several tints from one colouring matter by the use of different mordants.

The beautiful new dyes of the Aniline series, *Mauve*, *Magenta*, *Artificial Alizarine*, *Aniline Blue*, *Violet imperial*, *Aurine*, &c., obtained from coal-tar by various patented processes, are fully described, and formulæ given for their preparation in "Bleaching, Dyeing, and Calico Printing," vol. 2 of 'Churchill's Technological Handbooks.'

Mordants are earthy and metallic compounds, the bases of which unite both with the fibres of the material to be dyed and the colouring matter, thus rendering the dyes fixed. In calico printing, the mordants are formed into a paste with some gum or other thickening material, and printed with wooden blocks or by means of steam cylinders on to the cloth; which, after being dried, &c., is passed through the liquid dyes. The colouring matter combines

with the parts so printed, but is easily discharged from the other parts.

The principal mordants are the following:—

Alum Mordants. 1. Alum with one fourth its weight of tartar.

2. *Acetate of Alumina.* (See further back.) This is commonly prepared in solution for the purpose; 100 parts of alum in solution, with 150 parts of pyrolignite of lime of 20° B. density, is sometimes employed.

3. A solution of alum, with crystallized carbonate of soda, in the proportion of 1 oz. to each pound of alum.

4. **HAUSMANN'S.** This consists of a solution of alum with sufficient strong solution of caustic potash to redissolve the precipitated alumina; to which mixture a portion of linseed oil is added.

5. To 50 gallons of boiling water add 100 lbs. of alum; dissolve, and add slowly 10 lbs. of crystallized carbonate of soda. When the effervescence is over, add 75 lbs. of sugar of lead.

Tin Mordants. 1. *Protochloride of Tin.* To strong hydrochloric acid, add gradually small pieces of grain tin till no more is dissolved. It may be obtained in crystals by evaporation. In redissolving them, it is necessary to add to the water a few drops of hydrochloric acid.

2. *Nitro-muriate, or Perchloride of Tin.* Mix 1 measure of nitric acid with 4 of hydrochloric acid, and add tin in small quantities as long as any is dissolved. Or mix 4 oz. of hydrochloric with 1 of nitric acid and 1 of water; dissolve in it, by small portions at a time, 2 drs. of grain tin.

3. Aqua fortis (or equal parts of nitric acid and water) 8 parts, sal ammoniac 1 part; mix, and add gradually 1 part, or as much as it will dissolve, of grain tin.

4. Dr **BANCROFT'S** *Murio-sulphate of Tin.* Digest 2 parts of tin with 3 of strong hydrochloric acid for an hour. Add very cautiously 1½ part of oil of vitriol. Keep up the heat as long as hydrogen is evolved; on cooling, it crystallizes. Dissolve this salt in water, so as to form a solution containing 1 part of tin in 8.

5. *New Tin Crystals.* Add 3 lbs. of sal ammoniac to a gallon of solution of tin; evaporate, and crystallize.

6. *Mordant for Lac Dye.* Mix 27 lbs. of hydrochloric acid with 1½ lbs. of nitric acid (sp. gr. 1.19), put it into a

stone bottle, and add tin in small bits till 4 lbs. are dissolved.

7. *Stannate of Soda*. Digest litharge 36 parts, or minium 27 parts, in a metallic vessel, with a soda ley of 1.35 dens.; when dissolved, 8 parts of tin in grains are gradually added. The lead separates at once in a spongy state, and the solution of stannate of soda may be decanted.

Lac Spirit, used as a solvent for lac dye, in preference to hydrochloric acid alone, is thus made: Add gradually 3 lbs. of tin to 60 lbs. of hydrochloric acid. Digest $\frac{3}{4}$ lb. of this solvent in each pound of the dye for 6 hours. *Plum* or *puce spirit*, *peach spirit*, and *grain* or *scarlet spirit*, are names given by dyers to different solutions of tin employed in dyeing these colours. For scarlet, the nitro-hydrochloric solutions (Nos. 2 and 3, above) are used.

Iron Liquor. Scraps of iron are placed in casks or other vessels, and covered with commercial pyroligneous acid. There are usually a series of vessels, through which the solution is successively passed till it is fully saturated.

To produce good and permanent dyes, several successive processes are required, which cannot be described here. In domestic dyeing, for trifling articles, the dye and mordant are often boiled together, and the silk, &c., immersed in the liquid. The following are some examples: *Crimson*.—Boil 1 oz. Brazil-wood, with 1 dr. of alum and $\frac{1}{2}$ dr. of cream of tartar in a quart of water. *Purple*, or *Lilac*.—Archil 1 oz. (or cudbear 1 oz.), pearlsh 1 dr., hot water a quart. *Rose*, or *Flesh Colour*.—Pink saucers, with a little lemon-juice, will be found convenient. *Violet*.—Boil 4 oz. of logwood with 1 oz. alum and $\frac{1}{4}$ oz. of tartar in a quart of water. *Blue*.—Add to the water as much sulphate of indigo (chemic blue) as will give it the required colour. Or one of the other solutions of indigo (already described) may be used. *Yellow*.—Boil 2 oz. of turmeric, or 4 oz. quercitron, or a drachm of saffron, with $\frac{1}{2}$ oz. of alum, in a quart of water. *Green*.—Add to the yellow dye sufficient chemic blue to render it green. *Rose-red*, for silk.—Put your silk into a hot solution of alum and tartar; then into a hot decoction of madder. *Scarlet*, for silk.—Dye it yellow with saffron and alum; then put it into a decoction of cochineal and madder. *Black*.—Boil 1 oz. of logwood, $\frac{1}{4}$ oz. sumach, and 1 dr. of

copperas, in 4 pints of water. *Buff.*—Boil 1 oz. of fustic, 1 dr. of alum, in 4 pints of water.

EAU DE JAVELLE. See CHLORIDE OF POTASH.

EGGS, TO PRESERVE. JAYNE'S Liquid (expired patent) is thus made:—Take a bushel of lime, 2 lbs. of salt, $\frac{1}{2}$ lb. of cream of tartar, and water sufficient to form a solution strong enough to float an egg. In this liquid it is stated eggs may be preserved for two years. Eggs may be preserved for several months if rubbed over with linseed oil or poppy oil.

ELAINE. See OLEINE.

ELECTRIC TISSUE. Steep linen or cotton in a mixture of 1 part of strong sulphuric acid, and 3 of pure nitric acid, for an hour. Squeeze out the acid, wash with water until no sensible acidity remains, plunge it in a weak alkaline solution, then in water, and dry. By friction it yields a large quantity of resinous electricity.

ELECTRO-BRASSING. Mr WATT. Dissolve powdered acetate of copper, 5 oz., in $\frac{1}{2}$ gallon of water; add 1 pint of liquid ammonia; dissolve sulphate of zinc, 10 oz., in 1 gallon water, at 183° F., and when cool, add liquid ammonia 1 pint; dissolve potash, $4\frac{1}{2}$ lbs., in 1 gallon of water; lastly, cyanide of potassium, 8 oz., in 1 gallon of hot water. Now mix in the following order: add the copper solution to the zinc, and then the potash and cyanide, digest for an hour or so, add water to make up 8 gallons. Work with a brass *anode* and active battery power, adding occasionally a little more ammonia and cyanide.

ELECTRO-BRONZING BRUNEL'S Patent. Dissolve chloride of copper, 1 lb., in water $\frac{1}{2}$ gallon, then carbonate of potash, 25 lbs., in water 6 gallons; and sulphate of zinc, 2 lbs., in hot water $\frac{1}{2}$ gallon. Mix the solutions; add nitrate of ammonia, $12\frac{1}{2}$ lbs.; stir, and add water to 20 gallons. Work with an *anode* of brass and an active battery of two or more cells. Add occasionally some liquid ammonia and cyanide of potassium. These will prevent incrustations on the *anode*.

ELECTRO-COPPERING, for Iron or Zinc. The article must first be well cleaned, and "pickled" in dilute acid. Then dissolve sulphate of copper, 2 oz., in boiling rain-water; when cold add carbonate of potash, 4 oz., and liquid ammonia 2 oz. A precipitate forms, which is redissolved. Add cyanide of potassium (6 oz.) until all the blue colour

disappears. Make up to 1 gallon, and work with a two-cell battery. The surface obtained may be bronzed. (See back, BRONZING LIQUIDS.)

ELECTRO-GILDING. See GILDING, further on.

ELECTRO-SILVERING. See SILVERING, further on.

ELECTROTYPE MOULDS. These are sometimes made with fusible metals; sometimes with non-metallic compounds, having their surface covered with a conducting substance. The *fusible metal* is composed of 8 parts of bismuth, 3 of tin, and 5 of lead. The French *clichés* moulds consist of 8 parts of bismuth, 4 of tin, 5 of lead, and 1 of antimony. These are repeatedly melted together till perfectly mixed; and after being poured out on a suitable surface, are well stirred, and the metal forcibly pressed on the compound at the moment it is about to become solid. *Composition Moulds* are made with 8 oz. of spermaceti, 1½ oz. of white wax, and the same of strained mutton suet. These are melted together, and a little fine plumbago or flake-white stirred in. To copy plaster casts, the cast is set in a plate of hot water, with its face above the water, till it has absorbed water; it is then surrounded with a ring of cardboard, and the melted composition poured on it. The composition mould requires to be brushed over with finely powdered genuine black-lead.

ELECTROTYPE MOULDS. ELASTIC. *For copying medals when the figures are in high relief.* To 12 parts of carefully melted glue, add 3 parts of treacle, and incorporate them perfectly. *Gutta Percha Moulds* are made by softening a piece of gutta percha by warm water (150° to 160° F.) and pressing the metal into it by a screw. See SOLUTIONS.

ENGRAVINGS, PROCESS FOR CLEANING. FRANCIS. Put the engraving on a smooth board, and cover it thinly with common salt, finely powdered; pour and squeeze lemon-juice upon this salt, so as to dissolve a considerable portion of it. Now elevate one end of the board, that it may form an angle of about 45° with the horizon. Pour lastly on the engraving, boiling water from a tea-kettle, until the salt and lemon-juice are all washed off; the engraving will then appear perfectly clean and free from stains. It must be dried gradually, on the same board, or on some smooth surface.

ENGRAVING MIXTURE, FOR WRITING ON STEEL. Sulphate of copper 1 oz., sal ammoniac $\frac{1}{2}$ oz. Pulverize separately, adding a little vermilion to colour it, and mix with $1\frac{1}{2}$ oz. of vinegar. Rub the steel with soft soap, and write with a clean hard pen, without a slit, dipped in the mixture.

ESSENCE OF JARGONELLE PEAR. Acetate of amylene is sold under this name. It is made by distilling a mixture of 1 part of oil of grain, 2 of acetate of potash, and 1 of oil of vitriol. Wash the diluted liquid with alkaline water, agitate with dry chloride of calcium, and redistil it from litharge.

ESSENCE OF PINE APPLE. See BUTYRIC ETHER.

[For PERFUMED ESSENCES, see PERFUMERY. For CULINARY ESSENCES, see further back.]

ETCHING FLUIDS. CHEVALLIER. For *Lithography*. Dissolve 3 oz. of fused chloride of calcium in $9\frac{1}{2}$ oz. of water, add to the solution 2 oz. of gum arabic, and 1 oz. of pure hydrochloric acid.

For *Copper*. 1. Aqua fortis 2 oz., water 5 oz.; mix.

2. Iodine 2 parts, iodide of potassium 5 parts, water 5 to 8 parts.

3. *CALLOT'S Eau Forte for Fine Touches.* Dissolve 4 parts each of verdigris, alum, sea-salt, and sal ammoniac, in 8 parts of vinegar; add 16 parts of water, boil for a minute, and let it cool.

For *Steel*. 1. Iodine 1 oz., iron filings $\frac{1}{2}$ dr., water 4 oz.; digest till the iron is dissolved.

2. Pyroligneous acid 4 parts by measure, alcohol 1 part; mix, and add 1 part of double aqua fortis (sp. gr. 1.28). Apply it from $1\frac{1}{2}$ to 15 minutes.

3. Mix 10 parts of pure hydrochloric acid, 70 of distilled water, and a solution of 2 parts of chlorate of potash in 20 of water. Dilute before using with from 100 to 200 parts of water.

ETCHING VARNISHES. 1. White wax 2 oz., asphaltum 2 oz., melt the wax in a clean pipkin, add the asphaltum in powder, and boil to a proper consistence. Pour it into warm water, and form it into balls, which must be kneaded, and put into taffeta for use.

2. White wax 2 oz., Burgundy pitch and black pitch each $\frac{1}{2}$ oz.; melt together and add by degrees 2 oz. of asphaltum in powder and boil till a drop cooled on a plate becomes brittle.

3. Equal quantities of linseed oil and mastic, melted together.

FATS, TO OBTAIN RANCIDITY IN. Mr T. B. GROVES. Add oil of pimento or balsam of Peru, 2 drops to the ounce of lard or other fat.

FILTERING POWDER. Fullers' earth washed, dried, and coarsely powdered; mixed with prepared bone black (see CHARCOAL, ANIMAL) coarsely powdered.

FILTER FOR CORROSIVE LIQUIDS. Powdered glass or asbestos, or gun-cotton loosely packed in the neck of a funnel.

FININGS, FOR GIN. Carbonate of potash 4 oz., roche alum 8 oz. Brewers' finings consist of isinglass dissolved in stale beer.

FIRES, TO EXTINGUISH. Dr CLANNY'S *Solution* consists of 5 oz. of sal ammoniac to a gallon of water. The compound used in Phillips's Fire Annihilator is said to consist of dried prussiate of potash, sugar, and chlorate of potash.

FIRES, COLOURED. The ingredients for these compounds must be dry, not too finely powdered, and mixed very uniformly. The nitrate of strontian requires to be gently heated in an iron pan till it falls to powder. The ingredients should always be reduced to powder separately, and mixed very lightly with the other powders; the whole must then be passed through a sieve once or twice. Great caution is required when dealing with chlorate of potash.

Blue and Purple Fires. 1. Chlorate of potash 9, dried verdigris 2, sulphur 1; mix.

2. Nitre 12, sulphur 16, black antimony 4, charcoal $\frac{1}{2}$, orpiment $\frac{1}{2}$.

3. Chlorate of potash 9, sulphur 12, refiners' blue verditer 3; mix.

4. *Purple.* 1. Chlorate of potash 5, nitrate of strontian 16, realgar 1, sulphur 2, lamp-black 1; mix.

5. Nitre 5, sulphur 2, metallic antimony 1; mix.

6. *Purple.* Chlorate of potash 2 oz., sulphur 1 dr. oxide of copper 1 oz.; mix.

7. *Violet.* Chlorate of potash 1 dr., pure copper $\frac{1}{2}$ dr., sulphur a scruple, charcoal 16 grs.; mix.

Green Fires. 1. Sulphur $10\frac{1}{2}$, nitrate of baryta $62\frac{1}{2}$,

chlorate of potash $23\frac{1}{2}$, sulphuret of arsenic $1\frac{3}{4}$, charcoal or lamp-black $1\frac{3}{4}$; mix.

2. Sulphur 13, nitrate of baryta 77, chlorate of potash 5, metallic arsenic 2, charcoal 3; mix.

3. Nitrate of baryta 20, sulphur $1\frac{1}{2}$, sulphuret of antimony $\frac{1}{2}$, chlorate of potash 10, charcoal $\frac{1}{4}$; mix.

4. Nitrate of baryta $10\frac{1}{2}$ drs., sulphur 2 drs., chlorate of potash 162 grs., lamp-black 26 grs.; mix.

5. Dry nitrate of barium 12 parts, sulphur 4, dry and finely-powdered chlorate of potash 5 parts. The chlorate to be mixed by a horn spatula.

Lilac Fire. Chlorate of potash 49, sulphur 25, dry chalk 20, black oxide of copper 6 parts. For pans.

Red Fires. 1. Chlorate of potash 10, nitrate of strontian 80, sulphur 26, charcoal 6; mix.

2. Chlorate of potash 3, nitrate of strontian 24, sulphur 7, lamp-black 1, sulphuret of antimony 2. Half the lamp-black or charcoal only may be added at first; and if on trial it does not burn freely, add more.

3. Chlorate of potash 1, nitrate of strontian 5, sulphur 1, black sulphuret of antimony 1; mix.

4. Chlorate of potash $2\frac{1}{2}$, nitrate of strontian 20, sulphur $6\frac{1}{2}$, sulphuret of antimony 2, charcoal $\frac{1}{2}$; mix.

5. Chlorate of potash 5, nitrate of strontian 28, sulphur 13, sulphuret of antimony 4, lamp-black 1.

6. Chlorate of potash 35, nitrate of strontian 360, sulphur 93, sulphuret of antimony 60, charcoal 10.

7. Nitrate of strontian 72, sulphur 20, coaldust 2, gunpowder 6; mix.

8. 40 parts of nitrate of strontian, 13 of sulphur, 2 of lime-tree charcoal. Mix, and add by mixing with a horn spatula, 5 parts of dry and finely-powdered chlorate of potash.

9. Nitrate of strontian 9 parts, shell-lac in powder 3 parts, chlorate of potash $1\frac{1}{2}$ parts.

White Fires. 1. Nitre 30, sulphur 10, black antimony 5; mix.

2. Nitre 48, sulphur $13\frac{1}{4}$, black antimony 5 parts; mix.

3. Nitre 12, sulphur 16, black antimony 4, charcoal $\frac{1}{4}$, white arsenic $\frac{1}{4}$; mix.

4. Nitre $46\frac{1}{4}$, sulphur 23, meal powder $12\frac{1}{2}$, zinc filings 18.

5. *For stars.* Nitre 57, sulphur 28, zinc filings 15.

Yellow Fires. 1. Nitre 3 oz., meal powder 3 oz., flowers of sulphur 3 oz., dried salt 2 oz.; mix.

2. Nitrate of soda 6, sulphur 1, lamp-black 1; mix.

* * * To guard against the danger sometimes arising from the spontaneous combustion of coloured fires containing sulphur and chlorate of potash, Mr SAUNDERS recommends intimately mixing 120 grains of powdered bicarbonate of potash with each pound of sulphur before using it in the manufacture of any composition into which chlorates enter.

COLOURED FLAME PAPERS. Soak Swedish filtering paper for ten minutes in a mixture of 4 parts of oil of vitriol with 5 parts of strong nitric acid, both by measure. When the strips are removed from the acid they must be thoroughly washed, first with cold, then with hot rain, or distilled water, till the washings cease to be acid. Then make different solutions (not too strong) of such of the chlorates of the metals as give the desired flame reactions, make them slightly warm, and saturate the papers separately with them.

Dry the papers before the fire previous to lighting them. They show to best advantage when a slip is loosely crumpled up into a pellet, lighted quickly at one corner, and thrown into the air against a dark background. Paper prepared with a salt of potassa will give a violet flame; that with soda a yellow one, that with barium a green, and that with strontian a crimson.

COLOURED FLAMES. The flame of alcohol may be coloured by mixing certain salts with the spirit. A green colour is given by chloride of copper, or boracic acid; red by nitrate of strontian, nitrate of iron, or nitrate of lime; yellow by nitrate of soda.

FIRE-PROOFING. FOR PAPER, see PAPER. For dresses, &c. A strong solution of sulphate of ammonia. The dresses of stage dancers may be soaked in a weak solution of chloride of zinc. The tungstate of soda is said to be the only perfect fire-proofer.

FLINTS, LIQUOR OF. *Soluble glass.* Mix 70 parts of pearl-ash, 54 of washing soda, and 152 of siliceous sand, and fuse the mixture in a crucible. It is soluble in water, and the filtered solution evaporated to dryness leaves a transparent glass. It has been proposed to render wood, muslins, &c., incombustible by means of the solution. Dr

TURNER directs 3 parts of carbonate of potash, and 1 of silica. See GLASS, SOLUBLE.

FLOWERS, COMPOUND FOR PROMOTING THE BLOWING OF. Sulphate of ammonia 4 oz., nitre 2 oz., sugar 1 oz., hot water a pint. Keep it in a well-corked bottle. For hyacinth glasses add 8 to 10 drops of the liquid to the water, changing the water every 10 or 12 days. For flowering plants in pots, add a few drops to the water employed to moisten them.

FLOWERS AND PLANTS, TO PRESERVE. Any vegetable substance may be preserved moist in a solution of creosote, or in glycerin. The method of drying plants between sheets of paper needs no description. But the original form, and in many instances the colour, of a fresh flower may be preserved by carefully immersing it in some fine dry material, and then rapidly drying in a baking oven. Millet seed has been used for this purpose, and may answer well for coarse specimens. For fine ones white river sand in equal grains must be used. To separate large grains it should be passed through a sieve; to remove fine particles it is copiously washed with water. While drying, it is to be constantly agitated.

M. REVEL recommends further that 1000 parts of this sand be intimately mixed with 1 of stearic acid and 1 of spermaceti, before using. See next formula.

FLOWERS, TO PRESERVE IN THEIR NATURAL SHAPE AND COLOUR. Provide a vessel with a moveable cover. Fit to the top a piece of fine metallic gauze, and replace the cover. Pass through a sieve into an iron pot, sand sufficient to fill this vessel, and heat it with $\frac{1}{2}$ per cent. of stearin, carefully stirring. The vessel with its cover on, and the gauze beneath it, are then turned upside down, and the bottom being removed, place the flowers on the gauze, and pour in the sand and stearin, so as to cover and envelop them. Next place the vessel on the top of an oven for 48 hours. Remove the cover, invert the vessel, and the sand runs away through the gauze, leaving the flowers dried in their natural position. *Journ. Soc. Arts.*

Fresh flowers may be preserved for some time in glycerin.

FLOWERS, CUT, TO PRESERVE IN WATER. Add to the water a teaspoonful of salt, or a teaspoonful of charcoal. Flowers in pots may be watered with a weak solution of sulphate of iron.

FLUXES. In a general sense these are substances which promote the fusion of minerals, but particularly which cleanse a reduced metal, by assisting its separation from its impurities. They also serve to defend it from the action of the air, and some of them assist in the reduction of oxides.

Black Flux. Into an earthen crucible, heated sufficiently to cause feeble combustion, but not to fuse the flux, throw successive portions of a mixture of 1 part of nitre, and 2 of crude (or cream of) tartar. Keep the flux in a close bottle.

White Flux. Into a large earthen crucible, heated to redness, throw successive portions of a mixture of 2 parts nitre and 1 of tartar. Keep it as the last.

Crude Flux is the mixture of nitre and tartar, before deflagration.

Dr CHRISTISON'S Flux for reducing arsenic. Mix crystallized carbonate of soda with 1-8th of charcoal, and heat gradually to redness.

FRESENIUS'S Flux, for reducing sulphuret of arsenic. Dry carbonate of potash 3 parts, cyanide of potassium 1 part.

Cornish Flux. Cream of tartar 10 parts, nitre $3\frac{1}{2}$, borax 3.

MORVEAU'S Flux. Pulverized glass (free from lead) 8 parts, calcined borax $\frac{1}{2}$ part, charcoal $\frac{1}{2}$ part.

Mr TAYLOR'S Flux. Saturate a solution of tartaric acid with carbonate of soda, evaporate to dryness, and calcine in a covered platinum crucible.

Sal enixum (the acid sulphate of potash left in distilling nitric acid), sandiver or glass-gall, fluor spar, limestone, &c., are also used as fluxes.

FLY POISON. A common poison for flies consists of white arsenic, or King's yellow, with sugar, but the use of such compounds may lead to fatal accidents. A sweetened infusion of quassia answers the same purpose, and is free from danger. Pepper, with milk, is also used; and also some adhesive compounds by which the flies are fatally entangled. *Papier moure* contains a large quantity of arsenic.

FREEZING MIXTURES. The salts should be in a crystallized state, with as much water crystallization as possible without being damp. They should be coarsely pulverized at the time of using, and put into the water contained in a basin or other suitable vessel. The water to be frozen should be enclosed in a thin metallic vessel, and immersed in the

freezing mixture. To obtain extreme degrees of cold, the ingredients and vessels employed should be previously cooled by one of the freezing mixtures.

1. Sal ammoniac 5 oz., nitre 5 oz., water 16 oz.

2. Mix 4 oz. of nitrate of ammonia, 4 of crystallized carbonate of soda, and 4 of water. In 3 hours 10 oz. of water may be frozen.

3. Nitrate of ammonia and chloride of ammonium in equal proportions, water q. s.

4. Nitrate of ammonia 5 parts, nitrate of potash 5, sulphate of soda 8, water 16 parts.

5. Phosphate of soda 9 parts, dilute nitric acid 4 parts.

6. Sulphate of soda 8 parts, hydrochloric acid 5 parts.

7. Sulphate of soda 6 parts, nitrate of ammonia 5 parts, diluted nitric acid 4 parts.

8. Mix 1 part by weight of powdered sal ammoniac with 2 of powdered nitre. Reduce common washing soda to powder. Keep these powders in well-closed bottles, and when required for use take equal measures of each, and add an equal bulk of water, or sufficient to dissolve the salts.

9. Chloride of ammonium 11 drs., nitre 10 drs., sulphate of soda 2 oz. Powder separately, and mix in a tin vessel with 5 oz. of water.

With Ice.

1. Snow or pounded ice 2 parts, chloride of sodium 1 part.

2. Snow 2 parts, crystallized chloride of calcium 3 parts.

3. Snow 8 parts, hydrochloric acid 5 parts.

4. Snow or pounded ice 12 parts, chloride of sodium 5 parts, nitrate of ammonia 5.

5. Snow 7 parts, diluted nitric acid 4 parts.

6. Snow 3 parts, diluted sulphuric acid 2 parts.

7. Snow 1 and 1-10th part, oil of vitriol (containing 66 and 1-5th per cent. of oil of vitriol) 1 part.

FRENCH POLISH. This is an alcoholic solution of shell-lac; some of the softer resinous gums are usually added, but too much of them renders the polish less durable. Highly rectified spirit, not less than 60 over proof, should be used. Rectified wood naphtha is sometimes substituted, to which the unpleasant smell is the only objection. Methylated spirit is now almost invariably employed for the

manufacture of French polish, and it is advised to substitute it for the rectified spirit in the following formulæ.

1. Orange shell-lac 22 oz., rectified spirit 4 pints; dissolve.

2. Shell-lac 3 oz., gum sandarac $\frac{1}{2}$ oz., rectified spirit a pint.

3. Shell-lac 4 oz., gum thus $\frac{1}{2}$ oz., rectified spirit a pint; dissolve, and add almond or poppy oil 2 oz.

4. Shell-lac 5 oz., seed-lac 1 oz., gum juniper $\frac{1}{2}$ oz., mastic 1 oz., rectified spirit a pint.

5. Shell-lac 3 oz., seed-lac 3 oz., gum juniper $1\frac{1}{2}$ oz., mastic 1 oz., rectified spirit a quart.

6. Shell-lac 5 oz., oxalic acid $\frac{1}{2}$ oz., rectified spirit a pint; dissolve, and add linseed oil 4 oz.

7. Shell-lac 5 oz., gum benzoin 5 oz., oxalic acid 10 drs., rectified spirit a quart; dissolve, and add $\frac{1}{2}$ pint of linseed oil.

8. Shell-lac 8 oz., gum elemi $1\frac{1}{2}$ oz., rectified spirit 4 pints.

9. Shell-lac 10 oz., seed-lac 6 oz., gum thus 3 oz., sandarac 6 oz., copal varnish 6 oz., rectified naphtha 8 pints. Or dissolve 8 oz. each of seed-lac, gum thus, and sandarac, separately in a pint of naphtha; and 1 lb. of shell-lac in 8 pints of naphtha. Then mix 6 oz. of copal varnish, 12 oz. of the solution of seed-lac, 6 oz. of the solution of gum thus, 12 of the solution of sandarac, and $5\frac{1}{2}$ lbs. of the solution of shell-lac. Let the copal varnish be put into the tincture of shell-lac, and well shaken, and the other ingredients added. A correspondent speaks highly of this polish.

10. Copal $\frac{1}{2}$ oz., gum arabic $\frac{1}{2}$ oz., shell-lac 1 oz. Pulverise, mix, and sift the powders, and dissolve in a pint of spirit.

11. Shell-lac 5 oz., rectified naphtha a pint.

French polish is sometimes coloured with dragon's blood, turmeric root, &c. The general directions for preparing the polish are to put the gums with the spirit in a tin bottle, which should be placed in water of such a temperature as to keep it at a gentle heat, shaking it frequently. The cork should be loosened a little before shaking it, taking care that there is no flame near to kindle the vapour. When the gums are dissolved, let it settle for a few hours, and pour off the solution from the dregs. The method of using it is to have a roll of list

over the end of which five or six folds of linen rag are placed. The polish is applied to the linen with a sponge, and a little linseed oil is dropped on the centre of it.

FULMINATING COMPOUNDS. *Fulminating Powder.* Mix together in a warm mortar 3 parts of pulverized nitre, 2 of dry carbonate of potash, and 1 of sulphur. A small quantity heated on an iron shovel or ladle till it fuses, suddenly explodes with great violence. It should be used with great caution. Another kind of fulminating or detonating powder is made by mixing 3 grs. of chlorate of potash with 1 of sulphur: by strongly trituring it with strong pressure in a marble mortar, a succession of sharp explosions is produced. The same mixture, or 6 grs. of chlorate of potash, 1 of sulphur and 4 of charcoal, struck with a hammer on an anvil, gives a loud report.

Chloride, or Terchloride of Nitrogen, and Iodide or Teriodide of Nitrogen cannot be meddled with without extreme danger. *Fulminating gold*, and the *fulminating silver* prepared with ammonia, are also dangerous compounds, even in minute quantities. As they serve no practical use, the mode of preparing them is omitted.

Fulminating Mercury (HOWARD'S, as improved by Dr URE). Dissolve by a gentle heat 1 oz. of quicksilver in $7\frac{1}{2}$ fluid oz. (or 10 oz. by weight) of nitric acid, of 1.4 specific gravity, in a glass retort, the beak of which is loosely inserted into a large balloon or bottle. When the mercury is dissolved, the solution, at the temperature of 130° F., should be slowly poured through a funnel into 10 fl. oz. of alcohol of 0.830 specific gravity, contained in a vessel that will hold 6 times the quantity of ingredients. When the action ceases, pour the contents of the matrass on a double filter in a glass funnel; wash out any powder that may remain in the matrass with a little cold water, and throw this also on the filter; and wash the fulminate with more water till it passes free from acid. When sufficiently drained, lift the filter out of the funnel, and lay it open on a copper or earthen plate, and dry the fulminate 212° F. or under, by hot water or steam. Its manufacture requires great caution: some valuable lives having been sacrificed in its preparation.

Fulminating Silver (BRUGNATELLI'S). On 100 grs. of pulverized nitrate of silver, in an open glass vessel, pour first an oz. of alcohol, and then as much strong nitrous acid:

The mixture boils, and gives out ethereal vapours. When all the powdered nitrate has taken the form of white clouds, cold distilled water must be added to sustain ebullition, otherwise the fulminate will be dissolved. Collect the powder on a filter, and dry it at a low temperature. Dr TURNER directs 1 part of silver to be dissolved in 10 of nitric acid, at a gentle heat, 20 parts of rectified spirit to be added, and the mixture warmed. When it begins to boil, set it aside to cool, collect and wash the crystals on a filter, and carefully dry them. This is more violent and dangerous than fulminating mercury.

FUMIGATIONS. See FUMIGATIO, *Pocket Formulary*, for their medicinal uses. Though not strictly belonging to this place, it may be useful to give a few directions for the management of these important agents as disinfectants and purifiers.

Carbolic Acid Fumigation.—Steep rags in it, and then suspend them in various parts of the room, or dilute one ounce with a gallon of water, and sprinkle the mixture over the walls and floors.

Chlorine Fumigation. This is probably the most effective in destroying noxious effluvia and putrid odours, and in checking the spread of contagious diseases. But as the gas itself is deleterious, except in a very diluted state, it must be used with caution in occupied apartments. To disinfect rooms from which the occupants have been removed, mix common salt and black oxide of manganese in equal quantities. Mix also in an earthen basin equal weights of oil of vitriol and water, and when it has cooled put it into a bottle for use. Into a china or earthen basin put from 1 to 3 oz. of the powder, according to the size of the room, and pour over it twice or thrice as much of the mixed acid. Place it in the apartment, and close the doors and windows for a few hours; the doors and windows are then thrown open till the smell of chlorine disappears. Dr A. T. THOMSON directs a mixture of 1 oz. of salt and $\frac{1}{2}$ oz. of black oxide of manganese to be put into a china cup, and 6 fl. drs. of oil of vitriol poured on it, the cup being placed in a pipkin of hot sand. Instead of the above ingredients, some chloride of lime may be placed in a large jar or basin, and a mixture of acetic acid and water poured on it. When used in, or near the apartments of the sick, great care must be taken that the chlorine is

so diluted with air that it shall occasion no annoyance to the invalid. Some contrivances have been adopted to render the extrication of chlorine gradual and continual. SMITH'S Chlorine Fumigator, and the more simple one of Messrs HEATHFIELD & BURGESS, are very convenient. Another method was proposed by Mr SCANLAN in the *Pharmaceutical Journal*, vol. vii (1st series), p. 343. By such contrivances, chlorine may with care be safely employed in houses occupied by the sick (in the passages, stairs, &c.), to prevent the spread of infectious fevers; but chloride of lime, simply mixed with water, in the proportion of not more than 1 oz. to a quart, is usually sufficient to purify the chamber of the sick. It should be occasionally sprinkled on the floor, and also placed about the room in shallow dishes, or a linen cloth moistened with it, suspended on a line. The same method may be pursued in all places where unpleasant smells prevail.

Iodine. A little placed on a plate, diffuses into the atmosphere of a sick room, and is a useful disinfectant.

Nitric Fumigation. Put into a china cup equal measures of sulphuric acid and water, and add to it, from time to time, small quantities of powdered nitre; or put 2 or 3 drs. of powdered nitre into a cup, and pour over it about an equal quantity of oil of vitriol. Stir it with a glass rod or tobacco-pipe, and remove it from time to time to different parts of the apartment. For large rooms 2 or 3 cups may be required. It is often recommended to apply heat; but Dr BATEMAN, of the Fever Hospital, found this unnecessary and objectionable, especially in the apartments of the sick. No metallic or wooden stirrers, or vessels, must be used.

Muriatic Fumigation. Hydrochloric Fumigation. This is now almost disused, being less efficacious than the preceding. It is obtained by putting a few drachms of common salt into a cup, and pouring on it an equal quantity of oil of vitriol. The vapours are very injurious to the lungs.

Acetic Fumigation. The vapour of vinegar, and especially of strong acetic acid, is employed as a disinfectant, but its efficacy is now considered to be very limited. It may be used by keeping the vinegar boiling over a lamp. A coarser method sometimes used is to plunge a red-hot poker into a cup of vinegar. Aromatic vinegar, merely

held to the nose, may afford some slight protection to those who attend upon the sick.

Sulphur Fumigation. The fumes of burning sulphur may possibly have some effect in decomposing miasmata and noxious effluvia; but as they have no advantage over chlorine and are very disagreeable, and otherwise objectionable, they are not likely to be employed. Formerly the following powder was burnt to destroy contagious miasmata: flowers of sulphur, nitre, and powdered myrrh, of each 1 oz.

tar Fumigation. The vapour of boiling tar has been used as a disinfectant as well as a palliative in some affections of the respiratory organs. The usual plan is to keep the tar boiling over a lamp. See FUMIGATIO PICEA, *Pocket Formulary*.

Benzoin, styrax, and other odoriferous gums, cascarilla bark, coffee berries, and the compounds termed aromatic pastiles, are burnt as purifiers and disinfectants. But little confidence is now placed in them as prophylactics against infection. The same may be said of camphor and tobacco. They should not be depended on to the exclusion of more efficient means, nor be made a substitute for free ventilation and the removal of all sources of noxious effluvia, when practicable.

FUMIGATING PASTILES. See PERFUMERY further back.

FURNITURE CREAM. 1. Soft water a gallon, soap 4 oz., beeswax in shavings 1 lb.; boil together, and add 2 oz. of pearlash. To be diluted with water, laid on with a paintbrush, and polished off with a hard brush or cloth.

2. Wax 3 oz., pearlash 2 oz., water 6 oz.; heat them together, and add 4 oz. of boiled oil, and 5 oz. spirit of turpentine.

3. The name is sometimes given to a mixture of 1 oz. of white or yellow wax, with 4 oz. of oil of turpentine.

FURNITURE PASTE. 1. Melt 1 lb. of beeswax with $\frac{1}{2}$ pint of linseed oil, and add $\frac{1}{2}$ oz. of alkanet root; keep it at a moderate heat till sufficiently coloured; then remove from the fire, add $\frac{1}{4}$ pint of oil of turpentine, strain through muslin, and put it into small gallipots to cool.

2. Scrape 4 oz. of wax, and put it into a pipkin with as much oil of turpentine as will cover it, and $\frac{1}{2}$ oz. of powdered resin; melt with a gentle heat, and stir in sufficient Indian red to colour it.

3. Equal weights of beeswax, spirit of turpentine, and linseed oil.

FURNITURE OIL. 1. Linseed oil a pint, alkanet $\frac{1}{2}$ oz.; digest in a warm place till coloured, and strain.

2. The same with $\frac{1}{4}$ pint of oil of turpentine.

3. Linseed oil a pint, alkanet root 1 oz., rose pink 1 oz.; let them stand in an earthen vessel all night.

4. A quart of linseed oil, 6 oz. of distilled vinegar, 3 oz. of spirit of turpentine, 1 oz. of hydrochloric acid, and 2 oz. spirit of wine.

5. Linseed oil 8 oz., vinegar 4 oz.; oil of turpentine, mucilage, rectified spirit, each $\frac{1}{2}$ oz.; butter of antimony $\frac{1}{4}$ oz.; hydrochloric acid 1 oz.; mix.

6. Linseed oil 16 oz., black resin 4 oz., vinegar 4 oz., rectified spirit 3 oz., butter of antimony 1 oz., hydrochloric acid 2 oz.; melt the resin, add the oil, take it off the fire, and stir it in the vinegar; let it boil for a few minutes, stirring it; when cool put it into a bottle, and add the other ingredients, shaking all together. The last two are especially used for reviving French polish.

7. Linseed oil a pint, oil of turpentine $\frac{1}{2}$ pint, rectified spirit 4 oz., powdered resin $1\frac{1}{2}$ oz., rose pink $\frac{1}{2}$ oz.; mix.

8. Linseed oil $1\frac{1}{2}$ oz., vinegar $1\frac{1}{2}$ oz., hydrochloric acid $\frac{1}{2}$ oz.; mix.

FUSIBLE METAL. See ALLOYS, further back.

GALL, CLARIFIED. Ox-gall is prepared for the use of artists in the following manner: To a pint of fresh ox-gall boiled and skimmed, add 1 oz. finely powdered alum; leave it on the fire till the alum is dissolved, then let it cool, put it into a bottle, and cork it loosely. Treat another pint in the same way with 1 oz. of salt instead of alum. After standing more than 3 months, carefully decant from each bottle the clear portion, and mix them together. The colouring matter is precipitated, and a clear, colourless liquid is obtained by filtration. It is used for mixing artists' colours, and to prepare ivory, oiled paper, &c., to revive the colours. Also for taking out grease spots.

GALVANIC BATTERIES, SOLUTIONS FOR. See ACIDS, MIXED, FOR GALVANIC BATTERIES.

GANNAL'S SOLUTION. See ANATOMICAL INJECTIONS.

GANTEINE. (A composition for cleaning kid gloves; sometimes improperly termed *Saponine*). Dissolve 3 oz. of

soap by heat in 2 oz. of water, and when nearly cold add 2 oz. of eau de Javelle, and 1 drachm of water of ammonia; form a paste, which is to be rubbed with flannel over the glove until this is sufficiently clean.

GARANCINE. Madder (sometimes the spent madder of the dyer's bath) is mixed with its weight of oil of vitriol, added very gradually, so as to avoid overheating. The acid is then washed out.

GARROT'S COVERING FOR PILLS. Soak 1 oz. of purified gelatin in 2 or 3 drs. of water; keep it liquefied in a salt-water bath. The pills are stuck on long pins, and dipped in the solution; when cold the pins are withdrawn, after being heated by a small flame, which melts the gelatin and closes the hole. See *PILULÆ*, *Pocket Formulary*.

GASES. These are generated in gas bottles fitted, by grinding, with an S-formed tube; or in flasks to which a bent tube is adapted by means of a cork; in a common retort; or sometimes in iron bottles with a metal tube.* They are usually collected in vessels filled with water placed with their open end in a vessel of water. *Pepys's Gas Holder* is very useful for receiving, retaining, and transferring gases. The *pneumatic trough* consists of a vessel for holding water, with a shelf for sustaining the jars or bottles that are to be filled; these are filled by sinking them under water, and are then lifted on the shelf, above which the water rises, with their open end downwards. The beak of the retort, or bent tubes, are so placed that the gas issuing from them rises through the water into the vessel, and takes the place of the water in them. Some gases being very easily absorbed by water, are collected over mercury. Sometimes they are collected in dry bottles by displacement. For light gases, as ammonia, place a bottle in a vertical position with its mouth downwards over the tube from which the gas issues, which should touch the bottom of the bottle. When the bottle is filled with gas, and this escapes from the mouth, quickly withdraw the tube and close the bottle, still inverted, with a greased stopper. For gases heavier than air, as chlorine, or carbonic acid, the bottle must be placed with its mouth uppermost, and the tube delivering the gas must descend to the bottom of the bottle. When full of the gas,

* Whenever practicable, india-rubber tubing is now largely employed in place of tubes of glass and metal.

close it with the greased stopper. The tube connected with a flask in which a gas is generated should have a ball blown in it, into which asbestos may be introduced to arrest any particles thrown up by effervescence.

The following are the processes for procuring the principal gases :—

Ammoniacal Gas. This is obtained by mixing equal weights of slaked lime (previously cooled) and powdered sal ammoniac, and heating the mixture in a retort or flask. As water rapidly absorbs the gas, it must be collected over mercury, or in a dry bottle as described above. When required for laboratory use, it is most readily obtained by gently heating a strong solution of ammonia and collecting the escaping gas by displacement.

Carbonic Acid Gas. Carbonic Anhydride. This is obtained by acting on marble or chalk, or carbonate of soda or potash, by a diluted acid. For exact experiments it must be collected over mercury; otherwise it may be collected in a bottle. Mr BENSON states that a saturated solution of sulphate of magnesia may be used in collecting this gas, instead of mercury. See *Pocket Formulary*.

Carbonic Oxide is obtained by acting on binoxalate of potash with 6 times its weight of oil of vitriol at a gentle heat; or by strongly heating in an iron bottle or gun-barrel, equal weights of chalk and iron filings. The gas must be passed through water containing lime or caustic potash to absorb the carbonic acid gas.

Chlorine. This gas may be obtained from oxide of manganese, common salt, and sulphuric acid, as directed in making chloride of lime. Or from hydrochloric acid and oxide of manganese (see *CHLORINIUM, Pocket Formulary*); but more conveniently, on the small scale, by dissolving common salt in water, adding a sixth of its weight of nitric acid, and as much oxide of manganese. Apply a gentle heat, and the gas is abundantly produced without violent action.—*The Chemist*, vol. i.

Hydrochloric Acid Gas. It may be obtained by heating together in a flask equal weights of salt and oil of vitriol; or simply by heating strong hydrochloric acid. It must be collected over mercury; or otherwise conducted to the bottom of a dry bottle as described above.

Hydrogen Gas is readily procured by pouring on fragments of zinc, in a glass bottle, or flask with a bent tube, or

retort, some diluted sulphuric acid (1 measure of strong acid to 5 of water). It may be collected over water. If zinc be not at hand, fine iron wire, or the turnings or filings of iron, may be substituted for it. To procure gas of great purity, distilled zinc must be used.

HYDROGEN, PEROXIDE. See OXYGENATED WATER, further on, and HYDROGENII PEROXIDUM, *Pocket Formulary*.

Carburetted Hydrogen Gas. Light carburetted hydrogen is readily obtained by stirring the mud of stagnant pools. It may be artificially prepared by heating in a flask, a mixture of crystallized acetate of soda, and dry caustic potash of each 2 parts, and 3 parts of quicklime in powder. Heavy carburetted hydrogen is prepared by heating 1 part of alcohol with 6 or 7 of oil of vitriol, and conducting the mixed gases through milk of lime, which retains the sulphurous acid; and afterwards through oil of vitriol, which absorbs water, ether, and alcohol. Coal gas is a mixture of these gases, with other hydrocarbons, &c.

Phosphuretted Hydrogen. The spontaneously inflammable variety of this gas is made by boiling phosphorus with solution of potash in a small retort, the beak of which is kept under water; as each bubble of gas rises from the water, it inflames, and forms a ring of white smoke, which dilates as it ascends. The spontaneous inflammability of the gas when mixed with atmospheric air or oxygen renders caution necessary in its preparation. The other varieties of phosphuretted hydrogen have no special interest or application.

Sulphuretted Hydrogen. Hydrosulphuric Acid. Hydric Sulphide. This gas is best obtained from sulphide (sulphuret) of iron, broken into small lumps. On the sulphide, contained in a gas bottle, or other suitable apparatus, pour sulphuric acid previously diluted with 7 parts of water. If more acid be afterwards required, dilute the strong acid with only 4 of water. This gas is absorbed by water.

Nitrogen, or Azote. Atmospheric air may be made to yield an unlimited supply of nitrogen, by exposing it to the action of substances which combine with its oxygen. By burning phosphorus in a large bell-glass standing in water, and allowing it to stand over the water a few hours, nearly pure nitrogen is obtained, which may be further purified by agitating it with solution of pure potash.

Protoxide of Nitrogen. Nitrous oxide of DAVY. It is most conveniently made by heating nitrate of ammonia (formed by neutralizing pure nitric acid, diluted with 3 parts of water, with carbonate of ammonia, and boiling the solution till a drop let fall on a cold plate solidifies, adding a little ammonia towards the end to ensure neutralization) in a retort, at a heat not exceeding 500° Fahrenheit, till it is nearly all decomposed. It may be collected over warm water. This gas (principally, if not wholly, in the liquid form) is manufactured in enormous quantities for the use of dentists and for dental hospitals.

Oxygen Gas. Mix chlorate of potash with a third of its bulk of black oxide of manganese; put the mixture into a gas bottle, or clean flask, to which a bent tube is fitted by a cork, and apply gentle heat. The gas, which comes over freely, may be collected over water.

Ozone. This is supposed to be oxygen in an allotropic and more active state. 1. It is formed by putting into a wide-mouthed bottle pieces of clean phosphorus, with a little water, so that the phosphorus shall be partly in the water and partly uncovered. Close the bottle for some hours, when the air it contains will manifest the odour and effects of ozone.

2 (LINDER). Manganese dioxide, potassium permanganate, and oxalic acid p. æ. Two spoonfuls of this powder placed on a dish and gradually mixed with water, is sufficient for a room of medium size. More water is added in small portions from time to time, as the evolution ceases.

Sulphurous Acid Gas. Sulphurous Anhydride. It is procured in a nearly pure state by heating 2 parts of quicksilver with 3 of oil of vitriol, and collecting the gas over mercury. By passing the gas through a tube surrounded with a freezing mixture, it is condensed into a liquid. For ordinary purposes the gas may be obtained as directed under ACID, SULPHUROUS.

GAUZE, LISTER'S CARBOLIZED. Paraffin 7 parts, resin 5 parts, carbolic acid 1 part. Melt the paraffin and resin and add the carbolic acid. It is generally made by dipping the gauze into this mixture, and then passing it through rollers to remove the excess. *New Remedies.*

GELATIN, PUIFIED. *Grenetine.* It is made by various processes from gelatinous animal matters. NELSON'S

Patent Gelatine is made from cuttings of the hides of beasts. These, freed from hair, flesh, fat, &c., are washed and scored, then macerated for 10 days in a ley of caustic soda, and afterwards placed in covered vessels at a temperature of 60° to 70° F., until they become tender; then washed from the alkali, exposed to the vapour of burning sulphur until they become sensibly acid; dissolved in earthen vessels heated to 150° F., strained, put into settling vessels heated to 100° or 120° F. for nine hours, the clear liquor drawn off, and poured on the cooling slabs to the depth of $\frac{1}{2}$ an inch. When cold, the jelly is cut in pieces, washed till free from acid, redissolved at 85°, poured on slabs, cut up, and dried on nets.

Bone Gelatin. The bones are boiled to remove the fat, then digested in dilute hydrochloric acid till the earthy matter of the bone is dissolved. The gelatin, which retains the form of the bone, is washed in a stream of water, plunged in hot water, and again in cold, to remove all remains of acid, and sometimes put into a solution of carbonate of soda. When well washed, it is dried on open baskets or nets. By steeping the raw gelatin in cold water, dissolving it in boiling water, evaporating the jelly, and cutting it into tablets, it may be dried and preserved in that form.

GERMAN PASTE, for feeding insectivorous singing-birds.

Blanched sweet almonds 1 lb., pea meal 2 lbs., butter 3 oz., saffron a few grains, honey q. s. Form the whole into a paste, and granulate it by pressing it through a cullender. Some add the yolks of 2 eggs.

GILDING. Leaf gold is affixed to various surfaces, properly prepared by gold size or other adhesive medium. Metallic surfaces are coated with gold, by means of amalgam of gold and mercury, applied with a wire brush, wet with an acid solution of mercury (made by dissolving 10 parts of mercury in 11 of nitric acid, by a gentle heat, and adding $2\frac{1}{2}$ parts of water). The article thus coated is heated over charcoal till the mercury is dissipated, and afterwards burnished. To give it a redder colour, it is covered with gilders' wax (a compound of verdigris, ochre, alum, and yellow wax); again exposed to heat, and afterwards washed and cleaned by a scratch brush and vinegar. An inferior kind of gilding is effected by dis-

solving gold, with a fifth of its weight of copper, in nitrohydrochloric acid, dipping rags in the solution, drying and burning them, and rubbing the ashes on the metallic surface with a cork dipped in salt and water.

GILDING, by immersion. Dissolve teroxide or terchloride of gold in a solution of pyrophosphate of soda, and dip the article to be gilt in it.

Electro-Gilding, by Elkington's patent process, is thus performed:—A solution of 5 oz. of gold (see ACID, NITROHYDROCHLORIC, further back) is prepared, and boiled till it ceases to give out yellow vapours; the clear solution is mixed with 4 gallons of water, 20 lbs. of bicarbonate of potash added, and the whole boiled for 2 hours. The articles, properly cleaned, are suspended on wires, and moved about in the liquid from a few seconds to a minute, then washed, dried, and coloured in the usual way.

The solution used in gilding with the voltaic apparatus is made by dissolving $\frac{1}{4}$ oz. of oxide of gold, with 2 oz. of cyanide of potassium, in a pint of distilled water.

GINGER BEER. See BEVERAGES.

GINGER BREAD, PURGATIVE. See further back.

GLASS OF BORAX. Calcine borax with a strong heat till the water of crystallization is expelled, and the salt fuses into a clear glass.

GLASS, SOLUBLE. FUCHS. Mix 10 parts of carbonate of potash, 15 of quartz (or of sand free from iron and alumina), and 1 part of charcoal. Fuse together. The mass is soluble in 4 or 5 parts of water; and the filtered solution evaporated to dryness yields a transparent glass, permanent in the air.

GLASS. Glass differs considerably in composition, owing to the purposes for which it is intended; but it may be said to consist mainly of mixtures, in varying proportions, of silicates of potash, soda, lime, baryta, magnesia, alumina, and lead, coloured by the addition of small quantities of different metallic oxides, particularly those of iron, manganese, cobalt, uranium, and gold. Crown glass and green bottle glass contain a portion of lime. Green glass and some kinds of foreign white glass are free from lead, and should therefore be selected for chemical uses.

TOUGHENED GLASS. M. DE LA BASTIE'S process for toughening glass, consists in raising it to a very high tempera-

ture, and then plunging it while hot into a heated oleaginous bath.

The process can only be successfully carried out by attention to a number of minute details.

To mark on Glass. Glass may be written on for temporary purposes, by French chalk; pencils of this substance will be found convenient. Glass may be written on with ink, if the surface is clean and dry, and the pen held nearly perpendicular. The shell-lac ink (see INK) is the best for labels, as it resists damp, &c. *To scratch glass,* a scratching diamond is used; or a piece of flint, or crystal of quartz, or the point of a small 3-square file. *To engrave on glass,* fluoric (hydrofluoric) acid is used, either in the liquid state or in vapour. The glass must be warmed, and coated with wax, or engravers' cement, and the writing or design traced through the wax with a bradawl, or other pointed instrument. The liquid hydrofluoric acid is poured on it, and left to act on the uncovered portions of the glass; or the fluorspar may be powdered and made into a paste with oil of vitriol, and laid over the prepared surface, and covered with lead-foil or tea-lead; or bruised fluorspar is put in a Wedgewood evaporating basin, with sufficient oil of vitriol to form a thin paste, and the prepared glass laid over the basin, so that the vapours may act on the portions from which the wax has been removed. *To cut glass* (besides the usual method of dividing cut glass by a glazier's diamond), the following means may be used:—To divide glass tubes or rods, form a deep mark round them with the edge of a sharp 3-square file, then with a hand placed on either side of the mark, break the rod with a slightly stretching as well as bending motion. A diamond or sharp flint may be substituted for a file. Flasks, globes, and retorts may be divided by means of iron rings, having a stem fixed in a wooden handle. Make the ring red-hot, and apply it to the flask, &c. If the vessel do not break where it came in contact with the ring, wet the part, and it will generally separate. Another method is to twist together 2 or 3 threads of cotton, such as is used for wicks, moisten them with spirit of wine, and encircle the flask with them; then holding the flask horizontally, set fire to the wick, and turn the flask with the fingers, so as to keep the flame in the direction of the thread. If the separation does not take place

the first time, the process may be repeated after the glass has cooled. By these means a common oil flask may be divided into an evaporating dish and a funnel. By means of a stout iron rod, fixed in a wooden handle, and terminating in a blunt point, and heated to redness, broken retorts, globes, and flasks may be converted into useful evaporating dishes, &c. If any crack exist, it may easily be led in any direction, as it will follow the motion of the heated iron. If no crack exist, one must be produced by applying the point of the heated rod to any convenient spot on the edge of the broken glass, touching it afterwards with a moistened finger, if necessary. The edges of glass thus divided are rendered less apt to break by heating them in the flame of a blowpipe, or grinding them smooth with emery on a flat stone. See FARADAY'S Manipulations.

Glass, to Platinize. M. DULLO. This is recommended to prevent fusing of the end of the tube used in Marsh's test for arsenic. In drawing out the end of the tube leave the diameter slightly larger than is necessary; then roughen the narrow end with a file. Dip in a solution of bichloride of platinum, containing 1-20th of the metal; remove excess of the drop, and heat cautiously till the glass acquires a metallic appearance. Repeat this four or five times.

Glass, to Silver. 1. The term *silvering* is commonly applied to the process of coating the surface of the glass with amalgamated tinfoil, in forming mirrors. The tinfoil is rubbed over with quicksilver, and more of the latter poured over it; the plate of glass, perfectly clean and dry, is then applied to it in such a way as to exclude all air-bubbles, and to bring the glass and tinfoil into perfect contact. The plate, after being inclined, so as to allow the superfluous quicksilver to drain off, is loaded with weights, under which it remains till the adhesion is complete. To convex and concave mirrors, the amalgamated foil is applied by means of accurately fitting plaster moulds. The interior of globes is silvered by introducing a liquid amalgam (see AMALGAMS), and turning about the globe, till every part is covered with it.

2. PETTIJEAN'S process. 1540 grs. of nitrate of silver are treated with 955 grs. of strong solution of ammonia, and afterwards with 7700 grs. of water. To this solu-

tion, when clear, are added 170 grs. of tartaric acid dissolved in 680 of water, then 152 cubic inches more of water, with brisk agitation. When it has settled, the clear part is poured off; 152 cubic inches of water are added to the sediment, to dissolve as much as possible. The clear fluids are mixed, and again 62 cubic inches of water are added. This is Silvering Solution No. 1. No. 2 is prepared in the same manner, but with twice as much tartaric acid. The glass plate being cleaned, and laid horizontally, an even layer of Solution No. 1 is poured on it, to about the depth of 1-10th of an inch. Heat is then applied by means of a cast-iron water-bath beneath. Bright silver is soon deposited. When the stratum of fluid is exhausted it is poured off, and Solution No. 2 next applied in the same way. The silver surface may be polished, or coated with black varnish.

3. LIEBIG proposes to silver glass without mercury by a solution of ammonio-nitrate of silver, to which potash or soda is added, and which deposits silver at ordinary temperatures when brought in contact with an aqueous solution of sugar of milk.

4. Mr DRAYTON mixes 1 oz. of nitrate of silver, 3 oz. of water, 1 oz. of liquid ammonia, and 3 oz. of spirit of wine, and filters the solution after it has stood 4 hours. To every ounce of the solution he adds $\frac{1}{4}$ oz. of grape sugar, dissolved in equal quantities of water and alcohol. The surface to be silvered is covered with this liquid at a temperature of 160° F., maintained till the deposition of silver is complete. When quite dry, the coated surface is covered with mastic varnish. Other substances besides sugar occasion the deposition of silver from the ammoniacal solution; as oil of cassia, oil of cloves, and other essential oils, aldehyd, formic acid, &c. UNGER recommends a strong alcoholic solution of tannin. M. VOHL prefers an alkaline solution of gun-cotton. Dissolve gun-cotton in a solution of caustic potash, pour it into a solution of nitrate of silver, and add ammonia sufficient to redissolve the precipitate. The liquor, being slowly heated in a water-bath, becomes brown, effervesces, and deposits silver of superior brilliancy.

Glass, to Clean. The vessel to be cleansed is filled, or if large, rinsed with a moderately dilute solution of the permanganate of potash, contact being prolonged till a

film of hydrated manganic oxide has been deposited; the solution is then poured away, and the glass vessel rinsed with some strong hydrochloric acid. Concentrated sulphuric acid is used for the same purpose.

GLAZES. Common earthenware is glazed with a composition containing lead, on which account it is unfit for many pharmaceutical purposes. The following glaze has been proposed, among others, as a substitute:—100 parts of washed sand, 80 of purified potash, 10 of nitre, and 20 of slaked lime; all well mixed and heated in a blacklead crucible, in a reverberatory furnace, till the mass flows into a clear glass. It is then to be reduced to powder. The goods to be slightly burnt, placed under water, and sprinkled with the powder.

GLAZE FOR PORCELAIN. (ROSE.) Felspar 27 parts, borax 18, Lynn sand 4, nitre 3, soda 3, Cornwall china clay 3 parts. Melt together to form a frit, and reduce it to a powder, with 3 parts of calcined borax.

GLUE is made by boiling parings of ox-hides and other skins in water, evaporating the solution to a proper consistence, allowing it to gelatinize in wooden boxes, cutting it into layers with a wire, and drying the layers on nets stretched on wooden frames. *Bones* also yield a pale glue, described under **GELATIN**. Bank-note glue or mouth glue is made by dissolving 1 lb. of fine glue, or gelatin, in water, evaporating it till most of the water is expelled, adding $\frac{1}{2}$ lb. of brown sugar, and pouring it into moulds. Some add a little lemon-juice. It is also made with 2 parts of dextrin, 2 of water, and 1 of spirit.

GLUE, LIQUID. 1. Dissolve bruised orange shellac in $\frac{3}{4}$ of its weight of methylated or rectified spirit, or of rectified wood naphtha, by a gentle heat. It is very useful as a general cement and substitute for glue. 2. Another kind may be made by dissolving 1 oz. of borax in 12 oz. of soft water, adding 2 oz. of bruised shellac, and boiling till dissolved, stirring it constantly. 3. Dissolve 1000 parts of glue in 1000 parts by weight of water in a glazed pot, over a gentle fire. When it is melted, add nitric acid (sp. gr. 1.32) 200 parts, pouring it in very gradually. An effervescence is caused by the escape of hyponitrous acid. When all the acid is added, allow the mixture to cool. (This glue is found to remain unaltered on exposure to

the air. It is applied cold, and is recommended as very convenient in chemical operations.) M. DUMOULIN.

GLUE, MARINE. Cut caoutchouc into small pieces, and dissolve it, by heat and agitation, in coal naphtha. Add to this solution powdered shellac, and heat the whole with constant stirring, until combination takes place, then pour it while hot on metal plates, to form sheets. When used it must be heated to 248° F., and applied with a brush.

GLUE OF CASEIN. 1. BRACONNOT. Dissolve casein in a strong solution of bicarbonate of soda. 2. WAGNER. Dissolve casein in a cold saturated solution of borax. Superior to gum, and may take the place of glue in many cases. May be used for the backs of adhesive tickets.

GLUTEN, VEGETABLE. Form wheat flour into a stiff paste with cold water; then knead it under a stream of water till all the starch is washed away. What remains is impure gluten.

GOLDEN COMPOUND. Anhydrous tungstate of soda, or the salt obtained in fusing 2 equivalents of tungstic acid with 1 of carbonate of soda, is to be melted in a porcelain crucible, over a spirit lamp, at a temperature not more than sufficient; then add small pieces of pure tin to the melted mass. Cubes of a golden colour instantly form. The process should not be continued too long, or they acquire a purple hue. See AURUM MUSIVUM.

GUM, BRITISH. See DEXTRIN. It is also prepared by heating starch alone, or previously mixed with an acid. PINEL directs half a gallon of nitric and half a pint of hydrochloric acid to be mixed with 100 gallons of water, and as much potato fecula added as will form a paste. In 2 hours remove the paste in buckets, prepared for the purpose, to drain off all the water. Then place the paste in small lumps in a drying room till dry; pulverize it, and expose the powder the first day to the temperature of 100° F., the next day raise it to 150°, on the third day to 190°. It is then powdered, sifted, and heated from 300° to 350°. To give it the appearance of gum after it has gone through the stove, and is powdered and sifted, mix it to a paste with water to which 1 per cent. of nitric acid has been added, spread it on copper plates in layers $\frac{1}{4}$ of an inch thick, and heat it in an oven from 240° to 300° F., then remove it to the open air to cool.

GUM ARABIC, PURIFICATION OF. PICCIOTTO'S process. The gum is dissolved in water, and sulphurous acid gas passed into it. The sulphurous acid is sufficiently removed for common purposes by gently boiling the mucilage in a retort with a receiver attached. But to obtain the gum in a purer state, carbonate of barium is added, the mixture is filtered, afterwards agitated with gelatinous alumina, again filtered, and evaporated.

GUN BARRELS, TO STAIN. See **BROWNING LIQUIDS**, further back.

GUN COTTON. Mix $1\frac{1}{2}$ fl. oz. of each of the strongest nitric and sulphuric acid; put the mixture in a Wedgewood mortar, and when cool add 100 grs. of cotton wool. Stir it with a glass rod, and when it is fully soaked, squeeze out the acid with the pestle or a porcelain spoon, throw the cotton into a large quantity of water, squeeze it again, and wash it under a stream of water till quite free from acid. In the specification, the patentee directs 1 measure of nitric acid, sp. gr. 1.45 or 1.50 to be mixed with 3 measures of sulphuric acid, sp. gr. 1.85; the cotton to be soaked in the acid, then squeezed from it, and left in a covered vessel for an hour; and after washing, to be dipped in a solution of 1 oz. carbonate of potash in a gallon of water, then pressed, and partially dried; again dipped in a weak solution of nitre, then dried in a room heated to 150° F.

GUNPOWDER. A compound of nitre, charcoal, and sulphur. The nitre should be purified by recrystallization, the sulphur by distillation, and the charcoal selected of the best quality—that of the dogwood, alder, poplar, chestnut, or willow is preferred. The following is the composition of some of the most approved kinds:

	Nitre.	Charcoal.	Sulphur.
Royal Mills, Waltham Abbey	75	15	10
Marsh's Sporting	76	15	9
—— Mining	65	15	20
French (Government)	75	12.5	12.5
—— Sporting	78	12	10
Chinese	75	14.14	9.9

GUNPOWDER, WHITE. Well dried yellow prussiate of potash 1 part, white sugar 1 part, chlorate of potash 2 parts. Let the ingredients be separately reduced to a fine pow-

der, and the powders mixed by the hand, or by means of a leathern barrel turning on its axis. Or they may be moistened with water, and granulated by passing the paste through a wire sieve.

GYPNUM, TO HARDEN. KEATING'S patent process is to moisten calcined gypsum with a solution of 1 lb. of borax, 1 lb. of tartar, in 11 lbs. of water, it is then heated to redness for 6 hours, and pulverized. ERDEMANN recommends plaster figures, &c., to be soaked in a solution of FUCH'S soluble glass.

HAHNEMANN'S WINE TEST. See TESTS.

HARNES, JET. Take 4 oz. best glue, 1½ pints good vinegar, 2 oz. best gum arabic, ½ pint good black ink, 2 drs. best isinglass. Dissolve the gum in the ink, and melt the isinglass in another vessel in as much hot water as will cover it. Having first steeped the glue in the vinegar until soft, dissolve it completely by the aid of heat, stirring to prevent burning. The heat should not exceed 180° F. Add the ink and gum, and allow the mixture again to rise to the same temperature. Lastly, mix in the solution of isinglass, and remove from the fire. When used, a small portion must be heated until fluid, and then applied with a sponge, and allowed to dry on.

HARNES, WATERPROOF PASTE FOR. Put into a pipkin, black resin 2 oz., place on a gentle fire, and when melted add beeswax 3 oz. When this is melted, remove from the fire, and stir in ½ oz. fine lamp-black and ½ dr. Prussian blue, finely powdered. When completely mixed, add spirits of turpentine to form a thin paste, and let it cool. To be applied like blacking.

HEADING FOR BEER. GRAY. Equal parts of alum and sulphate of iron.

INCENSE. See PERFUMERY.

INDIA RUBBER COURT PLASTER. A stout frame of wood must be made, about 3 yards long and about 1½ yards wide. Within this frame must be placed two sides of another frame, running longitudinally and across, so fixed in the outer frame that the two pieces may slide independently of each other backwards and forwards about 6 inches. Tapes of canvas must be tacked round the inside of the inner frame and the corresponding sides of the outer frame, so as to form a square for the material to be sewn in, which when done the two loop frames must be drawn tightly to

the outer by means of a twine passed round each, in order to stretch perfectly free from irregularities, the silk or satin, previous to laying on the composition.

To make the plaster. Dissolve india rubber in naphtha or naphtha and turpentine, lay it on with a flat brush on the opposite side to that which is intended for the plaster. When the silk is perfectly dry and the smell in a great measure dissipated, it will be ready for the adhesive material; to make which take equal parts of Salisbury or fine Russian glue and the best isinglass, dissolve in a sufficient quantity of water over a water-bath, and lay on with a flat hogtoot while warm. It is requisite to use great caution to spread the plaster evenly and in one direction, and a sufficient number of coatings must be given to form a smooth surface, through which the texture of the fabric is not perceptible. Each coating should be perfectly dry before the succeeding one is given, after which the frame is to be placed in a situation free from dust, and where a draught of air would facilitate the drying. The quantity of water used and the weight of the two materials must be a little varied according to the season and the gelatin strength they possess. Lastly, the plaster being ready to receive the polishing coat, which gives also the balsamic effect to it, a preparation is made in nearly the same manner as the Tinct. Benz. Co. of the P. L., with the addition of more gums. This preparation must be laid on once only, and with a brush kept for the purpose. For making plasters on coloured silks it is only necessary to select the silk a shade deeper than the colour required, as the plaster causes it to appear a little lighter. See *EMPLASTRUM ICTHYOCOLLE*, *Pocket Formulary*.

INDIGO. (TAYLOR.) The principal preparations of indigo are described under **CHEMIC BLUE**, and **DYES**. Indigo may be purified by several methods, of which the following is the most simple:—Mix indigo with half its weight of Paris plaster and sufficient water to form a thin paste. Spread this evenly on an iron plate, about two inches wide, to the depth of 1-8th of an inch, and let it dry in the air. Then apply the flame of a large spirit lamp to the under side of the plate, beginning at one end and advancing it to the other as the sublimation proceeds. The violet vapour condenses on the surface in brilliant prisms or plates. Good indigo yields from 15 to 17 per cent.

Purified indigo is also obtained from the alkaline solution of reduced indigo, described under DYES; or by dissolving indigo in a mixture of 1 part of caustic soda, 1 of grape sugar and 20 of water. To the clear solutions thus obtained add hydrochloric acid to throw down the indigo, wash this perfectly with pure water, and finally with alcohol. If care be taken to exclude the air before and after adding the acid, and to wash it with recently boiled water, to drain it rapidly and dry it *in vacuo*, the indigo is obtained nearly white, but becomes blue on exposure to the air.

INKS. The following are specimens of the most useful kinds:

Black Writing Inks. 1. **BRANDE'S.** Bruised Aleppo galls 6 oz., soft water 6 pints; boil together, add 4 oz. of sulphate of iron and 4 oz. of gum arabic. Put the whole in a bottle, and keep it in a warm place, shaking it occasionally. In 2 months pour it off into glass bottles; and add to each pint a grain of corrosive sublimate, or 3 or 4 drops of creasote.

2. **Dr WOLLASTON'S.** Galls 1 oz., gum $\frac{1}{2}$ oz., cloves $\frac{1}{2}$ dr., sulphate of iron $\frac{1}{2}$ oz., water 8 oz. Digest with frequent shaking, till it has sufficient colour. A good durable ink, and will bear diluting.

3. *Prerogative Court Ink.* **GRAY.** Galls 16 oz., gum 6 oz., alum 2 oz., sulphate of iron 7 oz., kino 3 oz., logwood in powder 4 oz., water 8 lbs.

4. **Dr. URE'S Ink.** For 12 gallons of ink take 12 lbs. of bruised galls, 5 lbs. of gum Senegal, 5 lbs. of green sulphate of iron, and 12 gallons of rain-water. Boil the galls in a copper with 9 gallons of water for 3 hours, adding fresh water to replace what is lost by evaporation. Let the decoction settle, and draw off the clear liquor; add to it a strained solution of the gum; dissolve also the sulphate of iron separately, and mix the whole. Instead of boiling the galls, they may be macerated in a portion of hot water for 12 hours, then put into a percolator, and the rest of the water passed through it.

5. *Anti-corrosive Ink.* Aleppo galls 10 lbs., logwood 5 lbs., pomegranate-peel $2\frac{1}{2}$ lbs., cloves $2\frac{1}{2}$ oz., soft water 8 gallons. Let the whole boil gently for an hour or two, then cover the copper and leave it for 12 or 14 hours, stirring it now and then. Strain off the decoction and

add 2 gallons more water to the ingredients; simmer gently for an hour, and strain. Mix the liquors, and let them settle; draw off the clear liquid from the dregs; dissolve in a portion of it 2½ lbs. of gum arabic and ½ lb. of sugar candy; and in another portion 2½ lbs. of green sulphate of iron. Strain both solutions, and mix the whole together; then add 1 oz. of calcined borax, and ¼ oz. of creasote dissolved in ¼ pint of spirit of wine. Dr HARE recommends an ink free from acid, to be made with galls and "finery-cinder;" but we are not sure what is intended by this name.

6. RIBAUCCOURT'S *Ink*. Galls 1 lb., logwood ½ lb., gum 6 oz., sulphate of iron ½ lb., sulphate of copper 2 oz., sugar 2 oz., water 12 lbs. (or 5 quarts). This has the disadvantage of corroding the steel pens and the penknives with which it comes in contact.

7. Galls 3 oz., sulphate of iron 1 oz., logwood ½ oz., gum ½ oz., ale a quart. Let it stand in a loosely corked bottle in a warm place, for a week or more, shaking it daily.

8. Boil 4 oz. of logwood for an hour in 6 quarts of water; adding boiling water as it evaporates; then add 16 oz. of blue galls coarsely bruised, 4 oz. of dry sulphate of iron (*i.e.* heated till it becomes whitish and pulverulent), 3 oz. of brown sugar, 6 oz. of gum arabic, and ¼ oz. acetate of copper ground with a little of the decoction. Keep the whole in a bottle uncorked for a fortnight, shaking it twice a day.

9. READE'S *Patent*. This differs from common black ink, in containing a portion of soluble Prussian blue.

10. *Chrome Ink*. Extract of logwood ½ oz., gum ¼ oz., water a pint. Dissolve also in 12 oz. of water, ½ oz. of yellow chromate of potash (or ¼ oz. each of bichromate and bicarbonate of potash); and mix the two solutions. The ink is ready for immediate use.

11. Dr LEWIS'S *Writing Ink*. Powdered sulphate of iron 1 oz., powdered logwood 1 oz., powdered galls 3 oz., gum arabic 1 oz., white wine or vinegar 1 quart.

12. BERZELIUS recommends a solution of vanadate of ammonia in infusion of galls. Dr URE states that this forms the most perfect ink that can be desired; but the scarcity and high price of the vanadate prevents its use.

13. RUNGE'S *Black Writing Fluid*. Boil logwood

22 lbs., in enough water to yield 14 gallons of decoction. To 1000 parts of this decoction, when cold, add 1 part of chromate of potash. The mixture is to be well stirred. The proportions are to be carefully observed, and the yellow chromate, not the bichromate, employed. (This ink is said to possess some great advantages; to adhere strongly to paper, so that it can neither be washed off by water, nor even altered by weak acids; to form no deposit; and not to be in the least acted upon by steel pens.) Steel pens should be washed in an alkaline solution before being used with this ink. On exposure to the air, rapid decomposition of this fluid sometimes takes place. This may be prevented by adding a little carbonate of soda to the fluid.

14. *Alizarine Ink*. LEONHARDI. Digest 24 parts Aleppo galls with 3 parts of Dutch madder and 120 parts of warm water. Filter, and add 1.2 parts solution of indigo, 5.20 parts of sulphate of iron, and 2 parts crude acetate of iron solution. This ink contains no gum, and cannot get mouldy; the tannate of iron is prevented from separating by the sulphate of indigo. Alizarine ink may be evaporated to dryness and formed into cakes; 1 part with six of hot water will then form an excellent writing fluid.

Packers' Marking Ink is merely the dregs of black ink, for marking parcels with a brush.

Copying Ink. 1. Mr BRANDE directs 1 oz. of brown sugar to be added to No. 1, for copying. Another kind is made by dissolving $\frac{1}{2}$ oz. of gum, and 20 grs. of Spanish liquorice, in 13 drs. of water, and adding to it a drachm of lamp-black, previously mixed with a teaspoonful of sherry. If the lamp-black is greasy, it should be heated to redness in a covered crucible. Another published form is:—Black ink 3 oz., sugar candy 1 oz.

2. (*No press required*) PROFESSOR ATTFIELD. Reduce by evaporation 10 vols. of ordinary ink to 6, then add 4 vols. of glycerine.

Ink Powder. This consists of the dry ingredients for ink, powdered and mixed. 1. Powdered galls 4 oz., sulphate of iron (heated till it becomes white and pulverulent) 1 oz., powdered gum 1 oz., white sugar $\frac{1}{2}$ oz.; mix. To make a quart of ink, with water or beer.

2 (GRAY). Powdered galls 2 lbs., green vitriol 1 lb., gum 8 oz. In 2 oz. packets, each for 1 pint of ink.

Red Writing Inks. 1. Best ground Brazil wood 4 oz., diluted acetic acid a pint, alum $\frac{1}{2}$ oz. Boil them slowly in a covered tinned copper or enamelled saucepan, for an hour; strain, and add $\frac{1}{2}$ oz. gum. Some direct the Brazil wood to infuse for 2 or 3 days before boiling.

2. WEBER'S *Red Ink.* Boil 4 oz. of Pernambuca wood with 16 oz. of dilute acetic acid and an equal quantity of water, until 24 oz. remain. Add an ounce of alum, and evaporate again to 16 oz.; add gum arabic 1 oz., strain; and to the cold liquid add, lastly, 1 dr. of protochloride of tin. Said to be of a finer colour, and more permanent, than cochineal ink.

3. HENSELER. Boil 2 oz. of good Brazil wood, $\frac{1}{2}$ oz. of alum, and $\frac{1}{2}$ oz. of cream of tartar, in 16 oz. of rain-water, till reduced to half; strain, and dissolve in it $\frac{1}{2}$ oz. of gum arabic, and add a tincture made with $1\frac{1}{2}$ drs. of cochineal in $1\frac{1}{2}$ oz. of spirit of wine.

4. Brazil-wood 4 oz., alum 2 oz., water a quart. Boil for an hour, and strain; then add 1 oz. of gum.

5. Triturate 1 dr. of cochineal and 1 dr. of carbonate of potash, with a little boiling water; then add 1 dr. of burnt alum and 2 drs. of cream of tartar, and water to bring it to the desired colour. Not so permanent as the Brazil ink.

6. *Carminc Ink.* Heat a scruple of carmine with 3 oz. of water of ammonia for some minutes, a little below boiling, and add 15 to 20 grs. of gum. The inkstand must be kept well closed.

7. STEPHEN'S *Red Ink.* (Patent.) Take some common soda, potash, or carbonate of ammonia, and add to it, at intervals, twice its weight of crude argol in powder. When effervescence has ceased, pour off the solution, or filter it from insoluble matter. Add to it next, by measure, half the quantity of oxalate of alumina, prepared by adding to precipitated alumina in a damp state as much oxalic acid as will dissolve it. Into this mixture, when cold, put as much powdered cochineal as will give it a fine red colour, and, after letting it stand for 48 hours, strain it for use.

Blue Inks. 1. READE'S Patent. Prepare a solution of iodide of iron, from iodine, iron, and water; add to the solution half as much iodine as first used. Pour this solution into a semi-saturated solution of ferro-prussiate of potash, containing nearly as much of the salt as the

whole weight of iodine. Collect the precipitate, wash it, and finally dissolve it in water, to form the blue ink. The solution from which the precipitate is separated, evaporated to dryness, and the residue fused, redissolved, and crystallized, yields pure iodide of potassium. [This process being patented, ink must not be prepared by it for sale.]

2. Add a pint of a cold solution of persulphate of iron (prepared as directed, further on) to a solution of 6 oz. of ferro-cyanide of potassium in 2 pints of water. Collect the precipitate, wash it with distilled water until it begins to dissolve, then triturate it in a mortar with sufficient distilled water to form a blue ink.

3. Chemic, or Saxon blue (sulphate of indigo), diluted with water to the desired shade, with a little gum.

4. Pure Prussian blue, triturated with a sixth part of its weight of oxalic acid, with a little water, to a smooth paste, and more water added to bring it to the proper colour. A larger proportion of the acid is ordered in some recipes.

5. Dr NOEMANDY'S *Blue Ink*. Chinese blue (ferro-cyanide of iron) is ground in water with binoxalate of potash and gum arabic, in the following proportions:—7 oz. of water to 3 drs. of Chinese blue, 1 dr. of binoxalate of potash, and 1 dr. of gum arabic.

6. STEPHEN'S *Patent Blue Ink*. Common Prussian blue is first macerated in strong sulphuric acid, then repeatedly washed in water, and afterwards dried. This process is to render it more soluble in oxalic acid, which is now to be gradually added in the proportion of about 1 part to 6 of the Prussian blue (as before maceration), together with sufficient water to yield a dense blue solution.

7. Digest 2 oz. of the cuttings of tin plate with 4 oz. of nitrous acid, and add the solution to a gallon of water in which 2 oz. of prussiate of potash have been dissolved. [This requires frequent shaking to keep the precipitate (which is Prussian blue) suspended.]

Violet Inks. 1. Boil 8 oz. of logwood in 3 pints of rain or distilled water to 1½ pints. Strain, and add 1½ oz. of clean gum, and 2½ oz. of alum in fine powder. Agitate frequently till dissolved.

2. Cudbear 1 oz., pearlash 1½ oz., mucilage 2 oz., soft water to make a pint. Pour the water hot on the cudbear and pearlash, allow the mixture to stand for twelve hours,

then strain, and add the mucilage; 1 oz. of rectified spirit may also be added.

Purple Inks. 1. Add a little muriate (chloride) of tin to a strong decoction of logwood. A little gum may be added.

2. Dr NORMANDY'S *Purple Ink.* To 12 lbs. of Campeachy wood add as many gallons of boiling water, pour the solution through a funnel, with a strainer made of coarse flannel, on to 1 lb. of hydrate, or acetate of deutoxide of copper finely powdered (having at the bottom of the funnel a piece of sponge); then add immediately 14 lbs. of alum, and for every 340 gallons of liquid add 80 lbs. of gum arabic or gum Senegal. Let these remain for 3 or 4 days, and a beautiful purple colour will be produced.

Brown Ink. 1. Boil $\frac{1}{2}$ oz. of catechu with 8 oz. of water until dissolved, and strain. Dissolve 60 grs. of bichromate of potash in $1\frac{1}{2}$ oz. of water, and add it gradually to the solution of catechu until the desired shade is obtained. It requires no gum.

2. By adding to the violet ink finely powdered bichromate of potash, in the proportion of from 15 to 30 grs. to an ounce, various shades of brown and snuff colour are obtained.

Yellow Ink. 1. Gamboge triturated with water, and a little alum added.

2. Boil 8 oz. of French berries with 1 oz. of alum in a quart of water; strain, and add 1 oz. of gum.

Green Ink. WINCKLER. 1. Dissolve 3 drs. of bichromate of potash in 1 oz. of water; add to the hot solution $\frac{1}{2}$ oz. of alcohol, and decompose the mixture by a little strong sulphuric acid till it assumes a brown colour. Evaporate the liquid to half, let it cool, dilute with a sufficient quantity of water, and filter; add to the filtered liquid 4 drs. of alcohol, decompose with a few drops of sulphuric acid, and let it rest. After some time it assumes a fine green colour. A little gum may be added. There is danger of the paper and steel pens suffering from an excess of sulphuric acid.

2 (KLAPROTH). Distilled verdigris 2 oz., cream of tartar 1 oz., water 8 oz.; boil to half and filter.

3. Add to the yellow ink No. 2 sufficient sulphate of indigo.

4. Dissolve sap green in water with a little alum.

5. Rub $3\frac{1}{2}$ drs. of Prussian blue, and 3 drs. of gamboge, with 2 oz. of mucilage, and add $\frac{1}{2}$ pint of water.

Gold and Silver Ink. Fine bronze powder, or gold or silver leaf, ground with a little sulphate of potash, and washed from the salt, is mixed with water and a sufficient quantity of gum. Or, the gold leaf may be ground on a slab with honey, which is afterwards washed away.

Indestructible Writing Fluids. The common writing inks being liable to be obliterated by many chemical agents, several compounds more capable of resisting these agents have been proposed; of which the following appear deserving of notice:

1. *Carbon Ink.* Dissolve real Indian ink in common black ink; or add a small quantity of lamp-black previously heated to redness, and ground perfectly smooth, with a small portion of the ink.

2. *STEPHEN'S (patent) Carbon Ink.* Common soda of commerce is mixed with resinous matters (as shellac or resin), in about equal parts by weight. Water being added according to the strength required, the solution is boiled until the resin has become dissolved. Mix in a mortar with the requisite quantity of fine lamp-black, and add any suitable coloured solution.

3. *Shellac Ink, or COATHUPE'S Writing Fluid.* To 18 oz. of water add 1 oz. of powdered borax, and 2 oz. of bruised shellac, and boil them in a covered vessel, stirring them occasionally, till dissolved. Filter, when cold, through coarse filtering paper; add 1 oz. of mucilage; boil for a few minutes, adding sufficient finely-powdered indigo and lamp-black to colour it. Leave the mixture for 2 or 3 hours for the coarser particles to subside; pour it off from the dregs, and bottle it for use.

4. *Gluten Ink.* Dissolve wheat gluten, free from starch, in weak acetic acid of the strength of common vinegar, mix 10 grs. of lamp-black and 2 grs. of indigo with 4 oz. of the solution, and a drop or two of oil of cloves.

5. *HAUSSMANN'S.* Dissolve 1 part of genuine asphaltum with 4 parts of oil of turpentine, and sufficient lamp-black. If sufficient lamp-black be used to give it a suitable consistence, it may be used with types.

6. *BRACONNET'S Indelible Ink.* Take 20 parts of Dantzic potash, 10 of tanned leather parings, and 5 of sulphur; boil them in an iron pot with sufficient water to dryness;

then raise the heat, stirring the matter constantly, till the whole becomes soft, taking care that it does not ignite. Add sufficient water, and filter through cloth. It must be kept from the air. It flows freely from the pen, and resists many chemical agents; but it is not strictly indelible.

7. Dr NORMANDY'S *Indelible Ink*. Frankfort lamp-black 24 lbs., to be ground with mucilage, made by adding 24 lbs. of gum to 60 gallons of water, and the mixture filtered through a very coarse flannel; 4 lbs. of oxalic acid are then added, with as much decoction of cochineal and sulphate of indigo as will yield the shade of colour desired.

8. *Indian Ink*. Real lamp-black, produced by combustion of linseed oil, ground with gum, and infusion of galls. It is prepared both in a liquid and solid form, the latter being dried in the sun.

9. Gold is mixed with Indian ink, equal parts.

10. PUSCHER'S *Indelible Ink*. Dissolve 4 parts of aniline black in 16 parts by weight of alcohol, with 60 drops of strong hydrochloric acid, and dilute the solution with 90 parts by weight of water in which 6 parts of gum arabic have been previously dissolved. It is said not to act upon steel pens, nor be altered by alkalies or acids.

Indelible Ink, for printing Cotton and Linen fabrics intended for chlorine bleaching. 1 part of coal-tar mixed with 1 part of benzine, and 1-10th part of lamp-black. It can be made thicker or thinner by using more or less benzine.—*Pharm. Journal*.

Ink for writing on Zinc Labels. Horticultural Ink:

1. Dissolve 100 grs. of chloride of platinum in a pint of water. A little mucilage and lamp-black may be added.

2. Sal ammoniac 1 dr., verdigris 1 dr., lamp-black $\frac{1}{2}$ dr., water 10 drs.; mix.

Ink for writing on Steel or Tin Plate, or Sheet Zinc.

1. Mix 1 oz. of powdered sulphate of copper and $\frac{1}{2}$ oz. of powdered sal ammoniac with 2 oz. of diluted acetic acid; adding lamp-black or vermilion.

2. Dissolve 1 part of copper in 10 of nitric acid, and dilute with 10 parts of water.

White Marking for Black Bottles, in cellars. Grind flake white or sulphate of barium, with a little oil of

turpentine, and any light-coloured varnish, to a proper consistence.

Lithographic Ink. 1. LASTEYRIE'S;—Dried soap 1 oz., melt and add shell-lac 5 oz., then common soda 1 oz., mastic 1 oz., and lastly, lamp-black 3 drs. Melt, stir together, and, when completely melted, pour into moulds: to be used as Indian ink.

2. *Autographic.* White soap 100 parts, white wax 100, mutton suet 30, shell-lac 50, mastic 50, lamp-black 30 or 35. Melted as above.

3. *Lithographic Ink.* (WEISHAUPT.) Heat 40 parts of yellow wax until its vapour kindles on coming in contact with a burning match; then remove it from the fire, and add gradually, in small parts, Marseilles soap 22 parts, gum-lac 28 parts, and mastic 10 parts. Extinguish the flame, and incorporate perfectly with this mixture, lamp-black 9 parts. Then again heat until the vapour can be ignited, then remove it from the fire, and after the flame has been extinguished, pour it upon a stone. The mass is then cut into pieces.

4. *Crayons.* White wax 8 oz., white soap 2 oz., shell-lac 2 oz., lamp-black 3 tablespoonfuls. Melt the wax and soap with a brisk fire; stir in the lamp-black; allow the mixture to burn for half a minute, then extinguish the flame, and add the shell-lac by degrees, stirring continually. Put the mixture on the fire till it kindles, or nearly so. Extinguish the flame, let the mixture cool a little, and pour it into moulds.

Inks for Marking Linen. Some of these are used with types; others with a clean quill pen.

1. Sulphate of manganese 1 dr., water 1 dr., powdered sugar 2 drs., lamp-black $\frac{1}{2}$ dr. Triturate them together, and stamp in on the linen with types. When dry, wash the part with liquor potassæ; again dry, and wash with plenty of water.

2. Dr SMELLIE'S. Sulphate of iron 1 dr., linseed oil 1 oz.; vermilion $\frac{1}{2}$ oz.: grind perfectly smooth. Printers' ink is also used with type.

3. Heat to redness equal weights of black oxide of manganese and caustic potash, and mix it with an equal weight of pipe-clay, and sufficient water to give it a due

consistence. To be applied with types or stencils. It becomes brown and does not wash out. The following are used with a quill pen :

4. Nitrate of silver 100 grs., distilled water 1 oz., gum arabic 2 drs., sap green a scruple : dissolve. The linen is first to be wetted with the following *pounce*, dried and rubbed smooth, then written on by a clean quill or bone pen dipped in the ink. *Pounce* or *Mordant*. Subcarbonate of soda 1 oz., water 8 oz. [A great variety of recipes might be given, slightly differing from the above in the proportion of the ingredients, and in the colouring matter. GRAY directs 2 drs. of nitrate of silver, 6 drs. of water, and 2 of mucilage, and a pounce of 1 oz. of subcarbonate of soda in 16 of water, with a little sap green. Another form is—nitrate of silver 1 oz., distilled water 5 oz., powdered gum $1\frac{1}{2}$ oz., sap green sufficient to colour it. The linen to be first wetted with the following preparation:—Subcarbonate of soda 1 oz., water 6 oz., gum 1 oz. ; dissolve. Some add a little powdered bole to the preparation ; the object in colouring it being merely that the part which has been wetted may be more readily distinguished. The quantity of nitrate of silver should not be much less than 100 grs. in an ounce of ink ; the proportion of the other ingredients is of less importance. Some direct the addition of a drop or two of nitric acid.]

5. *Italian*. Moisten the linen with a solution of recently prepared chloride of tin, and write with a neutral solution of salt of gold.

Marking Ink, without Preparation. These inks merely require to have a hot iron passed over the part written on, and to be held pretty near the fire till the writing assumes a deep black colour.

1. Nitrate of silver 3 drs., water $1\frac{1}{2}$ oz. ; dissolve, and add as much strong liquid ammonia as will redissolve the precipitate formed by it ; add 2 drs. of mucilage, a little sap green, and water, if required, to make up the measure to 2 oz. A little ivory-black, Indian ink, or indigo, is sometimes used to colour it. Some recipes contain nitrate of copper in addition to nitrate of silver. Several recipes might be given, but they will all probably be superseded by PROFESSOR REDWOOD'S. In operating with ammonia and nitrate of silver, fulminating silver is sometimes unexpectedly formed, and may prove a source of danger.

In this respect, as well as others, PROFESSOR REDWOOD'S preparation claims a preference.

2. PROFESSOR REDWOOD'S. Rub together 1 oz. nitrate of silver, and 1 oz. of bitartrate of potash; add 4 oz. of liquor ammoniæ, and when dissolved mix in 6 drs. of white sugar, 10 drs. of powdered gum arabic, $\frac{1}{2}$ oz. of archil, and water to make up 6 oz. by measure. Instead of archil, $\frac{1}{4}$ oz. of sap green may be used to colour the ink; or 40 grs. of fine vegetable black, previously triturated with a little water or mucilage.

3. Rev. J. B. READE'S patent. This differs from the last in using tartaric acid instead of bitartrate of potash. The quantities may be 1 oz. of nitrate of silver, 3 drs. of tartaric acid, and the above quantities of the other ingredients. The use of tartaric acid he claims an exclusive right to.

4. Add to the last an ammoniacal solution of an oxide or salt of gold.—READE. This addition prevents its being acted on by cyanide of potassium, and some other agents which the silver ink fails to resist.

5. *Aniline Black Marking Ink.* The ink is prepared by means of two solutions, one of copper, the other of aniline, prepared as follows:

(1) *Copper Solution.* 8.52 grams of crystallized chloride of copper, 10.65 grams of chlorate of soda, and 5.35 grams of chloride of ammonium, are dissolved in 60 grams of water.

(2) *Aniline Solution.* 20 grams of hydrochlorate of aniline are dissolved in 30 grams of distilled water, and to this are added 20 grams of solution of gum arabic (1 part of gum to 2 of water), and 10 grams of glycerin.

By mixing in the cold, 4 parts of the aniline solution with 1 part of the copper solution, a greenish liquid is obtained, which can be employed directly for the marking; but as this liquid can only be preserved for a few days without decomposition it is advisable to keep the solution separately, until the ink is required for use. The ink may be used either with a pen or a stencil plate and brush; if it do not flow freely from the pen it may be diluted with a little water without fear of weakening the intensity of the colour. At first the writing appears of a pale green colour, but after exposure to the air it becomes black; or it may be changed to a black colour immediately by pass-

ing a hot iron over the back of the fabric, or heating it over the flame of a spirit lamp. As, however, a dry heat is apt to make the fibre saturated with the ink brittle, it is preferable to hold the marked fabric over a vessel containing water in full ebullition; the heat of the vapour is sufficient to determine almost immediately the reaction by which aniline black is formed. After the steaming the writing should be washed in hot soapsuds, which gives the ink a fine blue shade. The ink is not acted on by acids or alkalies, and if care be taken that the fibres are well saturated with it, there is no danger of its being removed by washing.—*Dingler's Journal*.

6. *Marking Ink without Silver*. Add a solution of caustic potash to a concentrated solution of chloride of copper in water as long as a precipitate is formed; allow this to subside, siphon off the clear fluid, and dissolve the precipitated oxide in the least possible quantity of water of ammonia, adding about 6 per cent. of dextrin to the solution. Articles marked with this ink require to be ironed before they are washed.

7. In addition to the above recipes, the following of M. HENRY may deserve attention in large establishments where economy is an object:—Take 1 oz. of iron filings and 3 oz. of vinegar, or diluted acetic acid. Mix the filings with half the vinegar, and agitate them continually until the mixture becomes thick, then add the rest of the vinegar and 1 oz. of water. Apply heat to assist the action; and when the iron is dissolved, add 3 oz. of sulphate of iron, and 1 oz. of gum previously dissolved in $\frac{1}{2}$ oz. of water; and mix the whole with a gentle heat. To be used with brush and stencil plates.

Crimson Marking Ink. Dissolve 1 oz. nitrate of silver and $1\frac{1}{2}$ oz. of carb. soda in crystals, separately in distilled water; mix the solutions, collect and wash the precipitate on a filter, introduce the washed precipitate, still moist, into a Wedgewood mortar, and add to it tartaric acid 2 drs. 40 grs., rubbing together till effervescence has ceased; dissolve carmine 6 grs. in liquor ammoniæ (882) 6 oz., and add to it the tartrate of silver, then mix in white sugar 6 drs., and powdered gum arabic 10 drs., and add as much distilled water as will make 6 oz.—*Pharm. Journal*.

Printing Ink. This is usually made by boiling linseed

oil in a large iron pot, setting fire to it, and letting it burn for half an hour or more. Various additions are made to it by some manufacturers, the use of which is not very evident. A viscid varnish is obtained, which is ground with lamp-black, vermilion, or other colouring matters, till perfectly smooth. $2\frac{1}{2}$ oz. of lamp-black are sufficient for each pound of varnish. See VARNISHES.

Printers' Ink from Resin Oil. Melt together 13 oz. of resin, 1 lb. of resin oil, and $1\frac{1}{2}$ oz. of soft soap; when cold, add lamp-black or other colouring matters.

Copper-plate Printing Ink. This is not rendered so viscid as the former, and is coloured with Frankfort black.

READE'S Patent Printing Inks. The *blue* consists of his soluble Prussian blue (see BLUE WRITING INK, further back) ground with oil as above. The *black*, by evaporating his black ink, and mixing the product with oil as usual. The *red* in the same manner, from his patent red ink.

Sympathetic or Secret Inks. The solutions used should be so nearly colourless that the writing cannot be seen till the agent is applied to render it visible.

1. Digest 1 oz. of zaffre, or oxide of cobalt, at a gentle heat, with $\frac{1}{2}$ oz. of nitro-hydrochloric acid till no more is dissolved, then add 1 oz. of common salt and 16 oz. of water. If this be written with, and the paper held to the fire, the writing becomes green, unless the cobalt should be quite pure, in which case it will be blue. The addition of a little nitrate of iron will then impart the property of becoming green. It is used in chemical landscapes for the foliage.

2. Put into a phial $\frac{1}{2}$ oz. of distilled water, 1 dr. of bromide of potassium, and 1 dr. of pure sulphate of copper. The solution is nearly colourless, but becomes brown when heated.

3. Boil oxide of cobalt in acetic acid. If a little common salt be added, the writing becomes green when heated; but with nitre it becomes a pale rose colour.

4. A solution of acetate of lead. Colourless, but becomes brown when exposed to sulphuretted hydrogen gas.

5. A weak solution of sulphate of copper. The writing becomes blue when exposed to the vapour of ammonia.

6. A solution of sulphate—or preferably, persulphate—of iron. It becomes black when washed with infusion of

galls; *blue*, by prussiate of potash. This constitutes colourless ink, which becomes visible when written with on paper containing galls, or tannin, or prussiate of potash.

7. Mix equal quantities of sulphate of copper and sal ammoniac, and dissolve in water. It becomes yellow when heated.

8. A weak solution of nitrate of mercury. Becomes black by heat.

9. Rice water, or any solution of starch. It becomes blue when washed over with an alcoholic solution of iodine.

10. Lemon-juice, milk, juice of onions, and some other liquids become black when the writing is held to the fire.

INK, to preserve from mouldiness. Add a small quantity of a solution of creasote in pyroligneous acid or rectified spirit, or of oil of cloves dissolved in spirit.

INSECTS, to kill. INSECT BITES, to cure. Camphor, and pepper, may be used to keep off moths. In Russia the powder of the flowers of a species of *Pyrethrum* is used as an insecticide. The powder of *Pyrethrum roseum*, or a diluted tincture, is used to prevent mosquitoes from biting in the East Indian islands. The powder of the root of *Acorus calamus* is also recommended. Liquid ammonia, sulphate of copper, or a mixture of toilet vinegar and glycerine are good as applications to bitten parts. Also a powder consisting of carbonate of lead 1 part, chalk 4 parts. Another method, said to be very effective, is to press the tube of a key firmly over the place stung; this prevents the immediate absorption of the poison, and presses the sting out.

IODATE OF POTASH. Fuse iodide of potassium in a capacious Hessian crucible. remove it from the fire and add to it, while still semi-fluid, successive portions of pulverized chlorate of potash, stirring after each addition, till no further action takes place. One part of iodide of potassium will require $1\frac{1}{2}$ of the chlorate. Wash the residuum in warm water, which leaves only iodate of potash.

IODIDE OF POTASSIUM. See **POTASSII IODIDUM, Pocket Formulary.**

IODINE. See *Pocket Formulary.* Other methods of obtaining it are the following :

To the mother liquor of kelp (after the crystallizable salts have been separated) add sulphuric acid to render the liquor sour. Introduce the acid liquor into a leaden still, heat to 140° F., add binoxide of manganese, and lute on with pipe-clay a leaden head, fitted to a series of spherical glass condensers, each having two mouths opposite each other, and inserted the one into the other. A stopper in the head of the still allows the contents to be occasionally inspected, and additions of acid or oxide made, if necessary. See WAGNER'S 'Chemical Technology' for a drawing of the apparatus. SOUBEIRAN proposes to add sulphate of copper to the ley, which precipitates half the iodine. He then decants the clear liquor, and adds more sulphate of copper with some iron filings. An iodide of copper is formed, which is separated from the iron filings and suspended in the liquor by agitation, collected on the filter, and heated with oxide of manganese and sulphuric acid. Glasgow is the principal seat of the Iodine manufacture.

IRON LIQUOR. See DYES, further back.

IRON, PERSULPHATE, SOLUTION OF. LIQ. FERRI PERSULPH.
See *Pocket Formulary*.

ISINGLASS. The air-bags, or sounds, of several kinds of fishes, washed, dried, and otherwise prepared. They are either dried without opening (purse, pipe, and lump isinglass), or opened and not folded (leaf and honeycomb isinglass), or folded (book isinglass), or twisted into the shape of a lyre or horse-shoe (short and long staple). The picked or cut isinglass of the shops consists of the lumps of staple isinglass, picked in shreds by women and children, or cut by machines. The leaf isinglass is sometimes rolled out into thin plates (ribbon and rolled leaf isinglass). The inner membrane, which is insoluble, is removed from the open air-bags in the best kinds. The Russian isinglass, which is most esteemed, is made from the air-bags of several species of ACIPENSER (*sturgeon*); particularly A. HUSO (the *Beluga*); A. GULDENSTADTII (the *Osseter*); A. RUTHENUS (the *Sterlet*); A. STELLATUS (the *Sewruga*); and also from the SILURUS GLANIS (the *Som*), which yields the Samovey isinglass. Brazilian and East Indian isinglass are of inferior quality; it is not certainly known from what genera or species of fish they are obtained. New York isinglass is the air-

bladder of the *common hake*, macerated in water and rolled out into ribbons. The sounds of the cod yield an inferior kind. Prepared sole skins are used as a chief substitute for isinglass. See Dr PEREIRA'S 'Elements' for the description of each variety.

IVORY, BLACK. Burn shavings and waste pieces of ivory from the ivory turners, in a covered crucible, till no more smoke issues. Cover it closely while cooling. It should be afterwards washed with diluted hydrochloric acid, then with water till no longer acid, dried, and again heated in a covered crucible. It is of a deeper colour than bone-black, and is used as a pigment, a tooth-powder, and to decolorize syrups and other liquids.

IVORY, FLEXIBLE. The pieces of ivory or bone, already manufactured into the shape required, are to be steeped for some time in dilute hydrochloric acid, until they have lost their earthy parts so far as to become yellowish, flexible, and elastic. When dry they become again inflexible, but their flexibility may at any time be restored by steeping them in water. In this manner flexible tubes, probes, bougies, &c., may be constructed.

IVORY, TO STAIN. Ivory is stained with the usual dyeing materials; it should be first steeped in the mordant and afterwards in the hot colour. Nitro-chloride of tin is the mordant for red, with decoction of brazil or cochineal; for yellow, with fustic; for violet, with logwood. After being plunged in hot liquor it should be placed in cold water. A black stain is given by nitrate of silver.

Ivory may be gilded by immersing it in a fresh solution of proto-sulphate of iron, and afterwards in solution of chloride of gold. It may be bleached by solution of sulphurous acid or peroxide of hydrogen.

IVORY AND BONE, TO BLEACH. 1. M. CLOEZ recommends the ivory or bones to be immersed in turpentine, and exposed for three or four days to sunlight. The object to be bleached should be kept an eighth or a fourth of an inch above the bottom of the bath by means of zinc supports.

2. M. PUSCHER says this may be easily and permanently effected by immersing the pieces in the following solution:—25 grammes of pure zinc white are covered with 40 cc. of water and 50 cc. of concentrated sulphuric acid gradually added; 150 cc. of hot water are then poured in, and with constant stirring, ammonia, in just sufficient quantity

to redissolve the precipitated zinc hydrate at first formed, avoiding, however, an excess. A solution of copper sulphate is then added until the yellow tint of the liquid has changed to a bluish white. The objects must remain in the liquid several days, after which they are to be well rinsed with water. Old discoloured bone or ivory bleached by this process can be polished.

IVORY, ARTIFICIAL. Let a paste be made of isinglass, egg-shell in very fine powder, and brandy. Give it the desired colour, and pour it while warm into oiled moulds. Leave the paste in the moulds until it becomes hard. See **IVORY, TO STAIN.**

JAPAN. See **VARNISHES.**

JELLIES. See **DIETETIC ARTICLES.**

KID-GLOVE CLEANER. Add 15 drops of strongest solution of ammonia to spirit of turpentine $\frac{1}{2}$ pint. Having fitted the gloves on wooden hands, apply this mixture with a brush. Follow up this application with some fine pumice powder. Rub with some flannel or sponge dipped in the mixture. Rub off the pumice, and repeat the same process twice or thrice. Hang in the air to dry, and, when dry, place in a drawer with some scent. Benzol is largely used as a glove cleaner. See **GANTEINE.**

KOUMISS. An alcoholic drink, made by fermenting the milk of mares, and beating it up with a whisk. It may be regarded as a somewhat coarse form of "rum and milk."

KYAN'S SOLUTION, for preventing the dry rot. Dissolve 1 lb. of corrosive sublimate in 5 gallons of water.

LABARRAQUE'S CHLORO-SODAIC LIQUOR is nearly identical with the **LIQUOR SODÆ CHLORATÆ** of the British Pharmacopœia. It is made by passing the chlorine gas from 2 oz. black oxide of manganese, and 8 oz. of hydrochloric acid, into a solution of 15 oz. of crystallized carbonate of soda in 3 pints of water; or sufficient to bring it to the density of 12° Baumé, or 1.09 specific gravity.

LABELS, INDESTRUCTIBLE, for acid bottles, &c. Paint the required label on the bottle with a camel-hair brush dipped in a solution of asphalt in benzol.

LABELS, GOLD, TO ATTACH. Use ordinary gum mucilage, in which a few pieces of gum tragacanth have been dissolved. After the labels have dried on they must be first sized over with a solution of isinglass in water (about $\frac{1}{2}$ dr. to an

oz.), and when that is dry, varnished over with a solution of Canada balsam in turpentine (1 dr. in an oz.). A thin coating should be applied, and when this is dry another coating, and so on.

LAC, PREPARATIONS OF. Stick-lac consists of the twigs of several kinds of trees encrusted with a resinous matter produced by the puncture of an insect (the *coccus lacca*). This, triturated with water and dried, forms seed-lac. The seed-lac heated and pressed in cotton bags forms shell-lac. Lac-dye is the colouring matter extracted from stick-lac by water, evaporated to dryness with the addition of earthy matters, and formed into square cakes. Seed-lac and shell-lac are chiefly used* in varnishes, dissolved in rectified spirit, methylated spirit, or rectified wood naphtha. The alcoholic solution is rendered paler, so that it may be used for polishing light-coloured woods, by digesting it in the sun, or near a fire for 2 or 3 weeks, with good animal charcoal, and then filtering it through paper in a funnel heated with hot water. Shell-lac may be bleached by dissolving it in a solution of potash or soda, and passing chlorine into the solution. The precipitated lac is collected and well washed. KASTNER directs 3 parts of carbonate of potash to be dissolved in 24 of water, and 3 of lime added, and the whole digested in a close vessel for 24 hours. The clear liquor is poured off, and boiled with 4 parts of shell-lac. When cold, dilute with four times its bulk of water, and filter; then add chloride of lime, and afterwards diluted hydrochloric acid.

LACQUERS. See VARNISHES.

LAKE LIQUOR. Boil 1 oz. each of cochineal and salt of tartar in 8 oz. of water, then add 1 oz. of cream of tartar, and the same of alum.

LAKES. These consist of vegetable colours in combination with alumina. Alum is usually added to an infusion or decoction of the colouring ingredient, and afterwards potash added, which throws down the colouring matter combined with alumina. Some of the lakes are noticed under PIGMENTS.

LEMON JUICE, FACTITIOUS. Dissolve 4 oz. of citric acid in 3 pints of water, with 8 drops of essence of lemon, rubbed with the acid, or dissolved in a little spirit. After stand-

* Methylated spirit is now almost invariably used for this purpose.

ing a few days, filter it, and preserve it in well-closed bottles.

LENSES, EXTEMPORANEOUS. FRANCIS. Procure a piece of thin platinum wire and twine it once or twice round a pin's point, so as to form a minute ring with a handle to it. Break up a piece of flint glass into fragments a little larger than a mustard seed; place one of these pieces on the ring of wire, and hold it in the point of the flame of a candle or gas-light, or better still, of a Bunsen burner. The glass will melt and assume a complete lens-like or globular form. Let it cool gradually, and keep it for mounting. Others are to be made in the same manner; and if the operation be carefully conducted, but very few will be imperfect. The smaller the drop melted, the higher in general will be its magnifying power. It may be mounted by placing it between two pieces of brass which have corresponding circular holes cut in them of such a size as to hold the edge of the lens. They are then to be cemented together.

LINSEED OIL, CLARIFIED, FOR VARNISHES.—Heat in a copper boiler 50 gallons of linseed oil to 280° F.; add 2½ lbs. of calcined white vitriol, and keep the oil at the above temperature for half an hour; then remove it from the fire, and in 24 hours decant the clear oil, which should stand for a few weeks before it is used for varnish.

LINSEED OIL, REFINED. (WILKS' PATENT.) Into 236 gallons of oil pour 6 lbs. of oil of vitriol, and stir them together for 3 hours; then add 6 lbs. of fullers' earth well mixed with 14 lbs. of hot lime, and stir for 3 hours. Put the oil into a copper boiler, with an equal quantity of water, and boil for 3 hours; then extinguish the fire, and when the materials are cold draw off the water, and let the oil stand to settle for a few weeks before using.

LIQUORICE, PURIFIED EXTRACT OF. Italian or Spanish juice may be purified by the following method:—Take a sugar-mould, close the vent-hole with a stopper, place inside it some coarse tow, and over this some clean straw, laid crossways in layers of an inch each, then the sticks of liquorice placed upright, and packed closely in the mould with chopped straw cut rather long. When this arrangement is completed to within an inch of the brim, pour water over the liquorice, allow it to remain for 24 hours,

then draw it off, and add more. The liquor, on evaporation, yields an extract perfectly soluble in water.

LITMUS. A preparation of some kind of lichen, probably *Lecanora tartarea*, or *Roccella tinctoria*, or both. It is prepared by a process similar to that used for archil; but it is moistened by a mixture of carbonates of ammonia and potash, or soda, the addition of the carbonates of ammonia, or of potash or soda, being essential to the development of the blue colour. It is made up with chalk or plaster of Paris into small cakes for the market. See TESTS.

LOZENGES. See TROCHISCI, *Pocket Formulary*, and LOZENGES, under PATENT MEDICINES, in this volume.

LUBRICATING COMPOUNDS. See ANTI-ATTRITION. The French compound term *Liard* is thus made:—Into 50 parts of finest rape oil put 1 part of caoutchouc cut small, and apply heat until it is nearly all dissolved.

MANKETRICK'S LUBRICATING COMPOUND consists of caoutchouc (dissolved in spirit of turpentine) 4 lbs., common soda 10 lbs., glue 1 lb., oil 10 gallons, water 10 gallons. Dissolve the soda and glue in the water by heat, then add the oil, and lastly the caoutchouc, stirring them until perfectly incorporated.

LUCIFERS. See MATCHES, further on.

LUMINOUS PHIALS. Nearly fill a bottle with olive or almond oil, and heat it in a water-bath. Drop into it small slices of phosphorus so long as it is dissolved. Let the solution cool, and pour off the oil from the undissolved phosphorus into clean dry phials, which should not be quite filled. When uncorked they emit light.

LUTES. See CEMENTS.

MANURES, ARTIFICIAL. These constitute a new and important branch of manufacture; but a few of the more simple and readily prepared kinds are all that can be noticed here.

Powder for Coating Seeds. Fine bone-dust 20 parts, gypsum 1 part. The seeds are steeped in water from the dunghill, then strewed over with the powder, so that each shall receive a layer of it. They are afterwards dried.

Sulphated Bones. See *Bones, Sulphated*. A usual proportion is 33 lbs. of sulphuric acid to 1 cwt. of bones.

Saline Mixture, as a top dressing for potatoes. Equal weights of nitrate of soda and dry sulphate of soda $1\frac{1}{2}$ cwt. to an acre.

Mr HUXTABLE'S *Mixture*. Bone-dust 4 cwt., gypsum 4 cwt., salt 2 cwt., ashes $\frac{1}{2}$ quarters, wood ashes 30 bushels.

Another Saline Mixture. Sulphate of ammonia 42 lbs., sulphate of lime 56 lbs., sulphate of potash 56 lbs., carbonate of magnesia 14 lbs., salt 56 lbs., to 1 acre.

Dr ANDERSON'S *Manure for Clover*. Sulphate of ammonia 98 lbs., gypsum 172 lbs., sulphate of potash 174 lbs., sulphate of soda 333 lbs., sulphate of magnesia 246 lbs., sulphuric acid 98 lbs., saltpetre 202 lbs., common salt 107 lbs., chloride of potassium 149 lbs.

Dr JOHNSTONE'S *Substitute for Guano*. Bone-dust 7 bushels, sulphate of ammonia 100 lbs., wood ashes 20 lbs., salt 100 lbs., dry sulphate of soda 11 lbs.

To Promote the Blowing of Flowers. See further back.

MARBLE TO CLEAN. Mix soft soap, solution of potash, and slaked lime, to a paste; spread it over the marble, and leave it for a day or two. Then wash it off.

MARBLE, TO STAIN. Make the marble hot, and pour on it the coloured liquid, also made hot. The stains usually employed are archil, solution of indigo, solution of verdigris, decoction of Brazil-wood, logwood, and sulphate of iron, tincture of dragons' blood, &c. But the most penetrating medium is wax, which may be coloured with alkanet, annotto, verdigris, &c.

MARINE GLUE. See GLUE.

MARINE SOAP. See SOAP, MARINE, further on.

MATCHES FOR INSTANTANEOUS LIGHT. 1. *Chlorate of Potash Matches (without sulphur)*. Chlorate of potash, separately powdered, 6 drs., vermilion 1 dr., lycopodium 1 dr., fine flour 2 drs.; mix carefully the chlorate with the flour and lycopodium, avoiding much friction, then add the vermilion, and mix the whole with a mucilage made with—1 dr. powdered gum arabic, 10 grs. of tragacanth, 2 drs. of flour, and 4 oz. of hot water; mix, add sufficient water to bring it to a proper consistence, and dip in it the wood, previously dipped in a solution of 1 oz. of gum thus, and $\frac{1}{2}$ oz. of camphor, in 6 oz. of oil of turpentine.

2. *With sulphur*. Chlorate of potash 9 grs., sulphur 2 grs., sugar 3 grs., vermilion 1 gr., flour 2 grs., spirit of wine q. s. The chlorate of potash, &c., must be separately reduced to powder, and the whole mixed with as little friction as possible. The wood should be previously

prepared as above, or with camphorated spirit. [These are ignited by dipping them in sulphuric acid, and instantly withdrawing them. The acid should be absorbed by abestos.] They are now become obsolete, having given place to—

Lucifer Matches. These contain phosphorus in a finely divided state, to which it is reduced by agitating it in some warm solution of gum or glue, then adding the other ingredients, so as to form a paste, into which the wood or card is dipped. It is said that urine and artificial urea have the property of readily dividing phosphorus when warmed and agitated together. The following are some of the published recipes :

1. Form 6 parts of glue into a smooth jelly, and rub with it 4 parts of phosphorus, at a temperature of 140° or 150° F.; add 10 parts of nitre, 5 of red ochre, and 2 of fine smalts. The matches are first dipped in melted wax to the depth of 1-10th of an inch, first rubbing their ends on a hot iron plate.

2. *Noiseless Congreves.* Dr BOETTGER. Triturate 9 parts of phosphorus with a solution of 16 parts of gum, and add 14 parts of nitre and 16 of vermilion.

3. Glue 6 parts, phosphorus 4, nitre 10, red lead 5, smalts 2; the glue is soaked in water for 24 hours, then liquefied in a warm mortar, and the phosphorus added, taking care that the temperature is not above 167° F.

4. Glue 21, phosphorus 17, nitre 38, red lead 24: proceed as before.

Promethean Matches. These consist of a composition similar to that of the chlorate of potash matches, inclosed at the end of a paper spill, with a minute glass bulb filled with oil of vitriol in the centre of the composition. When struck the vessel of acid is broken, and kindles the match.

METHYLATED SPIRIT. Spirit of wine mixed with 10 per cent. of rectified wood spirit is allowed to be sold under this name, free of the excise duty. It is forbidden to be used for drinking purposes, or for the preparations of medicinal tinctures. It is extensively employed for burning in lamps, and for the preparation of polishes, varnishes, and lacquers, as well as for many operations by the chemist. The law forbids methylated spirit or any preparation containing it to be used internally.

MICROSCOPE, PREPARATIONS REQUIRED IN THE
WORKING OF.

1. Alcohol, of various strengths.
 2. Arsenious acid.
 3. Benzol, pure.
 4. Bichloride of mercury.
 5. Ether, to dissolve oil-globules.
 6. Nitric acid, 1 part of strong acid to 5 of water.
 7. Sulphuric acid, 1 to 5.
 8. Hydrochloric acid.
 9. Acetic acid, glacial, and dilute (1 to 5).
 10. Chromic acid, very dilute, to harden tissues.
 11. Canada balsam, dissolved in chloroform or benzol.
 12. Carbolic acid.
 13. Carmine.
 14. Chloroform.
 15. Chromate of potash.
 16. Bichromate of potash.
 17. The coal-tar colours.
 18. Cloves, essential oil of.
 19. Dammar, gum, dissolved in chloroform or benzol.
 20. Glycerin.
 21. Glycerin jelly.
 22. Lemon, essential oil of.
 23. Logwood.
 24. Osmic acid.
 25. Picric acid.
 26. Prussian blue, soluble.
 27. Schultz's test.
 28. Solution of potash, saturated, and dilute (1 to 10).
 29. Solution of soda, 25 grs. of fused soda in 1 oz.
 30. Ammonia (1 part of the strongest solution to 3 of water).
 31. Nitrate of baryta, a cold saturated solution of.
 32. Nitrate of silver (120 grs. in 2 oz.). These two are tests for the mineral acids.
 33. Oxalate of ammonia, in solution. Test for lime.
 34. Solution of iodine saturated, *i. e.* 1 to 7000 water. Another solution is, 1 gr. of iodine, and 3 of iodide of potassium, in 1 oz. of distilled water.
- CEMENTS. 1. *Brunswick Black*. Boil together $\frac{1}{4}$ lb. of foreign asphaltum, and $4\frac{1}{2}$ oz. of linseed oil (previously

thickened with litharge), then mix to a proper consistence with oil of turpentine (about 1 pint).

2. *Gold Size*. Boil 25 parts of linseed oil with 1 of minium and 1-3rd part of umber for 3 hours; pour off the clear fluid, and mix with equal parts of powdered white lead and yellow ochre, added in small successive portions. Then boil well the whole again, and pour off the clear fluid. It dries slowly but firmly. Both this and the last are dissolved by turpentine.

3. *Goadby's Marine Glue*. Dissolve separately in coal naphtha equal parts of shell-lac and india rubber. Mix thoroughly with heat.

4. *Sealing-wax Varnish*. Dissolve the best sealing-wax in enough strong spirit of wine to make it the proper consistence. This is brittle.

5. *Canada Balsam*. This dries spontaneously.

Solutions of shell-lac, gum, and various other cements, and glues (which see), are employed by microscopic manipulators.

Preservative Fluids. Canada balsam, spirit and water, glycerin, solution of gelatin, saturated solutions of alum, chloride of zinc, and chloride of calcium, are all used to preserve microscopic objects. The following formulæ will be found useful:

1. *Goadby's Solution*. Bay salt 4 oz., alum 2 oz., corrosive sublimate 4 grs., boiling water 4 pints: mix and filter. It may often be more diluted.

2. *Thwaites' Fluid*. Mix spirit of wine 1 oz., with creasote sufficient to saturate it; rub up with chalk to form a thin paste, and mix gradually with water 16 oz. To this may be added an equal quantity of water saturated with camphor.

3. *Simple Creasote Solution*. Dissolve creasote 1 dr. in pyroligneous acid 1 dr., and mix gradually with cold water 1 pint.

4. *Passini's Solution*. For blood-globules, nerves, and white tissues generally. Chloride of mercury 1 part, chloride of sodium 2 parts, glycerin 13 parts, distilled water 113 parts.

Glycerin Jelly for Microscopic Mounting. Soak any quantity of good clear gelatin in cold water for three or four hours. Pour off the superfluous water, and melt the gelatin at a gentle heat; when melted, filter through

flannel, and to the filtrate add an equal quantity of PRICE'S glycerin. The above forms a good firm jelly, requiring little trouble in securing the cover.—*Pharm. Journal.*

MILK, CONDENSED. Cows' milk evaporated down *in vacuo*, and containing about one third its weight of sugar, when it is intended to be kept for any time. If required for early use it contains no sugar.

MILK TO PRESERVE. Dissolve 6 oz. of boracic acid in a gallon of warm water; and add half a gill of this mixture to every gallon of milk. By BETHEL'S patent the milk or cream is first scalded, and when cold strongly charged with carbonic acid gas by means of a soda-water machine.

MINERAL, CHAMELEON. See CHAMELEON MINERAL.

MODELLING, CLAY FOR. (BARRESWIL.) Knead up clay to the proper consistence with glycerin.

MOIRÉE MÉTALLIQUE. A method for ornamenting the surface of tin plate by acids. The plates are washed with an alkaline solution, then in water, heated, and sponged or sprinkled with the acid solution. The appearance varies with the degree of heat and the nature and strength of the acids employed. The plates, after the application of the acids, are plunged into water slightly acidulated, dried, and covered with white or coloured varnishes. The following are some of the acid mixtures used:—Nitro-hydrochloric acid, in different degrees of dilution; sulphuric acid with 5 parts of water; 1 part of sulphuric acid, 2 of hydrochloric acid, and 8 of water; a strong solution of citric acid; 1 part nitric acid, 2 sulphuric, and 18 of water. Solution of potash is also used.

MORDANTS. See DYES, further back.

MULTUM. A name given to a compound of liquorice and quassia, improperly sold by druggists to brewers.

NITRATE OF BARIUM. This may be made from the carbonate by dissolving it in dilute nitric acid, evaporating, and crystallizing; but more cheaply from the sulphate of baryta, by converting it into a soluble sulphide, by heating it with charcoal, and decomposing the filtered solution with nitric acid. M. WEISS recommends mixing the pulverized sulphate of baryta ("cawk or heavy spar") with 1-8th of charcoal and 1-4th of flour, heating it in a covered crucible, pulverizing the product and forming it into balls, with 1-8th of charcoal and a little water, and

again heating them placed between layers of charcoal. Hot water extracts the sulphide, which crystallizes from the filtered solution. By decomposing this by nitric acid (avoiding the gas which escapes) the nitrate is obtained. The other salts of baryta may be obtained in a similar manner.

NITRATE OF SILVER. It may be prepared from impure silver by the following process:—Dissolve the silver in nitric acid, add common salt till no more silver remains in solution. Wash the precipitate thoroughly; then add water and a very little hydrochloric acid, and introduce some pieces of zinc; let them remain together 24 hours, stirring frequently. Remove the zinc, and wash the reduced silver thoroughly. Again dissolve it in nitric acid, diluted with 2 or 3 parts of water; filter and evaporate, that it may crystallize. See ARGENTI NITRAS, *Pocket Formulary*.

NITRATE OF STRONTIAN. This may be obtained from the native carbonate of strontian, or more cheaply from the native sulphate, by the processes employed for Nitrate of Barium.

NITRIC AND NITRO-HYDROCHLORIC ACIDS. See ACIDS.

NITRITE OF POTASH. It is obtained mixed with a little nitre and potash by heating nitre to redness. To purify the residuum, dissolve it in boiling water, set aside for 24 hours, pour off the liquid from the deposited nitre, neutralize the free alkali with acetic acid, and add twice its volume of alcohol. In a few hours more nitrate crystallizes, and the liquid separates into two layers: the upper is alcoholic solution of acetate of potash, the lower is solution of nitrite of potash, which may be evaporated to dryness or kept in solution. CORENWINDER passes nitrous acid gas, formed by acting on 1 part of starch with 10 of nitric acid, through a solution of caustic potash, sp. gr. 1.38, until it becomes acid; then adds a little caustic potash, so as to render it distinctly alkaline.

NITRO-PRUSSIDE OF SODIUM. To 213 parts of powdered ferro-cyanide of potash, in a porcelain basin, add 450 parts of nitric acid of 1.42 density (or 337½ parts at 1.50), adding all the acid at once. When dissolved, transfer to a bolt-head, and digest in a water-bath, until the solution precipitates salts of protoxide of iron of a slate colour. Neutralize, when cold, with a cold solution of carbonate of soda; then boil, and separate the precipitate

by filtration. Evaporate the liquid again, filter, and allow the nitrates of potash and soda to crystallize out. Evaporate the liquid again, and remove the prismatic crystals of nitro-prusside as they form. They may be dissolved in hot water and recrystallized by cooling.

NOVARGENT. This is said to consist of a solution of freshly precipitated chloride of silver in hyposulphite of soda (or, according to the *Pharmaceutical Journal*, of oxide of silver in cyanide of potassium), mixed with prepared chalk.

OILS, PURIFICATION AND BLEACHING OF. Fish and other fat oils are improved in smell and colour by passing hot air or steam through them. **DUNN'S** method is to heat the oil by steam to 170° or 200° F., and force a current of air through it, under a chimney, till it is bleached and purified. **MR CAMERON'S** method of bleaching palm oil is to keep it at 230° F., with continual agitation by passing into it high pressure steam, through leaden pipes of two inches diameter. Four tons of oil requires 10 hours' steaming. Palm oil is bleached by chloride of lime. Take from 7 to 14 lbs. of chloride of lime, triturate it with 12 times the quantity of water, gradually added, so as to form a smooth cream. Liquefy 112 lbs. of palm oil, remove it from the fire, add the solution of chloride of lime, and stir well with a wooden stirrer. Allow it to cool, and when it has become solid, break into small fragments, and expose it to the air for 2 or 3 weeks. Then put it into a cast-iron boiler lined with lead, and add sulphuric acid in equal weight to the chloride of lime, diluted with 20 parts of water. Boil with a moderate heat till the oil drops clear from the stirrer; then let it cool.

To remove the fetor from fish oils, treat them in the same way (except the exposing to the air) using only 1 lb. chloride of lime to 112 lbs. of oil. It does not remove the natural smell of the oil.

Freshly burnt animal charcoal removes to some extent the colour and smell of most kinds of oil; but its effects are limited.

Calcined magnesia has been used to deprive oils of their rancidity.

MR GRIESELER states that the addition of a few drops of nitric ether will prevent rancidity in oils.

MR WATT'S patented method of bleaching oil is by chromic acid. For palm oil it is thus used:—The oil

is heated in a steam vessel, allowed to settle and cool down to 130° F., then removed into wooden vessels, care being taken that no water or sediment accompanies it. For a ton of palm oil, make a saturated solution of 25 lbs. of bichromate of potash; add 8 lbs. of sulphuric acid, and 50 lbs. hydrochloric acid (or an equivalent quantity of salt and sulphuric acid). Add a portion of the mixture to the oil, constantly stirring till the mixture becomes of a light-green colour. If not sufficiently decoloured, add more of the mixture. Let the oil settle for half an hour, then pump it into a wooden vat, boil it for a few minutes with fresh water by means of a steam-pipe, and let it settle. For linseed, rape, and mustard oil, a dilute solution of chromic acid is used, with a little hydrochloric acid; for olive, almond, and castor oil, no hydrochloric acid is required. Fish oils and fats are first boiled in a steam apparatus with a weak soda ley ($\frac{1}{3}$ lb. soda for every ton of fat) for half an hour; then $\frac{1}{2}$ lb. sulphuric acid, diluted with 3 lbs. of water, is added; the whole is boiled for 15 minutes, and allowed to settle for an hour or more, when the water and sediment are drawn off, and the oil further bleached by a solution of 4 lbs. of bichromate of potash and 2 lbs. of sulphuric acid properly diluted.

Mr DAVIDSON treats whale oil first with a solution of tan, next with water and chloride of lime, and lastly with diluted sulphuric acid and warm water. Rape and other seed oils are also refined by means of sulphuric acid and twice as much water. Mr GRAY directs 2 lbs. of oil of vitriol to 112 lbs. of oil. The oil should be carefully washed from the acid, and filtered.

Mr BANCROFT'S process for refining common olive oil, lard, oil, &c., for lubricating purposes, is to agitate them with from $3\frac{1}{2}$ to 8 per cent. of caustic soda ley, of 1.2 specific gravity. If on trial a small quantity of the ley be found to settle clear at the bottom, enough has been added. The oil is allowed to rest for 24 hours, for the soapy matter to subside; the supernatant oil is then filtered.

Another plan of purifying oils (especially lamp oils) is to agitate them with a strong solution of common salt.

The above methods of treating oil are of doubtful propriety in reference to such as are to be used as medicines. Oils which have been so carefully prepared from sound and fresh materials as to require no purification should be

selected for this purpose. This is especially important in reference to cod-liver oil. See LINSSEED OIL REFINED.

OIL FOR MACHINERY. Sperm oil, palm oil, olive oil, and petroleum heavy oil, are used. Care should be taken that they are not adulterated. For compound lubricants see ANTI-ATTRITION, and LUBRICATING COMPOUNDS.

OLEIN. This may be prepared by boiling fine olive oil with absolute alcohol, and evaporating the solution.

OXYGENATED WATER, OR DEUTOXIDE, OR PEROXIDE OF HYDROGEN. THENARD'S oxygenated water is thus made:—Expose fragments of perfectly pure baryta to a current of oxygen gas, in a green glass tube heated to a dull redness, to form a deutoxide of barium. To 7 oz. of water add as much pure hydrochloric acid as will dissolve 4 drs. of baryta; and carefully cool the mixture by means of ice, then add to it by degrees, 3 drs. of pulverized deutoxide of barium, and when this is dissolved, add sulphuric acid, drop by drop, till the baryta falls down in the state of sulphate. Then add more deutoxide, and precipitate by sulphuric acid as before. Then filter the solution; and repeat the solution and precipitation several times, till about 3 oz. of deutoxide of barium are used, filtering the liquid after every second repetition. Sulphate of silver is then added to remove the hydrochloric acid, and afterwards pure baryta, to throw down the sulphuric acid, and a few drops of diluted sulphuric acid to remove any excess of baryta. See HYDROGENII PEROXIDUM, *Pocket Formulary*.

This energetic compound must not be confounded with the oxygen water formed by impregnating water with oxygen gas; nor with the oxygenous aerated water of SEARLE, which is water strongly charged with protoxide of nitrogen.

OXYGEN GAS. See GASES.

PAPER, COPYING. Mix lard with black lead or lamp-black, into a stiff paste, rub it over writing paper with a flannel, and wipe off the superfluous quantity with a soft rag. These sheets alternated with writing paper and written on with a solid pen, produce 2 or 3 copies of a letter at once.

Lithographic Paper. Give the paper 3 coats of thin size, 1 of starch, and 1 of solution of gamboge. Each to be applied with a sponge, and allowed to dry before the next is applied.

Hydrographic Paper. This name has been given to

paper which may be written on with water. It may be made by rubbing paper over with a mixture of finely-powdered galls and sulphate of iron heated till it becomes white. The powder may be pressed into the paper by passing it between rollers or passing a heavy iron over it. A mixture of dried sulphate of iron and ferro-cyanide of potash may be used for blue writing. Or the paper may be imbued with a strong solution of one ingredient thoroughly dried, and the other applied in powder. Paper which has been wet with a solution of ferro-cyanide of potash also serves for writing on with a colourless solution of persulphate of iron.

Iridescent Paper. Nut-galls 8 parts, sulphate of iron 5, sal ammoniac 1, sulphate of indigo 1, gum arabic 1-8th. To be boiled in water, and the paper washed with it and exposed to ammonia.

Parchment Paper. Immerse blotting paper for a second or two in dilute sulphuric acid, then rinse repeatedly in several changes of water, and hang it up to dry. When dry, if it has a crumpled or wrinkled appearance, this can be removed by wetting it, and straining it upon a glass plate, the edges being made to wrap over the plate.

Photographic Paper. See PHOTOGRAPHY, further on.

Tracing Paper. Paper well wetted with Canada balsam and camphine, and dried. Another kind is made with nut oil, and oil of turpentine; the paper is moistened with it, and then rubbed with flour. A temporary tracing paper is made by moistening paper with pure alcohol; it must be used while wet.

Waxed Paper. Lay the paper on a clean hot iron plate, and rub it over with a piece of white wax inclosed in muslin.

Oiled Paper as a substitute for oiled silk. Boiled linseed oil is reboiled with litharge, acetate of lead, sulphate of zinc, and burnt umber, an ounce of each to a gallon. The sheet of paper being laid on a square board, it is well covered with the clear oil separated from the deposit. The first sheet is covered on both sides, the second, placed on this, receives one coating, and so on to 20, or 50. Separate and hang up to dry.

FIREPROOFING FOR PAPER. Dip in a strong solution of alum, and then dry it. Should the paper be extra thick, the same process may be repeated.

PAPER PASTE. Boil white paper in water for 5 hours; then pour off the water, and pound the pulp in a mortar; pass it through a sieve, and mix with some gum water, or isinglass glue. It is used in modelling by artists and architects.

PAPIER-MACHÉ. A plastic material, formed of cuttings of white or brown paper boiled in water, and beaten to a paste in a mortar, and then mixed with a solution of gum arabic in size, to give tenacity. It is variously manufactured by being pressed into oiled moulds, afterwards dried, covered with a mixture of size and lamp-black, and varnished.

PAPYRINE. Dip white unsized paper for $\frac{1}{2}$ a minute in strong sulphuric acid, and afterwards in water containing a little ammonia. When dried it has the toughness and appearance of parchment. See **PARCHMENT PAPER**, further back.

PARAFFINS. Liquid and solid paraffins are obtained by distillation of peat, petroleum, ozokerit, brown coal, Boghead shale, &c. By Young's process, in which Boghead coal is heated in tubes or retorts, a crude oil is first obtained, which, after purification by redistillation, followed by subsequent treatment with sulphuric acid and exposure to the action of caustic soda, is submitted to the process of fractional distillation. The first elevation of temperature drives over the lighter and more volatile portions, which, when purified by another distillation, yield the fluid known as "paraffin naphtha," a product used as a substitute for turpentine and as a solvent for india rubber. At a much higher temperature the burning oil (the paraffin oil of commerce) comes over. It is a perfectly safe lamp-oil. The third product in point of volatility is a comparatively heavy liquid (machinery oil), and from this, and others which come over at a very high temperature, the fourth commercial product is separated by the action of artificial cold, and is the solid paraffin now so much used in the manufacture of candles.

PASTES. See **BLACKING PASTE**, **FURNITURE PASTE**, &c., further back. For flour pastes, see **CEMENTS**. For almond paste, honey paste, and tooth pastes, see **COSMETICS**.

Paste for Cleaning Brass, &c. 1. Rotten stone in very fine powder 2 oz., soft soap 1 oz., oil of amber 1 dr.

2. Neats'-foot oil 16 oz., water of ammonia 1 oz., powdered rotten stone sufficient to form a paste.

3. Rotten stone $4\frac{1}{2}$ lbs., oxalic acid (dissolved in the water) 2 oz., soft soap 8 oz., sweet oil 8 oz.; oil of amber 1 oz., boiling water 1 lb. Some substitute oil of turpentine for oil of amber.

Paste for Razors. 1. Emery very finely levigated in the same manner as prepared chalk, mixed with lard or tallow; or a mixture of these with neats'-foot oil.

2. Equal parts of jewellers' rouge, black lead, and prepared suet.

3. PRADIER'S. Best putty powder 1 oz., jewellers' rouge 1 oz., scales of iron $\frac{1}{2}$ oz., levigated Turkey stone 3 oz., beef suet $1\frac{1}{2}$ oz.

4. Mix equal parts of dried sulphate of iron and salt, and apply a gradually increased heat in a closed vessel. Pulverize, elutriate, and mix with lard or tallow.

PASTILLES, AROMATIC. See PERFUMERY.

PAYNE'S PROCESS FOR RENDERING WOOD FIREPROOF.

The wood is introduced into a close vessel, which is exhausted of air; the liquid is then admitted and forced in by the pump till the pressure is from 110 to 140 lbs. to the square inch. The liquids employed are the liquid sulphides of calcium, or of barium; a solution of sulphate of iron is afterwards forced into the wood.

PERCUSSION CAPS, PRIMING FOR. 100 grs. of fulminating mercury are triturated, with a wooden muller on marble, with 30 grs. of water and 60 grs. of gunpowder. This is sufficient for 400 caps. Dr URE recommends a solution of gum mastic in turpentine as a medium for attaching the fulminate to the cap. The greatest caution is required in the mixing.

PHARAOH'S SERPENTS. The chemical toy sold under this name consists of the powder of sulphocyanide of mercury made up in a capsule of tin foil in a conical mass about an inch in height. Ignited at the apex, an ash is protruded, long and serpentine in shape. The fumes evolved are very poisonous.

PHARAOH'S SERPENTS, *non-poisonous*. Bichromate potash 2 parts, nitrate potash 1 part, and white sugar 3 parts; pulverize each of the ingredients separately, and then mix them thoroughly. Make small paper cones of the desired size, and press the mixture into them. They are now ready for use, but must be kept from light and moisture.

PHOSPHORESCENT OIL. Dissolve 1 gr. of phosphorus in 1 oz. of olive oil in a test tube by the heat of hot water, or add a larger quantity to some oil of lavender, in which it will dissolve spontaneously. Keep it in a close phial.

PHOSPHORUS. See *Pocket Formulary*.

PHOSPHORUS MATCHES. See **LUCIFERS.** The old phosphorus bottles with sulphur matches were made by melting phosphorus with a fourth part of wax in the bottles placed in warm water, and turning them about so as to coat the sides. **GRAY.**

PHOSPHORUS PASTE FOR VERMIN. Introduce 1 dr. of phosphorus into a Florence flask, and pour over it 1 oz. of rectified spirit. Immerse the flask in hot water, until the phosphorus is melted, then put a well-fitting cork into the mouth of the flask, and shake briskly until cold. The phosphorus is now reduced to a finely divided state. This after pouring off the spirit, is to be mixed in a mortar with 1½ oz. of lard. 5 oz. of flour and 1½ oz. of brown sugar, previously mixed together, are now added, and the whole made into a paste with a little water. Cheese may be substituted for sugar when the paste is intended for rats or mice.* (There is said to be no danger, whatever of spontaneous ignition, either during or after the preparation of this paste.)—*Pharmaceutical Journal*.

PHOTOGRAPHY. In all English photographic formulæ the solid and fluid measures of apothecaries' weight are used; but in buying or selling chemical articles, the avoirdupois weight is employed.

Positive Collodion. Pyroxilin and iodide of cadmium or ammonium 15 grs. of each; ether 3½ oz., alcohol 1½ oz. Place the two first in a dry bottle, then pour on the alcohol, shake the mixture well, then add the ether, shake again, and let it stand for 12 hours. Decant the clear portion into a wide-mouthed bottle, keep well stoppered and in the dark. Avoid shaking the bottle when about to use the collodion, and never use quite all the bottle contains, as the sediment, which will accumulate at the bottom, would spoil the picture.

Nitrate of Silver Bath (for positives). Recrystallized

* The following is an old rat-catcher's receipt for oils to attract rats:—Oil of aniseed 2 drs., nitrous acid 2 drops, musk 2 grs. Oil of rhodium is also used for the same purpose. Assafoetida is sometimes mixed with these oils.

nitrate of silver 5 drs., dissolved in 10 oz. of distilled water. Filter the solution until it is quite clear, then add 3 drops of nitric acid and 10 drops of collodion. Shake well together and filter. Blue litmus paper should slightly redden in this bath; should it turn very red, add a little ammonia, or oxide of silver; should it not redden at all, add a little acid carefully drop by drop. It is preferable to have a slight excess of acid.

Developing Solution (for positives). Protosulphate of iron 2 drs. dissolved in 8 oz. of distilled water, add $2\frac{1}{2}$ drs. of glacial acetic acid, $2\frac{1}{2}$ drs. of alcohol, and 5 minims of nitric acid. Filter, and pour into a well-stoppered bottle; do not expose to the air.

Fixing Solution (for positives). 50 grs. of cyanide of potassium dissolved in 5 oz. of distilled water; that is to say, for every fluid ounce of solution required, mix 10 grs. of cyanide of potassium in 1 oz. of distilled water. Filter and keep in a well-stoppered bottle, and label "Poison."

Positive Paper. Plain paper requires preparing or salting before it is ready for use, or it may be purchased already salted.

Procure some sheets of plain Saxe paper, and immerse them for five minutes (removing air-bubbles) in the following solution:

Chloride of ammonium	100 grs.
Chloride of barium	100 grs.
Citrate of soda	20 grs.
Water	20 oz.

Hang the sheets up to dry. For portraits and most other uses the paper is albuminized on one side. When photographs are printed to be afterwards coloured, unalbuminized paper is used.

Albuminized Paper. There are several well-known papers sold; RIVE'S, which is a French paper, has a high glaze and fine surface; the SAXE, which is more uniform in its texture, is made in Germany. Another maker is TO GOOD. Positive paper is albuminized by placing it in a mixture composed of white of eggs and salt. To the white of each moderate-sized egg use 15 grs. of common salt reduced to a fine powder; whisk until the albumin is all white froth. Leave this froth in a glazed earthen pan for about 12 hours, by which time most of it has settled into clear albumin; pour the clear portion into a flat

porcelain tray. This tray should be somewhat larger than the sheets of paper to be albuminized. Lift the paper up by the ends and lay it carefully on the albumin, keeping the side marked as "inferior" uppermost and dry.

The paper should be slightly damp before it is thus treated, as it then takes the albumin more regularly, and is not so liable to air-bubbles. The paper must be lifted at each end, and should any air-bubbles appear, brush them off with a card or small brush, replacing the paper in the bath. Wherever the albumin does not come into actual contact with the paper, a white mark will appear in the print. Remove the paper from the bath and place it to dry on a cardboard frame, or suspend it at the corners by clips. Paper glazed with pure albumin acquires too brilliant a glaze for portraits; the albumin may be diluted with from a $\frac{1}{4}$ to $\frac{1}{2}$ its bulk of water. It should be kept in tin or zinc cases.

Plain Paper. Albuminized paper may be used as plain paper, if, instead of sensitizing the glazed side, the plain side is placed in the sensitizing solution.

Plain Collodion. Mix in a bottle, gun-cotton 450 grs., ether 25 oz., spirits of wine 7 oz. Shake these well together, and leave to settle for several days. If well corked, the mixture will keep for a long time.

Sensitized Collodion. Add to 1 oz. of the plain collodion 6 drs. of spirits of wine, $1\frac{1}{2}$ oz. of ether, and 3 drs. of iodide and bromide solution. Shake the bottle well; the mixture is then ready, but is improved by being kept 4 or 5 hours before using. In hot weather a little more alcohol, and less ether, in very cold weather more ether, and less alcohol, must be used. As sensitized collodion does not keep well, it is better not to mix the plain collodion and the iodide and bromide solution until shortly before required for use.

Iodide and Bromide Solution. Iodide of cadmium 154 grs., bromide of cadmium 54 grs., spirit of wine $3\frac{1}{2}$ oz. Rub the iodide and bromide to fine powder in a mortar, add the spirit of wine gradually, and when the iodide and bromide are dissolved, filter through paper into a bottle. This solution will keep well in a closely-stoppered bottle.

Iodized Collodion may be made at one operation. It should be kept two days before using, but it is less reliable, if kept for any length of time, than sensitized

collodion, which has been made as above described. It is made as follows:—Place 16 grs. of gun-cotton in a bottle, add 18 grs. of iodide of cadmium, and 6 grs. of bromide of cadmium, in powder, and $1\frac{1}{2}$ oz. spirit of wine, sp. gr. 0.805. Shake the bottle till the iodide and bromide are dissolved, then add 3 oz. ether sp. gr. .720, and shake until the cotton is dissolved. After 24 hours decant the clear portion into small well-stoppered bottles.

Nitrate of Silver Bath for Negatives. Recrystallized nitrate of silver $\frac{1}{2}$ oz., distilled water 7 oz., collodion 7 drops. Shake well together until the crystals have dissolved, then filter.

The purity of the negative bath is a matter of great importance, hence the necessity of employing the very best nitrate of silver, and also of excluding foreign matters of every kind. When the bath gets out of order, it should be diluted with an equal bulk of distilled water, and exposed to the sun for a few days, in a white glass bottle, then filtered, and sufficient nitrate of silver added to restore the strength to 35 grains to an ounce, as indicated by the argentometer.

The Argentometer. This instrument is for ascertaining the strength of the nitrate of silver solution, which becomes weakened to a certain extent, after the immersion of every plate.

Developing Solution for Negatives. Protosulphate of iron 75 grs., glacial acetic acid 2 drs., alcohol 2 drs., distilled water 5 oz. Dissolve the crystals in the water, then add the acid and alcohol, and filter. In hot weather a little more acetic acid may be added, and if it does not flow readily, the alcohol may be increased.

Intensifying Solution, No. 1. Pyrogallic acid 10 grs., citric acid 10 grs., distilled water 5 oz. When it becomes brown it is useless. *No. 2.* Recrystallized nitrate of silver 40 grs., distilled water 1 oz.; dissolve and filter. This latter will keep for any length of time in the dark.

Another Intensifying Bath. A saturated solution of bichloride of mercury in water. Place the negative plate in a bath of the solution, remove when the film assumes a milky-white appearance, wash, and then plunge into a solution of 1 oz. of liquid ammonia to 10 oz. of water. Remove the plate, wash, and place to dry. This mode of intensifying may be regulated by leaving the

plate in the bichloride of mercury a shorter time, when it will require a weaker ammonia bath than that above given.

Fixing Solution for Negatives. Hyposulphite of soda, 5 oz., distilled water 5 oz.; dissolve and filter.

Sensitizing Solution, for Paper. Nitrate of silver 5 drs., distilled water 5 oz., nitric acid 2 drops, kaolin 1 oz. Dissolve the nitrate of silver in the water, and then add the acid and kaolin; the kaolin will not dissolve, its use being to prevent the solution becoming discoloured after using. This solution will not require filtering; it must be allowed to become quite clear, and when required for use must be carefully decanted. This solution should be occasionally tested with the argentometer, and sufficient nitrate of silver added to restore it to its original strength, which is 60 grs. to the ounce.

Another Negative Collodion. Ether $\frac{1}{2}$ oz., alcohol $\frac{1}{2}$ oz., gun-cotton 7 grs., bromide of cadmium $\frac{1}{2}$ gr., bromide of ammonium $1\frac{1}{2}$ grs., iodide of cadmium $2\frac{1}{2}$ grs., iodide of calcium 1 gr., iodide of potassium 1 gr., iodide of ammonium 1 gr. *For intensifying,* flood with chloride of gold 1 gr., water 15 oz., then wash, and flood with pyrogallic acid 2 grs., water 3 oz.

Toning Baths. 1. Chloride of gold 4 grs., acetate of soda $\frac{1}{2}$ oz., distilled water 10 oz.; dissolve and filter. 2. *To produce black to bright sepia tones, according to length of immersion.* Take carbonate of soda sufficient to cover a threepenny piece, dissolve it in a teaspoonful of cold water in a cup, add 2 grs. of chloride of gold, then add 3 oz. of boiling water; use in 15 minutes. This will suit HART'S albuminized paper.

Fixing Solution for Paper Prints. Hyposulphite of soda 8 oz., distilled water 1 pint. This solution must only be used once.

Stopping-out Negatives. Small, round, transparent spots are frequently found on glass negatives, which, if not stopped, occasion corresponding black spots on the print. Lay the plate on a slab of glass, having either direct or reflected light shining up through it. Then cover the spots with a mixture composed of 10 parts of ivory-black, 2 parts of saturated solution of gum arabic, 2 parts of white honey, 1 part of sugar-candy; well mix and apply with a fine camel-hair brush. Should the spots on the negative

be black or opaque, white spots will be formed on the print; these are easily tinted with a little water colour, to match the other portions of the print; it is seldom necessary, therefore, to alter the negative on this account.

Ferrotypes. In these, instead of a glass plate being used to receive the picture, as in a positive glass-photograph, a thin plate of black varnished iron is employed. Of course, no black backing is required; with this exception the photograph is produced in every particular by the same means and in the same manner as the glass positive is.

To Clean Glass Plates. Mr HUGHES.—The description of glass known as "flatted crown" is well suited for positives, but before using it requires careful cleaning. The sharp edges should be first removed with a "corundum" file, or by drawing the sharp edge of one piece over the sharp edge of another; then place the glass on a clean flat surface, or put it in a "plate-cleaning holder," and pour a few drops of the plate-cleaning solution in the middle. Rub it carefully over every part with a bit of clean soft rag; turn the glass over, and do the other side the same. Then polish each side with a clean cloth, and finish with a soft chamois leather kept expressly for this purpose. Now breathe on the glass, and if the breath deposits evenly, the plate is clean. If the plate, however, shows patches and marks, it must be recleaned. Let the edges be carefully wiped, and the plate is ready for use.

The following preparation makes a good plate-cleaning solution for glasses that require mechanical friction to make them clean:—Ordinary water 5 oz., alcohol 5 oz., iodide of potassium 15 grs., iodine 3 grs. When dissolved add tripoli, prepared chalk, whiting, or rotten-stone, in sufficient quantity to make a creamy paste.

This thin pasty solution is to be rubbed on the plates on both surfaces and polished off as already described. This amount of cleaning will generally be sufficient for new glasses, but when they have been used they require more labour.

They must then be well washed under the tap, to get rid of all collodion and chemicals, and be wiped on cloths kept expressly for the purpose. Should the plates have been varnished they must be soaked for some hours in a saturated solution of washing soda till the varnish and

film come freely off. The glasses must then be immersed for a few minutes in a solution composed of common nitric acid 2 oz., water 10 oz., and be well washed and treated as already described. It is a good plan when working to have a dish of water at hand, and to place the spoilt pictures in it at once while they are wet, and at the end of the day to wash the glasses and put them away clean. By thus not allowing the films to dry on the glasses, they are much easier cleaned, and fewer failures will arise from dirty glasses.

Collodion is a good material for cleaning glasses when they are not very dirty. Pour a few drops on the glass and well rub it with a clean cloth, and you will entirely remove all grease; a hint may thus be taken how to use up waste collodion.

Various Solutions used in the Dry Process. These are employed in cases where bromo-iodized collodion and the nitrate bath are used.

In all processes in which the bromo-iodized collodion is employed, two grains of bromide of cadmium should be added to each ounce of the collodion.

Mr Bartholomew advises diluted alcohol to be poured over the plate previous to developing.

Acid Pyro Developer. This developer is formed as follows:

Pyrogallic acid	1 gr.
Glacial acetic acid	30 minims.
Water	1 oz.

The plate, after being wetted with dilute alcohol and washed, has this solution flowed over it, to which has been added 2 or 3 drops of a ten-grain nitrate of silver solution.

Plain Pyro Developer. The strength of this may vary from 1 to 5 grs. in an ounce of water. Two grains may be taken as a medium. The dry plate being flooded with alcohol and water, and washed so as to well wet the film, this solution is floated over it.

Alkaline Pyro Developer.

No. 1.—Pyrogallic acid	96 grs.
Absolute alcohol	1 oz.
No. 2.—Carbonate of ammonia	96 grs.
Water	1 oz.
No. 3.—Bromide of potassium	10 grs.
Water	1 oz.

At the time of using make up the following solution :

Water	1 oz.
Solution No. 1	10 minims.
Solution No. 3	5 "

Pour this over the wetted plate, allow it to remain on a few seconds only, and then pour back into the developing cup, and add to it 5 minims of solution No. 2 and apply again.

The Collodio-Albumen Dry Process. Mr MUDD.—In this process the ordinary bromo-iodised collodion is employed. The plate being sensitized, is washed well, first with distilled, then with common water, and placed in a dish half filled with solution of iodide of potassium (3 grs. to the ounce), and allowed to remain while the next plate is being prepared. It is then removed from the solution and well washed with clean water, after which the following solution is poured over its surface :

Distilled water	2½ oz.
White of eggs	10 oz.
Iodide of potassium	50 grs.
Bromide of ammonium	10 grs.
Strongest solution of ammonia	120 minims.

Introduce these materials with some pieces of broken glass into a bottle capable of holding twice the quantity, and agitate till the whole forms a froth, and then, when settled, it is ready. A piece of camphor placed in the solution will help to preserve it. It must be filtered before using. After the plate has been coated with the above, it is finished by drying before the fire.

In this process all the above operations may be performed in ordinary white light. To render the plate sensitive, heat it as hot as the hand will bear, and when cool immerse it again in the following aceto-nitrate of silver bath for one minute, *using only a yellow light*, then wash thoroughly in clean water and dry in the dark :

Nitrate of silver	30 grs.
Distilled water	1 oz.
Glacial acetic acid	½ dr.

The development may be commenced by either plain or alkaline pyro ; Mr Mudd gives the preference to the plain, and intensifying after with acid silver.

Dr RYLEY's *Modified Collodio-Albumen Dry Process.* In this method the plate has to be sensitized as usual and

washed thoroughly. When the plate has been well drained, and while still wet, it is coated with the following solution:

Albumen	1 oz.
Water	2 oz.
Ammonia	30 minims.

The solution is beaten to a froth, allowed to settle, and filtered before using. Pour sufficient over the plate to cover it, letting it flow backwards and forwards so as to soak into the film. Pour the albuminous solution away and thoroughly wash the plate, the last rinsing being with distilled water. Let the plate dry; when perfectly dry moisten the plate with distilled water and pour over it the following solution:

Gallic acid	2 grs.
Water	1 oz.

Filter the solution before using. Pour it on and off the plate to well permeate the film, then set the plate up to drain, and dry without washing off the gallic acid solution. When the surface is dry finish by the heat of a dull fire.

Bromide of Silver, Wet Process. To every ounce of good collodion add 8 grs. of bromide of cadmium. The nitrate bath must be made 80 grs. to the oz., and slightly acidulated with nitric acid. The plate must remain in the bath the full time it requires to form a dense opaque film. When the plate is ready (it must not be removed from the bath until the film is much denser than in the ordinary wet process) it must be washed thoroughly to remove all silver. It must then have poured over it a 3-gr. solution of gelatine made slightly alkaline with carbonate of soda, or diluted albumen (albumen 1 oz., water 4 oz., well beaten together). When the plate is in this condition it may be exposed *wet*, or it may be allowed to dry. Prior to development it must be well washed, and the alkaline method must always be adopted. If the plates are used dry, a preliminary coating of dilute albumen is necessary, but if used wet, this is not necessary.

Bromide of Silver, Emulsion Process. By this method the nitrate bath is not necessary, as the sensitive material is contained in the collodion. The purchase of the material ready made for working this process is recommended in preference to its direct manufacture, as its preparation demands the use of considerable technical

skill, together with the employment of a gun-cotton not usually attainable. Mr HUGHES says the "Liverpool Dry Plate Company" supply an excellent emulsion. It is only necessary to pour the emulsion on to a plate and to allow it to dry, when the plate is ready for use. The development is by the alkaline process.

Gelatino-bromide of Silver Emulsion Process. In this process, the use both of the nitrate bath and of collodion are abolished.

The material employed is very troublesome to prepare, and on this account, as well as because of the risk of failure attending the use of the home-made article, it is far preferable it should be purchased. It may be obtained under the name of "KENNETT'S Sensitized Pellicle."

This pellicle consists of shreds of dry gelatin containing the sensitive salts.

Full directions accompany each packet of the "Sensitized Pellicle."

The Autotype Process. In this process the material employed consists of a layer of gelatin containing carbon, or some other permanent pigment, spread on paper.

In this condition the paper is not sensitive to light, but if it be treated with a solution of bichromate of potash, dried in the dark and afterwards exposed to sunlight under a negative, those portions of the paper which have been acted upon by the light will become insoluble, whilst those parts that have been protected from it, will be soluble. When, therefore, after sufficient exposure, the prepared paper is removed from the negative, a picture the reverse of the negative will have been formed, in which the pigmented gelatin remains and performs the part of the reduced silver in the ordinary photograph.

PHOTOGRAPHIC VARNISH. See VARNISHES.

The reader desirous of further information on the subject of photography cannot do better than consult Mr Ernest SPON'S valuable manual, entitled 'Workshop Receipts,' and Mr HUGHES' 'Principles and Practice of Photography,' to both of which works we are largely indebted. Captain ABNEY'S work on 'Photography' cannot be too highly commended.

PHOTOGRAPHY.—MISCELLANEOUS RECIPES.

To Clean Glass Plates. Mr MAYALL. Shake up together, alcohol 30 parts, strong liquid ammonia 10, water

40, and fine tripoli 30 parts. The plates are to be rubbed hard and evenly with balls of cotton wool dipped in this mixture. Rub again, when dry, with a clean ball of cotton; lastly, dust the back and edges with a clean hog's-hair brush.

To Clean off Collodion Pictures. This may be done, whether they have been varnished or not, by means of a tuft of cotton wool dipped in wood spirit.

To Colour Photographic Prints. This may be done variously in water and oils. A simple way is to rub in slowly with a small camel-hair brush a minute piece of dry colour laid upon the part, as of flesh tint for the face, &c. When properly distributed, the paper may be breathed upon, and the tint will not easily be rubbed off. Or it may be carefully coated with gelatin.

M. MINOTTO has described a plan of colouring on the back of the paper. The picture, being held up to the light, is first faintly outlined on the reverse side; colours are then laid on, of water or oil, as preferred, on this side. When dry, the paper is rendered transparent by a varnish, and the colours will then appear through it with all the delicacy and effect of a miniature on ivory. Good strong writing paper is best for this purpose; the colours must be vivid; and the varnish may consist of Canada balsam dissolved in turpentine, or a mastic varnish may be used, or turpentine and wax, or oil.

PIGMENTS. A few of these have been noticed before; see INDIGO, LAKES, PRUSSIAN BLUE, PURPLE OF CASSIUS. They generally constitute a distinct branch of manufacture, but a brief account of the composition of some of them may be useful. Those of which the colouring matter is derived from the animal and vegetable kingdoms will first be noticed; then the mineral colours.

Carmine. Several processes have been published for this beautiful pigment, but probably some minute precautions, not generally known, may be necessary to the production of the finest quality. The climate and state of the atmosphere are said to influence the result.

1. *Madame CENETTE'S process.* Into 6 pails of boiling, clear, soft water, in a copper vessel, throw 2 lbs. of powdered cochineal of good quality; boil for 2 hours, add 3 oz. of purified nitre, and, after a few minutes, 4 oz. of salt of sorrel. Remove the vessel from the fire, let the contents

settle for 4 hours, draw off the clear liquor with a syphon into flat plates, and leave it at rest for 3 weeks. Carefully detach the pellicle of mould from the surface, withdraw the liquid with a syphon and pipette, and dry the deposit in a stove.

2. Boil 4 quarts of soft water in a pewter kettle, add to it 4 oz. of finely powdered cochineal; boil for 5 minutes, adding 2 drs. of powdered cream of tartar; then add 8 scruples of Roman alum, and keep the whole on the fire for a minute longer. Let the decoction settle, decant it into cylindrical glasses, and cover them. When the carmine has subsided, pour off the clear liquor, and dry the sediment. By adding solution of tin to the liquid, more carmine is obtained.

3. Into a 14-gallon boiler of tinned copper, put 10 gallons of distilled water, or filtered rain-water; when it boils, sprinkle in, by small quantities, 1 lb. of powdered cochineal, and keep it boiling for half an hour. Then add 3½ oz. of crystallized carbonate of soda; in a minute or two, draw the fire, and add 1½ oz. of Roman alum in fine powder; stir with a glass rod till the alum is dissolved, leave it to settle for 25 minutes, draw off the liquor with a glass syphon, and strain the rest through a coarse linen cloth. Clean the boiler, return into it the clear-coloured liquor, and stir into it the whites of 2 eggs, previously well beaten with a quart of warm (not hot) water. Then light the fire, and heat the liquor till it begins to boil; separate the coagulum, collect the carmine which deposits, wash it on the filter with distilled water, spread it thinly on earthen plates, and dry it in a stove.

Inferior carmine may be improved by dissolving it in water of ammonia, and precipitating it by acetic acid and alcohol.

Cochineal Lake. Add 2 lbs. of pearlash to the red liquor from which the carmine has been separated in the last process, and return it to the boiler with the dregs of the cochineal; boil for half an hour, draw the fire, and when the sediment has subsided, draw off the clear liquid into an earthen vessel. Pour on the sediment a solution of 1 lb. of pearlash in 2 gallons of water; and boil for half an hour. Filter, and return both liquors into the copper. When as hot as the hand can bear, add to the liquor, by little and little, 3 lbs. of powdered Roman alum, and let

it simmer for 5 minutes. Allow it to settle, draw off the clear liquor, collect the sediment on a filter, wash it with clean rain-water, and leave it covered with a cloth for a few days, till half dry; form it into small lumps, and dry them in a stove.

Carthamin or Safflower Lake. Wash Safflower till the water comes off colourless; mix it with water holding 15 per cent. of carbonate of soda in solution, so as to form a thick paste; leave it for several hours, then press out the red liquid, and nearly neutralize it with acetic acid. Then put cotton into it, and add successive small portions of acetic acid, so as to prevent the liquid becoming alkaline. In 24 hours take out the cotton, wash it, and digest it for half an hour in water holding 5 per cent. of crystallized carbonate of soda in solution. Immediately on removing the cotton, supersaturate the liquid with citric acid, and collect the precipitate, which must be repeatedly washed in cold water. For pink saucers the liquor is allowed to deposit in the saucers. Mixed with the scrapings of French chalk it constitutes rouge.

Lakes are also obtained from Brazil-wood and madder, by adding alum to a concentrated decoction of the former, or to a cold infusion of the latter (made by triturating the madder, inclosed in a bag, with the water), and afterwards sufficient carbonate of potash or soda to throw down the alumina in combination with the colouring matter. The precipitate is to be washed and dried. A little solution of tin added with the alum improves the colour. Lakes may be obtained from most vegetable colouring matters by means of alum and an alkaline carbonate. *Yellow Lake* is made from French or Persian berries, by boiling them in water, with a little carbonate of soda or potash, and adding alum to the strained liquor as long as a precipitate is thrown down. Or by boiling weld, or quercitron bark, in water, and adding alum and chalk in a pasty state.

Rose Pink. Boil 6 lbs. of Brazil-wood and 2 lbs. of peachwood in water, with $\frac{1}{4}$ lb. of alum; and pour the strained decoction on 20 lbs. of sifted whiting.

Bistre. It is obtained from the soot of beech-wood.

Sap Green. The expressed juice of buckthorn-berries (and sometimes of other species of *rhamnus*, and also of

privet berries) is allowed to settle, and the clear liquid evaporated to dryness. A little gum arabic is sometimes added to the juice.

The beautiful colours of the *mauve* series are prepared from coal tar by patented processes. See "Dyeing, Printing, and Calico Printing," 'Churchill's Technological Handbooks.'

MINERAL PIGMENTS. *Azure Blue, or Smalts.* The common is made by fusing zaffre (roasted cobalt ore calcined with siliceous sand) with potash. A finer quality is obtained by precipitating a solution of sulphate of cobalt by a solution of silicate of potash. Another cobalt blue is obtained by adding a solution of phosphate of soda to a solution of nitrate of cobalt, and mixing the precipitate, washed, but not dried, with eight times its weight of fresh hydrated alumina. When dry, heat it to a cherry red.

Egyptian Azure. Carbonate of soda 16 oz., calcined flints 24 oz., copper filings 4 oz. Pulverise, mix, and fuse in a crucible for two hours. When cold, reduce to powder.

Blue Verditer. It is generally stated to be made by adding chalk to a solution of nitrate of copper produced in the process of refining silver; but Mr PHILLIPS did not succeed in making it by this means, and found no lime in the best samples.

New Blue. Mix equal parts of common arsenite of copper (see MINERAL GREEN, further on), and neutral arsenite of potash, fuse by heat in a large crucible, then add to the fused salt 1-5th of its weight of nitre. Effervescence takes place, and the salt becomes blue. Cool, pulverise, and wash.

Chrome Yellow. To a solution of bichromate of potash add a solution of nitrate of lead as long as a precipitate forms. Wash the precipitate, and dry it with a gentle heat. An inferior kind is said to be made by 4 lbs. of pure white lead, 1 lb. of bichromate of potash, and 20 lbs. of water, and boiling till the water becomes colourless. Or 75 parts of precipitated sulphate of lead may be acted on by a hot solution of neutral chromate of potash 25 parts. A mixed product of chromate and sulphate of lead is thus obtained, which is said to cover as well as the pure chrome yellow, and is much cheaper. RIOT.

Chrome Red. Melt saltpetre in a crucible heated to

dull redness, and add chrome yellow, by small portions, till no more red fumes arise. Allow the mixture to settle, then pour off the melted salt from the heavy sediment, and wash the latter with water, which should be quickly poured off, and dry the pigment. The liquefied salt poured off contains chromate of potash, and is reserved for making chrome yellow.

Orange Chrome is chrome yellow acted on by an alkali, which deprives it of part of the chromic acid.

King's Yellow. This is a yellow sulphide of arsenic, now almost superseded by chrome yellow, but occasionally used for killing flies.

Naples Yellow. Mix 12 parts of metallic antimony, 8 parts of red lead, and 4 of oxide of zinc, and calcine in a reverberatory furnace. The mixed oxides are rubbed together, fused, and the fused mass elutriated into a fine powder.—Dr URE. M. GUIMEL recommends 1 part of well-washed antimoniate of potash to be ground into a paste with 2 parts of red lead, and the powder exposed to a red heat for 4 or 5 hours, keeping the heat moderate.

Brighton Green. An inferior colour made with 28 lbs. of whiting, or white lead, 7 lbs. sulphate of copper, 3 lbs. sugar of lead, and $\frac{1}{4}$ oz. of bichromate of potash.

Brunswick Green. Pour a saturated solution of chloride of ammonium over copper filings in a close vessel placed in a warm situation; add more of the solution from time to time till 3 parts of the chloride have been used to 2 of copper. After standing for a few weeks the pigment is separated from the unoxidized copper by washing through a sieve. It is then to be well washed, and dried slowly in the shade. It is often reduced with white lead; some samples contain arsenic.

Arsenical Copper Greens. Of these there are several varieties.

Mineral Green, Scheele's Green, or Arsenite of Copper.

1. Dissolve 11 oz. of white arsenic and 2 lbs. of carbonate of potash, by heat, in a gallon of water. Dissolve also 2 lbs. of sulphate of copper in 3 gallons of water. Filter each solution separately, and add the former gradually to the latter as long as it occasions a precipitate. Wash the precipitate, drain it, and dry it.

2. Dissolve 50 lbs. of sulphate of copper and 10 lbs. of lime in 20 gallons of good vinegar, and add quickly the

boiling hot solution to 50 lbs. of white arsenic. Stir repeatedly, then allow it to settle; decant the clear liquor (which is reserved to dissolve the arsenic next time), and wash the precipitate and dry it.

3. *Emerald Green.* Mix 10 parts of pure verdigris with sufficient boiling water to form a soft pulp, and strain this through a sieve. Dissolve 9 or 10 parts of white arsenic in 100 parts of boiling water, and whilst boiling, let the verdigris pulp be gradually added, constantly stirring the mixture till the precipitate becomes a heavy, granular powder.

Green without Arsenic. Dissolve 48 lbs. of sulphate of copper, and 2 lbs. of bichromate of potash in water, and add to the clear solution, 2 lbs. of pearlash and 1 lb. of chalk.

Rinmann's Green Pigment. Dissolve together in sufficient water, 1 part of sulphate of cobalt and 3 of sulphate of zinc; precipitate with carbonate of soda, wash the precipitate and calcine it.

Chrome Green. A mixture of chrome yellow and Prussian blue. See also CHROME OXIDE, further back.

Barth's Green. A mixture of Prussian blue and yellow lake.

Ultramarine, Factitious. Take 70 parts of silica, or pure siliceous sand, in fine powder; 240 parts of recrystallized alum, calcined; 144 parts of sulphur; 48 parts of finely powdered charcoal; 240 parts of dry carbonate of soda. These are mixed together with the greatest care till the mixture appears of uniform colour under a powerful magnifier, and the mixture exposed to a moderate red heat in a closely covered crucible for an hour and a half. Wash the product with boiling water. When dry, mix the powder with its own weight of sulphur and $1\frac{1}{2}$ its weight of dried carbonate of soda, and burn as before; heat it again with sulphur and carbonate of soda, and wash it till the filtered fluid no longer colours acetate of lead. If a sample of the dried powder becomes blue when burnt with sulphur, it is ready for the last operation. Spread over a cast iron plate a layer of sulphur a line in thickness, and over it an equal layer of the dry powder after having passed it through a gauze sieve. Heat the plate so as to burn away the sulphur at the lowest possible temperature. Reduce the pigment to powder and repeat the

burning with sulphur and pulverization till the colour is perfect.

White Lead is carbonate of lead prepared by various processes. *Zinc White* is oxide of zinc, prepared by combustion. *Oxychloride of Zinc* is also used. *Antimony White* is oxide of antimony.

PINK SAUCERS. See **RED DYES**, further back.

PLATES, DAGUERRETYPE, are prepared by cleaning and polishing the silver surface, exposing it to the vapour of dry iodine, or tincture of iodine, or iodide of bromine, or bromide of calcium. After having the image thrown on them, they are exposed to the vapour of mercury. But the manipulations and precautions necessary to the success of the operation are too numerous to detail here. An exploded photographic process.

PLATE-BOILING POWDER. Equal parts of cream of tartar, alum, and common salt. A small quantity added to the water in which plate is boiled gives it a silvery whiteness.

PLATE POWDERS. 1. *Jewellers' Rouge.* Dissolve green vitriol in hot water, and add a solution of pearlsh as long as it throws down a precipitate. Wash the precipitate repeatedly with warm water, drain it on calico, and finally calcine it till it assumes a bright colour. It is sometimes made by calcining the sulphate of iron with a strong heat, till peroxide of iron only remains. Let it be triturated with water, and prepared in the same way as prepared chalk. See **POLISHING POWDER**, further on.

2. *French Plate Powder.* Mix 1 part of jewellers' rouge with 12 of carbonate of magnesia.

3. Finely prepared chalk, or burnt hartshorn. One way in which these are used is to boil them with water, with pieces of rag; the finer particles are entangled in the fibres of the rags, which are dried and kept for use.

4. Quicksilver with chalk 1 oz., prepared hartshorn 8 oz., prepared chalk 4 oz. Powders containing quicksilver, besides the necessary wearing of the surface, are supposed to render the plate more brittle. If used the quicksilver should not be in larger proportion than the above.

5. Finest putty powder 1 oz., levigated chalk 5 oz.; a little rouge may be added to colour it.

See **NOVARGENT, SILVERING POWDER, &c.**, for restoring the silver to plated goods.

- PLATINUM, BLACK (OXIFEROUS).** Dissolve protochloride of platinum in a boiling solution of potash, add alcohol in small portions till effervescence ceases. Boil the black precipitate successively with alcohol, hydrochloric acid, and potash, and finally 4 or 5 times with water.
- PLATINATED ASBESTOS.** Dr HARE. Dip asbestos in a solution of chloride of platinum, and heat it to redness. It causes the inflammation of hydrogen in the same manner as spongy platinum.
- PLATINIZED SILVER.** Silver plates for SMEE'S voltaic battery are covered with pulverulent platinum by adding a little bichloride of platinum to acid water, and decomposing the solution by the use of a platinum terminal in connexion with the copper of a battery, the silver plate to be platinized being in connexion with the zinc. Platinum itself is sometimes platinized in the same way. Sometimes the plates are platinized without the battery. The following solution is used by Dr Wright for the plates of his battery:—Saturated solution of chloride of platinum $\frac{1}{2}$ dr., sulphuric acid $1\frac{1}{2}$ drs., water 2 drs. Dip the plates in it for a few seconds, and wash them quickly.
- PLATINUM, CHLORIDES OF.** Dissolve platinum in nitrohydrochloric acid, and evaporate with a gentle heat to dryness. The red bichloride (tetrachloride) remains. Heated to 450° , the protochloride remains.
- PLATINUM SPONGY.** Dissolve separately in rectified spirit, chloride of platinum and sal ammoniac. Mix the solutions, and heat the precipitate to redness. For balls for hydrogen lamps, form the precipitate into balls while moist, and afterwards burn them.
- POISON.** See BEETLE WAFER, BUG POISON, RAT POISON; PHOSPHORUS PASTE; BLIGHTS, REMEDIES FOR, &c.
- POLISH.** See FRENCH POLISH.
- POLISH FOR BOOTS, &c.** See BLACKING.
- POLISHING POWDER FOR SPECULA.** LORD ROSS. Precipitate a dilute solution of sulphate of iron by ammonia in excess; wash the precipitate, press it in a screw press till nearly dry; then expose it to heat until it appears of a dull red colour in the dark.
- POT POURRI.** See PERFUMERY.
- POTASH, CARBONATE OF.** *Salt of Tartur.* *Subcarbonate of potash.* See POTASSÆ CARBONAS, and POTASSÆ CARBONAS PURUM, *Pocket Formulary.*

POTASH CHLORATE. See CHLORATE OF POTASH.

POTATO DISEASE, REMEDIES FOR. The following are some of the formulæ that have been proposed for the prevention of this terrible and well-known plant epidemic :

1. Some direct that the haulms and leaves should be entirely removed as soon as the least symptom of disease shows itself in them. The roots may grow on to full size without becoming affected.

2. The CHEVALIER CLAUSSEN'S Cure. The potatoes, before planting, are wetted with water acidulated with sulphuric acid (1 part to 500), and, before they are dry, powdered sulphate of lime is thrown over them. This plan has been found a very efficient one.

3. Mr T. HERAPATH'S plan. The roots, before planting, are allowed to become dry, and then dipped for a short time in a weak solution of sulphate of copper. The land is dressed with a mineral composition (2 parts of lime to 1 of salt) instead of farmyard manure. This process also has been attended with considerable success.

POUNCE. Powdered gum juniper is used under this name, for preparing parchment for writing on. For liquid pounce, see **INK, MARKING.**

POUDRE CLARIFIANTE. Beat together the whites and yolks of eggs, dry them with a very gentle heat, and reduce to powder. For clarifying wines and syrups.

POWDERS. See **TOOTH POWDERS,** and **HAIR POWDERS,** under **COSMETICS ;** **SCENT POWDERS,** under **PERFUMERY, &c.**

PRESERVATIVE LIQUIDS. See **ANATOMICAL SUBJECTS,** and **ANIMAL SUBSTANCES, TO PRESERVE.**

PRUSSIAN OF POTASH (YELLOW). What is known in commerce by this name is the ferro-prussiate of potash, or ferrocyanide of potassium. It is prepared by fusing in an egg-shaped iron pot a mixture of 2 parts of pearlsh and 5 parts of dry animal matters, such as horns, hoofs, tallow-chandler's greaves, &c., till fetid vapours cease to be produced. Iron filings are sometimes added, but usually the iron necessary to the formation of this salt is derived from the iron pots and stirrers. The fused mass (*prussiate cake*) is allowed to cool, dissolved in warm water, and the clear filtered or decanted solution evaporated, that crystals may form. These are dissolved in hot water, and the solution allowed to cool very slowly, that large crystals may form.

RED PRUSSATE OF POTASH. *Ferrid-cyanide of Potassium. Ferricyanide of Potassium.* Into a dilute solution of the yellow prussiate of potash, a current of chlorine gas is passed, till the solution ceases to give a blue precipitate with persalts of iron. It is then evaporated, crystallized, and recrystallized till quite pure. [M. POSSELT advises to add a few drops of solution of potash to the boiling liquor, to decompose the green matter that is formed; to filter the hot solution, to separate some peroxide of iron which is thrown down, and to let the liquor cool very slowly.] Or, boil yellow prussiate of potash with 12 or 15 parts of water, and while boiling add good chloride of lime until a filtered sample no longer yields a blue precipitate with persalts of iron. Filter quickly, and add carbonate of potash till the liquid has a faintly alkaline reaction, then evaporate for crystallization.

PRUSSIAN BLUE. *Berlin Blue. Percyanide, ferrocyanide, or ferro-prussiate of iron.* Commercial Prussian blue is made by adding to a solution of prussiate of potash, or of prussiate cake, a solution of 2 parts of alum and 1 of sulphate of iron, washing the precipitate repeatedly with water to which a little hydrochloric acid has been added, and exposing it to the air till it assumes a deep colour. A purer kind is made by adding a solution of persulphate or perchloride of iron to a solution of pure ferro-prussiate of potash. TURNBULL'S Prussian blue (ferrid-cyanide of iron) is made by adding a solution of red prussiate of potash to one of proto-sulphate of iron; or by adding proto-sulphate of iron to a mixture of yellow prussiate of potash, chloride of soda, and hydrochloric acid.

SOLUBLE PRUSSIAN BLUE. Add a solution of proto-sulphate of iron to a solution of prussiate of potash, and expose the precipitate to the air till it becomes blue, and wash it till the soluble salts are washed away. By continuing the washing, the blue itself dissolves, forming a deep blue solution, which may be evaporated without decomposition. Or add a solution of persulphate of iron to a solution of ferro-prussiate of potash, keeping the latter in excess; wash the precipitate until it begins to dissolve, and dry it. See *INK, READE'S Patent Blue*, for another method.

PURPLE OF CASSIUS. See *AURUM STANNO PARATUM, Pocket Formulary*. Many other processes have been proposed,

of which the following is one:—Dissolve 3 grs. of gold in aqua regia, avoiding excess, and dilute with 3 oz. of water. Mix 30 grs. of pink salt (the bichloride of tin with sal-ammoniac) with $3\frac{1}{4}$ grs. of tin filings and 2 drs. of water till the tin is almost entirely dissolved: add 7 drs. of water, and add this solution to the gold solution, slightly warmed. Wash the precipitate, and dry it.

PYROLIGNEOUS ACID, }
 PYROXYLIC SPIRIT, } By the destructive distillation
 PYROACETIC SPIRIT, &c. } of dried wood, chiefly that of
 the beech and birch, in iron
 cylinders, an acid liquor and tar are produced. These are received in proper reservoirs, and are afterwards separated. The tar is subjected to distillation, and yields *oil of tar* (containing creosote, eupion, &c.), and leaves a residuum of pitch, or English asphalt. The acid liquor, separated from the tarry deposit, is also distilled: the first portion which comes over contains the pyroxylic spirit, which is rectified by one or more distillations. It may be further purified by distilling it with dried chloride of calcium, and finally with quicklime. This constitutes one of the articles sold under the name of wood naphtha, and is regarded by chemists as a hydrated oxide of methyl. After the pyroxylic spirit has come over, the crude pyroligneous acid distils, which still holds some tar and empyreumatic oil in solution. It is purified by saturating it either directly with common soda, or first with lime, or rather chalk, and when the neutral solution has become clear, evaporating it to 1.114 sp. gr., and adding sufficient saturated solution of sulphate of soda to decompose the impure acetate of lime. The clear solution obtained by either process is then evaporated, that the acetate of soda may crystallize. This is afterwards roasted at a temperature of about 500° Fahrenheit, to destroy the tar, and again dissolved and crystallized. The purified acetate is then distilled with sulphuric acid. See ACIDUM ACETICUM, *Pocket Formulary*. For some manufacturing purposes, an impure acid is obtained by merely saturating the crude pyroligneous acid with lime, evaporating to dryness, and distilling with sulphuric acid.

If acetate of lime or acetate of lead be distilled without addition, and the liquid which comes over be rectified over lime, *pyroacetic spirit* is obtained: this is also termed *acetone*.

PYROPHORUS. This name is given to several compounds, prepared by calcination, which take fire when exposed to the air, especially when breathed upon. The following are perhaps some of the best :

1. Heat tartrate of lead in a tube of hard glass, and securely close the tube before the charred residuum becomes cold. A little poured out and breathed upon takes fire. The tartrate of lead is made by dissolving separately 2 drs. of tartaric acid and 5 drs. of crystallized acetate of lead in sufficient water, mixing the solutions, and collecting, washing, and drying the precipitate.

2. Calcine tartar emetic in a similar manner, or in a closed crucible.

3. Mix 11 parts of lamp-black with 2 of powdered sulphate of potash, and heat the mixture strongly in a closely covered crucible. The product is so combustible that it can scarcely be transferred to a bottle without danger.

4. Mix 3 parts of powdered alum with 1 of flour, and calcine the mixture in a common phial coated with clay or placed in sand, till it ceases to emit a blue flame. Before it is cold, close it securely with a sound cork or glass stopper.

5. Mix neutral chromate of lead with 1-6th its weight of sulphur; triturate them with water sufficient to form a paste, and make it into pellets; dry these perfectly, then heat them in a tube till the sulphur is all driven off, and secure as the last.

RENNET. The stomach of a calf, washed, salted, and dried.

RENNET LIQUID. ESSENCE OF RENNET. Fresh rennet 12 oz., salt 2 oz., proof spirit 2 oz., white wine a quart; digest for 24 hours and strain. A quart of milk requires 2 or 3 teaspoonfuls. **WISLIN** directs, 10 parts of a calf's stomach, salt 3 parts. The membrane of the stomach is to be cut with scissors, and kneaded with the salt and with the rennet found in the interior of that organ; the whole left in a cool place in an earthen pot till the cheesy odour is replaced by the proper odour of rennet, which will be in 1 or 2 months. Then add 16 parts of water and 1 of spirit. Filter and colour with burnt sugar.

RUST, TO PREVENT AND REMOVE. Steel goods are rubbed over with a mixture of lime and oil, to preserve them from rusting. Mercurial ointment has been recommended for the same purpose. **M. PAYEN** recommends plunging the

articles into a solution of common soda. Spots of rust are removed by rubbing them with very fine emery and sweet oil; as a chemical means of removing them, the ammoniacal chloride of zinc may be found useful. Vaseline rubbed over steel goods preserves them from rusting. To remove iron-mould spots from linen, oxalic acid or binoxalate of potash is commonly used. RUNGE recommends prussiate of potash. Dissolve a little in water, and just acidulate with sulphuric acid. The stained linen macerated in this water turns blue. Its whiteness is now restored by a solution of carbonate of potash.

SAXON BLUE. See **CHEMIC BLUE.** The solution of indigo in sulphuric acid, diluted with twice its weight of water, is so termed.

SCOURING DROPS FOR REMOVING GREASE. 1. Alcohol 6 oz., camphor 2 oz., rectified essence of lemon 8 oz.

2. Rectified oil of turpentine 3 oz., essence of lemon 1 oz.; mix. Some direct them to be distilled together.

3. *French.* Camphine 8 oz., alcohol 1 oz., sulphuric ether 1 oz., essence of lemon 1 dr.

4. Spirit of wine a pint, white soap 3 oz., ox-gall 3 oz., essence of lemon $\frac{1}{4}$ oz.

5. Benzol, or ether, or chloroform.

SEALING-WAX. 1. *Blue.* Shell-lac 2 parts, dammar resin, 2 parts, Burgundy pitch 1 part, Venice turpentine 1 part, artificial ultramarine 3 parts.

2. *Light Blue.* As the last, with 1 part of dry sulphate of lead.

3. *Dark Blue.* Venice turpentine 3 oz., finest shell-lac 7 oz., clear amber or black resin 1 oz., Prussian blue 1 oz., carbonate of magnesia $1\frac{1}{2}$ drs. The last two to be made into a stiff paste with oil of turpentine, and added to the melted shell-lac and Venice turpentine.

Black. 1. Venice turpentine $4\frac{1}{2}$ oz., shell-lac 9 oz., colophony $\frac{1}{2}$ oz., lamp-black mixed to a paste with oil of turpentine q. s.

2. *Inferior.* Venice turpentine 4 oz., shell-lac 8 oz., 3 oz. of colophony, and sufficient lamp-black mixed with oil of turpentine to colour it.

3. Shell-lac 8 oz., Venice turpentine 4 oz., lamp-black 6 oz.

4. *Common, for bottles.* Resin 6 oz.; shell-lac 2 oz., Venice turpentine 2 oz., lamp-black q. s.

Brown. 1. *Light Brown.* Venice turpentine $\frac{1}{2}$ oz., shell-lac $7\frac{1}{2}$ oz., brown earth (English umber?) $\frac{1}{2}$ oz., cinnabar $\frac{1}{2}$ oz., prepared chalk $\frac{1}{2}$ oz., carbonate of magnesia moistened with oil of turpentine $1\frac{1}{2}$ drs.

2. *Light Brown.*—Second quality. Venice turpentine 4 oz., shell-lac 7 oz., resin 3 oz., English umber 3 oz., cinnabar $\frac{1}{2}$ oz., prepared chalk 1 oz., magnesia as the last.

3. *Dark Brown.* Venice turpentine $\frac{1}{2}$ oz., fine shell-lac $7\frac{1}{2}$ oz., English umber $1\frac{1}{2}$ oz., magnesia as before.

4. *Dark Brown.*—Second quality. Venice turpentine $\frac{1}{2}$ oz., shell-lac 7 oz., colophony 3 oz., English umber $1\frac{1}{2}$ oz., magnesia as before.

Green. Venice turpentine 2 oz., shell-lac $\frac{1}{2}$ oz., colophony $1\frac{1}{2}$ oz., King's yellow $\frac{1}{2}$ oz., Prussian blue $\frac{1}{2}$ oz., magnesia as for brown.

Gold. 1. Venice turpentine $\frac{1}{2}$ oz., fine shell-lac 8 oz., leaf-gold 14 sheets, bronze powder $\frac{1}{2}$ oz., magnesia (made into a paste with oil of turpentine) $1\frac{1}{2}$ drs.

2. GRAY. Use gold talc instead of gold leaf and bronze.

Marbled. GRAY. Melt each coloured wax separately, and just as they begin to grow solid, mix together.

Red. 1. *Fine Carmine Wax.* Venice turpentine 2 oz., finest shell-lac 4 oz., colophony 1 oz., Chinese vermilion $1\frac{1}{2}$ oz., magnesia (moistened with oil of turpentine) $1\frac{1}{2}$ drs.

2. *Finest Red.* Venice turpentine $\frac{1}{2}$ oz., shell-lac 7 oz., cinnabar $\frac{1}{2}$ oz., carbonate of magnesia (with oil of turpentine) $1\frac{1}{2}$ drs.

3. As the last, with only $3\frac{1}{2}$ oz. of cinnabar.

4. Venice turpentine $\frac{1}{2}$ oz., shell-lac $6\frac{1}{2}$ oz., colophony $\frac{1}{2}$ oz., cinnabar $2\frac{1}{2}$ oz., magnesia (with oil of turpentine) $1\frac{1}{2}$ drs.

5. Venice turpentine $\frac{1}{2}$ oz., shell-lac 6 oz., colophony $\frac{3}{4}$ oz., cinnabar $1\frac{3}{4}$ oz., magnesia as before.

6. As the last, but use colophony and cinnabar each $1\frac{1}{2}$ oz.

7. Venice turpentine $\frac{1}{2}$ oz., shell-lac $5\frac{1}{2}$ oz., colophony $1\frac{1}{2}$ oz., cinnabar $1\frac{1}{4}$ oz., magnesia as before.

8. *English.* Venice turpentine 2 oz., shell-lac 4 oz., vermilion 1 oz.

9. *Spanish.* Venice turpentine 8 oz., shell-lac 2 oz., colophony 4 oz., vermilion 1 oz. Remove from the fire, and add $\frac{1}{2}$ oz. of rectified spirit.

Yellow. Venice turpentine 2 oz., shell-lac 4 oz., colophony $1\frac{1}{4}$ oz., King's yellow $\frac{1}{4}$ oz., magnesia as before.

Perfumed Wax. Add to any of the above a small quantity of fine benzoin.

Common Bottle Wax. 1. Dark resin 18 oz., shell-lac 1 oz., beeswax 1 oz. Mix together, and colour with red lead, Venetian red, or lamp-black.

2. Resin 19 oz., beeswax 1 oz., colour as before.

SEA-WATER, ARTIFICIAL. See AQUARIUM.

SHELL-LAC, TO BLEACH. See LAC, further back.

SILK CLEANER. Mix well together $\frac{1}{4}$ lb. of best soft soap, a teaspoonful of brandy, $\frac{1}{2}$ pint of methylated spirit and $\frac{1}{2}$ pint of water. It is to be spread with a sponge on each side of the silk without creasing it; the silk is then rinsed out two or three times, and ironed on the wrong side.

SILVER, TO CLEAN. See PLATE POWDER. To clean silver utensils, blackened by sulphuretted hydrogen, BOTTGER recommends a boiling saturated solution of borax, or a solution of caustic potash, with some fragments of metallic zinc.

SILVERING POWDER, &c., for silvering copper, covering the worn parts of plated goods, &c. 1. Nitrate of silver 30 grs., common salt 30 grs., cream of tartar $3\frac{1}{2}$ drs. Mix. Moistened with water and rubbed on dial plates or other copper articles, it coats them with silver.

2. Silver precipitated from its nitric solution by copper 20 grs., alum 30 grs., cream of tartar 2 drs., salt 2 drs.

3. Precipitated silver $\frac{1}{2}$ oz., common salt 2 oz., chloride of ammonium 2 oz., corrosive sublimate 1 dr., make it into a paste with water. Copper utensils are previously boiled with tartar and alum, and rubbed with this paste, then made red-hot, and afterwards polished.

4. Dissolve chloride of silver in a solution of hyposulphite of soda, and mix this with prepared hartshorn or other suitable powder.

SILVERING PASTE. Nitrate of silver 1 part, cyanide of potassium (Liebig's) 3 parts, water sufficient to form a thick paste. Apply it with a rag. A bath for the same purpose is made by dissolving 100 parts of sulphite of sodium, and 15 of nitrate of silver, in water, and dipping the article to be silvered, into it.

ELECTRO-SILVERING. Mr WATT. 1. To the crystallized nitrate of silver obtained by dissolving 1 oz. of fine silver

in nitric acid, add 3 pints of cold distilled water to dissolve it. Precipitate with common salt. Wash well. Dissolve the chloride in just enough of the strong solution of cyanide of potassium. Filter several times, then add distilled water to 1 gallon. This is excellent for a dead white, or to plate cleaned figures, or clock-dials, which only require to be scratch-brushed first. Or if worked weaker, with a feeble battery, and a small surface of *anode* exposed, a thick layer is obtained, which will bear bur-nishing.

2. Dissolve as much nitrate in the same quantity of distilled water. Add strong solution of cyanide of potassium till there is no further precipitate—(not enough to re-dissolve this). Wash the precipitated cyanide several times with pure water. Add now enough strong solution of cyanide of potassium to dissolve it, and make up the solution to 1 gallon. Filter before using. When it is desired that the articles should come out with a bright appearance, a little bisulphide of carbon is added to the solution.

SILVER, OXIDIZED. Ornaments to which this name is given have a surface of silver which has been acted on by some chemical liquid. A solution of sal ammoniac will give a brownish tint; a still better one is obtained by using equal parts of sulphate of copper and sal ammoniac dissolved in vinegar. A fine black tint may be produced by a slightly warm solution of sulphide of potassium. Solutions of the chlorides of platinum and gold are sometimes adopted in these processes.

SILVERING GLASS. See **GLASS.**

SILVER, TO PURIFY AND REDUCE. Silver, as used in the arts and coinage, is alloyed with a portion of copper. To purify it, dissolve the metal in nitric acid slightly diluted, and add common salt, which throws down the whole of the silver in the form of chloride. To reduce it to a metallic state several methods are used:—1. The chloride must be repeatedly washed with distilled water, and placed in a zinc cup; a little diluted sulphuric acid being added, the chloride is soon reduced. The silver when thoroughly washed is quite pure. In the absence of a zinc cup, a porcelain cup containing a zinc plate may be used. The process is expedited by warming the cup.

2. **HORNUNG.** Digest the washed chloride with pure copper and ammonia. The quantity of ammonia need not

be sufficient to dissolve the chloride. Leave the mixture for a day, then wash the silver thoroughly.

3. Boil the washed and moist chloride in solution of pure potash, adding a little sugar; when washed it is quite pure.

SILVER, SOLVENT FOR. See further on.

SIZE. *Oil Size* is made by grinding yellow ochre or burnt red ochre with boiled linseed oil, and thinning it with oil of turpentine. *Water Size* (for burnished gilding) is parchment size ground with yellow ochre.

SMALTS. See **PIGMENTS.**

SKELETON LEAVES, to make. Steep the leaves, seed-vessels, or other parts of the plant to be dissected, in rain-water, until the whole of the soft matter is decomposed. Some require a few weeks, others several months. The rotted parts are now to be carefully removed by a fine brush, under the surface of water, or in a stream of water. A syringe is sometimes required. To bleach the skeletons, soak them for some hours in a mixture of 1 oz. of strong solution of chloride of lime and a quart of distilled water, or in a solution of peroxide of hydrogen. Lastly, wash thoroughly in cold water, and dry by exposure to air.

SMOKING FLUID. One drop of creasote in a pint of water imparts a smoky flavour to fish or meat dipped into it for a few minutes.

SOAP. FOR PERFUMED AND TOILET SOAPS, see **SKIN COSMETICS**, further back. For the manufacture of soaps generally, see Dr. URE's 'Dictionary of the Arts,' WAGNER'S 'Chemical Technology,' 'Spon's Cyclopædia of the Industrial Arts,' and other similar works. Hard soaps are made by boiling oils or fats with a ley of caustic soda. Soft soaps consist of oil and potash; and as they do not separate from the ley like the hard soaps, they generally contain an excess of caustic alkali. Silica soap has silicate of soda incorporated with it. Soap is adulterated by earthy matters, as pipe-clay, &c.; these and other impurities remain when soap is dissolved in alcohol.

SOAP, MARINE. *Patent.* This is made by substituting cocoa-nut oil for the fats and oils used in the manufacture of common soap. It has the advantage of forming a lather with salt water.

SODA. For its medical and pharmaceutical compounds, see *Pocket Formulary*.

SODA, HYPOSULPHITE OF. Dissolve 1 lb. of crystallized carbonate of soda in a quart of boiling water. Slake $\frac{1}{2}$ lb. of lime in another quart of water. Mix the solutions, let them stand in a covered vessel until cold, pour off the clear liquid, and boil it with more sulphur than it will dissolve. Pour off the clear solution into a deep vessel, and pass sulphurous acid gas through it until it becomes nearly colourless. While still a little yellow, filter, and evaporate it quickly in an earthen vessel to a syrupy consistence. Shake this with half its bulk of rectified spirit, and allow the lower layer to crystallize under the alcoholic solution which floats on it. It must be kept from the air and light.

SOLVENTS FOR INDIA RUBBER. *Ether* for this purpose should be agitated with water and decanted. *Benzol* will dissolve caoutchouc with warmth and long digestion. *Rectified coal naphtha* forms an imperfect solution employed in MACINTOSH'S waterproof fabrics. *Oil of turpentine*, rendered pyrogenous by absorbing it with bricks of porous ware, and distilling it without water, and treating the product in the same way, is also used for this purpose. It is stated that the solution on evaporation does not leave the caoutchouc in a sticky state. Another method is to agitate oil of turpentine repeatedly with a mixture of equal weights of sulphuric acid and water; and afterwards to expose it to the sun for some time. *Bisulphide of carbon* is a good solvent, dissolving the gum without heat. *Crude PETROLEUM* also dissolves it. This constitutes PARKES' Patent Solvent. *Chloroform* is an excellent but rather expensive solvent.* *Caoutchoucine* has also been employed as a solvent. It is prepared by distilling india rubber without addition, increasing the heat to 600° Fahr. The product is rectified by distilling it with one third of water. It is then a colourless fluid of 0.680 specific gravity. Its smell is improved by agitating it with 5 oz. of nitro-hydrochloric acid to each gallon. India rubber is rendered more readily soluble by first digesting it with a solution of carbonate of soda, or water of ammonia.

SOLVENTS FOR GUTTA PERCHA. Benzol readily dissolves it. So do chloroform and bisulphide of carbon.

* Methylated chloroform is much cheaper than that made from rectified spirit.

SOLVENT FOR OLD PUTTY AND PAINT. Soft soap mixed with solution of potash or caustic soda; or pearlash and slaked lime mixed with sufficient water to form a paste. Either of these laid on with an old brush or rag, and left for some hours will render it easily removable.

SOLUBLE GLASS. See GLASS.

SOLUTIONS USED IN ELECTROTYPE MANIPULATIONS, &c.

For the Decomposing Cell. 1. Saturated solution of sulphate of copper 2 parts, sulphuric acid 2 parts, water 6 or 8 parts. 2. Mr. WALKER directs 2 measures of a saturated solution of sulphate of copper, and 1 measure of acidulated water (1 part of sulphuric acid to 9 of water). 3. ROBELL'S solution consists of 2 parts of a saturated solution of sulphate of copper, and 1 part of a saturated solution of Glauber's salt, to which as much sulphate of copper has been added as it will take up.

2. *Gold Solution.* Dissolve 2 oz. of cyanide of potassium (prepared by LIEBIG'S method) in a pint of warm distilled water, add $\frac{1}{2}$ oz. of oxide of gold, and agitate together.

3. *Silver Solution.* Dissolve 2 oz. of LIEBIG'S cyanide of potassium in a pint of distilled water; add $\frac{1}{2}$ oz. of moist oxide of silver (precipitated by lime-water from a solution of the crystallized nitrate), and agitate together till the oxide is dissolved.

4. *Solution into which Steel Articles are dipped before Electroplating them.* Nitrate of silver 1 part, nitrate of mercury 1 part, nitric acid (sp. gr. 1.384) 4 parts, water 120 parts.

5. *Solution, or Pickle, for Immersing Copper Articles in before Electroplating.* Sulphuric acid 64 parts, water 64, nitric acid 32, hydrochloric acid 1: mix. The article, free from grease, is dipped into the pickle for a second or two. See GILDING, SILVERING, ELECTRO-BRASSING, &c.

SOLUTION FOR MULLINS' VOLTAIC BATTERY. *In contact with the Zinc:* 1 part of sal ammoniac to 5 of water. *In contact with the Copper:* a saturated solution of sulphate of copper. M. BACHOFFNER used a saturated solution of common salt and a saturated solution of sulphate of copper.

SOLUTION FOR SOLDERING. Dissolve zinc in hydrochloric acid to saturation, add pulverized sal ammoniac, and boil for a short time. Applied with a sponge or feather it facilitates the flow of the solder.

SOLUTIONS FOR THE WATER-BATH. Various salts dissolved in water materially raise the boiling-point, and thus afford the means of obtaining a steady temperature at different degrees above 212° F. The following are some of the most useful :

A saturated solution of nitrate of soda boils at	246°	Fahr.
" " Rochelle salts at . . .	240°	"
" " nitre at	238°	"
" " chloride of sodium at	224°	"
" " sulphate of magnesia at	222°	"

SOLVENT FOR SILVER. *Nitro-sulphuric Acid.* Dissolve 1 part of nitre in 10 parts of oil of vitriol. Used for dissolving the silver from plated goods, &c. It dissolves silver at a temperature below 200° F., and scarcely acts upon copper, lead, and iron, unless diluted. The silver is precipitated from the solution, after moderately diluting it by common salt, and the chloride reduced as directed, under **SILVER, TO PURIFY AND REDUCE.**

SPIRIT OF NITRIC ETHER. See *Pocket Formulary*. It need only be added here, that its acidity is removed and prevented, by rectifying it from neutral tartrate of potash.

SPONGE, BLANCHED. Soak the sponge for several days in cold water, renewing the water and squeezing the sponges occasionally. Then wash them in warm water, and place them in cold water to which a little hydrochloric acid has been added. Next day take them out and wash them thoroughly in soft water; then immerse them in aqueous sulphurous acid (sp. gr. 1.034) for a week. They are afterwards washed in plenty of water, squeezed, and allowed to dry in the air.

2. Soak the sponges, previously deprived of sand and dirt by bleaching and washing, in a 1 per cent. solution of permanganate of potash. Then remove them, wash them thoroughly with water, and press out the water. Next put them into a solution of a half pound of hyposulphite of soda in one gallon of water, to which 1 oz. of oxalic acid has been added and leave them in the solution for 15 minutes. Finally, take them out and wash them thoroughly. *New Remedies.*

For burnt, prepared, and waxed sponge, see **SPONGIA**, *Pocket Formulary*.

SPONGES, SLIMY, TO CLEAN. (**VATTA.**) Press the sponge as dry as possible, place it on a plate, and sprinkle pow-

dered chloride of calcium on it, and allow to deliquesce on the sponge. After half an hour the sponge may be washed in water and dried.

STAINS FOR WOOD, IVORY, &c. See **IVORY, TO STAIN**; **WOOD, TO STAIN**; **BOOKBINDERS' STAINS, &c.**

STAINS, TO REMOVE. Stains of *iodine* are removed by rectified spirit. *Ink* stains by oxalic acid or superoxalate of potash. *Iron moulds* by the same; but, if obstinate, it has been recommended to moisten them with ink, then remove them in the usual way. See **RUST, TO PREVENT AND REMOVE**, further back.

Grease Spots may be removed either by chloroform, liquid ammonia, benzol, oil of turpentine, or ether. The three last are very inflammable. See also **SCOURING DROPS**.

Red Spots on black cloth, from acids, are removed by spirit of hartshorn, or other solutions of ammonia.

Stains of Marking Ink, or Nitrate of Silver, to remove.

1. Wet the stain with fresh solution of chloride of lime, and after 10 or 15 minutes, if the marks have become white, dip the part in solution of ammonia, or of hypsulphite of soda. In a few hours wash with clean water.

2. Stretch the strained linen over a basin of hot water, wet the mark with tincture of iodine.

3. They may also be most readily removed by cyanide of potassium; but this should be done by the druggist, and not entrusted to anyone else, since the cyanide is a potent poison.

STARCH. Starch is procured from various roots and seeds. Its varieties are numerous, but a few of the most important only can be noticed here.

Arrowroot (West Indian). The fecula of the tubers of the *Maranta arundinacea*. The fresh tubers are washed and beaten to a pulp, which is well stirred in a large tub of cold clean water, and the fibrous part wrung out by the hands, and thrown away. The water in which the fecula is suspended is passed through a hair sieve or coarse cloth, allowed to settle, and the water poured off. After being repeatedly washed, the wet starch is drained, and afterwards dried in the sun. The other varieties of arrowroot (see **DIETETIC ARTICLES**) are prepared by analogous processes from the roots which yield them.

Potato Starch. The tubers are washed and peeled, usually by machinery, rasped by a revolving grater, and

the pulp washed on hair sieves till freed from the starchy matter. Successive portions of the pulp are thus treated till the vessels over which the sieves are placed are sufficiently full. The starch held in suspension in water, falls to the bottom; the water is then drawn off, and the starch stirred up with fresh water, and again allowed to subside. This is repeated several times till the starch is sufficiently pure. The fibres and the washing waters are used as manures. The washed fibres have also been recommended as an ingredient in bread for diabetic patients.

Wheat Starch. Wheat flour is steeped in water for a week or two and allowed to ferment. The acid liquor is drawn off, and the residue washed on a sieve; what passes through is allowed to settle, the sour liquor drawn off, and the starch thoroughly washed from the slimy matter. It is then drained in perforated boxes, cut up into square lumps, placed on bricks to absorb the moisture, and dried in a stove.

Various means are adopted to free the starch from gluten and other impurities. In the patent rice starch, and probably other kinds of starch, alkaline solutions are used. Ammonia has been recommended, as it does not, like potash and soda, dissolve any portion of pure starch.

The various kinds of fecula are distinguishable by the form of their particles or grains. By a microscopical examination of these, the mixture or substitution of potato starch with the more expensive kinds is readily detected. Figures of the different kinds of starch grains are given in the *Pharmaceutical Journal*, vol. iv, in Dr PEREIRA'S 'Elements,' in ATTFIELD'S 'Chemistry, General, Medical and Pharmaceutical,' and in the microscopic manuals. M. GOBLEY has proposed to distinguish them by the coloration produced when the several kinds are exposed to the vapour of iodine; but the effect seems to depend greatly on the relative dryness of the samples.

STEARIC ACID. Fat is saponified, and the soap decomposed by an acid, with a large quantity of water, the mixture being kept warm and well stirred. The water being drawn off, the stearic acid is well washed, allowed to cool, and submitted to strong pressure.

STORM GLASS. Take $2\frac{1}{2}$ drs. of camphor, 38 grs. of nitre, and 38 grs. of sal ammoniac; dissolve them in 9 drs. of water, and 6 drs. of rectified spirit, with a gentle heat.

This is placed in a glass tube covered with a brass cap, with a small hole to admit air. Or it may be put in an eau de Cologne or other long bottle, tied over with bladder. Its various changes are supposed to indicate changes of weather, but the indications are not to be relied on.

STUFFING BIRDS AND ANIMALS, PREPARATIONS FOR. 1.

Camphor 1 oz., corrosive sublimate $\frac{1}{2}$ oz., alum $\frac{1}{2}$ oz., sulphur 1 oz.; all finely powdered and mixed.

2. Tanner's bark dried and powdered 2 oz., burnt alum 1 oz., snuff 1 oz.; mix, and add arsenic $\frac{1}{4}$ oz., camphor $\frac{1}{4}$ oz., sulphur 1 dr.

3. BÉCŒUR'S *Arsenical Soap*. Camphor 5 drs., arsenic 4 oz., white soap 4 oz., carbonate of potash 12 oz., air-slaked lime 4 oz.; make a stiff paste with a little water.

STYROL. Mix 20 parts of storax with 7 of carbonate of soda, and put them into a retort with water, and apply heat. A limpid fluid distils, which becomes when heated to a certain point, a transparent solid.

SUGAR RESIN. Mix 16 parts of strong sulphuric acid with 8 of the strongest nitric acid; when cooled to 70° Fahr., stir in 1 part of finely powdered sugar. In a few seconds, when the sugar has become pasty, take it out of the acid and plunge it into cold water. Add more sugar to the acid, and proceed as before. Wash the resinous matter carefully, and dissolve it in alcohol or ether. Evaporate the solution with a gentle heat. It is very combustible. Its solution may be used to render gunpowder, lucifer matches, &c., waterproof.

SULPHIDE. A sulphide was formerly termed a *sulphuret*, which latter name has been almost invariably retained in the present edition.

SULPHITE OF COPPER. To a concentrated solution of bisulphite of potash add a cold solution of sulphate of copper, filter, and heat gently.

SULPHOCYANIDE OF AMMONIUM. Neutralize hydrosulphocyanic acid with ammonia and gently evaporate the solution to dryness over a water-bath.

SULPHURET OF CARBON. See BISULPHURET OF CARBON.

SYRUP OF MILK. Evaporate, with constant stirring, 6 lbs. of skimmed milk to 3 lbs.; add 4½ lbs. of sugar; dissolve with a gentle heat, and strain. It may be flavoured with the addition of 1 oz. of cherry-laurel water. [For other Syrups see *Pocket Formulary*.] Milk may be preserved

by first heating it, and when cold, charging it with carbonic acid gas. Boracic acid is now very largely used for preserving milk.

TANNIN. See ACID, TANNIC, further back.

TERPINE. Leave oil of turpentine for a long time in contact with a mixture of nitric acid and alcohol. Crystals of terpine form. By boiling an aqueous solution of terpine with a small quantity of sulphuric or other acid, *terpinol* is formed, and may be separated by distillation. It has the odour of hyacinths.

TEST LIQUORS, TEST PAPERS, &c. For TEST LIQUORS see *Pocket Formulary*. In preparing the papers, the liquid should be placed in an earthenware plate or dish, and the paper carefully immersed in it so as to be uniformly wetted, then dried out of reach of acid, ammoniacal, or other vapours likely to affect it; and afterwards kept in bottles, jars, or cases. Dr FARADAY recommends unsized paper, but Mr PARNELL and other good authorities direct good letter paper to be used.*

Brazil Paper. Dip paper in a strong decoction of Brazil-wood and dry it. It is rendered purple or violet by alkalis; generally yellow by acids.

Cabbage Paper. Make a strong infusion of red cabbage leaves, strain it, and evaporate it by a gentle heat till considerably reduced. Then dip the paper in it and dry it in the air. This paper is of a greyish colour; alkalis change it to green, acids to red. It is a very delicate test; if rendered slightly green by an alkali, carbonic acid will restore the colour.

Dahlia Paper. From the petals of violet dahlias, as cabbage paper.

Elder-berry Paper. This is merely paper stained with the juice of the berries. Its blue colour is changed to red by acids, and to green by alkalis.

Indigo Paper. Immerse paper in sulphate of indigo, wash it with water rendered slightly alkaline, then with pure water, and dry it in the air.

Iodide of Potassium and Starch Paper. Mix starch paste with solution of iodide of potassium, and moisten bibulous paper with it. It becomes blue when exposed to ozone. Chlorine has the same effect.

Lead Paper. Paper dipped in a solution of acetate of

* Dr Fresenius recommends unsized paper.

lead. When moistened it detects sulphuretted hydrogen, which renders it black.

Blue Litmus Paper. Bruise 1 oz. of litmus in a mortar, and add boiling water; triturate together, put them in a flask and add boiling water to make up to half a pint; when cool, strain it, and dip paper in it. More colour may be extracted from the litmus by hot water, but the liquid will require to be concentrated by evaporation. Acids change the colour to red, but it does not become green with alkalis.

Red Litmus Paper. As the last, adding to the strained infusion a few drops of nitric acid, or of pure acetic acid. Dr FARADAY recommends holding blue litmus paper over a large jar, into which a few drops of hydrochloric acid have been introduced, till sufficiently reddened.

Rose Paper. Make a strong infusion of the petals of the red rose, and dip unsized paper in it. Dipped in an alkaline solution, so weak as not to affect turmeric paper, it assumes a bright green colour.

Manganese Paper. Dip paper in a solution of sulphate of manganese. It becomes black in an ozonized atmosphere.

Rhubarb Paper. Dip paper in a strong infusion of rhubarb and dry it. Alkalis render it brown. It is not, like turmeric paper, affected by boracic acid.

Starch Paper. This is merely paper imbued with starch paste. Cotton cord is sometimes used instead of paper. As a test for iodine, which it turns blue.

Turmeric Paper. Boil 1 oz. of coarsely powdered turmeric root in half a pint of water for half an hour, and strain: dip paper in the liquid, and dry it. It is rendered brown by alkalis, and also by boracic acid and borates.

TEST SOLUTIONS, &c. The vegetable preparations are here placed first.

Tincture and Infusion of Red Cabbage. Digest red cabbage with rectified spirit in a warm place for a few days; strain, distil off most of the spirit, and evaporate what remains to the consistence of syrup. It will keep for years. When required for use, dilute it with a little water; or the concentrated infusion directed above for the paper may have a little spirit added to it. If the cabbage leaves be well dried, they may be kept in a close vessel for use, and a strong infusion made when wanted.

Acid Infusion of Red Cabbage. FARADAY directs one or more red cabbages to be cut up in strips, and boiling water poured on them, and a little dilute sulphuric acid (equal to $\frac{1}{2}$ oz. of oil of vitriol to a large cabbage) to be added, and the whole kept hot for an hour or two in a copper or earthen vessel. It is then strained, the cabbage infused in a little more water and acid, and the mixed infusion evaporated to one third its first bulk, allowed to settle, and put into bottles. When required for use, the acid is neutralized by caustic potash or soda. Another plan is to dry the leaves at 120° F.; and when required for use to make a strong infusion, adding a drop of sulphuric acid, to neutralize the strained infusion with marble, filter, and add a little spirit, if required to be kept.

Infusion or Tincture of Litmus. This is made as directed above for litmus paper. Or an ounce of powdered litmus may be triturated with 6 oz. of boiling water, digested near the fire for an hour, and mixed, when cool, with 2 oz. of spirit. Or digest 1 oz. of powdered litmus in a pint of proof spirit for 7 days. If required red, a few drops of acetic acid are added to either of these. The next day decant the clear liquor. Dr PEREIRA directs 1 part of litmus to 25 of water. When made very strong, it must be diluted when used.

Tincture of Galls, Infusion of Galls, &c. Fresh powdered blue galls 1 oz., proof spirit 8 oz.; digest in a close vessel for a week, and filter. A watery infusion of galls may be made in the same proportion with boiling water, for immediate use. PETTENKOPFER directs 1 oz. of powdered galls to be infused in 3 or 4 oz. of boiling water for several hours, and 2 oz. of salt added. After filtration, it retains its transparency and power of precipitating gelatin for years. [This is used to detect iron, with the persalts of which it produces a bluish-black precipitate; for gelatin, which it precipitates in brownish-white flocks; and several of the organic alkaloids.]

MARSH'S Dahlia Test. Make a strong infusion of the petals of dark dahlias; strain, and add to every pint $\frac{1}{2}$ oz. of strong sulphuric acid; stir with a glass rod, and when cold, add to each pint 2 grs. of corrosive sublimate.* Filter through coarse cloth, and bottle. When required

* The corrosive sublimate is an undesirable ingredient.

for use, neutralize it carefully with ammonia, and use the liquid by dipping paper in it.

Syrup of Violets. On 4 oz. of fresh petals of violets pour half a pint of water at 104° F., stir them together, and in a minute or two strain off the water with gentle pressure, and pour 8 oz. of boiling distilled water on the flowers. In 12 hours strain through linen, let the infusion settle, and decant. A delicate test for acids and alkalies.

Dr CLARK'S Test for Hardness of Water. Dissolve 1 oz. of Hawes's best white soap in a gallon of proof spirit. If not of such strength that it requires 32 measures to be added to 100 measures of solution of chloride of calcium of 16 degrees of hardness (see below) before it lathers, it must be adjusted to that strength. The chloride of calcium solution is thus made:—Dissolve 16 grs. of pure carbonate of lime (Iceland spa) in a small quantity of pure hydrochloric acid, avoiding loss from effervescence; evaporate the solution to dryness, and dissolve the residue in water, and again evaporate till a neutral solution is obtained; then dissolve in a gallon of water. This forms the standard solution of 16 degrees of hardness. One measure of this solution with 15 of distilled water constitutes a solution of one degree of hardness; and so on up to 16 degrees. The degree of hardness expresses the number of grains of carbonate of lime per gallon contained in the water. For the mode of using this test, see Dr CLARK'S pamphlet, or any of the manuals on "Water Analysis."

HAHNEMANN'S Wine Test, for detecting lead in wine. Sulphuret of lime 3 oz., tartaric acid 3 oz., water 2 lbs., mix, decant, and add 4 oz. of tartaric acid. Or, simple sulphuretted hydrogen water 4 oz., tartaric acid 1 dr. For the detection of other adulterants in wine, see Article "Wine and Wine-making" in vol. i of 'Churchill's Technological Handbooks.'

TROMMER'S Test for sugar in urine. Put some of the suspected urine into a large test-tube, and add a few drops of solution of sulphate of copper, then sufficient solution of potash to render it strongly alkaline. If sugar be present, the precipitated oxide dissolves into a blue liquid, and on boiling, red suboxide of copper is precipitated.

"FEHLING found that one equivalent of grape-sugar or 180 parts, decomposed exactly ten equivalents, or 1246·8

parts, of sulphate of copper. Accordingly he prepared a solution of copper of standard strength and applied it to fluids containing grape-sugar; and the quantity of these required to decompose a fixed volume of the standard solution furnished an exact measure of the sugar they contained.

"Fehling's standard solution is prepared according to the following prescription:—Sulphate of copper $90\frac{1}{2}$ grs., neutral tartrate of potash 364 grs., solution of caustic soda (sp. gr. 1.12) 4 fl. oz., add water to make up 6 fl. oz. 200 grs. of this solution are exactly decomposed by 1 gr. of sugar."—Dr ROBERTS.*

Dr ROBERTS gives the following directions for the quantitative examination of urine for sugar:—"Pour some of the prepared test-liquor ("Fehling's," as above) into a narrow test-tube to the depth of $\frac{3}{4}$ of an inch; heat until it begins to boil, then add 2 or 3 drops of the suspected urine. If the sugar be abundant a thick yellowish opacity and deposit of yellow suboxide are produced (and this changes to a brick-red at once if the blue colour of the test remain dominant). If no such reaction ensue, go on adding the urine until a bulk nearly equal to the test employed has been poured in; heat again to ebullition, and, no change occurring, set aside without further boiling. If no milkiness is produced as the mixture cools, the urine may confidently be pronounced free from sugar."

Picric Acid Test for sugar. Consult Dr GEORGE JOHNSON'S paper in the *Pharmaceutical Journal* for June 9th, 1883. The length of the paper precludes its insertion here.

Nitric Acid Test for albumen in urine. Fill a test-tube to the depth of about an inch with the urine, then incline the tube and pour in strong nitric acid, so that it may trickle down along the side of the tube to the bottom and form a stratum some quarter of an inch thick below the urine. If the urine contain albumen three layers will be perceptible—one, perfectly colourless, of nitric acid at the bottom; immediately above this an opalescent zone of the coagulated albumen, and on the top the unaltered urine.

Heat Test for albumen in urine.—"If the urine have its usual acid reaction it becomes turbid on boiling when it

* For the method of performing the analysis, &c., consult Dr Roberts' work, 'Urinary and Renal Diseases.'

contains albumen, and this turbidity persists after the addition of an acid. There are two points to be remembered on using heat alone as a test for albumen:—First, that albumen is not coagulated by heat when the urine is alkaline; in such cases, therefore, it is necessary before boiling to restore the acidity by a few drops of acetic acid. Secondly, when the urine is neutral, or very feebly acid, it may become turbid on heating, from precipitation of the earthy phosphates; but turbidity from this cause is easily distinguished from albumen by a drop of acetic or nitric acid, which instantly causes the phosphates to disappear.”—Dr ROBERTS.

Picric Acid Test for albumen in urine. Into a test-tube about six inches long, the urine is poured to within two inches of the top, then the tube being held in a slanting position, about an inch of a saturated aqueous solution of picric acid is gently poured on the surface of the urine, when in consequence of its low sp. gr. (1003), it only partly mixes with the upper layer of the urine, and as far as the yellow colour of the picric solution extends there will be more or less turbidity from coagulated albumen, contrasting with the pellucid unstained urine below. If, then, the tube be placed in a stand, the coagulated albumen will gradually subside, and form a delicate horizontal film at the junction of the coloured and the unstained stratum of urine, the yellow liquid above, and the unstained urine below being quite free from turbidity.

If the urine should be turbid with urates, it must be cleared by heat before the addition of the picric acid solution. If the urine be highly alkaline it must be acidulated with a vegetable or mineral acid before adding the picric acid solution.—Dr GEORGE JOHNSON.*

PETTENKOFER'S Test for bile in urine, &c. Put a small quantity of the suspected liquid into a test-tube, and add to it, drop by drop, strong sulphuric acid till it becomes warm, taking care not to raise the temperature above 122° Fahr. Then add from 2 to 5 drops of syrup, made with 5 parts of sugar to 4 of water, and shake the mixture. If the liquid contain bile, a violet coloration is observed. Acetic acid, and those substances which are converted into sugar by sulphuric acid, may be substituted for sugar.

* Consult Dr Johnson's paper in the *Pharmaceutical Journal* for June 9th, 1883.

Another test consists in placing a little of the suspected urine in a test-tube, and adding to it a few drops of tincture of iodine, when, if bile be present, the fluid becomes distinctly green.

TOBACCO WATER. See WASHES for vermin on plants.

TOUCH PAPER. Dip a piece of white blotting-paper, or printing-paper, in a solution of 1 oz. of nitre in 8 oz. of water. Dry it perfectly.

TRAUMATICINE AUSPITZ. Gutta percha 1 part, chloroform 10 parts.

TREES, METALLIC. *Lead Tree.* Dissolve 1 oz. of sugar of lead in a quart of distilled or filtered rain-water, adding a few drops of acetic acid. Filter, and put the clear solution into a decanter or bottle. Suspend in it a piece of zinc, and set it aside.

Silver Tree. Dissolve 20 grs. of crystallized nitrate of silver in an ounce of distilled water; put it into a phial, and add about $\frac{1}{2}$ a drachm of pure quicksilver.

Tin Tree. Dissolve 3 drs. of chloride of tin in a pint and a half of water, with 10 or 15 drops of nitric acid; and suspend in it a rod of zinc.

TURPENTINE, VENICE (*factitious*). It is usually made by dissolving black resin in oil of turpentine. Dr PEREIRA states the proportion to be 5 fluid oz. of the oil to 16 oz. of resin; but some makers put as much as 8, 10, or even 12 oz. of oil of turpentine to each pound of resin. We have introduced this factitious preparation, because no genuine Venice (or larch) turpentine is rarely, if ever, to be obtained.

TURPENTINE, OIL OF. Common turpentine, chiefly American, is distilled with water; the oil comes over with the water and is found floating on it. It is rectified by distilling it again with water. See CAMPHINE and SOLVENTS for INDIA RUBBER, for further modifications of this oil.

URN POWDER. Peroxide of iron, crocus, or jeweller's rouge.

VANILLIN. A crystalline substance obtained from coniferin, (the glucoside contained in the cambium of coniferous woods), by digesting it with water and a small quantity of emulsin. It has been shown to be identical with the aromatic principle of vanilla. It must be kept for a twelvemonth before it is fit for use.

VARNISHES. These constitute a distinct branch of manufacture, and many of them can be advantageously or

safely made only on the large scale on premises adapted for the purpose. A few of the most easily prepared and useful varnishes have been selected for insertion. For fuller information, see Dr URE'S 'Dictionary of Arts,' DUMAS' 'Chimie appliquée aux Arts,' &c. Some practical information on this subject will be found in Mr REDWOOD'S edition of GRAY'S 'Supplement,' and in the 49th vol. of the 'Transactions of the Society of Arts;' also in SPONS' 'Workshop Receipts.'

Spirit Varnishes. The spirit* employed should not be less than 60° overproof. In preparing and using varnishes, they should be kept at a distance from a candle or other flame. Respecting the gum-resins employed, it may be useful to mention that shell-lac is rendered more soluble by being powdered and exposed for a long time to the air; sandarach gives hardness to varnishes; mastic gives a gloss to solutions of other gums; benzoin still more, but its colour is objectionable; anime readily dissolves, but renders the varnish long in drying; copal and amber are scarcely soluble in spirit, but are rendered partially so by other gums, and also by being previously fused by heat. Shell-lac gives a durable varnish, objectionable only on account of its colour, which may be rendered paler by charcoal. See LAC.

1. *White Spirit Varnish.* Methylated spirit 2 gallons, gum sandarach 5 lbs. Put them into a tin bottle, cork securely, and agitate frequently, placing the tin occasionally in hot water till the gum is dissolved, then add a quart of pale turpentine varnish.

2. *Brown.* Methylated spirit 2 gallons, sandarach 3 lbs., shell-lac 2 lbs., pale turpentine varnish a quart. Proceed as the last.

3. Sandarach 2 oz., shell-lac $\frac{1}{2}$ oz., methylated spirit 16 fluid oz.

4. *White.* Gum sandarach $1\frac{1}{2}$ oz., mastic $\frac{1}{2}$ oz., elemi $\frac{1}{4}$ oz., foreign oil of lavender $\frac{1}{2}$ oz., methylated spirit 8 oz.

Copal Spirit, or Drying Varnish. Copal, fused and pulverized, 3 oz., sandarach 6 oz., mastic 3 oz., Venice turpentine $2\frac{1}{2}$ oz., methylated spirit a quart, powdered glass 3 oz. Mix the powdered glass and resins, and sift them; introduce them into a matrass with the spirit, and heat to

* Methylated spirit is now universally employed in making spirit varnishes, being equal to, as well as very much cheaper than, duty-paid spirit.

boiling, constantly agitating till the gums are dissolved; then add the turpentine. Heat the varnish for half an hour, and when removed from the fire, agitate till cold.

Brilliant Amber Spirit Varnish. Fused amber 4 oz., sandarach 4 oz., mastic 4 oz., methylated spirit a quart. Expose to the heat of a sand-bath, with occasional agitation, till dissolved. [The amber is fused in a close copper vessel, having a funnel-shaped projection, which passes through the bottom of the furnace by which the vessel is heated.]

Amber Varnish for Photographs. Dissolve 3 to 4 grs. of amber in 1 oz. of chloroform.

1. *Colourless Varnish for Photographs.* Dissolve shell-lac by heat in 8 parts of water and 1 part of pearlsh. Precipitate by chlorine, and dissolve in rectified spirit.—
Dr HARE.

2. 1 oz. white lac, dissolved in 10 oz. of warm methylated spirit of wine. Let it settle for several weeks, then carefully decant for use.

Chinese Varnish. Mastic 2 oz., sandarach 2 oz., rectified spirit a pint. Close the matrass with bladder, with a pin-hole for the escape of vapour; heat to boiling in a sand- or water-bath, and when dissolved, strain through linen.

Crystal Varnish. Picked mastic 4 oz., methylated spirit a pint, animal charcoal 1 oz. Digest and filter.

French Polish and Lacquers are varieties of spirit varnishes. The former has already been noticed. A few formulæ for the latter are here added.

Pale or Gold Lacquers. To a pint of methylated spirit add as much gamboge as will give it a bright yellow colour, then add 12 oz. of seed-lac in fine powder, and set it in a sand-bath till dissolved. Or a tincture of annatto (1 part to 8 of spirit) may be added to give the desired colour.

Dark Lacquer. Clear seed-lac 1 lb., dragon's blood 1 oz.; pulverize together, and add them to a pint and a half of methylated spirit. Set in a warm place till dissolved.

Lacquer for Brass Work. Turmeric 1 oz., saffron $\frac{1}{4}$ oz., Spanish annatto $\frac{1}{4}$ oz., methylated spirit a pint. Digest at a gentle heat for several days; strain through coarse linen, put the tincture into a bottle, and add 3 oz. of good seedlac coarsely powdered. Put in a moderately hot place and shake

frequently till dissolved; if wanted of a redder shade, increase the quantity of annotto, or add a little dragon's blood. [Some makers prepare a strong tincture of the various colouring ingredients, and add them to the lacquer to produce the required shade.

Oil of turpentine, and other essential oils, are used as solvents, forming essence varnishes, as the following :

Mastic Varnish. Clean mastic 5 oz., rectified oil of turpentine a quart. Digest in a warm place, shaking frequently till the solution is complete, then strain.

Picture Varnish. For Oil Paintings. Chio turpentine $2\frac{1}{2}$ oz., mastic 12 oz., camphor $\frac{1}{2}$ dr., powdered glass 4 oz., rectified oil of turpentine 3 pints.

Canada Varnish. For Maps, Drawings, &c. Clear balsam of Canada 4 oz., rectified oil of turpentine 8 oz.; warm gently, and shake together till dissolved. The drawings are first sized over with a solution of isinglass, taking care that every part is covered; when dry, the varnish is brushed over it.

TINGRY'S Essence Varnish. Mastic in powder 12 oz., pure turpentine $1\frac{1}{2}$ oz., camphor $\frac{1}{2}$ oz., powdered glass 5 oz., rectified oil of turpentine a quart.

Common Turpentine Varnish. This is merely clear pale resin dissolved in oil of turpentine; usually 5 lbs. of resin to 7 lbs. of turpentine.

Oil Varnishes. These consist of copal and other gums dissolved by heat in boiled linseed oil; generally with the addition of oil of turpentine.

Cabinet Varnish. Fuse 7 lbs. of African copal, and pour on it 4 pints of hot clarified linseed oil (see OILS); in 3 or 4 minutes, if it feels stringy, take it out of the building, where there is no fire near, and when it has cooled to 150° F., mix in 3 gallons of oil of turpentine of the same temperature, or sufficient to bring it to a due consistence. [Various qualities of copal varnish are made for different purposes; inferior gums are often substituted for, or mixed with, copal.]

Amber Varnish. Amber 16 oz.; melt in an iron pot, and add $\frac{1}{2}$ pint drying linseed oil, boiling hot, and then 3 oz. of resin and 3 oz. of asphalt, each in fine powder. Stir till they are thoroughly incorporated; remove from the fire, and add a pint of warm oil of turpentine.

Common Oil Varnish. Resin 3 lbs., drying oil $\frac{1}{2}$ gallon; melt together, and add, when removed from the fire, 2 quarts of warm oil of turpentine.

Varnish for Printers' Ink. To every 10 lbs. of clarified linseed oil (see further back) add 5 lbs. of clear black resin and $\frac{1}{2}$ lb. oil of turpentine. It is then ready for mixing with lamp black or other colouring matter. A twelfth part of Canada balsam is sometimes added for the finer sorts.

A rapidly-drying Varnish is prepared, according to Mr Dauner, as follows—Mix intimately colophony with thick milk of lime. After 24 hours dry by heat and powder. Melt 100 parts of pine resin, add, with constant stirring, 10 to 15 parts of the above powder. Continue the heat for thirty minutes, remove from the fire, and add 25 to 50 parts of linseed oil, and 35 to 90 parts of turpentine, according to the thickness desired.

A few miscellaneous varnishes are added.

Varnish for Engraving on Copper. Yellow wax 1 oz., mastic 1 oz., asphaltum $\frac{1}{2}$ oz.; melt, pour into water, and form into balls for use. A softer varnish for engravers is made with 1 part of tallow and 2 of yellow wax; or with 2 oz. of wax, 1 dr. of common turpentine, and 1 dr. olive oil. See ETCHING VARNISHES, further back.

Varnish for Engraving on Glass. 1. Wax 1 oz., mastic $\frac{1}{2}$ oz., asphaltum $\frac{1}{2}$ oz., turpentine $\frac{1}{2}$ dr.

2. Mastic 15 parts, turpentine 7 parts, oil of spike 4 parts.

LE BLOND'S Varnish. Keep 4 lbs. of balsam of copaivi warm in a sand- or water-bath, and add 16 oz. of copal, previously fused and coarsely powdered, by single ounces daily, and stir it frequently; when dissolved add a little Chio turpentine.

BESSEMER'S Varnish, for Metallic Paint. This is made with 8 lbs. of copal, $2\frac{1}{2}$ gallons of drying oil, and 25 gallons of oil of turpentine. These are made into a varnish nearly as directed for CABINET VARNISH; and afterwards mixed with a gallon of slaked lime, and left for 3 days to settle. The clear portion is then drawn off, and 5 parts of varnish mixed with 4 parts of bronze powder.

MACINTOSH'S Caoutchouc Varnish. Dissolve 1 lb. of india rubber cut in shreds in a quarter of a pint of rectified coal naphtha. [Caoutchouc varnishes may be made

with any one of the solvents for it, noticed before. The following are also used:]

India-rubber Varnish, for boots. Dissolve $\frac{1}{2}$ oz. of caoutchouc in 2 oz. of mineral naphtha. Dissolve also $\frac{1}{2}$ oz. of asphaltum in 1 oz. of oil of turpentine. Mix the solutions.

Balloon Varnish. Melt india rubber in small pieces with its weight of boiled linseed oil, and thin it with oil of turpentine.

Varnish for Frames for Hot Beds. Mix 4 oz. of pulverized white cheese, 2 oz. of slaked lime, and $\frac{1}{2}$ oz. of boiled linseed oil. Mix, and add $\frac{1}{2}$ oz. each of whites and yolks of egg, and liquefy the mixture by heat. This curious mixture is said to produce a pliable and transparent varnish.

Coloured Varnishes. Oil varnishes are coloured by grinding with them the most transparent colours, as distilled verdigris for green, &c. Spirit varnishes are also coloured with dragon's blood, gamboge, &c.

Sealing-Wax Varnish. Black or coloured sealing-wax broken small, and sufficient rectified spirit to cover it, digested till dissolved. An article called black lac is sold as an economical substitute for black sealing-wax.

Black Japan for Leather, &c. Boil together a gallon of boiled linseed oil, 8 oz. of umber, and 3 oz. of asphaltum. When sufficiently cool, thin in with oil of turpentine.

Japan for Tin-ware. 1. Common copal varnish. 2. Dissolve copal 2 oz., and camphor 1 dr., in oil of turpentine 8 oz.

Brunswick Black. Melt 4 lbs. of asphaltum and 2 lbs. of hot boiled linseed oil together, and when sufficiently cool add a gallon of oil of turpentine.

Varnish for Gun Barrels, after browning them. Shell-lac 1 oz., dragon's blood $\frac{1}{2}$ oz., methyated spirit a quart. Dissolve and filter.

Transfer Varnish. Alcohol 5 oz., pure Venice turpentine 4 oz., mastic 1 oz.

Hair Varnish. Dissolve 1 part of clippings of pig's bristles, or of horsehair, in 10 parts of drying linseed oil by heat. Fibrous materials (cotton, flax, silk, &c.), imbued with the varnish and dried, are used as a substitute for haircloth.

Glass Varnish. Used to protect Wood, &c., from fire. This is a solution of soluble glass, and should be thus made:—Fuse together 15 parts of powdered quartz (or of fine sand), 10 parts of potash, and 1 part of charcoal. Pulverize the mass, and expose it for some days to the air; treat the whole with cold water, which removes the foreign salts, &c. Boil the residue in 5 parts of water until it dissolves. It is permanent in the air, and not dissolved by cold water.

Label Varnish. Sandarach in coarse powder 100 parts, oil of turpentine 40 parts, copaiba 15 parts, alcohol (*absolute*) 90 parts, mastic 40 parts, Venice turpentine 30 parts, alcohol 90 parts. Macerate until dissolved.

1. *Best pale Carriage Varnish.* 8 lbs. of 2nd sorted African copal, 2½ gallons of clarified oil; boil till very stringy. ¼ lb. of dried copperas, ¼ lb. of litharge, 5½ gallons turpentine; strained. 8 lbs. of 2nd assorted gum anime, 2½ gallons of clarified oil, ¼ lb. of dried sugar of lead, ¼ lb. of litharge, 5½ gallons of turpentine; mix with the former part while hot. This varnish will take 4 hours boiling in summer and 6 hours in winter—if well boiled will dry hard. It is intended to be used for varnishing the wheels, springs, and carriage parts of coaches, it has a strong gloss and dries quickly.

2. *Oak Varnish.* Clear pale resin 3½ lbs., oil of turpentine 1 gallon. Dissolve. If it is required of a darker colour, add a little fine lamp-black.

3. *Dammar Varnish.* Gum dammar 10 parts, gum sandarach 5 parts, gum mastic 1 part. Digest at a low heat, with spirit of turpentine 20 parts, occasionally shaking. Add sufficient turpentine to bring it to the consistence of syrup.

For gilt articles. Gum lac, gamboge, dragon's blood, and annotto, of each 125 parts, saffron 32 parts. Dissolve each resin in 1000 parts by measure of absolute alcohol. Two separate mixtures must be made with the dragon's blood and annotto in the above proportions of alcohol; and a proper proportion of each should be added with the gamboge to the varnish according to the shade of colour required.

For Polished Metal. 1. Take shell-lac pounded in a mortar, place the bruised fragments into a bottle of alcohol until some shell-lac remains undissolved; agitate

the bottle and contents frequently, and let the whole stand till clear; pour off the clear fluid. This forms the varnish. Warm the metal surface, and coat with a camel-hair brush. If not perfectly transparent, warm the varnished surface before a fire or in an oven until it becomes clear.

2. Digest 1 part of bruised copal in 2 or 3 parts of absolute alcohol and 1 part of oil of rosemary; this gives a clear varnish as limpid as water. It should be applied hot, and when dry it will be found hard and durable.

3. Mix equal quantities of Canada balsam and very clear spirit of turpentine until of the consistence of ordinary varnish. This may be applied without heating either the metal or the varnish.

For Silver. Gum elemi 30 parts, white amber 45 parts, charcoal 30 parts, spirit of turpentine 375 parts, used in a heated condition, both metal and varnish.

For Iron and Steel. Dissolve 10 parts of clear grains of mastic, 5 parts of camphor, 15 parts of sandarach, and 5 parts of elemi, in a sufficient quantity of alcohol, and apply without heat.

Violin Varnish. Coarsely powdered copal and glass, of each 4 oz., alcohol (64 O. P.) 1 pint, camphor $\frac{1}{2}$ oz.; heat the mixture with frequent stirring in a water-bath, so that the bubbles may be counted as they rise, until solution is complete, and when cold decant the clear portion.

Wainscot Varnish. Gum anime 8 lbs., clarified linseed oil 3 gallons, litharge $\frac{1}{2}$ lb., acetate of lead $\frac{1}{2}$ lb., sulphate of copper $\frac{1}{2}$ lb. These materials must be carefully but thoroughly boiled together until the mixture becomes quite stringy, and then $5\frac{1}{2}$ gallons of heated turpentine stirred in. If the colour is not sufficiently deep add a little gold size.

VASELIN. A proposed substitute for lard in the preparation of ointments, &c. See *Pocket Formulary*.

VEGETABLE OILS. Vegetable oils are divided into two classes; the drying and the non-drying. To the first class belong linseed, hempseed, and poppy oils. The second includes olive, palm, almond, and cocoanut oils. According to the mode of obtaining oils, they are distinguished as oils of the first and second pressure. Those of

the second pressure are mostly in demand by the soap manufacturers, for although less liquid and often mucilaginous, they contain more stearin, which gives rise to the production of a harder soap.

VEGETABLE PARCHMENT. *De la Rue's patent.* Strong unsized paper is immersed for a few seconds in oil of vitriol, diluted with half its volume of water. It is then washed in pure water. It strongly resembles animal parchment, and is used for the same purposes. The acid solution must be exactly of the strength indicated, and not warmer than the air around.

VINEGAR. Vinegar may be made from wine or ale, by keeping it for some weeks or months in a warm place, with access of air. In this country it is usually made from malt, or a mixture of malted and unmalted barley, which is mashed as for beer, and fermented with yeast. The fermented liquor is then placed in a warm room for many weeks in unclosed casks, and finished by transferring it into large vessels with false bottoms, on which are placed the refuse raisins, &c., from which wine has been prepared. A much quicker method of acetification is sometimes employed: the fermented liquor is made to pass in drops into tubs filled with beech chips, so as to expose an extended surface to the action of the air. In Germany it is also made by the direct acetification of spirit by means of platinum black. The method of preparing wood-vinegar has already been noticed. (See **PYROLIGNEOUS ACID.**) The following is one of the processes followed in making vinegar from sugar:—Boil 10 gallons of water for 10 minutes with a quart of bran; run it into a tub through flannel, and put into it 12 lbs. of coarse brown sugar, and when cooled to 70° F. add a quart of yeast at three different times. Let it work for four days, then take off the yeast, and run the liquor into a clean tub. Fill the tub nearly with the liquor, leaving room for 2 lbs. of bruised crab apples and 1 lb. of raisins. If it ferments, add a little reserved liquor, or water boiled with sugar, till the fermentation ceases. Then place the cask upon a plank fronting the sun in summer, and near the fire in winter. Put into it 1 oz. of isinglass well beaten up with a quart of old vinegar, cover the bung-hole with a piece of hop-bag (fastened to the edge of the hole by pitch), and lay a tile over it. Leave it in this state till it becomes fit for use.

On a small scale, Dr TURNER states that vinegar may be made from 120 parts of water, 12 of brandy, 3 of brown sugar, 1 of tartar, and $\frac{1}{2}$ of sour dough, left some weeks in a warm place. For Perfumed Vinegar, see PERFUMERY.

CARBOLIC VINEGAR. QUESNEVILLE. Crystallised carbolic acid 100 parts, acetic acid 900 parts, powdered camphor 5 parts. Sprinkled in infected cabins on board ship.

VULCANITE AND EBONITE. The only difference in these two articles is in the colouring materials employed. Vulcanite and ebonite are compounds of india rubber and sulphur, exactly the same as the well-known elastic bands, the only difference being in the time and heat required to vulcanize or harden the product.

To prepare vulcanite or ebonite combs, the india rubber is put into a masticator, along with a proper proportion of sulphur, and when thoroughly mixed, a sufficient quantity is put into a mould of the right shape made of plaster of Paris, or other material which will not combine with sulphur, and exposed in a steam boiler to a heat of 315° , and a pressure of about 12 lbs. to the inch for two hours. It is then removed from the mould and finished and polished in the same manner as ivory. The application of heat as above without a steam pressure is sufficient to vulcanize or harden the product, but the result is not always as satisfactory, as the material is liable to be porous if not compressed whilst hardening.

WAFERS, GELATIN. Dissolve fine glue or isinglass in such a quantity of water as that the solution, when cold, may have a firm consistence. Pour it hot on a plate of mirror glass (previously warmed with steam and slightly greased), which is fitted in a metallic frame, having edges just as high as the wafers should be thick. Lay on the surface a second glass plate, also hot and greased, so as to touch every point of the gelatin while resting on the edges of the frame. By its pressure the thin cake is rendered uniform. When the glass plates have cooled, the gelatin will be solid, and may be removed. It is cut into discs of different sizes by means of proper punches.

WASHES FOR VERMIN IN PLANTS. 1. *Tobacco water.* Infuse 1 lb. of tobacco in a gallon of boiling water, in a covered vessel, till cold.

2. *For Lice in Vines.* Boil $\frac{1}{2}$ lb. of tobacco in 2 quarts

of water; strain and add $\frac{1}{2}$ lb. of soft soap and $\frac{1}{4}$ lb. of sulphur. Mix.

3. *For Aphides.* Boil 2 oz. of lime and 1 oz. of sulphur in water, and strain.

4. *Poison for Plant Lice, and other insects.* Boil $3\frac{1}{2}$ oz. of quassia chips, and 5 drs. of powdered stavesacre seed, in 7 pints of water, to 5 pints. Strain when cold, and use with a watering pot or syringe.

5. *For Red Spiders.* A teaspoonful of salt in a gallon of water. In a few days wash the plant with pure water. See BLIGHTS, remedies for.

6. *For Mildew and Insects.* Sulphide of potassium $\frac{1}{4}$ oz., water 1 gallon.

WASHING POWDERS. These consist of soda-ash combined with gelatinous substances as a solution of glue, linseed jelly, &c., dried and powdered. Another form, by KILNER, is: carbonate of soda in powder 9 lbs., hyposulphite of soda 1 lb., borax $\frac{1}{2}$ lb. Add a teaspoonful to each gallon of water when the clothes are boiled.

WASHING LIQUIDS are chiefly solutions of caustic soda and silicate of soda.

WATCHMAKER'S OIL. 1. Put some lead shavings into neat-foot oil, and allow it to stand for some time, the longer the better. The lead neutralizes the acid, and the result is an oil that never corrodes or thickens.

2. Stir up for some time best olive oil with water kept at the boiling point; then after the two fluids have separated, decant the oil, and shake it up with a little freshly burnt lime. Let the mixture stand for some weeks in a bottle exposed to the sunlight and air, but protected from wet and dirt. When filtered the oil will be nearly colourless, perfectly limpid, and will never thicken or become rancid.

WATER FOR MARINE AQUARIUM. See AQUARIUM, MARINE, WATER FOR.

WATER, HARD (Dr CLARK'S *patent for softening*). This consists simply in adding milk of lime to the water in the reservoir. It combines with free carbonic acid, which it precipitates as carbonate of lime, and at the same time causes the deposition of the carbonate of lime previously held in solution by that gas.

WATERPROOFING COMPOUNDS. *For Boots, &c.* (ROOME'S patent.) Suet 8 oz., linseed oil 8 oz., yellow beeswax

6 oz., neatsfoot oil $1\frac{1}{2}$ oz., lamp-black 1 oz., litharge $\frac{1}{2}$ oz. Melt together, and stir till cold.

2. Linseed oil 8 oz., boiled ditto 10 oz., suet 8 oz., yellow wax 8 oz. Melt.

3. Dr HARVARD'S. Wax 8 oz., resin 4 oz., mutton suet 4 oz.; boil together, and apply warm to new boots.

4. Col. HAWKER'S. Drying oil 1 pint, wax 2 oz., Burgundy pitch 1 oz., oil of turpentine 2 oz. Melt over a slow fire, and add a few drops of oil of lavender or thyme. Brush the boots repeatedly with the composition before the fire, till they appear fully saturated.

5. *For Leather, &c.* Cut 3 drs. of india rubber into small pieces, soak them for 24 hours in a solution of common soda; dissolve this and 3 oz. of asphaltum in 12 oz. of camphine, then add $\frac{1}{2}$ oz. of boiled linseed oil.

For Cloth. It is alternately dipped in a solution of acetate of lead with a little gum, and solution of alum.

For Hats. Boil 8 lbs. of shell-lac, 3 lbs. frankincense, and 1 lb. borax in sufficient water.

For Canvas, &c.—Mr CASTLEY. Gutta percha 3 parts is dissolved in resin spirit 9 parts, at a heat of 120° to 140° Fahr., stirring occasionally.

WAX. Dr PEREIRA. Yellow beeswax is bleached by pouring the melted wax in a divided state on a revolving cylinder partly immersed in water, so as to form it into fine ribbons, which are exposed to air and moisture till bleached, and subsequently refined by melting with water containing sulphuric acid. M. INGENHOL. It has been proposed to bleach wax by adding to each pound of melted wax 2 oz. of powdered nitrate of soda, and afterwards stirring in, by little at a time, 1 oz. of sulphuric acid diluted with 10 parts of water, keeping the mixture warm, and constantly stirred with a glass rod in a capacious earthen vessel, till all the acid is added. It is then allowed to become somewhat cool, and the vessel filled with boiling water, well agitated, and set aside. The cake of wax is removed into boiling water, till this no longer produces a precipitate with chloride of barium. [We have not found this render wax perfectly white.]

WAX FOR MODELLING. Lead plaster 8 oz., beeswax 8 oz., Burgundy pitch 8 oz.; melt together, stir in sufficient chalk to form a paste, and form it into small sticks for use. **For SEALING WAX,** see further back.

WELDING COMPOSITION. Mix borax with 1-10th of sal ammoniac, fuse the mixture, and pour it on an iron plate. When cold, pulverize it, and mix it with an equal weight of quicklime, sprinkle it on iron heated to redness, and replace it in the fire. It may be welded below the usual heat.

WHEAT, STEEPS FOR. A pound of sulphate of copper in sufficient water, for each sack of seed. Arsenic is also used; sulphate of zinc has been recommended; so has quicklime, which is thus used:—Soak the seed in a warm mixture of 36 to 48 oz. of quicklime to 6 or 7 gallons of water. This is for $4\frac{1}{2}$ bushels of wheat; the solution should be sufficient to cover the seed 3 or 4 finger-breadths deep, and it should lie in it 24 hours. M. BOUSSINGAULT has proposed to sprinkle the grain, first with water, then with a mixture of 10 parts of lime with 1 of white arsenic. This was with the double object of preserving the grain, and destroying a plague of field mice. Sulphate of copper has seemed to give satisfactory results. It would be desirable, however, to find an innocuous substitute, as traces of copper have been found in wheat grown from the steeped seed. This desirable substitute appears to have been discovered in the use of a solution of sulphate of soda with lime, which has proved more successful in France than either arsenic or sulphate of copper. DOYÈRE recommends for grain infested with weevils, a small quantity of disulphide of carbon to be enclosed in an air-tight chamber with the grain; in a few hours both the larvæ and the eggs are killed, and the grain is not injured, as on exposure to air the disulphide quickly evaporates. See **BLIGHTS, REMEDIES FOR.**

WHITEWASH. Whiting is made into a milk with water, and a small quantity of melted size or dissolved glue added. It is applied to walls or ceilings with a broad, flat brush, worked in a uniform direction. Should the surface have been previously whitewashed, it is requisite first to remove the dirt by washing it with a brush and abundance of clean water.

“*Lime-washing* is, from the cleansing action of the quicklime, much the more effectual mode of purification, but is less frequently had recourse to, from the general ignorance respecting the proper mode of preparing the lime-wash. If glue be employed it is destroyed by the corrosive action

of the lime, and in consequence the latter easily rubs off the walls when dry. This is the case also if the lime be employed, as is often absurdly recommended, simply slaked in water, and used without any fixing material. Lime-wash is prepared by placing some freshly burned quicklime in a pail, and pouring on sufficient water to cover it. Boiled linseed oil should then be immediately added, in the proportion of a pint to a gallon of the wash. For coarser work any common refuse fat may be used instead of the boiled oil. The whole should then be thinned with water to the required consistency, and applied with a brush. Care should be taken not to leave the brush in the lime-wash for any length of time, as it destroys the bristles."—TEGETMEIER.

WOOD, TO BRONZE. The wood is first covered with a uniform coating of glue, or of drying oil, and when nearly dry, the bronze powder contained in a small bag, is dusted over it. The surface of the objects is afterwards rubbed with a piece of moist rag. Or the bronze powder may be previously mixed with the drying oil and applied with a brush.

WOOD, TO STAIN. 1. *Mahogany colour (dark).* Boil $\frac{1}{2}$ lb. of madder and 2 oz. of logwood in a gallon of water; and brush the wood well over with the hot liquid. When dry, go over the whole with a solution of 2 drs. of pearlsh in a quart of water.

2. (*Light.*) Brush over the surface with diluted nitrous acid, and when dry apply the following with a soft brush:—Dragon's blood 4 oz., common soda 1 oz., methylated spirit 3 pints; let them stand in a warm place, shaking frequently, then strain. Repeat the application until the proper colour is obtained.

3. (*To Stain Maple a Mahogany colour.*) Dragon's blood $\frac{1}{2}$ oz., alkanet $\frac{1}{2}$ oz., aloes 1 dr., spirit of wine 16 oz. Apply it with a sponge or brush.

4. *Rosewood.* Boil 8 oz. of logwood in 3 pints of water until reduced to half; apply it boiling hot two or three times, letting it dry between each. Afterwards put in the streaks with a camel-hair pencil dipped in a solution of copperas and verdigris in decoction of logwood.

5. *Ebony.* Wash the wood repeatedly with a solution of sulphate of iron; let it dry, then apply a hot decoction of logwood and nutgalls for two or three times. When

dry, wipe it with a wet sponge; and when again dry, polish with linseed oil.

6. *Oak.* Vandyke brown $2\frac{1}{2}$ oz., carbonate of ammonia 1 oz., bichromate of potash $\frac{1}{2}$ oz., common soda $\frac{1}{2}$ oz., water 1 quart. Boil together for 6 or 8 minutes.—*Pharmaceutical Journal.*

7. *To Stain Wood Red.* Use a strong decoction of Brazil-wood and alum.

Woods may be stained with the various dyes before described. See DYES.

A fine stain can be imparted to furniture made of beech or pine in a very simple manner. Dissolve 3 oz. of permanganate of potash and 3 oz. of sulphate of magnesia in 2 quarts of hot water. Apply this to the surface of the wood with a brush, and repeat if necessary. The manganese salt is decomposed in contact with the fibre of the wood, and a fine permanent stain is produced. If the objects are small a more dilute bath can be prepared, and the wood immersed in it for from one to five minutes, until it is thoroughly stained.—*The Laboratory.*

YEAST, ARTIFICIAL. 1. Honey 5 oz., cream of tartar 1 oz., malt 16 oz., water at 122° F. 3 pints; stir together, and when the temperature falls to 65° cover it up, and keep it at that temperature till yeast is formed.

2. Wheat flour is to be mixed with water into a thick paste, which is to be slightly covered in a moderately warm place; about the third day it begins to emit a little gas, and to exhale a disagreeable sour odour; about the sixth or seventh day the smell changes, much gas is evolved, accompanied by a distinct and agreeable vinous odour, and it is then in a state to excite either vinous or panary fermentation, and may be either at once employed for that purpose, or formed into small and very thin cakes, dried in the air, and preserved for future use.

Wort fermented with it in the ordinary way yields a large quantity of yeast, of excellent quality, which is found at the bottom of the vessel.

3. *With a ferment.* Take of bean flour $\frac{1}{4}$ lb., water 6 quarts, boil for half an hour, pour the decoction into any suitable vessel, add of wheat flour $3\frac{1}{2}$ lbs., stir the whole well together, and when the temperature reaches 55° F. add of beer yeast 2 quarts, mix well, and in 24 hours after the commencement of the fermentation

add of barley or bean flour 7 lbs.; make a uniform dough by thorough kneading, form it into small cakes as above, and then preserve these in a dry situation.

For use, one of these cakes is to be broken into pieces, laid in tepid water, and set in a warm place during twelve hours, when the soft mass will serve the purpose of beer yeast.

Patent Yeast. Take of hops 6 oz., water 3 gallons; simmer for three hours, strain, and in ten minutes stir in of ground malt $\frac{1}{2}$ peck. Next reboil the hops in water as before, and let the strained liquor run into the first mash, which must then be well stirred up, covered over, and left for four hours; after that time drain off the wort, and when the temperature has fallen to 90° F., set it to work with yeast (preferably patent) one pint; after standing for 20 to 24 hours in a warm place, take off the scum, strain it through a coarse hair sieve, and it will be fit for use. One pint is said to be enough for one bushel of bread.

YEAST. *Barm.* Yeast which consists almost entirely of minute vegetable cells, called *Torula cerivisiæ*, is either the froth or the deposit of fermenting worts, according to the character of the fermentation.

Ordinary beer yeast may be kept fresh and fit for use for several months by placing it in a close canvas bag, and gently and gradually squeezing out the moisture in a screw press, until the remaining matter acquires the consistence of clay or soft cheese, in which state it must be preserved in close vessels, or wrapped in waxed cloth. This is the method usually adopted for the best Flanders or German yeast.

YEAST, GERMAN. Ordinary beer yeast from which the moisture has been squeezed out by strong pressure. It is preserved in close vessels.

ZINC, AMMONIO-CHLORIDE OF. By dissolving equivalents of chloride of zinc and sal ammoniac, a crystallizable salt is formed, which dissolves oxides of copper and of iron, and is used in tinning or zincing those metals.

ZINC, AMALGAMATED (*for voltaic plates*). Put a little mercury on the zinc plate, and pour on it dilute sulphuric acid; then rub the mercury over the surface by means of a piece of linen. Another method, which is said to give a more permanent coating, is that of Mr WALENN. Having cleaned the plates by emery, and by immersion in diluted

sulphuric acid, and then in clean water, dip them into a mixture of equal parts of a saturated solution of corrosive sublimate and a similar solution of acetate of lead; then rub them with a cloth.

ZINC, PLATINIZED, for Dr Wright's Battery. Saturated solution of chloride of platinum $\frac{1}{2}$ dr., sulphuric acid $1\frac{1}{2}$ drs., water 2 drs. Mix; dip the zinc plates into the solution for a few seconds, and wash them quickly.

ZINC, OXIDE OF. M. J. LEFORT. It may be prepared from the purified sulphate, by precipitating it from a hot solution, by carbonated or bicarbonated alkalies and subsequent ignition. It cannot be obtained pure by caustic ammonia. Mr MIDGLEY prepares it on a large scale by the combustion of zinc in a muffle, heated by a furnace of peculiar construction; the zinc is introduced into the muffle from time to time, as the combustion proceeds; he is thus able to prepare one or two hundredweights at a time, by a continuous process.

ZINC, PURIFICATION OF. Granulate zinc by melting it, and pouring it while very hot into a deep vessel filled with water. Place the granulated vessel in a Hessian crucible, in alternate layers, with one fourth its weight of nitre, with an excess of nitre at the top. Cover the crucible, and secure the lid; then apply heat. When deflagration takes place, remove from the fire, separate the dross, and run the zinc into an ingot mould. It is quite free from arsenic.

APPENDIX.

WEIGHTS AND MEASURES

AVOIRDUPOIS WEIGHT. (B. P.)

lb. Pound.	oz. Ounces.	grs. Grains.	French Grammes.
1	= 16	= 7000	= 453·592
	1	= 437·50	= 28·349

Avoirdupois weights can be made available as substitutes for Troy weights by bearing in mind that 42·5 grains added to the avoirdupois ounce will make it equal to the Troy ounce, and that 1240 grains deducted from the avoirdupois pound will reduce it to the Troy pound.

MEASURES OF CAPACITY. (B. P.)

1 Minim	min.	
1 Fluid Drachm	fl. dr.	= 60 minims.
1 Fluid Ounce	fl. oz.	= 8 fluid drachms.
1 Pint	O.	= 20 fluid ounces.
1 Gallon	C.	= 8 pints.

The *old wine gallon* was adopted in the London Pharmacopœia before 1836, and the Dublin Pharmacopœia before 1850. Its use in this kingdom is no longer legal.

MEASURES OF LENGTH. (B. P.)

1 inch	in.	
12 inches	=	1 foot.
36 „	=	3 feet = 1 yard.

RELATION OF MEASURES TO WEIGHTS. (B. P.)

1 Minim	is the measure of	0.91 grains of water.
1 Fluid Drachm	„	54.68 „
1 Fluid Ounce	„ 1 ounce or	437.5 „
1 Pint	„ 1.25 pounds or	8750.0 „
1 Gallon	„ 10 pounds or	70,000.0 „

To find the weight of any given measure of a liquid, multiply the weight of the water it will contain by the specific gravity, water being 1.000. The weight of a gallon of any liquid, in avoird. lbs. and decimal parts, is at once seen from its density, merely removing the decimal point one place to the right. Thus, a gallon of ether at .750 weighs 7.50 (7½) lb. A gallon of nitric acid at 1.500 weighs 15 lbs.

APOTHECARIES' WEIGHT. (L. P.)

lb.	3	3	9	Gr.	Minims
Pound.	Ounces.	Drachms.	Scruples.	Grains.	of water.
1	= 12	= 96	= 288	= 5760	= 6319.54
	1	= 8	= 24	= 480	= 526.62
		1	= 3	= 60	= 65.82
			1	= 20	= 21.94
				1	= 1.09

The Troy pennyweight, 24 grains, is not used in compounding medicines.

APOTHECARIES' MEASURE.

C.	O.	f 3	f 3	℥	Grains of
Congius.	Octarii	Fluid	Fluid	Minims.	water.
Gallon.	Pints.	Ounces.	Drachms.		
1	= 8	= 160	= 1280	= 76,800	= 70,000
	1	= 20	= 160	= 9,600	= 8,750
		1	= 8	= 480	= 437.5
			1	= 60	= 54.7
				1	= 0.9

IMPERIAL MEASURE.—(Common Divisions.)

Quarter.	Bushels.	Pecks.	Gallons.	Quarts.	Pints.	Gills.
1	= 8	= 32	= 64	= 256	= 512	= 2048
	1	= 4	= 8	= 32	= 64	= 256
		1	= 2	= 8	= 16	= 64
			1	= 4	= 8	= 32
				1	= 2	= 8
					1	= 4

WEIGHTS AND MEASURES OF OTHER COUNTRIES.

The working formulæ of the United States Pharmacopœia are so constructed, that in their practical application any system of weights, or in certain cases measures, may be used, except in the preparation of fluid extracts, when metric weights and measures are recommended. The weights and measures referred to by physicians in prescribing, and used by pharmacists in dispensing, are either those of the apothecaries' or Troy system of weights, and the wine measure, or those of the metric system.

The Troy weights and the old wine measure are subjoined :

WEIGHTS.

The ounce,	} contains	{	eight drachms, ʒ.
The drachm,			three scruples, ʒ
The scruple,			twenty grains, gr.

MEASURES.

These are derived from the *wine gallon*, and are given in the following table with their signs annexed :

The fluid ounce,	} contains	{	eight fluid drachms, f ʒ.
The fluid drachm,			sixty minims, m.

The unit of the British India ponderary system is the *tola*, equal to 180 Troy grains. 32 *tolas* are equal to 1 lbj Troy. The *maund* is equal to 100 Troy ounces.

In France the metrical or decimal system is now the only legal one. The following table shows the correspondence of the French metrical weights with English grains.

	Troy grains.		Troy grains
Milligramme =	·0154	Decagramme =	154·32
Centigramme =	·1543	Hectogramme =	1543·20
Decigramme =	1·5432	Kilogramme =	15432·00
Gramme =	15·4320	Myriagramme =	154320·00

The measures of capacity in France are multiples and divisions of the *LITRE*, which is the measure occupied by a kilogramme (15432 Troy grains) of distilled water at its greatest density. It exceeds the old Paris *pinte* by $\frac{1}{4}$ th, and is equal to 35 fluid ounces and 103 minims, or 1·7608 imperial pints, or 61·028 English cubic inches. $4\frac{1}{2}$ litres make an imperial gallon, within about fʒxij.

The following table will show the relations between the litre and the imperial gallon of 277·2738 c. inches :

Litres.	Cubic Inches.	Gall.	Pts.	Fl. 3.	Fl. 3	Min.
$\frac{1}{1000}$ = Millilitre	·061028					16·9
$\frac{1}{100}$ = Centilitre	·61028				2	49
$\frac{1}{10}$ = Decilitre	6·1028			3	4	10·36
1 = Litre	61·028		1	15	1	43·69
10 = Decalitre	610·28	2	1	12	1	16·9
100 = Hectolitre	6102·8	22	0	1	4	49
1000 = Kilolitre	61028·	220	0	16	6	40
10000 = Myrialitre	610280·	2201	(or 175½ bushels).			

French Measures of Length.

The standard unit is the *mètre*, equal to 39·371 English inches, or 1 yard, 3 inches, and $\frac{37}{100}$ ths. The *kilomètre* (1000 mètres) is 4 furlongs, 213 yards, 1 foot, 11 inches.

The litre, with its divisions and multiples, is the measure now used. It contains 1000 grammes of water at the temperature of 4° C.; the number of grammes of other liquids corresponds with their specific gravity; water being 1000.

The metric system of weights and measures is also in general use in Germany, Austria, Italy, Spain, Norway, Sweden, Holland, Switzerland, Greece, and British India. It was legalized in Great Britain in 1864, and in the United States in 1866.

Comparison of Thermometric Scales.

To convert the degrees of centigrade into those of Fahrenheit, multiply by 9, divide by 5, and add 32.

To convert degrees of Centigrade into those of Réaumur, multiply by 4 and divide by 5.

To convert degrees of Fahrenheit into those of Centigrade, deduct 32, multiply by 5, and divide by 9.

To convert degrees of Fahrenheit into those of Réaumur, deduct 32, divide by 9, and multiply by 4.

To convert degrees of Réaumur into those of Centigrade, multiply by 5 and divide by 4.

To convert degrees of Réaumur into those of Fahrenheit multiply by 9, divide by 4, and add 32.

In DE LISLE'S thermometer, used in Russia, the graduation begins at boiling point, which is marked *Zero*, and the freezing point is 150.

EFFECTS OF TEMPERATURE

Degrees
of Fahr.

- 2786 Cast iron melts (Daniel).
 2016 Gold melts (Daniel).
 1996 Copper melts (Daniel).
 1873 Silver melts (Daniel).
 1750 Brass (containing 25% of zinc) melts (Daniel).
 1000 Iron bright cherry red (Poillet).
 980 Red heat, visible in daylight (Daniel).
 773 Zinc melts (Daniel).
 644 Mercury boils (Daniel), 662 (Graham).
 620 Sulphuric acid boils (Marignac), 620 (Graham).
 630 Whale oil boils (Graham).
 617 Pure lead melts (Rudberg).
 507 Bismuth melts (Gmelin).
 442 Tin melts (Crichton).
 424 (about) Arsenious acid volatilizes.
 312 to 324 Commercial oil of turpentine boils.
 256 Sat. sol. of acetate of soda boils.
 257 „ sal ammoniac boils (Taylor).
 239 Sulphur melts (Miller), 226 (Fownes).
 238 Sat. sol. of nitre boils.
 221 „ salt boils (Paris Codex).
 220 „ alum, carb. soda, and sulph. zinc boil.
 218 „ chlorate and prussiate of potash boil.
 216 „ sulph. of iron, sulph. of copper, nitrate of
 lead boil.
 214 „ acetate of lead, sulphate and bitartrate of
 potash boil.
 213 Water begins to boil in glass (or 213·5).
 212 Water boils in metal, barometer at 30°
 211 Alloy of 5 bismuth, 3 tin, 2 lead, melts.
 212 „ 8 bismuth, 5 lead, 3 tin, melts (below).
 207 Sodium melts (Regnault).
 185 Nitric acid 1·50 begins to boil.
 176 Rectified spirit boils.
 173 Alcohol (sp. gr. ·796 to ·800) boils.
 140 Beeswax melts, not under.
 150 Pyroxylic spirit boils (Scanlan), (156 Fownes).
 141·8 Chloroform boils.
 554 Crystallized glycerin boils (Mendlejeff).

Degrees
of Fahr.

- 144·5 Potassium melts (Bunsen).
 132 Acetone (pyroacetic spirit) boils (Kane).
 122 Mutton suet melts.
 116 Bisulphuret of carbon boils (Graham).
 100 Spermaceti melts, not under.
 111 Phosphorus melts (Miller).
 98 Temperature of the blood.
 95 Ether (·713) boils.
 95 Carbolic acid crystals become an oily liquid.
 64·4 Oil of anise liquefies.
 59 Gay Lussac's *Alcoomètre* graduated at.
 39 (about). Olive oil becomes partially solid.
 32 Water freezes.
 5 Cold produced by snow 2 parts and salt 1 part.
 —37·9 Mercury freezes.

SPECIFIC GRAVITIES

1. Solids. Water = 1·000.

Platinum, 21·53; Gold, 19·34; Thallium, 11·9; Lead, 11·350, Silver, 10·500; Bismuth, 9·822; Copper, 8·95; Cadmium, 8·604; Nickel, 8·82; Cobalt, 8·538; Iron, 7·844; Tin, 7·291; Zinc, 7·146; Antimony, 6·720; Aluminium, 2·56; Glass, 2·540 to 2·953; Sulphur (crystals of native), 2·07; Magnesium, 1·75; Calcium 1·58; Rubidium, 1·52; Gum arabic, 1·355; Scammony, 1·210; Amber, 1·078; Resin, 1·072; Camphor (laurel), ·996; Sodium, ·972; Beeswax, ·962; Spermaceti, ·943; Caoutchouc, ·933; Potassium, ·865.

2. Liquids. Water = 1·000.

Mercury, 13·590; Sulphuric acid 1·843; Nitric acid (monohydrated), 1·517; Nitric acid, P. B., 1·420; Commercial nitric acid, 1·380 to 1·390; Double aqua fortis, 1·360; Single aqua fortis, 1·220; Hydrochloric acid (strongest), 1·210; Hydrochloric acid, P. B., 1·160; Solution of caustic potash, P. B., 1·058; Solution of ammonia, P. B., ·959; Stronger solution of ammonia, P. B., ·891; Saturated solution of alum, 1·033; Saturated solution of common salt

1·200; Saturated solution of sulphate of copper, 1·150; saturated solution of sulphate of magnesia, 1·218; Sea-water, 1·027; Milk, from 1·018 to 1·045; Alcohol, P. B., ·795; Rectified spirit, P. B., ·838; Proof spirit, P. B., ·920; Glycerin, P. B., 1·25; Chloroform, P. B., 1·49; Bisulphide of carbon, 1·272; Syrup, P. B. 1·330; Spirit of nitrous ether, P. B., ·845; Ether (pure), ·713; Acetic ether, P. B., ·900; Caoutchoucine, ·680; Oil of turpentine, ·869 to ·876; Olive oil, ·9175; Castor oil ·950 to ·970; Spermaceti oil, ·875; Southern whale oil, ·920; Almond oil, ·917; Creasote, 1·071; Oil of wine, 1·05; Essential oil of anise, about ·976; of caraway, about ·920; of cinnamon, about 1·040; of cloves, about 1·055; of cajeput, about ·925; of lemon, distilled, about ·847; of rosemary, about ·897.

3. *Gases and Vapours.* Atmospheric Air = 1·000.

Hydrogen, ·0692; Nitrogen, ·972; Oxygen (Graham), 1·1056; Carbonic oxide, ·972; Carbonic acid (carbonic anhydride), 1·524; Light carburetted hydrogen, ·5595; Olefiant gas, ·981; Chlorine, 2·470 (2·421 Graham); Vapour of ether, 2·582; v. of water, ·622; v. of sulphur 2·218 (Graham); v. of phosphorus, 4·351; v. of iodine, 8·716.

TABLE OF CHEMICAL ELEMENTS MENTIONED IN THE BRITISH PHARMACOPŒIA (1885), WITH THEIR SYMBOLS AND EQUIVALENT NUMBERS.

ELEMENTARY BODIES.	SYMBOLS AND EQUIVALENTS.
Aluminium	Al = 27
Antimony (Stibium)	Sb = 120
Arsenium.....	As = 75
Barium.....	Ba = 137
Bismuth	Bi = 209
Boron	B = 11
Bromine	Br = 80
Cadmium.....	Cd = 112
Calcium	Ca = 40
Carbon.....	C = 12
Cerium.....	Ce = 141
Chlorine	Cl = 35.5
Chromium	Cr = 52.5
Copper (Cuprum)	Cu = 63.4
Gold (Aurum).....	Au = 196.5
Hydrogen	H = 1
Iodine	I = 127
Iron (Ferrum).....	Fe = 56
Lead (Plumbum).....	Pb = 207
Lithium	L = 7
Magnesium	Mg = 24
Manganese	Mn = 55
Mercury (Hydrargyrum) ...	Hg = 200
Nitrogen	N = 14
Oxygen.....	O = 16
Phosphorus	P = 31
Platinum	Pt = 195
Potassium (Kalium)	K = 39
Silver (Argentum)	Ag = 108
Sodium (Natrium)	Na = 23
Sulphur	S = 32
Tin (Stannum).....	Sn = 118
Zinc	Zn = 65

TABLE OF THE SOLUBILITY OF CHEMICALS IN WATER AND
IN ALCOHOL.

(From the United States Pharmacopœia.)

Abbreviations.—s. = soluble; ins. = insoluble; sp. = sparingly; v. = very; alm. = almost; dec. = decomposed.

Chemicals.	Water.		Alcohol, sp. gr. 0·820.	
	At 59° Far. = 15° Centg.	Boiling.	At 59° Far. = 15° Centg.	Boiling.
One Part is soluble in :	Parts.	Parts.	Parts.	Parts.
Acidum Arseniosum . . .	30—80	15	sp.	sp.
„ Benzoicum . . .	500	15	3	1
„ Boricum . . .	25	3	15	5
„ Carboicum . . .	20	—	v. s.	v. s.
„ Chromicum . . .	v. s.	v. s.	dec.	dec.
„ Citricum . . .	0·75	0·5	1	0·5
„ Gallicum . . .	100	3	4·5	1
„ Salicylicum . . .	450	14	2·5	v. s.
„ Tannicum . . .	6	v. s.	0·6	v. s.
„ Tartaricum . . .	0·7	0·5	2·5	0·2
Alumen	10·5	ins.	0·3	ins.
„ Exsiccatum . . .	20	ins.	0·7	ins.
Aluminii Hydras . . .	ins.	ins.	ins.	ins.
„ Sulphas	1·2	v. s.	alm. ins.	alm. ins.
Ammonii Benzoas . . .	5	1·2	28	7·6
„ Bromidum	1·5	0·7	150	15
„ Carbonas	4	dec.	dec.	dec.
„ Chloridum	3	alm. ins.	1·37	alm. ins.
„ Iodidum	1	0·5	9	3·7
„ Nitras	0·5	v. s.	20	3
„ Phosphas	4	ins.	0·5	ins.
„ Sulphas	1·3	1	sp.	sp.
„ Valerianas	v. s.	v. s.	v. s.	v. s.
Antimonii et Potassii Tartaras	17	3	ins.	ins.

TABLE OF THE SOLUBILITY OF CHEMICALS, ETC.—*continued.*

Chemicals.	Water.		Alcohol, sp. gr. 0·820.	
	At 59°Far. = 15° Centg.	Boiling.	At 59°Far. = 15° Centg.	Boiling.
One Part is soluble in :	Parts.	Parts.	Parts.	Parts.
Antimonii Oxidum . . .	alm. ins.	alm. ins.	ins.	ins.
„ Sulphidum . . .	ins.	ins.	ins.	ins.
„ Sulphidum Purificatum . . .	ins.	ins.	ins.	ins.
Antimonium Sulphura- tum	ins.	ins.	ins.	ins.
Apomorphinæ Hydro- chloras	6·8	dec.	50	dec.
Argentii Cyanidum . . .	ins.	ins.	ins.	ins.
„ Iodidum	ins.	ins.	ins.	ins.
„ Nitras	0·8	0·1	26	5
„ Fusus	0·6	0·5	25	5
„ Oxidum	v. sp.	v. sp.	ins.	ins.
Arsenii Iodidum	3·5	dec.	10	dec.
Atropina	600	35	v. s.	v. s.
Atropinæ Sulphas . . .	0·4	v. s.	6·5	v. s.
Bismuthi Citras	ins.	ins.	ins.	ins.
„ et Ammonii Citras	v. s.	v. s.	sp.	sp.
„ Subcarbonatis . . .	ins.	ins.	ins.	ins.
„ Subnitras	ins.	ins.	ins.	ins.
Bromum	33	—	dec.	dec.
Caffeina	75	9·5	35	v. s.
Calcii Bromidum	0·7	v. s.	1	v. s.
„ Carbonas Precipi- tatus	ins.	ins.	ins.	ins.
„ Chloridum	1·5	v. s.	8	1·5
„ Hypophosphis	6·8	6	ins.	ins.
„ Phosphas Precipi- tatus	ins.	ins.	ins.	ins.
Calx	750	1300	ins.	ins.
Camphora Monobromata	alm. ins.	alm. ins.	v. s.	v. s.

TABLE OF THE SOLUBILITY OF CHEMICALS, ETC.—*continued.*

Chemicals.	Water.		Alcohol, sp. gr. 0·820.	
	At 59°Far. = 15° Centg.	Boiling.	At 59°Far. = 15° Centg.	Boiling.
One Part is soluble in :	Parts.	Parts.	Parts.	Parts.
Cerii Oxalas	ins.	ins.	ins.	ins.
Chloral	v. s.	v. s.	v. s.	v. s.
Chrysarobinum	alm. ins.	alm. ins.	sp.	sp.
Cinchonidinæ Sulphas	100	4	71	12
Cinchonina	alm. ins.	alm. ins.	110	28
Cinchoninæ Sulphas	70	14	6	1·5
Codeina	80	17	v. s.	v. s.
Creta Preparata	ins.	ins.	ins.	ins.
Cupri Acetas	15	5	135	14
„ Sulphas	2·6	0·5	ins.	ins.
Elaterinum	ins.	ins.	125	2
Ferri Chloridum	v. s.	v. s.	v. s.	v. s.
„ Citras	s.	v. s.	ins.	ins.
„ et Ammonii Citras	v. s.	v. s.	ins.	ins.
„ „ Sulphas	3	0·8	ins.	ins.
„ „ Tartras	v. s.	v. s.	ins.	ins.
„ Potassii Tartras	v. s.	v. s.	ins.	ins.
„ Quininæ Citras	s.	v. s.	ins.	ins.
„ Strychninæ Ci- tras	v. s.	v. s.	ins.	ins.
„ Hypophosphis	sp.	sp.	ins.	ins.
„ Lactas	40	12	alm. ins.	alm. ins.
„ Oxalas	sp.	sp.	ins.	ins.
„ Oxidum Hydra- tum	ins.	ins.	ins.	ins.
„ Phosphas	v. s.	v. s.	ins.	ins.
„ Pyro Phosphas	v. s.	v. s.	ins.	ins.
„ Sulphas	1·8	0·3	ins.	ins.
„ „ Precipitatus	1·8	0·3	ins.	ins.
„ Valerianas	ins.	dec.	v. s.	v. s.
Hydrargyri Chloridum Corrosivum	16	2	3	1·2

TABLE OF THE SOLUBILITY OF CHEMICALS, ETC.—*continued.*

Chemicals.	Water.		Alcohol, sp. gr. 0·820.	
	At 59°Far. = 15° Centg.	Boiling.	At 59°Far. = 15° Centg.	Boiling.
One Part is soluble in :	Parts.	Parts.	Parts.	Parts.
Hydrargyri Chloridum	ins.	ins.	ins.	ins.
Mite .	12·8	3	15	6
Cyanidum .	alm. ins.	alm. ins.	130	15
Iodidum Ru- brum	alm. ins.	alm. ins.	ins.	ins.
" Viride	ins.	ins.	ins.	ins.
Oxidum Fla- vum	ins.	ins.	ins.	ins.
" Rubrum	ins.	ins.	ins.	ins.
Subsulphas Flavus .	ins.	ins.	ins.	ins.
Sulphidum Rubrum .	ins.	ins.	ins.	ins.
Hydrargyrum Ammonia- tum	ins.	ins.	ins.	ins.
Hyoscyaminæ Sulphas .	v. s.	v. s.	v. s.	v. s.
Iodoformum	ins.	ins.	80	15
Iodum	sp.	—	11	—
Lithii Benzoas	4	2·5	12	10
" Bromidum	v. s.	v. s.	v. s.	v. s.
" Carbonas	130	130	ins.	ins.
" Citras	5·5	2·5	sp.	sp.
" Salicylas	v. s.	v. s.	v. s.	v. s.
Magnesia	alm. ins.	alm. ins.	ins.	ins.
" Ponderosa	alm. ins.	alm. ins.	ins.	ins.
Magnesii Carbonas	alm. ins.	alm. ins.	ins.	ins.
" Sulphas	0·8	0·15	ins.	ins.
" Sulphis	20	19	ins.	ins.
Manganesii Oxidum Ni- gram	ins.	ins.	ins.	ins.
" Sulphas	0·7	0·8	ins.	ins.
Morphina	v. sp.	500	100	36

TABLE OF THE SOLUBILITY OF CHEMICALS, ETC.—*continued*

Chemicals.	Water.		Alcohol, sp. gr. 0·820.	
	At 59°Far. = 15° Centg.	Boiling.	At 59°Far. = 15° Centg.	Boiling.
One Part is soluble in :	Parts.	Parts.	Parts.	Parts.
Morphinæ Acetas . . .	12	1·5	68	14
„ Hydrochloras . . .	24	0·5	63	31
„ Sulphas . . .	24	0·75	702	144
Phosphorus	ins.	ins.	v. sp.	v. sp.
Physostigminæ Salicylas	130	30	12	v. s.
Picrotoxinum	150	25	10	3
Pilocarpinæ Hydrochloras	v. s.	v. s.	v. s.	v. s.
Piperina	alm. ins.	alm. ins.	30	1
Plumbi Acetas	1·8	0·5	8	1
„ Carbonas	ins.	ins.	ins.	ins.
„ Iodidum	2000	200	v. sp.	v. sp.
„ Nitras	2	0·8	alm. ins.	alm. ins.
„ Oxidum	ins.	ins.	ins.	ins.
Potassa	0·5	v. s.	2	v. s.
Potassii Acetas	0·4	v. s.	2·5	v. s.
„ Bicarbonas	3·2	dec.	alm. ins.	alm. ins.
„ Bichromas	10	1·5	ins.	ins.
„ Bitartras	210	15	v. sp.	v. sp.
„ Bromidum	1·6	1	200	16
„ Carbonas	1	0·7	ins.	ins.
„ Chloras	16·5	2	v. sp.	v. sp.
„ Citras	0·6	v. s.	v. sp.	v. sp.
„ Cyanidum	2	1	sp.	sp.
„ et Sodii Tartras	2·5	v. s.	alm. ins.	alm. ins.
„ Ferrocyanidum	4	2	ins.	ins.
„ Hypophosphis	0·6	0·3	7·3	3·6
„ Iodidum	0·8	0·5	18	6
„ Nitras	4	0·4	alm. ins.	alm. ins.
„ Permanganas	20	3	dec.	dec.
„ Sulphas	9	4	ins.	ins.
„ Sulphis	4	5	sp.	sp.

TABLE OF THE SOLUBILITY OF CHEMICALS, ETC.—*continued.*

Chemicals.	Water.		Alcohol, sp. gr. 0·820.	
	At 59°Far. = 15° Centg.	Boiling.	At 59°Far. = 15° Centg.	Boiling.
One Part is soluble in :	Parts.	Parts.	Parts.	Parts.
Potassii Tartras	0·7	0·5	alm. ins.	alm. ins.
Quinidinæ Sulphas	100	7	8	v. s.
Quinina	1600	700	6	2
Quininae Bisulphas	10	v. s.	32	v. s.
„ Hydrobromas	16	1	3	1 or less
„ Hydrochloras	34	1	3	v. s.
„ Sulphas	740	30	65	3
„ Valerianas	100	40	5	1
Saccharum	0·5	0·2	175	28
„ Lactis	7	1	ins.	ins.
Salicinum	28	0·7	30	2
Santoninum	alm. ins.	250	40	3
Soda	1·7	0·8	v. s.	v. s.
Sodii Acetas	3	1	30	2
„ Arsenias	4	v. s.	v. sp.	60
„ Benzoas	1·8	1·3	45	20
„ Bicarbonas	12	dec.	ins.	ins.
„ „ Venalis	12	dec.	ins.	ins.
„ Bisulphis	4	2	72	49
„ Boras	16	0·5	ins.	ins.
„ Bromidum	1·2	0·5	13	11
„ Carbonas	1·6	0·25	ins.	ins.
„ Chloras	1·1	0·5	40	43
„ Chloridum	2·8	2·5	alm. ins.	alm. ins.
„ Hypophosphis	1	0·12	30	1
„ Hyposulphis	1·5	0·5	ins.	ins.
„ Iodidum	0·6	0·3	1·8	1·4
„ Nitras	1·3	0·6	sp.	40
„ Phosphas	6	2	ins.	ins.
„ Pyrophosphas	12	1·1	ins.	ins.
„ Salicylas	1·5	v. s.	6	v. s.
„ Santoninas	3	0·5	12	3·4

TABLE OF THE SOLUBILITY OF CHEMICALS, ETC.—*continued.*

Chemicals.	Water.		Alcohol, sp. gr. 0·820.	
	At 59°Far. = 15° Centg.	Boiling.	At 59°Far. = 15° Centg.	Boiling.
One Part is soluble in:	Parts.	Parts.	Parts.	Parts.
Sodii Sulphas	2·8	0·4	ins.	ins.
„ Sulphis	4	0·9	sp.	sp.
„ Sulphocarbolas	5	0·7	132	10
Strychnina	6700	2500	110	12
Strychninæ Sulphas	10	2	60	2
Sulphur Lotum	ins.	ins.	ins.	ins.
„ Precipitatum	ins.	ins.	ins.	ins.
„ Sublimatum	ins.	ins.	ins.	ins.
Thymol	1200	900	1	v. s.
Veratrina	v. sp.	v. sp.	3	v. s.
Zinci Acetas	3	1·5	30	3
„ Bromidum	v. s.	v. s.	v. s.	v. s.
„ Carbonas Precipitatus	ins.	ins.	ins.	ins.
„ Chloridum	v. s.	v. s.	v. s.	v. s.
„ Iodidum	v. s.	v. s.	v. s.	v. s.
„ Oxidum	ins.	ins.	ins.	ins.
„ Phosphidum	ins.	ins.	ins.	ins.
„ Sulphas	0·6	0·3	ins.	ins.
„ Valerianas	100	—	40	—

Composition, Equivalent Numbers, and Formulae of some of the more important Compounds employed in Pharmacy and the Arts.

[Fractions are omitted.]

	OLD SYSTEM.		NEW SYSTEM.	
	Formulae.	Equiv.	Formulae.	Equiv.
Acetone	$C_6 H_6 O_2$	58	$C_3 H_6 O$	58
Acid, Acetic (anhydrous)	$C_4 H_4 O_4$ (or \bar{A})	60	$C_2 H_4 O_2$	60
— Arsenious	$As_2 O_3$	99	$As_2 O_3$	198
— Arsenic	$As_2 O_5$	115	$As_2 O_5$	230
— Benzoic	$C_{14} H_5 O_3 HO$	122	$H C_7 H_5 O_2$	122
— Boracic (anhydrous)	$B_2 O_3$	35	$B_2 O_3$	70
— — Crys.	$B_2 O_3, 3HO$	62	$H B O_2, H_2 O$	62
— Carbolic	$HO, C_{12} H_5 O$	94	$H C_6 H_5 O$	94
— Carbonic	$C O_2$	22	$C O_2$	44
— Chromic	$Cr O_3$	51	$Cr O_3$	101
— Citric (anhydrous)	$C_{12} H_5 O_{11}$	165	$C_6 H_5 O_7$	192
— — (crystals)	$\bar{C} 2HO$	183	$H_3 C_6 H_5 O_7, H_2 O$	210
— Gallic	$3 HO, C_{14} H_3 O_7 + 2 HO$	160	$H_3 C_7 H_3 O_5, H_2 O$	188
— Hydrochloric	$H Cl$	37	$H Cl$	37
— Hydrocyanic	$C_2 N H$ (or $H Cy$)	27	$H Cy$	27
— Hydrosulphuric	$H S$	17	$H_2 S$	34
— Iodic (anhydride)	$I O_5$	167	$I_2 O_5$	334
— Lactic	$C_6 H_6 O_6$	90	$\bar{H} C_3 H_5 O_3$	90
— Nitric	$N O_5$	54	$N_2 O_5$	108
— — sp. gr. 1.5	$N O_5, HO$	63	$H N O_3$	63
— Oxalic (anhydrous)	$C_4 O_6$ (or \bar{O})	72	$H_2 C_2 O_4$	92
— — Crys.	$C_4 O_6, 4HO$	108	$H_2 C_2 O_4, 2 H_2 O$	126
— Phosphoric (anhydride)	$P O_5$	71	$P_2 O_5$	142
— Salicylic	$C_{14} H_6 O_6$	138	$H C_7 H_5 O_3$	138
— Sulphuric (anhydride)	SO_3	40	SO_3	80
— — hydrated	SO_3, HO	49	$H_2 SO_4$	98
— Tartaric, Crys.	$\bar{C}_8 H_4 O_{10} 2HO$	150	$H_2 C_4 H_4 O_6$	150
Alcohol	$C_4 H_6 O_2$	46	$C_2 H_6 O$	46
— Amylic	$C_{10} H_{12} O_2$	88	$C_5 H_{12} O$	88
Alumina	$Al_2 O_3$	52	$Al_2 O_3$	103
Alum	$Al_2 O_3, 3 (SO_3); KO, SO_3 + 24 HO$	474	$K, Al (SO_4)_2 12 HO_2$	474
Aloin	$C_{22} H_{18} O_{14}$	322	$C_{11} H_{18} O_7$	322
Ammonium	NH_4	18	NH_4	18
Amidogen	NH_2	16	NH_2	16
Ammonia	NH_3	17	NH_3	17
— Hydrochlorate*	$NH_3, H Cl$	54	$NH_4 Cl$	54

* Ammonium Chloride.

	OLD SYSTEM.		NEW SYSTEM.	
	Formule.	Equiv.	Formule.	Equiv.
Ammonia Sulphate, crys.	NH ₃ SO ₃ HO.....	66	(NH ₃) ₂ SO ₄	132
— Carbonate	2NH ₃ O, 3CO ₂	118	N ₂ H ₁₂ C ₃ O ₃	236
Amyl, Nitrite	C ₁₀ H ₁₁ O, NO ₃	117	C ₅ H ₁₁ NO ₂	117
Antimony, Teroxide ..	Sb O ₃	146	Sb ₂ O ₃	292
— Tersulph.....	Sb S ₃	170	Sb ₂ S ₃	340
— Potassio-tartrate ..	KO, Sb O ₃ , T, 2 HO	343	K, Sb, C ₄ H ₄ O ₇ , H ₂ O...	343
Apomorphine Hydro- chlorate	C ₃₄ H ₁₇ NO ₄ , HCl ...	303	C ₁₇ H ₁₇ NO ₂ , HCl ...	303
Atropia	C ₃₄ H ₂₃ NO ₆	289	C ₁₇ H ₂₃ NO ₃	289
Baryta	Ba O.....	77	Ba O.....	153
— Carbonate	Ba O, CO ₂	99	Ba CO ₃	197
— Sulphate	Ba O, SO ₃	117	Ba SO ₄	233
Barium, Chloride ...	Ba Cl.....	104	Ba Cl ₂	208
Bismuth, Oxide	Bi O ₃	234	Bi ₂ O ₃	468
— Subnitrate	Bi O ₃ , NO ₂ , 2 HO.....	306	Bi ₂ NO ₃ , H ₂ O.....	306
Borax	Na O, 2 BO ₃ , 10 HO	191	Na ₂ O, 2 B ₂ O ₃ , 10 H ₂ O	382
Caffeine	C ₁₆ H ₁₀ N ₄ O ₄ H ₂ O ₂ ...	212	C ₈ H ₁₀ N ₄ O ₂ H ₂ O...	212
Calcium, Chloride ...	Ca Cl.....	55	Ca Cl ₂	111
— — crys.	Ca Cl, 6 HO.....	110	Ca Cl ₂ , 6 H ₂ O.....	219
— Oxide (Lime).....	Ca O.....	28	Ca O.....	56
Cerium, Oxalate	2 Ce O, C ₄ O ₆ + 6 HO	234	Ce C ₂ O ₄ , 3 H ₂ O.....	234
Chloral, Hydrate.....	C ₂ H Cl ₃ O ₂ , 2 HO...	166	C ₂ H Cl ₃ O, H ₂ O.....	166
Chloroform	C ₂ H Cl ₃	120	C H Cl ₃	120
Cinchonina	C ₂₀ H ₁₂ NO.....	154	C ₂₀ H ₂₄ N ₂ O.....	308
Cocaine Hydrochlor- ate	C ₃₄ H ₂₁ NO, HCl.....	339	C ₁₇ H ₂₁ NO ₄ , HCl ...	339
Copper, Oxide.....	Cu O.....	41	Cu O.....	82
— Dioxide	Cu ₂ O.....	72	Cu ₂ O.....	144
— Sulphate (anhyd.)	Cu O, SO ₃	80	Cu SO ₄	160
— — crys.	Cu O, SO ₃ , 5 HO.....	125	Cu SO ₄ , 5 H ₂ O.....	249
— Nitrate	Cu O, NO ₃	94	Cu (NO ₃) ₂	188
— Acetate, crys.....	Cu O, A, HO.....	100	Cu, 2 C ₂ H ₃ O ₂ , H ₂ O...	200
Cyanogen	C ₂ N (or Cy).....	26	CN (or Cy).....	26
Ether	C ₄ H ₈ O.....	37	C ₂ H ₅ O.....	74
— Acetic	C ₄ H ₈ O, C ₄ H ₃ O ₃ ...	88	C ₂ H ₅ C ₂ H ₃ O ₂	88
Ethyl	C ₄ H ₈	29	C ₂ H ₅	29
Glycerin	C ₆ H ₈ O ₆	92	C ₃ H ₅ O ₃	92
Iron, Protoxide	Fe O.....	36	Fe O.....	72
— Black Oxide	Fe ₃ O ₄	116	Fe ₃ O ₄	232
— Peroxide	Fe ₂ O ₅	80	Fe ₂ O ₅	160
— Chloride	Fe Cl.....	64	Fe Cl ₂	128
— Perchloride.....	Fe ₂ Cl ₃	163	Fe ₂ Cl ₆	326
— Iodide	Fe I.....	155	Fe I ₂	310
— Sulphate (anhyd.)	Fe O, SO ₃	76	Fe SO ₄	152
— — crys.	Fe O, SO ₃ , 7 HO.....	139	Fe SO ₄ , 7 H ₂ O.....	278
Lead, Acetate, crys.	Pb O, A, 3 HO.....	190	Pb, (C ₂ H ₃ O ₂) ₂ , 3 H ₂ O	379
— Protoxide	Pb O.....	112	Pb O.....	223
— Iodide	Pb I.....	231	Pb I ₂	461
— Chloride	Pb Cl.....	139	Pb Cl ₂	278
— Sulphate	Pb O, SO ₃	152	Pb SO ₄	303
— Nitrate, crys.....	Pb O, NO ₃	166	Pb (NO ₃) ₂	331
Lime, Carbonate.....	Ca O, CO ₂	50	Ca CO ₃	100

	OLD SYSTEM.		NEW SYSTEM.	
	Formule.	Equiv.	Formule.	Equiv.
Lime, Hydrate.....	Ca O, HO.....	37	Ca H ₂ O ₂	74
— Hypophosphite ...	Ca O, PO, 2 HO	85	Ca 2 PH ₂ O ₂	170
— Sulphate.....	Ca O, SO ₃	68	Ca SO ₄	136
— Phosphate (bone-earth)	3 Ca O, PO ₅	155	Ca ₃ (PO ₄) ₂	310
Lithia, Carbonate ...	LO, CO ₂	37	L ₂ CO ₃	74
Magnesia.....	Mg O.....	20	Mg O.....	40
— Carbonate	(Mg O, CO ₂) ₂ ; MgO; 5 HO.....	191	(Mg CO ₃) ₂ , MgO, 5 H ₂ O	382
— Sulphate (anhyd.)	Mg O, SO ₃	60	Mg SO ₄	120
— — crys.	Mg O, SO ₃ , 7 HO	123	Mg SO ₄ 7 H ₂ O	246
Manganese, Binoxide	Mn O ₂	44	Mn O ₂	87
— Chloride.....	Mn Cl	63	Mn Cl ₂	126
— Sulphate.....	Mn O, SO ₃	76	Mn SO ₄	151
Mannite	C ₆ H ₇ O ₆	91	C ₆ H ₁₄ O ₆	182
Mercury, Ammoniated	NH ₂ Hg ₂ Cl.....	251	NH ₂ Hg Cl	251
— Chloride (Subchloride, B.)	Hg ₂ Cl	236	Hg Cl	236
— Bichloride (Perchloride, B.)	Hg Cl ₂	171	Hg Cl ₂	271
— Protoxide (suboxide)	Hg O.....	108	Hg ₂ O	416
— Binoxide (red oxide)	Hg O.....	216	Hg O	216
— Iodide	Hg ₂ I.....	327	Hg I	327
— Biniodide	Hg I	227	Hg I ₂	454
— Sulphate	Hg O ₂ , 2 SO ₃	196	Hg SO ₄	296
— Bisulphuret*	Hg S ₂	132	Hg S	262
Morphia	C ₃₄ H ₁₉ NO ₆	285	C ₁₇ H ₁₉ NO ₃	285
— Acetate, crys.....	C ₃₄ H ₂₉ NO ₆ , A, HO	345	C ₁₇ H ₁₉ NO ₃ , C ₂ H ₄ O ₂	345
— Hydrochlorate, { crys.	C ₃₄ H ₁₉ NO ₆ , H Cl, 6 HO.....	375	C ₁₇ H ₁₉ NO ₃ , H Cl, 3 H ₂ O	375
Physostigmine.....	C ₂₀ H ₂₁ N ₃ O ₄	275	C ₁₅ H ₂₁ N ₃ O ₂	275
Platinum, chloride ...	Pt Cl ₂	170	Pt Cl ₂	339
— Ammonio-chloride	Pt Cl ₂ , NH ₃ , H Cl	223	(H ₄ N Cl) ₂ , Pt Cl ₄	446
Potash (anhyd.)	KO.....	47	K ₂ O.....	94
— Hydrate	KO, HO	56	K H O	56
— Acetate	KO, A	98	KC ₂ H ₃ O ₂	98
— Carbonate	KO, CO ₂	69	K ₂ CO ₃	138
— Bicarbonate, crys.	KO, 2 CO ₂ , HO	100	K H CO ₃	100
— Chlorate	KO, Cl O ₅	123	K Cl O ₃	123
— Chromate	KO, Cr O ₃	98	K ₂ Cr O ₄	195
— Bichromate	KO, 2 Cr O ₃	148	K ₂ O (Cr O ₃) ₂	295
— Nitrate	KO, NO ₃	101	K NO ₃	101
— Sulphate.....	KO, SO ₃	87	K ₂ SO ₄	174
Potash, Bisulph., crys.	KO, HO, 2 SO ₃	136	K H SO ₄	136
Potash, Tartrate.....	2 KO, T.....	226	K ₂ C ₄ H ₄ O ₆	226
— Bitartrate	KO, T, HO	188	K H C ₄ H ₄ O ₆	188
— Permanganate ...	KO, Mn ₂ O ₇	158	K Mn O ₄	158
Potassium, bromide.	K Br.....	119	K Br	119
— Chloride	K Cl	75	K Cl	75

* Mercuric Sulphide.

	OLD SYSTEM.		NEW SYSTEM.	
	Formulæ.	Equiv.	Formulæ.	Equiv.
Potassium Iodide ...	K I	166	K I	166
Quinia	$C_{20} H_{12} NO_2$	162	$C_{20} H_{24} N_2 O_3$	324
— Sulphate, crys....	$2 (C_{20} H_{12} NO_2)_2 SO_3$	436	$(C_{20} H_{24} N_2 O_3)_2 H_2$	872
— Neutral sulphate.	$C_{20} H_{12} NO_2 SO_3$	274	$C_{20} H_{24} N_2 O_3 H_2$	548
Santonin	$C_{30} H_{18} O_6$	246	$C_{15} H_{18} O_3$	246
Silver, Oxide	Ag O	116	$Ag_2 O$	232
— Chloride	Ag Cl	144	$Ag Cl$	144
— Iodide	Ag I	235	$Ag I$	235
— Cyanide	Ag Cy	134	$Ag Cy$	134
— Nitrate	Ag O, NO_2	170	$Ag NO_3$	170
Soda	Na O	31	$Na_2 O$	62
— Carbonate (exsic- cat.).....	Na O, CO_2	53	$Na_2 CO_3$	106
— Carbonate, crys...	Na O, CO_2 , 10 HO	143	$Na_2 CO_3$, 10 $H_2 O$	286
— Bicarbonate	Na O, 2 CO_2 , HO	84	$Na HCO_3$	84
— Sulphate (anhyd.)	Na O, SO_3	71	$Na_2 SO_4$	142
— — crys.	Na O, SO_3 , 10 HO	161	$Na_2 SO_4$, 10 $H_2 O$	322
— Phosphate	2 Na O, PO_5 , 25 HO	358	$Na_2 HPO_4$, 12 $H_2 O$	358
— tartarated crys. ...	Na O, KO, T, 8HO	282	$Na K C_4 H_4 O_6$, 4 $H_2 O$	282
— Salicylate	$(Na C_{14} H_5 O_3)_2 (HO)_2$	338	$(Na C_7 H_5 O_3)_2 H_2 O$	338
Sodium, chloride ...	Na Cl	59	Na Cl	59
Starch	$C_{12} H_{20} O_{10}$	162	$C_6 H_{10} O_5$	162
Strychnia	$C_{42} H_{22} O_4 N_2$	334	$C_{21} H_{22} N_2 O_2$	334
Sugar, Cane.....	$C_{12} H_{11} O_{11}$	171	$C_{12} H_{22} O_{11}$	342
Sugar of Milk	$C_{12} H_{12} O_{11} HO$	180	$C_{12} H_{22} O_{11} H_2 O$	360
Thymol	$C_{20} H_{14} O_2$	150	$C_{10} H_{13} HO$	150
Water	HO	9	$H_2 O$	18
Zinc, oxide	Zn O	41	$Zn O$	81
— Carbonate	Zn O, CO_2	63	$Zn CO_3$	125
— Sulphate	Zn O, SO_3 , 7 HO	144	$Zn SO_4$, 7 $H_2 O$	287
— Chloride	Zn Cl	68	$Zn Cl_2$	136

Table of the Neutralizing Proportions of some of the Acids and Alkaline Carbonates. The best commercial preparations are intended.

100 parts of.	Anhydrous Carb. of Potash.	Bicarbonate of Potash.	Carbonate of Soda Exsiccated.	Carbonate of Soda Xialized.	Bicarbonate of Soda.	Carbonate of Ammonia.	Carbonate of Magnesia.
Citric Acid require .	98	145	75	200	120	84	6·5
Lemon Juice „	7	10	5	14	8·5	6	4·6
Tartaric Acid „	9·2	13·3	70	190	112	78	60
Vinegar (containing 4·5 parts of abso- lute acid) require .	5	7·5	3·9	10·5	6·3	4·3	3·4

It must be borne in mind that lemon juice varies in strength. The above quantities are sufficient when it contains from 7 to 7·5 per cent. of citric acid.

Specific Gravities corresponding to Degrees of BAUMÉ'S Hydrometer for Liquids heavier than Water (POGGIALE).

Degrees.	Specific Gravity.						
0	1.000	19	1.152	38	1.357	57	1.652
1	1.007	20	1.161	39	1.370	58	1.671
2	1.014	21	1.171	40	1.383	59	1.691
3	1.022	22	1.180	41	1.397	60	1.711
4	1.029	23	1.190	42	1.410	61	1.732
5	1.036	24	1.199	43	1.424	62	1.753
6	1.044	25	1.210	44	1.438	63	1.774
7	1.052	26	1.221	45	1.453	64	1.796
8	1.060	27	1.231	46	1.468	65	1.819
9	1.067	28	1.242	47	1.483	66	1.846
10	1.075	29	1.253	48	1.498	67	1.872
11	1.083	30	1.264	49	1.514	68	1.897
12	1.091	31	1.275	50	1.530	69	1.921
13	1.100	32	1.286	51	1.546	70	1.946
14	1.108	33	1.297	52	1.563	71	1.974
15	1.116	34	1.309	53	1.580	72	2.000
16	1.125	35	1.320	54	1.597	73	2.031
17	1.134	36	1.332	55	1.615	74	2.059
18	1.143	37	1.345	56	1.634		

Specific Gravities corresponding to Degrees of BAUMÉ'S Hydrometer for Liquids lighter than Water (FRANCEUR).

Degrees.	Specific Gravity.						
10	1·000	23	0·918	36	0·849	49	0·789
11	0·993	24	0·913	37	0·844	50	0·785
12	0·986	25	0·907	38	0·839	51	0·781
13	0·980	26	0·901	39	0·834	52	0·777
14	0·973	27	0·896	40	0·830	53	0·773
15	0·967	28	0·890	41	0·825	54	0·768
16	0·960	29	0·885	42	0·820	55	0·764
17	0·954	30	0·880	43	0·816	56	0·760
18	0·948	31	0·874	44	0·811	57	0·757
19	0·942	32	0·869	45	0·807	58	0·753
20	0·936	33	0·864	46	0·802	59	0·749
21	0·930	34	0·859	47	0·798	60	0·745
22	0·924	35	0·854	48	0·794		

The temperature at which these instruments were originally adjusted by BAUMÉ was 12·5° Cent. (54·5° Fahr.) They are now commonly adjusted in this country at 58° or 60° Fahr.

The degrees of TWADDLE'S hydrometer may be converted into the corresponding specific gravities by multiplying them by 5 and adding 1000.

INDEX

TO THE

VETERINARY FORMULARY

See also pages 489 to 507

[THE VETERINARY MATERIA MEDICA being alphabetically arranged, it is not considered necessary to include the Drugs, whose uses and doses are there stated, in this Index.

Abbreviations employed in this list:—*c*, Cattle; *s*, Sheep; *d*, Dogs; *sw*, Swine. The Horse Medicines have no mark of distinction.]

	PAGE		PAGE
Absorbent Powders	62	BALLS, for horses	43—64
Administration of balls	43	for dogs	143—150
Ægyptiacum	94	alterative	44, <i>d</i> . 144
Aloes, solution of	77	alt. tonic	57
balls	54	anodyne	58
Alteratives	3, 44	antimonial	44, 47
Alterative balls	44, 45	antispasmodic	58
drenches	<i>sw</i> . 141	for appetite	58
powders	62	astrigent	45, <i>d</i> . 144
Alum, solution of	93	bronchitis	47
mouth-wash	89	chewing	59
collyrium	92	common mass for	44
whely	119	cordial	48
Anodyne ball	58	cough	46, <i>d</i> . 144
drench	76	diuretic	49
fomentation	91	diaphoretic	58
poultice	95	for distemper	<i>d</i> . 145, 146
tincture	77	farcy	51
Antimonial balls	44, 47, 50	fever	50
Antiseptic drench	76	for fits	<i>d</i> . 146
Antispasmodic balls	58	glanders	51
drenches	67, <i>c</i> . 113	grease	53, 59
Appetite, balls for	59	hydrophobia	58, <i>d</i> . 150
Apoplexy	134	inflam. of lungs	47
Aromatic powder	61	inflammatory disorders	<i>d</i> . 143
Arsenical powder	64	iodine	58
balls	52	for jaundice	53, <i>d</i> . 149
Astringent balls	45, <i>d</i> . 144	laxative	55
drenches	<i>c</i> . 116	Leicester red	58
electuaries	61	masticatories	59, 120
glyster	79	merc. alterative	45
lotions	87, 89, <i>d</i> . 151	mixed	48
paste	105	miscellaneous	58
powders	107, <i>c</i> . 130, <i>s</i> . 137	mode of administering	43
caustics	<i>s</i> . 139	nauseating	55

	PAGE		PAGE
Balls, parturificient	<i>d.</i> 150	Caustic for foot-root	<i>s.</i> 139
physic or purging	53, <i>d.</i> 143	liquid, 89—91, <i>c.</i> 128; solid	91
poultry	155	Charcoal poultice	95
sedative	59	Charges	106, <i>c.</i> 130
stomachic	55	Chewing balls	59
tonic, mineral	56	Chloride of lime lotion	89
tonic, vegetable	56	poultice	96
tonic, diuretic	50	Citrine ointment	103
St. Vitus's dance	<i>d.</i> 149	Clark's physic balls	54
worm	57, <i>d.</i> 148, 149	conglutinum	88
zinc and valerian	59	embrocation	86
for yellows	53	lotion	86
Barley water	65	pulvis utilis	61
Basilicon	98	Cleansing drench	<i>c.</i> 112
Bites of vipers	<i>c.</i> 119, <i>s.</i> 134	Clysters, <i>see</i> Glysters.	
Black oils	84	Colchicum drench	72
Blain in the mouth	<i>c.</i> 126	Colic in dogs	146
Blanche water	65	Colic drenches	68
Bleeding, 8; to stop	130	tincture	77
Blistering	97	Collyria	92, <i>c.</i> 126, 136, <i>d.</i> 152
Blistering liniments	80	Common ball-mass	44
ointments	96, <i>c.</i> 128	poultice	94
Blown or blast	<i>c.</i> 113, <i>s.</i> 133	Composition for sand-cracks	106
Bran mash	66	Condition powders	63
Braxy	134	Confections, <i>see</i> Electuaries.	
Bog spavin	82	Conglutinum	88, 93
Bronchitis, balls for	47	Convulsions	<i>d.</i> 146, <i>ducks</i> 157
Bruises, lotions for	<i>c.</i> 125	Cooling drinks	66
		Copper, sulphate solution	93
		Cordial balls	48
		diuretic balls	49, 50
CALVES, medicines for	123	drenches	69, <i>c.</i> 113
calving, drench after	112	calves'	123
cordial, &c.	123	powders	<i>c.</i> 120
Canker liniment	83	electuaries	60
Canquoin's caustic	91	Cough balls	46, <i>d.</i> 144
Cantharides ball	52	electuaries	60
drench	74	Cow spice	120
liniments	81	Cow-pock, lotion for	125
oil of	81, 83	Cracked heels, ointment,	99, 100
ointments	96, 97	Creasote ointment	103
tincture compound	94	liniment	83
Carminative drenches	67, <i>c.</i> 119, <i>sw.</i> 141	Cropsick poultry	156
glyster	79	Croton ball	61
tincture	77	drenches	16, <i>c.</i> 110
Catarrh, drenches for	69, 70, 71	electuary	61
Cataplasms, <i>see</i> Poultices.		liniment, <i>c.</i> 127; tincture	77
Catechu lotion, 89; tincture	94	DARBY'S oils	84
Cattle medicines, various	109—130	Decoction of linseed (compound)	66
drenches	109—119	Demulcent drinks	65
powders	119	electuary	60
outward applications	125—130	Deobstruent drenches	74
plague, medicines for	121	Detergent liniments	83
Caustics, Canquoin's and Filho's	91	lotions	88
Caustic, Duville's	139	ointments	98
for farcy buds and poll evil	90		

	PAGE		PAGE
Detergent powders	107	Drenches, cough	70, c. 116
Diabetes, drench for	67	cordial	69, c. 113
Diapente	63	cow-pox	s. 133
Diaphoretic ball	58	demulcent	69—71
Diaphoretic powders	62	deobstruent	74
Diarrhœa, balls (astringent) for	45, 46, <i>d.</i> 144	for diabetes	66
drenches 66, c. 116, s. 132, 134		for diarrhœa	66, c. 116, s. 132, 134
Digestive ointment	98, c. 129	for diarrhœa in lambs	132
Distemper	118	diuretic	71, c. 119
Diuretic balls	49	dropsy	71
cordial balls	49	drying	c. 119
drenches	71, c. 119	for dysentery	66, s. 134
drink	66	emollient	69
electuary	60, 61	for enteritis	67
powders	62	for epidemics	c. 117, s. 133
Divine stone	93	expectorant	c. 116
Dogs, medicines for	143	farcy	72
alterative balls, &c.	144	fever	72, c. 110, s. 132
canker in ear	151	glanders	72
clysters	154	hoose, in calves	124
distemper	145—146	hoven	c. 113
dropsy	150	inflammation of liver	c. 111
emetic powders	147	influenza	72, s. 133, 135
enlarged glands	150	iodine	76
epilepsy, or fits	146, 147	laxative	72, c. 109
eye-water	152	locked jaw	c. 110, 113
hydrophobia	150	murrain	c. 118
inflammatory disorders	147	parturifacient	76, s. 134
lotions, various	151—154	pectoral	69, c. 116
mange, washes, &c.	152	for poisoning by yew	76
ointments	152	purging	73, c. 109, s. 131
piles	153	putrid fever	75
purging balls	143	for rabies	76
rheumatism	148	for red water	c. 115, s. 132
scirrhous tumours	153	rheumatism	c. 113
St Vitus's dance	149	stomach staggers	73
tonic	148	strangury	c. 112
worms	148, 149	tonic	74, s. 132
yellows	149	tonic febrifuge	c. 111
Drenches 65—76, c. 109—119		worm	75, calves 124
alum	c. 124	yellow, or jaundice	c. 111
anodyne	76	Driffield oils	c. 127
antiseptic	76	Drying drench	c. 119
antispasmodic	67, 113	Dysentery	67
astringent	66, c. 116	EAR, canker of	<i>d.</i> 151
for bites of vipers	119	sore	<i>sw.</i> 142
blown or hoven	c. 113, s. 133	Eggs, to promote laying	157
bloody urine	c. 114	Electuaries	60, 61
bulling	119	Electuary, cough and laxative, 60, 61	
for calves, various	123	cordial	60
after calving	112	demulcent and pectoral	60
carminative	67, c. 119, <i>sw.</i> 141	diaphoretic	61
for catarrh	69, 70, 71, c. 116	diuretic	61
for cleansing	112	opiate	60
colic	68		

	PAGE		PAGE
Electuary, purgative	61	Glanders, balls for	51
tonic and astringent	61	drenches	72
Embrocations	85, c. 127	poultry	155
Clark's	86	Glysters	c. 120
Embrocatio frigidifera	86	anodyne	78
excitans	86	astringent	79
Embrocation, cooling	86	carminative and stimulant	79
Embrocation croton	c. 127	cooling	79
for garget	c. 127	diuretic	79
mustard	85	dogs	154
for saddle-galls	85	emollient	78
strains	85	for gripes	79
Emetics for dogs	147	irritable bladder	79
Emulsions	69, 70	laxative	78, c. 120
Enteritis, drench for	67	nourishing	80
Epidemic, affecting mouth and		purgative	78
feet	c. 117, s. 133	uterine	79
malignant	75	worm	79
pleuro-pneumonia	c. 120	Goulard liniment	83
Epilepsy in dogs	146	ointment	99
Euphorbium ointment	99	poultice	95
tincture of	94	water	86, 93
Expectorant balls	46, 59	Grease, balls for	53, 59
External applications, <i>see</i> Lo-		lotions	87
tions, Liniments, Oint-		ointments	99
ments, &c.		poultice	96
Eye-waters	92, c. 126, s. 136	Gripe tincture	77
ointments	99, d. 152	drenches	67
powders	s. 137	Gripes powders	64
FARCY balls	51	Gruel	65
caustic for	90	HAIR, to restore	100
drenches	72	Healing ointment	c. 129, d. 153
lotions	87, 89	Heels, scurfy and cracked	99
ointment	101	greasy	53, 87, 99, 100
Fattening powder	c. 124, <i>svo.</i> 142	Hellebore ointment	101
Febrifuge tonic drench	c. 111	Hepatitis, balls for	53
Feet, stopping for	105	Hind's sweating balls	59
Fever balls	50	Hoof ointment	101, 106
drenches	72, c. 110, s. 132	Hoose	c. 116, <i>calves</i> 124
powders	63	Horse spice	61
medicines for swine	141	medicines	1-108
Fleas	d. 154	Hoven, drench for	c. 113, s. 133
Fly-powder	s. 137	Hydrophobia, remedies for	58, 76
Flux, <i>see</i> Diarrhœa.		INFLAMMATION of lungs	47, s. 135
Fomentations	91	liver	c. 111
Foot-and-mouth disease	117	Inflammatory disorders	
Foot-rot	s. 139	47, c. 124, s. 135, d. 147, <i>svo.</i> 142	
Formulae, veterinary	43-157	Influenza	c. 110, s. 133-135
Foul in foot	c. 125, 130	Infusion of foxglove	77
Foxglove tincture	77	Iodine balls, 58; drench	76
GARGET ointment	c. 129	liniment	84
liniment	c. 127	ointment	101, c. 129
Giddiness	73, <i>poultry</i> 157	tincture	77

	PAGE		PAGE
Iodide of mercury ointment . . .	101	Lotions for bull-burnt	c. 126
Itching, to allay . . .	d. 151	for canker . . .	c. 125
JAUNDICE, balls for . . .	53, d. 149	catechu . . .	89
Joint wounds . . .	c. 129	chloride of lime . . .	89
KNEES, broken . . .	105	Clark's . . .	86
LAMBS, medicines for . . .	131—140	for cow-pox . . .	c. 125
cordial . . .	133	detergent . . .	88
Lapis divinus . . .	93	discutient . . .	86, c. 125
Lapis mirabilis . . .	93	farcy . . .	89
Laxative balls . . .	55	foul in foot . . .	c. 125
and purgative drenches		mange . . .	88, c. 126, d. 152
c. 72, 109, 112		mercurial . . .	88
clysters . . .	78, c. 120	for the mouth . . .	89, c. 126
electuaries . . .	60, 61	nitric acid . . .	88
Leicester red balls . . .	see 58	refrigerant . . .	86
Licc, ointment for . . .	101, c. 128	for strains . . .	87, c. 125, d. 154
liniment for . . .	82	styptic . . .	89
lotions . . .	c. 126	for vermin . . .	89, 126
Liniments, various . . .	80—85	wounds c. 126, d. 154, poultry	157
Liniments of ammonia . . .	84	yellow . . .	88
anodyne . . .	84	Louse-water . . .	89
for sore backs . . .	82	Low, ointment for . . .	c. 130
blistering . . .	80	Lungs, inflammation of . . .	47
creosote . . .	83	MADNESS, canine . . .	150
detergent . . .	83	Materia medica . . .	1—42
Goulard . . .	83	Malignant epidemics . . .	75
for grease . . .	83	Mallenders . . .	102
iodine . . .	84	Malt mash . . .	66
for itching humours . . .	82	Mange, lotions for, 88, c. 126, d. 152	
lime-water . . .	84	liniments and oils for 83, c. 127	
mange . . .	82, c. 127	ointments 101, c. 128, see 142	
marshmallow, narcotic, &c.	83, 84	Marshall's oils . . .	84
saturnine . . .	83	Marshmallow ointment . . .	102
for sore throats . . .	c. 127	liniment . . .	83
stimulating . . .	81, 82	Mashes, bran . . .	66
suppling . . .	106	linseed . . .	66
sulphate of copper . . .	94	malt . . .	66
sweating . . .	80	Masticatories . . .	59, c. 120
turpentine . . .	82	Meadow sickness . . .	s. 133
verdigris . . .	94	Medicated provender . . .	64
Linseed mash . . .	66	Medicine for horses . . .	1—108
tea . . .	65	cattle . . .	109—130
Liquid blister . . .	80	for cattle plague . . .	121—123
caustics . . .	89—91	sheep . . .	131—140
snuff . . .	c. 128	swine . . .	141—142
Lockjaw, drench for . . .	c. 110, 113	dogs . . .	143—154
Lotions, various 86—89, c. 125, 126,		poultry, &c. . .	155—157
d. 151		Mercury, perchloride solution . . .	93
Lotion, astringent . . .	87, d. 151	Mercurial balls . . .	45, 51
black . . .	88	alterative . . .	44
for blain in mouth . . .	c. 126	lotion . . .	88
bruises . . .	c. 125, d. 154	ointment . . .	103, d. 153
		Mild drinks . . .	65
		caustics . . .	90

	PAGE		PAGE
Mineral tonics	56	Ointment, verdigris	98, 105
Miscellaneous balls	58, 59	for vives	105
Mixed balls	48	warts	105
Mixture, smearing	<i>s.</i> 137	Opiate electuary	60
Moulting	157		
Murrain drinks	<i>c.</i> 118	PARTURITION, to promote, 76, <i>d.</i> 150	
lotion	118	Paste, astringent	105
Mustard embrocation	85, <i>c.</i> 127	for broken knees	105
poultice	96	styptic	130
Myrrh and aloes tincture	77	Pectoral electuary	60
		balls	46, 47
NAUSEATING balls	55	drenches	69, <i>c.</i> 116
Newmarket oils	85	powder	64
Nine oils	85	Physic balls	53
Nitre balls	49	Pacumonia, epizootic	<i>c.</i> 120
powder	62, 63	Poisoning by yew	76
OILS, various	83, 85	Pot belly	157
black	84	Poultices, common	94
Darby's	84	anodyne	95
egg	84	charcoal	95
for mange	82, 84	chlorine, cleansing, drawing	95, 96
Marshall's nine, Radley's	84	Goulard	95
Stamford's, Lord	85	for grease	96
Ward's	84	mustard, rubefacient	96
white	84	resolvent	95
Ointments	96—106, <i>c.</i> 128—130	yeast	95
Ointment for blemished knees	100	Poultry, medicines for	155—157
blistering	96, <i>c.</i> 128	Powders, absorbent	62
for canker	104	alterative	62, <i>c.</i> 119, <i>d.</i> 144
citric and creasote	103	aromatic	61
detergent	98	arsenical	64
digestive	98, 129	bitter	64
eye	99	condition	63
euphorbium	99	escharotic	107
farcy	101, 105	fever	63, <i>c.</i> 120
foot	129	farcy	63
grease	99	fly in sheep	137
healing	<i>c.</i> 129, <i>d.</i> 153	tor grease	62
hellebore	101	for gripes	64
hoof	101, 106	pectoral	64
iodine	101, <i>c.</i> 129	purgative	64
for mallenders	102	for outward use	107
mange	101, <i>c.</i> 128, <i>sv.</i> 142	astringent	107
marshmallow	102	diaphoretic	62
mercurial	103, <i>s.</i> 142	detergent	107
nitrate of silver	153	diuretic	62
scab or shab	<i>s.</i> 137	styptic	108
simple	96	sneezing	108
for sore backs	104	veterinary	61
for sore teats	<i>sv.</i> 142	worms	64
for sitfasts	104	Provender, medicated	64
spavins	104	Purging balls	53, <i>d.</i> 143
sulphuric acid	104	drenches	72, 73, <i>c.</i> 109, <i>s.</i> 131
tar	104	electuaries	61
for thrush	104		

	PAGE		PAGE
Purgatives for swine	141	Sturdy, drink for	134
sheep	131	Styptic stone	93, 108
dogs	143	powder	108
for poultry	156	paste	c. 130
Purgative powders	64	Sulphuric acid ointment	104
Putrid fever, drenches for	75	caustic	91
RABIES, drench for	76	Sulphate of copper liniment	94
Rabbits, medicines for	155, 157	balls, <i>see</i> Farcy balls.	
Radley's oils	85	Sweating oils	80
Red water drench	c. 115, s. 132	TAR ointment	104
Résolvent ointment	103	Thriving powder	c. 124, <i>see</i> 142
Rhenmatic drench	c. 113	Thrush, ointment for	104
Rinderpest, treatment of	121—123	Tincture of aloes and myrrh	77
Roaring, ball for	58	anodyne, carminative	77
Rot, foot	134, 135	of cantharides	94
in rabbits	157	colic	77
in sheep	134, 135	of croton	77
Roup, poultry	155	euphorbium	94
SADDLE galls	85, 104	alkaline	94
Sallender's ointment	102	foxglove	77
Sanderack, composition for	106	gripe	77
Sedative balls	59	of iodine	77
Setons	35, c. 130	compound	77
Sheep and lambs, medicines, 131—140		myrrh	77
drench for apoplexy	134	opium, ethereal	77
braxy	134	of pimento	77
diarrhœa	134	styptic	94
purging	131	Tonic balls	56
for red-water	132	drenches	74, c. 111, s. 132
epidemic distemper	133	electuary	61
sturdy	134	Tumours, ointment for, c. 130, d. 153	
skit	133	Turpentine liniment	82
ointment for scab	137	ointment	105
wounds and sore heads	139	ULCERS, lotions for	88
caustic	139	ointments	98, c. 129
Sitfasts, ointment for	104	VETERINARY Materia Medica, 1—42	
Sneezing powder	108	formulæ	43—157
liquid	128	powders	61—64
Solutions, various	77, 78	Verdigris ointment	98, 105
Sore backs, ointment	104	liniment	94
Softening ointment	103	Vermin, to destroy	
Spavins, ointment for	104	89, c. 126, s. 136, d. 154, <i>poultry</i> 157	
Spice, cow	130	Viper, bites of	c. 119, s. 134
horse	61	Vitus's (St) dance	d. 149
Stamford's oils	85	WARD's white oils	84
Staggers, stomach	73	Warts, ointment for	105
Stimulating drench	69	caustic for	d. 154
Stomachic balls	56	Washes	86—89
Stone, styptic	93, 108	black and yellow	88
wounds	93	for scab, &c.	s. 136
mirabilis	93	for maggots	s. 136
divine	93		
Strains, lotions for, 87, c. 125, d. 154			
Strangury, drench for	c. 112		

	PAGE		PAGE
Washes, mercurial . . .	88	Wounds, applications to	
various, <i>see</i> Lotions.		107, 108, c. 129, s. 139. p. 157	
Waters, eye, <i>see</i> Collyria.		&c., <i>see</i> Ointments and Lotions.	
White's solution of potash . . .	124	YELLOWS, or jaundice, c. 111, d. 149	
White egg oils . . .	84	Yew, poisoning by . . .	76
Wind galls . . .	104	ZINC and valerian balls . . .	59
Worms, balls for . . . 58, d. 148, 149		sulphate solution . . .	93
Worm drenches . . . 75, <i>calves</i> 124			
Worm powders . . .	64		

INDEX

*Observe : For Medicines for Horses, Cattle, &c., see the
VETERINARY INDEX, pages 481 to 488.*

	PAGE		PAGE
ABERNETHY'S Pills	158	Acton's solution for baldness	239
Abernethy Biscuits	284	Ægyptiacum	94
Acetate of alumina	298	Acrated chalybeate	204
iron	298	lemonade	269
lime	299	lemonade powders	269
of amylene	344	orangeade	270
soda	299	orangeade powder	270
Acetic acid	299	sherbet	270
perfumes	220, 221	sherbet powder	271
fumigation	354	waters	198
Acetimetry	299	waters, chalybeate	204
Acid, acetic	299	lithia water	199
boracic	300	magnesia water	198
carbolic	300	saline waters	199
carbonic	358	Ague drops	158
chromic	300	Aix-la-Chapelle water	203
cinnamic	300	Albespeyres paper	189
fluoric	300	Albuminous size	304
hippuric	300	Alcohol	304
hydrochloric	301	Alibert's cosmetic	228
iodic	301	Alkalimetry	304
nitric	301	Alkaline lotion for the mouth	260
nitro-hydrochloric	301	Alkaline solution, Brandish's	164
oxalic	301	Allcock's porous plaster	158
phosphoric	302	Allen's anti-fat	158
picric	302	world's hair restorer	158
pyrogallic	302	Alloys, various	305, 306
salicytic	302	Almond flavour	290
sulphuric	302	paste	230
sulphuric, dry	302	powder	231
sulphurous	303	Almond honey paste	231
tannic	303	soap	234
Acidimetry	304	bloom	233
Acids for galvanic batteries	303	Alum	307
Acids and alkaline carbonates, table of the neutralising pro- portions of some	478	mordants	340
Acidulated kali	270, 304	Aluminium, frosted	307
effervescing powders	271	bronze	307
lemonade	268	Amalgams	306
lemonade powder	269, 270	for teeth	264
syrup of raspberries	272	Amalgamated zinc	457
syrup of strawberries	273	Amadou	307
		American shampoo	240
		tooth-powder	252

	PAGE		PAGE
Ammonia, carbonate	307	Azote, or nitrogen	359
sulphate	307	Azure blue	416
Ammoniacal gas	358	BACHER'S pills	161
Ammoniated perfumes	219, 220	Baden water	199
Eau de Cologne	219	Baking powder	161, 286
Anatomical injections	308	Bailey's itch ointment	161
preparations, to preserve	308	Baillie's pills	161
subjects to preserve	308	Balaruc water	202
Anchovies, essence	292	Baldwin's phosphorus	311
paste	293	Baldness, cures for	237-240
Anderson's pills	159	Balm of Gilead	161, 193
Angelica, spirit of	213	of Rakasiri	161
Animal charcoal	328	of Syriacum	161
substances, to preserve	308, 309	Balls, ash	311
Animals, to preserve	309	blackening	312
Annotto, purified	349	breeches	311
solution	309	camphor	321
Anodyne necklaces	159	for copying	311
Anti-attribution	309	wash	235
Antibilious pills	159	heel	311
Anti-fat	158	scouring	311
Antiferment	309	furniture	312
Antipertussis	159	Balsam of honey	181
Antiscorbutic elixir	258	of horehound	177
Antiseptic tooth-powder	252	Bancroft's solution of tin	340
Antispasmodic drops, Grindrod's	180	Bandoline	247
Aperient pills, various	159	Barclay's pills	162
Apothecaries' measure	460	Bareges water	203
weight	460	Baregian balls	162, 187
Apparatus, luting for	326	Barium, peroxide of	312
Aqua cosmetica	227	Bark, essential salt of	162
fortis	309	Barker's tooth-tincture	162
Aquarium, marine, salts for	310	Barley water	275
Arabine	310	Barth's green	418
Arbor Diane	310	Baryta, chlorate of	329
Argentum musivum	310	Baster's farina	280
Armenian cement	323	Bate's eye-water	162
Aromatic pastils	311	anodyne balsam	162
tooth-powder	252	styptic wash	162
vinegar	159, 220	unguentum crinisum	238
Arquebusade (acid)	159	Bateman's drops	162
aromatic	209	itch ointment	162
Arrow-root	274, 433	Bathing spirits	162, 178
Artificial manure	390	Bath digestive pills	162
sea-water	310	lozenges	162
Asiatic tooth-powder	252	Battle's vermin powder	163
Asses' milk, artificial	275	Battley's senna powder	163
Atkinson's infant preservative	160	Baume de vie	163
Atropine paper	160	Bay rum	163
Aurum musivum	311	Baynton's plaster	163
Australian moss	277	Bean flower water	206
Avoirdupois weight	459	Bears' grease	239, 242
table for converting into Troy	459	factitious	242
Axle-grease	309	green	242
		Beavan's cholera remedy	168

	PAGE		PAGE
Béccur's arsenical soap	435	Blue pigments	416
Beddoe's pills	163	Prussian	422
Belloste's pills	163	Saxon	425
Beef tea	281	smalts and azure	416
Beer, ginger	266, 267	Turnbull's	422
spruce	271	verditer, &c.	416
treacle	272	Bochet's syrup	164
Beetle wafers	163, 312	Boerhaave's odontalgic	261
poison	312	Boilers, to prevent incrustations in	317
Bell metal	305	Bones, sulphated	317
Bengal chitni	289	Bookbinders' stains	318
Benzine	312	Boot powder	318
collas	312	Boot-top liquid	317
Benzol	312	Boots, to render waterproof	452
Benzoline	312	Borax, glass of	362
Bestucheff's nervous tincture	163	Borated tincture of myrrh	258
Betton's British oil	163	Bories' odontalgic	259
Bewley's chalybeate	163	Boroglyceride	317
Beverages	266—271	Bottle cement	325
Bird-lime	312	Bouchardat's chalybeate purgative	205
Biscuits, Abernethy	284	Boudet's depilatory	251
aperient	163	Braden's farina	280
meat	282	Brande's toothache tincture	261
sponge	284	ink	371
Bismuth white	233	Brandish's alkaline solution	164
Bisulphide of carbon	313	alkaline tincture of rhubarb	164
Bistre	415	Brandreth's pills	165
Black, Brunswick	320	Brass	305
draught	164	Brazil paper	436
drop	163	Bread, acrated	482
flux	349	unfermented	283
ink	371	gluten	285
dyes	339	Breakfast powder	280
japan	315	Bright's nutritious farina	280
oils	84	Brighton chalybeate water	204
pomatum	250	green	417
reviver	315	Brilliantine	246
Blackening ball	315	Brine for pork	297
liquid	313	British herb tobacco	165
paste	314	oils	165
for harness	315	Brodum's cordial	165
Blackening, various	313—315	Brocchieri's styptic	165
Blackening without polishing	314	Broma	279
patent india rubber	314	Bronze (alloys)	305
Blaine's distemper powder	164	for figures	318, 319
Blake's toothache remedy	261	powder	318
Blanc-mange	278	Bronzing liquids	318
Bleaching of oils	397, 398	Bronzing ball	320
of wax	453	pastes	320
liquid	315	Brosse de corail	320
Blights, remedies for	316	Brown pomatum	250
Blistering paper	164	Browning for gun-barrels	320
tissue	164	Brown's bronchial troches	164
Blood mixture	169	Brunatelli's fulminating silver	352
Blue for linen	316		
ink	375		

	PAGE		PAGE
Brunswick black	320	Catsup, mushroom	293
green	417	walnut	293
Bug poison	330	Cattell's hair dye	249, 250
Bullock's semola	280	Caustic for corns	171
Burnett's (Sir W.) disinfecting fluid	165, 321	Cayenne, soluble essence	290 291
Busang water	204	Celery, essence of	291
Butter of antimony	321	Cements, various	323—326
to preserve	321	Armenian	323
Butyric ether	321	for apparatus	326
		Cement, bottle	325
CABBAGE paper	436	cap	324
Cachou aromatisé	225	coppersmiths'	325
Cadet's tooth-powder	252	curd	324
Cajeput liniment	165	diamond	323
Calvert's disinfecting powder	336	engineers'	325
Camphine	321	for cracks in iron	325
Camphor, artificial	321	for india rubber	326
ball	231	for jewels	323
fluid	188	for leather and metals	326
ice	231	liquid	323
liniment	165	for marble	324
washballs	235	French	326
Camphorated chalk	252	hydraulic	325
tooth-powder	253	Maissiatt's	325
Candies	322	oxychloride of zinc	325
Candy, caraway	186	Roman	325
live-long	186	shell-lac	323
Cantharidin blister	322	soft	326
Canton's phosphorus	322	for teeth, various	263—265
Caoutchouc, solvents for	322	for wood, porcelain, and glass	323
Cap cement	324	for fixing labels to tin	326
Capillaire syrup	272	Singer's	324
limoniated	322	Turner's	325
Capsules, gelatinous	186	Cephalic snuff	165
Caraway candy	291	Ceylon moss jelly	276
essence of	291	Chalybeate, Palmer's aerated waters	189 204
Carbon, <i>see</i> Charcoal.		Chamberlaine's pills	166
writing ink	377	Chameleon mineral	328
bisulphide of	313	Chamomile pills	166
Carbonic acid gas	358	drops	166
oxide	358	Chapman's wheat flour	280
dentifrice	253	Charcoal, animal	328
Carburetted hydrogen	359	aluminized	328
Carlsbad water	199	from coal tar	328
Carmine	232, 413, 414	vegetable	327
Carminative, Dalby's	171, 172	Charcoal tooth-powder	253
Carabelli's tooth-powder	253	tooth-paste	257
Carrara water	199	Chelsea pensioner	166
Carron oil	165	Cheltenham salts	166
Carthamine lake	415	Chemic blue	329
Cartwright's tooth-powder	253	Cherry paste for teeth	257
Case-hardening powders	322	Chevallier's hair dye	248
Cassareep	294	Chilblains, remedies for	166, 167, 184
Castillon's powders	165		
Castor-oil pomade	243		

	PAGE		PAGE
Chinese depilatory	251	Colours for maps	333
Ching's lozenges	167	for confectioners	333
Chirayta pills	168	for druggists' show bottles	331, 332
mixture	168	liqueurs	333
Chitni, Bengal	289	various	331
Chloralum	329	vehicles for	333
Chlorate of barium	312, 329	Coloured fires	345, 346, 347
of potash	329	flames	347
Chloride of lime	329	coloured flame papers	347
potash	330	Colouring for brandy, &c.	333
soda	330	Comparison of thermometric	
Chlorine gas	358	scales	462
fumigation	353	Composition for electro-moulds	343
lozenges	226	Condiments, various	288—290
Chlorodyne	167	Condy's fluid	333
Chloroform	330	Consumption, remedies for	169
Chocolate and cocoa	278	Copaiva, solution of	194
Cholera, remedies for	167, 168	Copland's pills	160
Chromate of lead	416	Copper, black and red oxide	333, 334
of potash	330	nitrate of	334
Chrome red, &c.	416	Coral dentifrice	253
yellow	416	tooth-paste	256
Chromic acid	300	Corns, caustic for	171
oxide	330	plasters	170
Cinnamic acid	300	plasters, mechanical	170
Cinnamon, essence of	290	solvents for	170
water	206	Cosmetics, hair	237
tincture of	290	skin	227
Circassian cream	243	teeth	252—260
tooth-powder	253	for chapped nipples	184
Citrate of magnesia	169	Cosmetic gloves	232
Clark's blood mixture	169	Cosmétiques, in sticks for hair	250
Clichés moulds	306	Cottereau's toothache drops	261
Clothes, powder to keep moths		Cotton, gun	368
from	330	Cough lozenges	170
Clutton's febrifuge	169	linctus	169
Coathupe's writing fluid	377	Court plaster	171, 369
Cochineal colouring	330	Crayons, lithographic	379
lake	414	for writing on glass	334
liquid	388	Cream, Circassian	243
Cochrane's cough mixture	169	cold	229, 230
Cod-liver, Soyer's preparations of	287	lemon	228
with cod roe	287	wash-balls	235
and potatoes	287	Crystalline cream	243
and rice	287	castor-oil pomade	243
and tapioca	287	Cucumber pomade	229
preparations of	287, 288	Culinary essences, &c.	290—294
Cod-liver oil sauce	287	Currant jelly	274
Coffee, dandelion	172	Curry powder	288, 289
Cold cream	229, 230	Custard powder	286
granulated	230		
Collier's wine of quinine	169	DAFFY'S elixir	171
Collins's disinfecting powder	169, 335	Daguerreotype plates	419
Collodion	330	Dalby's carminative	171
elastic	330	Dahlia test	436, 438
styptic	169		

	PAGE		PAGE
Dandelion coffee	172	Easton's tonic syrup	173
Darby's oils	84	Eaton's styptic	173
Darceet's lozenges	172	Eau d'ambre	211
Dawson's lozenges	172	d'ange	209
De Handel's toothache paste	262	de Bottot	259
Delamott's golden drops	172	de bouquet	211
Denham's food	280	des carmes	209
Dentifrices, various	252—256	de Cologne	208
Deodorizing compounds	335, 336	collante	247
Deodorizing fluid, Ellerman's	335	dentifrice de Stahl	259
Depilatories, various	251	d'Egypte	249
Derbyshire's embrocation	172	d'élégance	210
Deschamps's tooth-powder, 253, 254	254	de fée	173
Desforges's tooth-powder	254	d'hélotrope	210
extract of pellitory	258	de javelle	174
Deshler's salve	172	d'Ispahan	210
Detergent tooth-powder	254	de luce	219
Deutoxide (peroxide) of hydrogen	399	de mareschale	210
Dextrin	334	de miel	209
Diapente	63	de millefleurs	210
Diamond cement	323	de magnanimité	173
Diastase	334	médicinale d'Husson	173
Dictamia	279	de O'Meara	259
Dietetic articles	274—286	de Naphe	206
Digestive pills	172	de Portugal	210
Dinneford's fluid magnesia	199	Romaine	210
Dinner pills	172	de Rabel	174
Disinfectants	335	sanspareille	211
Disinfecting fluid, Burnett's, 165, 321	321	sedative	174
Ellerman's	335	de Rosières	211
Ledoyen's	183, 335	de Madame Vrillière	259
compounds	335, 336	Ebony pomatum	250
Distemper powders, Blaine's	164	Edinburgh lozenges	186
Dixon's pills	173	ointment	174
Daver's powders	173	Effervescing aperient	188
Drayton's silvering process	365	citrate of magnesia	174
Drinks, summer	266	powders	267—271
Dryers, for painters	336	lemonade	269
Drying oil	336	powder	269
Dubbing, curriers'	336	orangeade	270
Duffin's Harrogate salts	181	orangeade powder	270
Duncan's lactuc. lozenges	173	sherbet	270
fl. ext. of senna	173	sherbet powders	271
gout remedy	173	chalybeate water	204
Dupasquier's gaseous water	205	Seidlitz powders	193
Dupuytren's eye-salve	173	Eger water	199
pomade	237	Eggs, to preserve	342
Dutch drops	173	Elderflower water	206
Dyers' compounds	337—341	Electric tissue	342
Dyes, various	337—339	Electro-gilding	343, 362
for the hair	248—250	solutions for	362
for ivory	386	brassing	342
for wood	455	bronzing	342
Dyon's tooth paste	257	coppering	342
EARACHE, simple cure for	265	silvering	343, 428
		Electrotype moulds	343

	PAGE		PAGE
Electrotype solutions	431	Essence, mignonette	217
Electuary, Mahomed's	186	millefleurs	217
Elixir, Daffy's	171	mint	176
de Garus and Haller's	174	moss rose	217
Longæ Vitæ	174	myrtle	217
Radcliffe's	191	musk	217
of roses	258	mustard	176
Squire's	194	peppermint	176
Stoughton's	194	pennyroyal	176
for the teeth	258	patchouli	215, 217
of vitriol	174	perfumery	214—219
Ellerman's deodorizing fluid	174	of pine-apple	291
Emerald green	418	of rennet	424
Ems water	200	of roses	212
Embrocation, Roche's	192	rondoletia	217
Embrocations for the hair	239—241	royale	217
Emulsions, cosmetics	228	of sarsaparilla	176
Engraving, liquid	344	for scenting pomatums	218
on glass	363	for smelling bottles	220
Engravings, to clean	343	sweet briar	217
Engravers' varnish	446	sweet pea	217
Epicurean sauce	293	of Tyre	249
Equivalents, table of	466	of vanilla	214
Ervalanta	280	of verbena	218
Eschalot wine	292	of vetiver	214
Esprit de suave	211	of violets	212, 219
rose, &c.	212	Westphalian	297
Ess bouquet	213	wood violet	215
Essence of ambergris	216	white lilac	218
anchovies	292	Essences, culinary	290, 291
d'ambrette	214	compound	215
of bark	175	conc. infusions	177
of bitter almonds	290	vinous	177
Cambrian	297	perfumed	215—219
of caraway	291	Etching fluids	344
of cayenne	291	varnish	344
cedrat	216	Extrait de bouquet	215
of celery	291	de mareschale, &c.	215
chamomile	176	Extract of liquorice	389
cinnamon, cloves, &c.	290, 291	of meat	282
of clove pink	216	of malt	177
Cologne	216	Extracts (<i>Extraits</i>) of flowers	215
cubeb	175	FAIRBORN'S pills	177
flavouring	290	Fards, or face paints	232, 233
ginger	175	Farinaceous food, Hard's, &c.	280
soluble	175	Farina, <i>see</i> Starches	433, 434
heliotrope	216	Bright's nutritious, &c.	280
honeysuckle	216	Fats, to obviate rancidity in	345
hovenia	216	Feculum Saxonica	279
jargonelle pears	291, 344	Filter for corrosive liquids	345
jessamine	215, 219	Filtering powder	345
jonquil	216	Finings for spirits	345
lavender	207	Fireproofing	347, 402
lily of the valley	216	Fires, coloured	345, 346
Madden's	186	to extinguish	345
magnolia	216		

	PAGE		PAGE
Fixateur	247	Furniture cream	355
Flames, coloured	347	oil	356
Flame papers coloured	347	paste	355
Flavouring essence	290	Fusible metal	305
Flint, liquor of	347	GAD'S cement	325
Floating soaps	235	Galvanic battery, acids for	303, 304
Florentine dentifrice	254	tooth-powder	254
Florida water	211	Gall, clarified	356
Flowers, to expand	348	Galls, tincture of	438
to preserve	348	Gants cosmétiques	232
Fluide de Java	246	Garancine	357
Fluxes, black and white	349	Gardiner's alimentary prepara- tions	280
Flux, Christison's	349	Gannal's solution	308
crude	349	Ganteine	356
Cornish	349	Garrot's covering for pills	357
Morveau's	349	Gases, to collect	357
Fresenius's, &c.	349	Gas, ammoniacal	358
Taylor's	349	carbonic acid	358
Fly poison	349	carbonic oxide	358
Food for infants, &c.	280	chlorine	358
Foot powder	336	hydrogen	358
Ford's balsam, horehound	177	hydrochloric acid	358
laudanum	177	carburetted hydrogen	359
Foreign weights and measures	461, 462	phosphuretted hydrogen	359
Forges water	204	sulphuretted hydrogen	359
Fothergill's pills	177	nitrogen	359
Fox's cream	244	protoxide	360
Frangipanni bouquet	218	oxygen	360
Franks' solution	177	ozone	360
Freckles, lotion for	228	sulphurous acid	360
Freeman's bathing spirit	178	Gauze, Lister's carbolized	360
Freezing mixtures	349, 350	Gelatin	361
French polish	350, 351	capsules	322
lip-salve	232	bone	361
mustard	292	wafers	451
tooth-powder	254	Gelée pour le goître	178
Friedrichshall water	200	Gentil's gluten flour	286
Fryar's balsam	178	German lip-salve	232
Frigorific mixtures	349	paste	361
Fruit lozenges	186	pomade	242
Fulminating compounds	352	silver	305
cotton	368	tooth-powder	254
powder	352	Gilding	361
mercury	352	electro	362
silver	352	by immersion	362
Fumigating pastils	224	Ginger beer	266, 267
Fumigations, acetic	354	beer powders	267, 268
carbolic acid	353	essence of	175
chlorine	353	aromatic tincture	291
iodine	354	Gingerbread, common	285
muriatic	354	cake	285
nitric	354	purgative	178
sulphur	355	Girambing	267
tar	355	Glaes	362
for toothache	262		

	PAGE		PAGE
Glass, to cut	. 363	Gum, British	. 367
to mark or write on	. 363	Arabic, purified	. 368
to silver	. 364, 365	Gun barrels, browned	. 320
of borax	. 362	cotton	. 368
flux	. 324	powder	. 368
to platinize	. 364	white	. 368
soluble	. 362	Guthrie's black ointment	. 180
toughened	. 362	eye ointment	. 180
to clean	. 365	Gutta-percha moulds	. 343
Glazes	. 366	solvents for	. 430
Gloves, to clean	. 387	Gypsum, to harden	. 369
Glue	. 366		
liquid	. 366	HÆMLEM drops	. 173
marine	. 367	Hahnemann's hair dye	. 249
Gluten, vegetable	. 367	wine test	. 369
bread	. 285	Hair cosmetics	. 237
granulated	. 286	dyes, various	. 248—250
Glycerine jelly	. 178	oils	. 244—246
jelly for microscopic		pomades	. 241—244
mounting	. 394	powder	. 237
and lime cream	. 244	restorer	. 249
paste	. 178	restorer, Allen's	. 158
Goody's solutions	. 308	washes	. 246, 247
Godbold's veg. balsam	. 178	to remove	. 251
Godfrey's cordial	. 178, 179	to promote	. 237—240
smelling salts	. 179, 220	to stiffen	. 247
Gold, aluminium	. 306	Hands, bad odours to remove	
common	. 306	from	. 232
factitious	. 306	Halford's (Sir H.) pills	. 180
mosaic	. 306	Hamilton's pills	. 160
Golden compound	. 367	Harcees blacking	. 315
hair dye	. 249	jet	. 369
ointment	. 180	waterproof paste	. 369
spirit of scurvy grass	. 180	Harrogate salts	. 181, 203
Gout, specific, Murray's	. 188	water	. 203
paper	. 180, 189	Hartshorn jelly	. 277
medicine, Duncan's	. 173	Heading for beer	. 369
pills, Lartigue's	. 183	Helvetius' styptic	. 181
tincture, Wilson's	. 197	Hemet's dentifrice	. 254
Granulated gluten	. 286	Henbane fumigation	. 262
Granville's (Dr) lotions	. 180	Henry's magnesia	. 181
Grape lip-salve	. 232	Herb soup-powder	. 289
Graves's (Dr) chilblain remedy	. 166	tobacco	. 165
gout preventive	. 180	Hill's balsam of honey	. 181
Grecian water	. 249	Holloway's pills	. 181
Gregory's powder	. 180	ointment	. 181
Green dyes	. 339	Honey paste	. 231
pigments	. 417, 418	soap	. 234
Greenough's tincture	. 180, 259	water	. 209, 210
Griffin's tincture	. 180	Hoffman's pills	. 181
Griffith's mixture	. 180	Hooper's pills	. 181
Grindrod's antispasmodic	. 180	Hooping-cough remedies	. 182
Grosvenor's tooth powder	. 254	Horse-radish powder	. 290
Gruel, oatmeal	. 281	Holton's laudanum	. 181
Guarana	. 279	Hudson's tincture	. 259
Guestonian embrocation	. 180		

	PAGE		PAGE
Huiles antiques	244	Ink, Redwood's	381
rose, jessamine, &c.	244, 245	Stephens's	375
macassar	246	Inks, various	371—384
Huiles de phénix	245	writing, black	371
philcome and comogene	245	anti-corrosive	371
Hungary water	209	copying	373
Hunyadi Janos water	200	packers'	373
Huxham's tincture of bark	182	Insects, to kill	384
Hydrochloric acid	301	bites, to cure	384
gas	358	Iodate of potash	384
Hydrogen gas	358	Iodide of potassium	384
deutoxide (peroxide)	399	Iodine	385
carburetted, &c.	359	Iodized test	436
Hypochlorite of potash	330	Iridescent paper	400
Hyposulphite of soda	430	Irish moss, preparations of	276
.	276	Iron, acetate of	298
chocolate	276	liquor	298, 341
Imperial drink	268	persulphate of	385
pop	267	cement	325
measure	460	Isinglass	385
Incense	224, 369	plaster, Liston's	185
India pickle	297	Issue peas	182
rubber, solvents for	430	Ivory, artificial	387
court plaster	369	black	386
Indigo purified	370	flexible	386
dyes	337	to stain	386
solutions	329	to bleach and gild	386
test paper	436	JAMES's analeptic pills	182
Infants' preservative	160	powder	182
Infusions concentrated	177	Japan	387
Ink, to keep from mouldiness	384	black	315
alizarine	373	Japanese cement	326
aniline black, marking	381	Jarave, Spanish	182
powder	373	Jelly, currant	274
red	374	gelatin	277
blue	374	hartshorn	277
green	376	Irish moss	276
yellow	376	carrageen	276
gold and silver	377	Iceland moss	276
indestructible	377	of Ceylon moss	277
Indian	378	salep	277
horticultural	378	isinglass	278
for writing on steel	378	Jessamine, esprit	212
lithographic	379	perfumed oil of	244
for marking linen	379	extract	215
without preparation	380	water	213, 219
for printing cotton fabrics, &c.	378	Jesuits' drops	182
crimson marking	382	Jockey club bouquet	212
printers'	382, 383	Jones's patent flour	284
Runge's	373	KALYDOR lotion	227
sympathetic	383	Keating's lozenges	183
Weber's	374	Kemmerer's tooth-powder	255
violet	375	Kennedy's corn-plaster	183
Normandy's	375, 378	Keyser's pills	183
Reade's	372		

	PAGE		PAGE
Kid glove cleaner	387	Lemon embrocation	228
King Cup	268	cream	229
King's yellow	417	wash balls	235
cordial	183	Lemonade, acidulated	268
Ketchup, mushroom	293	acerated	269
walnut	293	dry	269
Kirkland's cerate	183	powders	269, 270
Kittoe's freckle lotion	228	milk	269
Kitchener's essence of cayenne	291	Lentil powder	280, 281
of ginger	291	Leroy's purgative	184
of spices	291	Lenses, extemporaneous	389
pease powder	289	Liebert's nipple cosmetic	184
peristaltic persuaders	183	Liebig's food for infants	282
soup herb-powder	289	soup for children	282
spirit of soup herbs	291	Lignum's drops	184
savoury spices	289	Lime juice and glycerine	244
tincture of cinnamon	290	Limoniated ginger-beer	267
Koumiss	387	Linctus, for coughs	169
Kyan's solution for wood	387	Liniment, cajeput	165
		camphorated	165
LABARRAQUE'S liquid	387	St John Long's	185
Labels, indestructible	387	Linseed tea	276
gold to attach	387	oil, refined	359
Lac, stick, seed, and shell	388	boiled	356
to bleach	388	clarified	359
cement	323	Lip-salve, rose	232
spirit	341	Peruvian	232
Lacquers	388	German	232
Lactucarium lozenges	186	grape	232
Lait de fraicheur	228	French	232
virginal	228	Liqueur dorée	184
Lake, cochineal	414	Liquid amalgam	306
Brazil, &c.	415	Liquids for baldness	239, 240
liquid	388	blister	184
yellow, &c.	415	disinfecting	335
Lakes	388, 415	glue	366
Lamplough's pyretic saline	183	Liquids for the teeth	258-260
Landerer's cure for baldness	240	tables of weights and	
Lane's, Dr, vinous essences	177	measures of	459-462
Lardner's tooth-powder	255	Liquorice, purified	389
Lartigue's gout-pills	183	Lisbon water	210
Lavender tooth-powder	255	Lister's carbolized gauze	360
water	207, 208	Liston's plaster	185
Leath's alimentary farina	280	Lithographic paper	399
Lee's pills	184	ink	379
Ledoyen's disinfecting fluid	183, 335	Litmus	390
Leech-bites, astringent for	183	paper	437
Lefoulon's elixir	259	tincture	438
tooth-powder	255	Live-long candy	186
Lefandinière's elixir	259	Locatelli's balsam	188
Lejeune's balsam	184	Locock's lotions for the hair	259
Lemazurier's odontalgic	261	wafers	185
Lemon juice, factitious	268, 388	Long's, St John, liniment	185
flavour	290	Lotions for the mouth	260
kali	270	of chloride of soda	260
Lemon, pickle	293	for chilblains	166, 167

	PAGE		PAGE
Lotions for the skin	227	Metallic cements	264
Lozenges	390	Metge's tooth-powder	255
absorbent	185	tooth-paste	257
aperient	185	Methylated spirit	392
Ching's	167	Mexican hair renewer	187
black currant	185	Mialhe's dentifrice	255
and ipecac.	186	chalybeate water	205
cough	170, 186	ioduretted water	205
digestive	186	Mice and rat poison	420
Edinburgh	186	Microscope, formulæ for	393, 394
fruit	186	Milk of cucumbers	227
marshmallow	186	of houseleek	228
peppermint	186	condensed	395
Lubricating compounds	390	of roses	227
Lucifer matches	392	preserved	395
Luminous phials	390	Mineral green	417
Lutes, various	326, 327	waters	198-205
Lynch's embrocation	186	chameleon	328
		Miscellaneous preparations, 298-458	
MACASSAR oil	246	Mistura odorata	218
McDougal's disinfecting powder	336	Modelling, clay for	395
Madden's essence	186	Mohr's lute for retorts	327
Magnesia, citrate of	169	Moirée métallique	395
fluid	187, 199	Montein's balls for baths	187
Magnesian aperient	188	Mont d'Or water	204
Mahomed's clectuary	186	Mordants	339-341
Mahy's plaster	187	alum	340
Moissiat's cement	325	tin	340
Manures, artificial	390, 391	Morphia collodion	188
Maredant's Norton's drops	187	Morrison's paste	188
Marble, to clean	391	pills	188
to stain	391	Morton's drops	187
Marienbad water	200	Morveau's flux	349
purging salts	200	Moseley's pills	188
Marine glue	367	Moss, Australian	277
soap	429	Ceylon	277
Marking ink for linen	379	Iceland	276
without preparation	380	Irish	276
Marmoratum	264	Moth powder	330
Marmalade	273	Moulds, electrotype	343
Marsden's drops	187	Mouth cosmetics	252-260
Marsh's dahlia test	438	Moxon's magnesian aperient	188
Marshall's cerate	187	Munro's cough medicine	188
eye-drops	187	Mucilage for minerals	324
Marrow oil	246	Multum	395
pomatum	242	Muriatic acid	301
Mastic varnish	445	Muriate of tin	340
Matches, lucifer, &c.	391, 392	Murray's fluid magnesia	187, 199
Mathieu's vermifuge	187	camphor	188
Maury's tooth-powder	255	gout specific	188
Meat biscuit	282	Mushroom catsup	293
extract of	282	Musk water	207
Medals, to bronze	319	Mustard for the table	292
Medallions, perfumed	223, 224	French	292
Mercury, fulminating	352	leaves	188, 192
		tissue	188

	PAGE		PAGE
Myrrh dentifrice	255	Orangeade powder	270, 271
tincture, odoriferous	258	Orange chrome	417
NANKEEN dye	339	juice, factitious	268
Naples water	203	marmalade	273
yellow	417	Orfila's hair-dye	248
Neligan's, Dr, pills	160	Or-molu	306
Nelson's gelatin	361	Ormskirk medicine	189
Neuraline	188	Ostermaier's cement	263
Nepenthe	188	Oxygen gas	360
Neutralizing proportions, table of	478	Oxygenated water	399
New-mown hay	212	Ozone	360
Nipples, lotion for	184	PALAMOUD	279
Nitrate of baryta	395	Palm oil, to bleach	397
of silver	396	Palmer's tooth-powder	255
strontian	396	aerated chalybeate	189
Nitrite of potash	396	vitaroborant	280
Nitrogen gas	359	Panada	275
protoxide of	360	Paper, Brazil, cabbage, &c.	436, 437
Nitro-prusside of sodium	396	copying	399
Norris's drops	189	fire-proofing for	400
Nouffer's worm medicine	189	hydrographic	400
Novargent	397	tests, various	436, 437
ODONTALGIC tinctures	261	lithographic	399
elixir	259	oiled	400
Odor delectabilis	212	parchment	400
Odoriferous tincture of myrrh	258	paste	401
lavender water	207	photographic	400
Oil, boiled and drying	336	tracing	400
furniture	356	iridescent	400
Macassar	246	waxed	400
marrow	246	Papier épispastique	189
Oils of ambergris, musk, &c.	245	Fayard	189
roses, violets, &c.	245	maché	401
perfumed, various	244—246	Papyrine	401
for the hair	245	Paraguay-roux	261
to purify	397, 398	Paraffin	401
to bleach	397, 398	Parfum des rois	211
for machinery	399	Passy water	204
Carron	165	Paste	326
nine	85	almond	230
vegetable	449	flour	326
watchmakers'	452	furniture	355
Ointments, Bailey's	161	for the hands	236
Bateman's	162	honey	231
Edinburgh	174	Regnauld's	191
golden	180, 193	Swediaur's	166
Holloway's	181	tooth, various	256, 257
Plunket's	190	for toothache	262
Smellome's	193	Ward's	196
Oleine	399	for cleansing brass	401
Ollivier's biscuits	189	for razors	402
Opodeldoc	189	blacking	314
Orangeade	270	Pastils for burning	224
aerated	270	for the mouth	225, 226
		Pâte divine de Venus	229

	PAGE		PAGE
Payne's preparation for wood	402	Pills, Belloste's	163
Pearl dentifrice	255	Brandreth's	165
white	233	Chamberlaine's	166
Pease powder	209	chamomile	166
Pelletier's quinine dentifrice	255	digestive	172
odontine	257	dinner	172
Pensioner, Chelsea	166	Dixon's	173
Percussion caps, priming for	402	Fairthorn's	177
Peristaltic persuaders	183	Fothergill's	177
Perfumes, various	206—226	Halford's, Sir H.	180
acetic	220	Hamilton's	160
ammoniated	219, 220	Holloway's	181
Persian insect powder	190	Hooper's	181
Perry's drops	187	James's analeptic	182
Peruvian lip-salve	232	Johnson's	160
Peter's pills	190	Kitchener's	183
Pettenkofer's test	441	Lartigue's	183
Pharaoh's serpents	402	Lee's	184
non-poisonous	402	Lynn's	160
Philocomo	245	Morrison's	188
Phosphorescent oil	403	Moseley's	188
Phosphoric acid	302	Neligan's	160
Phosphorus	403	Reece's chirayta	191
Baldwin's	311	Robinson's	192
Canton's	322	Scudamore's	160
matches	403	Smith's (Hugh)	193
bottles	403	Speediman's	193
paste	403	Vance's	160
*Photography and preparations	403—413	Ward's red	196
Phosphuretted hydrogen	359	Whitehead's	197
Pickles, various	294—297	covering for	190, 357
Pickled beet	295	Pink saucers	419
cabbage, cucumber, &c.	295	Pivots for teeth	265
mangoes and mushrooms	296	Plasma	190
nasturtiums and onions	296	Plaster, Bayton's	163
walnuts	296	court	171, 370
piccalilli	297	isinglass	185
Pickle for pork, &c.	297	Liston's	185
Pigments, various	413—419	Mahy's	187
vegetable	413—416	Plate powder	419
mineral	416—419	for boiling	419
Piles, remedies for	190	Plates, Daguerreotype	419
Pills, Abernethy's	158	Platinum, black	420
Anderson's	159	chlorides	420
aperient	159, 160	sponge	420
antibilious	159, 160	Platinated asbestos	420
Bacher's	161	Platinized silver	420
Baillie's	161	Plombières water	201
dinner	161	Plumbe's farinaceous food	280
Barclay's	162	Plunket's ointment	190
Bath	162	Poison for beetles	163, 312
Beddoe's	163	for bugs	320
		for rats and mice	420

* Pages 403 to 413 include directions for the preparation of the various chemicals, solutions, papers, &c, employed in this art.

	PAGE		PAGE
Polish, French	351	Powell's balsam	191
furniture	355, 356	Preserving animals	308, 435
Polishing powder	420	Prince of Wales's food	280
Pomades	241—243	Prometheans	392
for baldness	237, 239	Provin's water	204
Pomatums	241—243	Prussiate of potash	421
Pomatum, common	241	red	422
coloured, &c.	243	Prussian blue	422
crystalline	243	soluble	422
cucumber	229	Turnbull's	422
marrow	242	Pullna water	200
hard or roll	242	salts	201
rose, &c.	242	Purging drink, Dr Young's	197
Pommade contre l'alopecie	238	Purple of Cassius	422
castor oil	243	Puscher's indelible ink	378
castor oil and glycerine	243	Pyrmont water	204
Cattell's, Dr	238	Pyro-acetic spirit	423
Cazenave's	238	Pyrogallic acid	302
collante	247	Pyroligneous acid	423
d'Hèbe	229	Pyroxylic spirit	423
de jeunesse	250	Pyrophorus	424
divine	190, 230	QUEEN of Hungary's water	191, 209
Dupuytren's	237	Quillai bark	244
for beautifying hair	242	Quin sauce	293
for the skin	229	Quinine dentifrice	255
de beauté	229	wine	169
de Ninon, &c.	229	RACAHOUT des Arabes	279
Pond's extract	191	Radcliffe's elixir	191
Pot-pourri	222	Ragout spice	289
Potash, carbonate	420	Raspberry vinegar	273, 292
Pop, imperial	267	syrup, acid	272
Portland's powder	191	Raspail's cigarettes	191
Potato-disease, remedies for	421	Rat poisons	420
Poudre clarifiante	421	without arsenic	403
métallique	264	Redwood's marking ink	381
à la mousseline	223	depilatory	251
de Chypre, &c.	223	Reece's chirayta pills	191
Pounce for parchment	421	Regnault's dentifrice	255
marking ink	380	pectoral paste	191
Powder, almond	231	Rennet	424
baking	161, 286	liquid	424
filtering	345	wine	286
fulminating	352	Reynolds's gout specific	191
ginger-beer	267, 268	Revalenta	191, 280, 281
gun	368	Rhatany tooth-powder	255
hair	237	Rice cake	284
hand	236	Riga balsam	192
lemonade	269, 270	Righini's tooth-powder	255
orangeade	270, 271	Rigollot's mustard leaves	192
spruce-beer	272	Robinson's pills	192
soda	193	tooth-cement	263
soap	236	Roche's embrocation	192
Scidlitz	192, 193	Rogé's magnesian purgative	192
for rats, &c.	420	Rose lip-salve	232
plate	419		
Powders for the teeth	252—256		

	PAGE		PAGE
Rose tooth-powder	255	Scott's plaster	192
pink	415	pills	192
soap	234	Scouring drops	425
tooth-paste	257	Scudamore's pills	160
water	219	Scurf, wash for	247
Rosemary tooth-paste	257	Sea water	202, 310
Rossiter's hair restorer	192	salts for	427
Rouge	233	Sealing-wax	425—427
jewellers'	419	Seidlitz water	201
Rousseau's drops	192	salts	201
Rubini's camphor	192	powders	192, 193, 201
Ruspini's tincture	260	Seidschutz water	201
tooth-powder	256	Seltzer water	201
styptic	192	Semolina	280
Russian tooth-powder	256	Shaving liquid	236
Rust's toothache pill	262	paste	236
Rust, to remove and prevent	425	powder	236
Ryan's essence of coltsfoot	192	Shampoo liquor	240
Rymer's tincture	192	Shell-lac	388
		to bleach	427
SACHETS	222	cement	323
Sago, to prepare	275	Sherbet	270
milk	275	powders	270, 271
posset	275	Siemmerling's cosmetic	228
Saint Galmier water	201	Silica cement	264
Safflower	331	Silk-cleaner	427
Salep	277	Silver, to clean	427
Saline solutions	432	fulminating	352
tooth-paste	257	oxidized	428
waters	199—202	to purify and reduce	428
powders, Stevens'	167	tree	442
Salts, Cheltenham	166	Silvering	427
Harrogate	181	glass	364, 365
Marienbad	200	silvering paste	427
Pulna	200	powder, &c.	427
Seidlitz	201	Sinapine tissue	193
smelling	230	Singer's cement	324
Vichy	202	Singleton's ointment	193
Sanitas	336	Sirope d'orgeat	272
Sap green	415	Size, oil, &c.	429
Sauce, anchovy	292	Skeleton leaves	429
Epicurean	293	Skin cosmetics	227
Quin	293	Smalts	429
superlative	294	Smelling salts	220
Waterloo	293	Smellome's eye-salve	193
Saunders' dentifrice	256	Smith's (Hugh) pills	193
Savon au bouquet	234	Smoking fluid	429
Savonettes	235	Snuff, cephalic	165
Savoury spices	289	Soap	429
Scent bags	222	arsenical	435
balls	223	carbolic	234
for pomatums	218	marine	429
powders	223	powder	236
for snuff	218	paste	236
Scented soaps	233—236	toilet, scented, various	233—236
		floating	235

	PAGE		PAGE
Soap, transparent	235	Stopping for teeth	263—265
Windsor	234	Storey's worm cakes	195
Soda, hyposulphite of	430	Storm-glass	434
powders	193, 271	Stoughton's elixir	194
Soft cement	326	Strawberry syrup	273
Solders, various	306	spirit	215
Solomon's anti-impetigines	193	water	206
balm of Gilead	193	Stuffing birds, &c.	435
Solvents for gutta percha	430	Styptic, Brocchieri's	165
india rubber	430	Eaton's	173
corns	170	Heivetus's	181
old putty	431	Rubini's	192
silver	432	Styrol	435
Soluble glass	347, 362	Struve's lotion	195
cayenne	290	Sugar resin	435
Solution of copaiva	194	Sulphide	435
for electro-plating	431	Sulphite of copper	435
Gannal's	308	Sulphocyanide of ammonium	435
for soldering	431	Sulphuretted hydrogen	359
for voltaic battery	431	waters	203
for water-baths	432	Swaim's vermifuge	195
Soubiran's tooth-cement	263	Sydenham's laudanum	195
Soup for children	282	Syrup of almonds	272
Soup-powder	289	capillaire	272
Soy	294	limoniated	272
Spa water	204	milk	435
Specific gravities	464	pine-apple	272
corresponding to degrees of		raspberries	272
Baumé's hydrometer	479, 480	strawberries	273
Specula, polish for	420	Syrups, concentrated	271
Speedman's pills	193	for flavouring effervescing	
Spices, mixed	289	draughts	271
ragout	289		
savoury	289		
sausage	289		
essences of	291		
Spiced vinegar	294		
Spilsbury's drops	194		
Spirit of nitric ether	432		
savoury spices	291		
soup herbs	291		
Spirits, perfumed	207—213		
Sponge, blanched	432		
to clean	432		
Spruce beer	271		
Squire's elixir	194		
Stains, to remove	433		
Standert's mixture	194		
stomachic candy	194		
Starch	434		
potato, &c.	433		
Steam-pipe cement	325		
Stearic acid	434		
Steedman's soothing powders	194		
Steer's opodeldoc	194		
Stiffening for the hair	247		
		TABLE of weights and measures	459—462
		chemical elements	466
		of solubilities	467—473
		of equivalent numbers and	
		formule of various chemical	
		compounds	474—477
		Tuffetas vesicant	164
		Tamara	289
		Tannin	303
		mouth lotion	268
		Tanjore pills	195
		Tapioca	275
		milk	275
		pudding	275
		Taveare's tooth-cement	263
		Teeth cosmetics	252—260
		stopping for	263—265
		Temperature, effects of	463
		Terpine	436
		Test papers	436
		solutions	437, 438
		Clark's, for water	439

	PAGE		PAGE
Tests, Hahnemann's wine . . .	439	Vanilla, essence of . . .	214
Pettenkofer's for bile . . .	441	tooth-paste . . .	257
picric acid . . .	440, 441	Vanillin . . .	442
Frommer's . . .	439	Varnish, amber . . .	444
Thibaut's balsam . . .	195	Bessemers . . .	446
Tin, solutions of . . .	340	copal . . .	443
mordants . . .	340	cabinet . . .	445
tree . . .	442	Canada . . .	445
Tincture of ambergris . . .	214	etching . . .	334
civet . . .	214	Le Blond's . . .	446
red cabbage . . .	437	Macintosh's . . .	446
Tincture balsam of Peru . . .	214	india rubber . . .	447
galls . . .	438	for hotbeds . . .	447
litmus . . .	438	sealing-wax . . .	447
musk seed . . .	214	hair . . .	447
musk . . .	214	coloured . . .	447
myrrh, odoriferous . . .	258	colourless, for photographs . . .	444
borated . . .	258	glass . . .	448
patchouli . . .	215	Japan . . .	447
of rhodium . . .	214	for gun barrels . . .	447
of vanilla . . .	214	various . . .	442-449
vetiver . . .	214	spirit . . .	443
Tinctures, culinary . . .	290, 291	(lacquers) . . .	444
for the teeth . . .	258-260	essence . . .	445
perfumed . . .	214	oil . . .	445
Tissot's purgative . . .	195	Vaselin . . .	449
Tissue, blistering . . .	164	Vegetable extract for the hair . . .	246
Tobacco water . . .	442	gluten . . .	367
Toilet vinegar . . .	221	parchment . . .	450
Tooth powders . . .	252-256	Venlo's syrup . . .	195
pastes . . .	256, 257	Verditer . . .	416
tinctures . . .	258-260	Vermifuge, Matthieu's . . .	187
Toothache remedies . . .	260-262	Nouffleur's . . .	189
Touch paper . . .	442	Swaim's . . .	195
Tous les mois . . .	274	Vermil powder, Battle's . . .	163
Trade chemicals . . .	298-458	Vichy water . . .	201, 202
Transparent soap . . .	235	salts . . .	202
Traumaticine . . .	442	Victoria perfume . . .	212
Treacle beer . . .	272	Vinaigre de Bully . . .	221
Trees, metallic . . .	442	de Cologne, &c. . .	221
Trophazome . . .	282	de quatre voleurs . . .	221
Trousseau's chalybeate . . .	205	Vinegar (wine, malt, &c.) . . .	450
Turlington's balsam . . .	195	wood . . .	423
Turner's cement . . .	325	curry . . .	292
Turpentiné, oil of . . .	442	camp . . .	292
as a solvent . . .	430	garlic, shallot, &c. . .	291-293
Venice . . .	442	raspberry . . .	273-292
Tyre, essence of . . .	249	spiced . . .	294
UNFERMENTED bread . . .	283	Tarragon, &c. . .	291
Urine, tests for . . .	439-441	carbolic . . .	451
Urn-powder . . .	442	Vinegars, culinary . . .	291, 292
Ultramarine, factitious . . .	418	perfumed . . .	220, 221
VALANGIN's mineral solvent . . .	195	Vinous preparations, Dr Lane's . . .	177
Vance's pills . . .	160	Violet powder . . .	223, 237
		Violet tooth-powder . . .	256
		Violets, essence of . . .	215

	PAGE		PAGE
Vogler's pill for toothache	. 262	Weights and measures, tables	459—462
Vrillière's mouth-wash	. 259	of	. 454
Vulcanite and ebonite	. 451	Welding composition	. 454
WACAKA des Indes	. 279	Wheat, steep for	. 273
Wafers, gelatin	. 451	Whey	. 273
Walnut catsup	. 293	powder	. 273
Warburg's fever tincture	. 196	Whitewash	. 454
Ward's dropsy powder	. 196	Whitelaw's tinct. of lobelia	. 197
essence for headache	. 196	Whitehead's essence of mustard	. 197
paste	. 196	pills	. 197
red pill	. 196	White flux	. 349
sweating powder	. 196	bismuth	. 233
washing powder	. 196	zinc	. 419
white drops	. 196	lead	. 419
white oils	. 84	antimony	. 419
Walker's Jesuits' drops	. 195	Wilson's gout tincture	. 197
Warner's cordial	. 196	lotion for baldness	. 239
Warren's hair dye	. 249	Windsor soap	. 234
Warts, to cure	. 196	Winslow's soothing syrup	. 197
Warwick's powders	. 196	Wyndham's pills	. 184
Wash powder, for the hands	. 232	Wine of shallots	. 292
balls	. 235	quinine	. 169
carbolic	. 247	Wine's, Lane's medicated	. 177
Washes for the hair	246, 247	Wirth's tooth cement	. 264
for vermin in plants	. 451	Wisdom's eye-water	. 197
Washing powders	. 452	Withering's cosmetic	. 228
liquids	. 452	Wright's pearl ointment	. 197
Water, hard, to soften	. 452	Writing fluids, <i>see</i> Inks	. 377
Waters, mineral	198—205	Wood, to bronze	. 455
aerated	. 198	to stain	455, 456
chalybeate	. 204	Worm lozenges, Ching's	. 167
distilled (perfumed)	. 206	Storey's	. 195
saline, &c.	. 199	YEAST	. 457
perfumed	. 206, 207	artificial	. 456
spirituous (perfumed)	207—213,	German	. 457
	219	Ylang-Ylang	. 213
Water-bath, solutions for	. 432	Young's purging drink	. 197
Waterloo sauce	. 293	ZINC, ammonio-chloride	. 457
Waterproofing compounds	. 452	amalgamated	. 457
Wax, to bleach	. 453	oxide	. 458
sealing	425—427	platinized	. 458
modelling	. 453	purified	. 458
Waxed paper	. 400		
Webster's diet drink	. 196		
(Lady) pills	. 197		



NATIONAL LIBRARY OF MEDICINE



NLM 00107175 2