

Selection for sucrose + mutants.

Oct. 26, 1947.

Pupae 10ml. 2% sucrose 1% peptone, ~~at 1% yeast~~ with
 Branched Purple Indicator. broc. as indicated, in series.

	2P27	5P28
A. Y53. 10/26/47. (1) 5P.	st. discoloration funding ±	-
2.	+++	±
3.		
4.		
5.		
6.		
B. 58-161. (1) 10/26/47. 5P.	+++	+++
(1)	-	-
2.		-
3.		
4.		
5.		
6.		

(-)
 1A3 was mov. into 1A4 P30
 and left overnight at room temperature. Later turned alkali. Not suc +

P4. (1A4) was +. Struck out on sucrose EM13. $\left. \begin{matrix} 25^\circ \\ 37^\circ \end{matrix} \right\}$ both Suc -

Y105 - *
 Y106. *

Abandoned Ab.

Selection for d-methyl glucoside mutants.

Oct. 26, 1947.

1% d-methyl glucoside (EK) broth, autoclaved together. BCP media.

growth only seen at 24 hrs

	2P27	5P28.
A. 453. (1) 5P	- ↓	- ↓
B. 58-161 (1) 5P.	- ↓	- ↓

abandoned A 6.

"Transmission" of alkaligenesis.

Oct. 26, 1942.

Glucose + 2% peptone. Vary glucose in methyl-red indicator to establish the critical level. K-12. 5P26. In duplicate.

Glucose: 2% 1% .5% 2% 1%.
Methyl red destroyed.

3P27. Growth + fermentation +. No color changes. Test pH in 1 series.

pH:
acid to MR ++ ++ ++ ±
" " BCP. ++ ++ ++ ±

Use glucose .3%; Brilliant Purple. Prepare series of tubes c 2% peptone. Inc. each with K-12. p28.

11A29. pH B. = ca. 6.8. pH A. < 5.2.

(B) (-glucose).

Add ca 8ml B to A1. pH ca. 6.0.

(A) 1-10.

Retain sample in refrigerator for comparison

to detect alkali production.

(A1) A30. alkali produced. Transmit to A2.
i.e. mix cultures.

(A2) A3. alkali produced. Add to (A3)

(A3) A6. alkali produced. ... Start A4

(4) A10. alkali produced.

end. Expt.

10/25/47.


Swab 453 from 2x25ml / 125 ft. YP broth and from 3x10ml YP agar slants 20 hrs. old. Suspended in 10ml M/10 phosphate buffer at pH. 6.0. Add 20mg HN2 nitrogen mustard [$\text{Cl}(\text{CH}_2)_2\text{NCH}_2\text{Cl} \cdot \text{HCl}$] in 10 ml buffer. Let stand 30 minutes. Sediment cells and replace with peptone 10ml. 1% noc 1ml samples into 25ml YB broth and incubate to prepare inoculum for mutant detection. Squab 1ml as check on survival \rightarrow ca. 10^4 . Estimate.

$PS = \log \frac{10^{10}}{10^5} = 5.$

Many mucoids noted!

Plates	Colonies/plate	Total colonies	Suspected mutants				
10/28/47. 40	250	10,000.	-1 not Mal- -2 (mucoid). -3				
	T1	T4	T6	T7	Lac	Nutr.	
-1 (W1)	S✓	R	S	S	-	T-L-B ₁ ±	∴ mutant!
-2	R✓	R	S	S ²	-		
-3 (W3)	S✓	R	S	S	-	?	Mal - papillae on first streaks.
4105	R	R	R?	R	+		Resolute.
4106	R	R	R?	R plaque	+		
K-12	S	R	R?	S	+		

10/29/47. Squab similar population on galactose EMB plates. 11AM.

(colony dimorphism noted •  S/R?)

Of ca. 40 x 250 = 10,000 colonies, no galactose - were noted. However, one colony which was unusually light purple was noted + picked for further study - 4-4. ∴ Not Mal-.

Oct. 28, 1947. Test the cultures indicated on the following EMB:

	Y53	Y87	58-161	Y53	Y87	58-161	
Lactose	++	18 hrs. ++	++	→	→	→	✓ + + + ^{***} _{peptid!}
Rhamnose	-	-	-	-	-	alk	✓
Inulin	-	-	-	-	-	-	✓ *
Melibiose	+	+	+	+	+	+	✓
α-methyl glucoside	-	-	-	-	-	-	✓
Cellobiose	-	-	-	alk	alk	alk	✓ - ± ± **
Dulcitol	-	-	-	-	-	-	✓
Sorbitol	++	±	+	++	+	+	✓
Sucrose	-	-	-	alk	alk	alk	✓
Ethyl Butyrate	inhibited			→	→	→	produce alk
Lactose	-	-	+	-	-	++	

on CaCO₃ (.1%) peptone agar:

Lactose Minor clearing around lact colonies. Not distinct enough!
 Ethyl Butyrate (add "sterile" to hot medium.)

Cellobiose α-methyl glucoside Sucrose Melibiose

w-1

w-3.

* streaks not uniformly dark: lighter in center. However, this is rather a thin plate.

** streak out the papilla.

P 31. Y53 is Dulcitol negative; Y87 + 58-161 Dulcitol +.

[dulcitol is related to lactose!]

~~w-1~~ w-1. Melibiose slow +
 α-methyl glucoside -
 Sucrose -
 Cellobiose -

Papilla (?) as 58-161 and Y87?

Streak out again

Carbon Source Utilization

T(m) + indicated carbohydrate + BM for 58-161

G = glucose 1/2%
g = glucose .05%

(1/2%)

+ TLB₁ for Y53...

16 hrs. 37°

	O	G	g	Maltose	+ d.m.g.	d.m.g.	+g	+G	Cellulose	+g	+C
W-1	-	++	++	+	+	±	++	++	+	++	++
Y53	-	++	++	±	++	±	++	++	+	++	++
W-1	±	++	++								
Y53	±	++	++								

	-BM	+BM	G	Lac	Suc
58-161	-	-	++	++	±

48 hours:

	O	B	g	M	M+d.m.g.	d.m.g.	+g	+G	C	C+g	C+G	Suc	+G	+g
W-1	-	++	++	±	+	±	++	++	+	++	++	±	++	++
Y53	-	+++	++	++	±	±	++	+	++	++	±	++	++	

	BM	Gl	Lac	Suc
58-161	-	++	++	±
Y87	-	++	+	±

Use .05% in testing sugar utilization in future growth experiments.

P 29. Inoc 58-161 into YP broth. 2 P 30 Harvest and resuspend in 1 ml
 M/10 phosphate pH 6.0. Add 10 mg. to each of two tubes -
MP2.

a. 2:28 PM - 3:02 + 8 min centrif. 34 min.
~~7:48 PM.~~ b. 3:02 - 3:17 + min centrif 15 min.

at 3:02 PM Centrifuge 10 min. ^{dilute peptone} Resuspend in peptone
 Inoc 1 ml each into 25 ml YP. Spread 1 ml samples to
 assay killing.

Killing a) $\gg 10^4$

b) $> 10^4$ survived. $ps < 5$. Squad out +

Retreat survivors also!
 Spread on Yac, Gal. and Mal.

Galactose. 83 plates. Sharp division into large + small colonies
 noted. ca. 300 large + 900 small noted. Noted
 noted on Yac or Malt plates Yac + Malt plates are v. crowded &
 "uniform" colonies - ca. 1000/plate. \therefore are some cells inhibited by
 galactose?? Only large could be scored: 25,000 colonies. Ca 7
 most likely possibilities. Also isolate S^R and S^S colonies +
 test as contaminations? 7:1-7.

Lactose: 37 plates. Ca 800 each plate scoreable = 30,000.
 Almost 1 lac - per plate noted. 12 selected for further study. 7:11-30

Maltose ^(1/1000) 63 plates ca 55,000 colonies scoreable.

8 apparent Mal - noted.

7:41-50.

Total tests: 110,000

Reversion of Lac + Mal in W-1 and W-3.

Streak out papillae of W-1, W-3 and ^{cultures of} 4530 on Lac and on

Nov. 2, 47 Malt. EMB agar. Note that on original plate, W-3 had some papillated, some non-papillated colonies.

W-1. Lac. all Lac-. Papillae in streaks?

Mal. all Mal-. All Mal-.

W-3. Lac All -.

Mal All -!

} Hold for restructing



Restruct A6.

W-1. Reverted for Lac found. Verify and number W-33.

W-3- (Malt) all -; papillae are not Mal+!

W-3- (Lac) + and -. Verify. ✓ 1034

Mustard killing: 58-161.

9

Add 10 mg mustard to 10 ml 58-161, assay = 3×10^9 , in phosph.
M/10 pH 6.0. Dilute 1:100 at intervals and spread (on sucrose EMB which
is available). 0.1 ml. Colony count = survivors/ml $\times 10^{-3}$

0 time assay = 3×10^9

60 mins assay = 2×10^6 . pS = 3+. for 60 mins.

assume 10% survive for each 20 mins. at .1% in phosphate buffer.
Use .2% and 30 mins. treatment

Nov. 2, 1947. S^S S^R

P1 - Inoc small and large colonies of 58-161 on gelatinose ETMB plates of Expt. 7 into YP 25ml. Inc. overnight.

P2. Treat \bar{c} .2% H₂O₂ in phosphate buffer 40 mins. Add 1ml susp. to 25ml YP for further incubation.

Incubate S^S on left (sinistral) side of incubator.

Gal. Inc. 70 pl. x 20 col. → 1400 tests → 12 colo. fragments. inc. all bac +.

ca 50% of colonies are bac - (As inoculum!).

Pick to sl. 10-1. Pick colonies from a Gal plate inc. with the untreated suspension. Pick 3 colonies to slants W30, W31, W32.

Test mutation: DM ✓

Malt. 16 x 40 = 640 tests.

1:1 demultiplex of very dark and less dark colonies. Both are + and have a sheen. (Corresponds to above??)

Gal. 23 x 30 = 700. No mutants. some contaminated

S^S Inc. 63 x 100 = 6300. No mutants.

Malt. 17 x 80 = 1360 No mutants.

Gal. 13 x 50 = 650 No mutants.

Nov. 3, 1947

Characterization of Mutants of Exp. 7.

A. Galactose mutants - 1st. streak on galactose EMB.

- 7-1. Majority of light purple colonies \odot with a few typical Galactose +. Pick to glucose slants. BM W-2.
- 7-2. Gal \pm as in 7-1. No Gal + BM W-4
- 7-3 do. BM W-5
- 7-4. Majority of allelic, v. small colonies. In agar of heavy streak, a few Gal +, larger colonies. Test on bac: bac - allelic. throw out. BM W-6
- 7-5 As 7-1. BM W-6
- 7-6. Do. BM W-7.
- 7-7. As 7-2. BM. W-8.

B. Lactose mutants.

- 7-11. Typical bac - . ^{n.g. in 16h.} not B-M-! W-9
- 7-12. Do. Papilloquin W-10
- 7-13 Do. Pap. W-11
- 7-14. Do. + two lac + colonies. W-12
- 7-15 Do. W-13
- 7-16. Do. W-14
- 7-17 Do. W-15
- 7-18 Do. Colonies smallest. lac + in heavy streak W-16
- 7-19 Do. Weak utilization? W-17
- 7-20 Do. W-18.

7-21. Do.
7-22 do.
7-MALTOSE

what is the mutation of these creatures?

w-19
w-27

7-41.	Majority are light purple. Some Mal+.	BM	w-20
7-42.	Mal+.		
7-43.	As 41.	BM	w-21
7-44.	all Mal+.	BM	w-22
7-45	do.	BM.	w-23
7-46.	do.	BM	w-24
7-47	do.	BM	w-25
7-48	do.	BM.	w-26.

Test units. by comparison of $T(B+M) \bar{c} T(o)$.

58-161. $\frac{S^R}{S^S}$ w-28
w-29

Galactose Inhibition

Nov. 3. Demargination on galactose was noted in 7.

$G^S + G^R$ were streaked out on EMB.

	G^S	G^R
lac	++	++
Gal	++ only d. smaller than.	++
Mal	++	++
Lac.	± (slow)	± (slow).

The galactose effect was not reproduced here, as in the plates streaked from cultivated G^S, G^R in expt. 10. These were, however, a different batch. Test on:

galactose 1%	peptone 1%
galactose 1%	Neamine 1%
galactose 1%	N_2 Tone 1%.

Dulcitol.

Nov. 3 '47.

K-12 ferments dulcitol only weakly. Inc into broth
+ compare. In 16 hours, dulcitol broth is vigorously fermented
by 58-161.

A. from broth

B. from plate (a slow colony).

} Both form only "weak" colonies.

Streak A. again. - slow + as before!

?? Reductions of Methylene Blue??

Nov. 4, 1947.

EMB - Gal 1% - agar 1 1/2%

- A. Peptone 1%
- B. " 0.3%
- C. N-2-Amino "B" 1%
- D. N-2-Tare 1%.

Streak out $G^R + G^S$ colonies. P4.

A5.

- A No dimorphisms
- B large + small colonies. Not as marked as C.
- C large + small colonies.
- D. no dimorphisms.

4
Nov. 24, 1947.

Mutant Rem. Treat a single colony culture of
58-161. E HN2 .2% for A) 5 minis. B) for 30
Lac mutant plating. Lo. W-1 for galactose mutant plating.
Incubate 24 hours before plating per cocurrent technique

A. 58-161 (treated) - 50 pl x 70 cols. on lac = 3500 tests

No mutants

B. W-1.

35 pl x 100 cols on Gal = 3500 tests

No mutants.

Reduced sugar utilization.

4
Nov. ~~18~~, 1947.

See 14A. for Treatments.

Add ~~plate~~ ca 10^{10-9} cells (survivors of mustard) to:

		A5	A6	A7	A9
58-161 :	1. 100 ml T(m) + (B.M.)	±	'		
	2. 100 ml T(m) + glucose .05%	+++			
	3. do. cellobiose	±			
	4. do sucrose	±			
	5. do sucrose 200ml.	±			
	6. do. sucrose 200ml.	±			
	7. do. 1-methyl glucoside.	±			
w-1.	8. do. sucrose TLB, 300ml.	±			

-No change

[Handwritten mark]

ditto.

abandoned experiment on Nov. 9.

Nov. 5, 1947.

U.-V. Killing Rate.

(10^8 cells)

Hanovia lamp. 6" from plates. Spread 1 ml of grown cultures of 58-161 on EMB lactose plates. Irradiate as indicated.

t.	colonies
0	$> 10^5$
5 sec.	$> 10^3$
10 sec.	112
15 sec.	ca 100.
20 sec	50
30 sec	28
1 M.	20
2 M	7
6 M	3.

There is certainly a break in the killing curve between 5 and 10 seconds, or else, the survivors of higher doses represent large clumps.

10 sec. is a convenient irradiation time which has a $pd = \log 10^8 / 10^2 = 6$.

One "weird" vac colony noted on 10 second plate.

Streak + compare with subs.

β -methyl glucoside

Nov. 4, 1947.

Sample from H A Laidy. (mp 106-108°)

1) 1% in EMB.	W-1	—	A6.
	Y53	—	dark spot - ketone. →
	Y10	—	all —.
	58-161	—	
	W-22	—	

PS

2) .05% as C-source in T(m) + BM. 58-161 mol. PS.

	A6.	A7	A10.
1) β -methylgl.	—		+
2) α -methylgl.	—		—
3) $\alpha + \beta$	—	<i>auto</i>	+
4) glucose	++		++
5) β +glucose	++		+++

Nov. 6, 1947.

W-30 represents a series of Lac - colonies which constituted 50% of the colonies found upon spreading mustard-treated 58-161 on lactose EMB. 12 colonies derived from the untreated inoculum were all Lac+. The inoculum was derived from a single large colony (GR) of a type constituting about 1/4 the colonies found on a galactose - N2 Annie B-EMB plate (see. Expt. 7) In Expt. 10, the ratio of

Lac+ : Lac - was :


	+	-
plate 1.	18	10
plate 2.	10	4
plate 3.	8	6
plate 4	12	16
	48	36

A large fraction of both these types was found on each of the 70 plates of the experiment. (10)

- A). Pick 5 colonies from plate Expt. 7 in attempt to reproduce the expt.
- B). Pick + streak out 5 ^{Lac+} colonies from expt. 10 to determine stability of these Lac+ : all plates show all Lac+
- C). Set up selection Expts. between these Lac+ and Lac - in broth.

On the Lac plates, W-30 forms broad flat colonies. These seem to engulf Lac+ colonies which they may contact.

On maltose two colony types are seen. Initially, they differ only in that some are less intensely colored on EMB than the others. Later they develop large lobes of gummy material which projects from the surface.



On galactose no peculiarities had been noted.

C. Selection: Expt.

A. Broc Lac+ colonies 2P6 into YP broth 25ml.

B. Do. + 1 loopful of Lac- (W30) suspension.

Streak out duplicate lac plates for initial assay.

A: all colonies lac+ (Lac- < 1/100)

B: ditto.

		P7.	
C. Streak out A	A 6:	all+	all+
D. B	A 6:	all+	all+
A2 Streak out P7.	A10	all+	all+
B2 " "		all+	all+

The combination: W-30 and the population of Lac+ found on these plates does not seem to satisfy selective hypothesis. The Lac+ may be resistant to the (hypothetical) inhibitor. 58-161 (18-A) should be used instead.

D. Selection: Expt. 11/10/47. Mix 1ml broth culture of 18A1 with .05ml similar suspension of W-30 from 5 lamb in YP broth. Streak out initially, etc.

①. 18D-A1	Streak 18A1 on EMB-Lac	A 11	all+ all+
- B1	18A1 + W-30 on EMB-Lac.	A 12	ca 100:1, +: -
N12	A2 ✓		all+ all+
A12	B2 -		ca 20:1, +: -
	B3		ca 20:1 +: -

⤴ This selection expt. does not explain original findings.

Nov. 6, 1947.


Spread .1 ml 58-161 culture on EMBS-Malt and Lac plates.
 Irradiate each for 10 secs. \bar{E} Dr. Smith's Hanovia lamp.
 ca. 120 / plate

Lac (10) Nothing noted on 9 plates. See below for 10th. *

Malt (10) No mutants or sectors noted.

~~No mutants or sectors noted.~~

* A colony was found of the following appearance.



This suggests further a delayed effect of the mutagenic agent.

Picks from each of the 4 distinguishable zones to YP Broth + incubate. When grown spread on ~~and streak out.~~

1	W-35	W-38	38
2	W-36	W-35	35
3	W-37	W-37	37
4	W-38	W-36	36

Lac EMBS.

(A10).

50:

1. Ca 50:1 lac+ : lac-. No evidence

of sectoring of the +.

2. Ca 50:1 lac- : lac+ . No sectoring

3. Ca 100:1 lac+ : lac-. No sectoring.

4. All lac-.

Streak out and transfer to slants.

Nov. 6, 1947.

P3. Streak out Y53 on EMB-lactose - 3/10% peptone.

A6. Papillae well developed. Pick papillae to small H₂O and streak out.

#P8. All of 20 plates pure Lac⁺ exc. for 1 which was streaked from a mixed colony. ∴ Lac⁺ is a stable reversion in most cases.

Nov. 8, 1947.


Squad 10^8 cells of a single cell culture of 58-161 from 4P on maltose + lactose EMB plates. Irradiate each plate ca 10 secs. \bar{c} Hanovia u.v lamp at 6". Score at 36 h.

Maltose. 24 pl x 60 cols. = 1440 colonies tested. All colonies +.

Lactose 59 pl. x ca. 150 scoreable colonies = 9000 tested.


12 plates showed 14 likely colonies. Some of these are clearly simply mucoid.


①, ② Mucoid.



- 4 clearly sectored
- 3 not sectored
- 5 mucoid
- 2 ??

4. Small white colony ✓

5. a  c Streak out 3 compe. b


6 M 7.  ✓

8.  ✓ 9. M

10.  ✓

11. a  ✓ b

12.  → no lac. in streaking

13.  ✓  b. all $\frac{1}{2}$ mucoid.

14. 0 mucoid.

3a 1:1 +: - (+) W50
 3b almost pure(-) W48
 3c 1:1 \neq and - (-) W49

4. pure - . W39.

5. a. 20:1 $lac^+ : Lac^-$ Purify + : W41.
 b. 2:1 +: - \rightarrow
 c. pure - . W40.

7. pure - W42. (slight utilization?) ✓

8. 1:1 +: - The lac^- is W43.

10. a. 10:1 +: - +, W44
 b. pure - - W45

11. a. 2:1 +: - (+) W46

b. pure - W47.

12. all +. Not mutant.

Nov. 10, 1947.

Compare 58-161 and 17-1.

	glucose	β -Me Gluc.	cellobiose	
(maltose) 17-4.	\pm $+++$	- -	+ \pm	- -
(m β -methyl) 17-1.	++ $+++$	- \pm	\pm \pm	- -
(Maltose Neg.) W-20.	\pm $+++$	- -	- -	- -

def. diffusive
maltose content?

24 hrs., 36 h.

A15. do. only glucose is $+++$. others from - to \pm .

\therefore 17-1 is only slow utilization.

5 A16: P17

	β -Me Gluc.	Cellobiose.
17-41	\pm \pm ✓	\pm ✓
17-4	\pm ✓	\pm ✓
W-20.	\pm ✓	\pm ✓
17-1A.	\pm +	\pm

Transfer 17-~~4~~ to similar series. \rightarrow much slower m β -methyl than α glucose.

Nov. 14, 1947.

Irradiate 50 ml of 24 hr. ~~10-37~~ 10-37 conc. to 10 ml. with washing. 3 ml / 25 ml quantity flask. 40 secs. at 6 in. from Hanovia Lamp., with manual twisting. Spread 1 ml on EMB.

30 Lac, 20 Dal plates. All too dense (ca. $10^5 - 10^6$ / plate).
(Autoabsorption, rather, masked!)

1 ml samples irradiated 5 secs. in vitro synthetic minimal.

- glucose

sucrose

α-methyl glucoside

cellobiose

} all neg. after 4 days.

Attempts at induced utilization
of sucrose & other sugars.

Dec. 10, 1947.

Purple suspensions, ca 10¹⁰/ml, of Y10. Irradiate 3ml at a time
5 secs. in quartz flask rotated at hood of Hanovia uv lamp. 1/2 ml
inocula into 50 ml / 125 ml flasks containing (1m) + sugars
at .05% (except. sucrose .15%) ± glucose .005% (g.)
ca. 4P 10.

	9A11.	2P11.	P611	9A12	P13	P16.
1. Sucrose	±	±				
2. "	±					
3. " g.	++		- + Not	* Not.		⊕
4. " g.	++					✓
5. Ref.	±					
6. Ref.	±				* Not.	
7. " g.	++			++		
8. " g.	++±	* Not		+++		⊕
9. Cellulose	±					
10. " g.	++		- * Not	* Not.		do. w55.) ⊕
11. d. He gluc	-		* Not.			✓
12. " g.	+					✓
13. Trehalose	+++					
14. "	+++					
15. Glucose	+++	+++				✓
16. g.	+	+				✓
17. --	±		- berline.			

Experiment terminated 12/24, without the recovery of any plus-variations in this series.
Y55 (Salicin-plus mutant) should be tried on the beta-glucosides.

* Strals out on corresponding medium.

[.005% glucose is apparently an excessive "boost". Use .001%.] Also these sugars were
evidently somewhat broken up by this prolonged autoirradiation.