

Fermentation tests.
Mutants.

434

Spread out mustard-treated 440 (see 426) on
EMB-lactose + a few maltose, sucrose, glycerol plates
and examine for mutants ca 400 per plate.

bacteria: 36. # of plates. (mucoid found)

∴ ca 15,000 colonies examined.

no fermentation mutants.

~~Haltose~~
M. amital 2 plates + 2 streak plates.
ca 1000 examined.

2 very small M- colonies noted. streak out on fresh medium.
all M+.

Glycerol - slow utilization. - compare passage on glycerol
& peptone.

Sucrose - very slow, but definite utilization

Fermentation contents - enrichment cultures.

435

Feb. 25, 1947.

Incoc 50 ml NB + 2% sugar + Bromoresol Purple A25. Y53.

	A26	A27	A3	A4	A5	symbols	acid.	acid.
glucose	+++							
glycerol	+	+++	Y81.					growth heavier than in glucose
lactose	-	-	+	++	+	+	+	no % black on EMB.
sucrose	-	-	-	-	-	-	-	* growth mic. on standing at room temp.

Streak out some apparently gly^(a)- colonies from 434 +

compare with atypical gly^(b) ±. — (b).

- a) Y84. — ~~Y80.~~ Y80.
- b) +

Compare a, b + gly+ enrichment culture above.

	EMB
Y80	-
(from glucose above) Y53 gly ±	+
Y53 gly +	+ or ++

1/2 evidence of papillae.

different from Y53?

on BCP —, medium is not changed in color, cells show slightly different shades (+, ± in pinkish, — more translucent or violet.)

on gly-BCP broth — gly-, gly ± and gly+ all show slow acid + gas.

(see over)

1. Enrichment for gly+

A5 - streak from gly tubes to new gly EMBA

A8 - scoring OK, as before!

Resistance mutants - cross test.

436

Streak out susp. of

in MAgar + MalZum. 10^{-4} .

Streptomycin 5 u/ml

	M.G.	St.	B.G.
Y77	++*	-	-!
Y78	-	++	

* dye is decolorized.

appeared on the streak.

after 3 days, several hundred colonies
Y79.

streak Y77 over Y78/M.G. to determine if decolorization reverse
effect of dye. No evidence of stimulation of previously streaked

(Y79³) culture. Probably due to pH change.

Inversion Detection

437

See 437a for summary.


1 MAR 1961

Streak out Y40, Y53 ^(T from 426) on NA plates.

P1.

Pick single colony and streak on NA, overlapping a streak of the other type in the center of the plate, and mix well in center.

This can then be picked after growth and inoculated onto minimal.



P3.

Test combinations by suspending growth from center in ^{1 ml} water, and streaking on minimal plates. (i.e. 84 tests!) ca 10⁹ microbium.

Tests: log prototrophs

1	1+	13	1+	26	2
2	2	14	1+	27	1+
3	2	15	1+	28	2
4	1	16	1+	29	2
5	1.5	17	1+	30	1
6	1.5	18	1+	31	1+
7	2	19	1	32	1+
8	1	20	1.5	33	1
9	1+	21	1+	34	1
10	1+	22	1+	35	1
11	1	23	1+	36	1
12	1	24	1+	37	1
		25	2	38	1

39	1
40	1
41	1
42	1

1-6 also tested by streaking (hopeful on T(0) plates).
colonies

1	0
2	3
3	1
4	2
5	7
6	10
7	1
8	1

84 tests - no inversions

Streak technique not as reliable as desired!

Inversion Tests : Summary.

437A.

exp.	material.	tests:		Cumul. Yield Tests.
426	MN ₂ fit.	Y40 x Y53T; Y53 x Y40T, by st. tech., 0 + 0,	20 tests.	0
433.	MN ₂ treat.	do.	17	37. 0
437.	"	Y40T x Y53T. in only, X mag; 2 x 42 =	84	121 0
508	X-ray	Y40T x Y53T. in only. mag. 2 x 14 =	28.	149 0

Trace of prototroph initiation.

438

Pour 440x453 plates in T(0). To sup. add also 10^{-6} R-12 cultured + washed similarly in order to compare rates of colony development.

See 445.

Y65, } test for inversions
Y68 }

439.

Plate Y65 x Y40

Y68 x Y68.

in $T(0) + T(B_1)$.

no prototrophia shell!

\therefore not due to suppression of X in B_1 - B_4 region.

Segregation of Gly-

March 4, 1947 -

Y80 x Y81. cf. 435.

$B_1^- \times B_1^+$ as usual. Pick & plate on EMB lac + EMB gly agar.

Lac-R	Lac-S	Lac+ R	Lac+ S				
				Gly-	Lac-R - S	+ R	+ S.
				Gly+			

Lac, V
segregation.

March 6, 1947

Lac-V^R Lac-V^S Lac+V^R Lac+V^S

B₁⁺ ~~15~~ 15 ~~9~~ 9 0 1 25 Faulty segregation of Lac+?

all Gly+.

B₁⁻
Gly- 2 19 5 0

Gly+ 1 (1?)

This scoring of Gly, done at 2 da. is not borne out by readings at 3 days, when in both groups, all were Gly+, ¹⁵ c some variation in intensity.

Note: 24 Lac- : 1 Lac+ in B₁⁺ (95%!!)

46 Lac- : 5 Lac+ in B₁⁻.

1) B₁⁻/B₁⁺

25	24	2	1	25	χ ² = .04 .02 .04 .03 <u>.13</u>
41	40	4	5	45	
	64	6		70	

compare standard.

18 24 7 1 25.
51- 64 19+ 6 70

$$\chi^2 = \frac{36}{18} + \frac{36}{7} = 7$$

$$\chi^2 = \frac{13^2}{51} + \frac{13^2}{19} = \frac{3.3}{8.9} = 12.2$$

- a) All data: 72% lac-
- b) new data 64% lac-

efficiency of Lac+

Peculiar segregation may explain peculiarities of Gly segregation.

Consider:



if vac. eq. is disturbed

Segregation of drug-resistance

- A. ~~Y79 x Y80~~
Y77 x Y78. - in T(0) OK! Streptomycin 5 u/ml
Molacitate St. 100 u/ml
- B. Y53 x Y78 off! #6 Lac- 1 Lac+.
no streaks just 12h
- C. Y40 x Y77 OK! no #

A. Y77 x Y78.

	Lac		V	str	M.G.		Lac		str	M.G.
	+		R	↓	R		-		↓	R
	+		R		R		-			R
	+		S		R		-			S
# peculiar dup blue color	+		S		R?		-			S
	+		R		R?		-			R
	+		R		R?		+			R
	+		R		R		+			R
	-		R		R		+			R?
	-		R		R		-			R?
	-		R		R		-			R.

? probably S.

scoring uncertain
due to selection of
resistant residue

do.

C.

	Lac+MR	Lac+Ms	Lac-MR	Lac Ms.	
	2	1	8 8	2	
	0	2	16	3	<u>scoring not certain</u>

total.

Lac+S x 2 3 24 5 ind. cont.

Lac-R.

no. of resistants. valid, indicates linkage to B4.

should score on minimal plates to avoid selection for resistant contaminants.

Selection of recombinants with drugs.

442

P3. - mix YB cultures of 477 (Mg^r) and 478 (Sti_5^r). 30°
 also use mixture of anti. plating technique of 441.

- | | | |
|------------------------------|-----------|--|
| 1. Mix culture in $Mg + Sti$ | A6
0,0 | A8.
1 colony; |
| 2 477 | " | 2 colonies; ^{also} as 3.; 4 cols. |
| 3 478 | " | 1 colony
ca 100 minute clumps, ^{mini} cols. |
| 4 477+478 | " | ca 10 v. small A - as 3.
B > 10 large colonies
innumerable minute clumps |

A6 - incubate at 35° .

Indeterminate whether the mult. resistant colonies represent recombinations.

Compare 4 (cross) \bar{c} $\underline{=}$ 2 mutations of $Sm^S Mg^R$ to $Sm^R Mg^R$

There may be some synergism in view of the long lag before colonies are detectable!

[Use 484 x 479. and plate in brilliant green.
 + streptomycin.]

March 4, 1947.

Recd. 127,000 u. streptomycin from Woodruff, 9 units.
non-streak ampules

Suspend in 2.7 ml 95% alcohol for 3 hours. Add
10.3 ml sterile H₂O → ca 10,000 ^u/ml in 20%

alcohol. Dilute further as required.

u/100 ml = 1 ml
100 u 1
478 T
453 T

500 5
478 T
453 1-200

streak 478 on 5u agar for -
482 - streak out on 5u agar

1000 10
478 50
453 1

483 - streak on 10u agar
484 - do.

10,000 100
478 0
453 0

478 Streptomycin
on 5u agar gives colonies but not so large
as 482.

483, 484 OK on 10u agar.

Segregation of B⁻, etc.

445

See 452.

Y40 x Y53. Mix cells in agar pour 5 ml each plate.

A) O:
 11
 15
 9
 10
 15

 60
 m = 12.0

B: 12
 11
 14
 12
 9
 13
 6
 12

 m = 11.1

B₁: 124
 136

 m = ~~10~~
 130
 more turbid.

B) 20 B tested: 18 B⁺
 2 B⁻ (Lac⁻R; Lac⁺R)
~~1 R~~

5/20	364
2/20	445
5/50	452

12/90	13%

C. Seg. of Lac, U^R in O, B₁:

O: ~~-R -S +R +S~~
~~24 ~~11~~ 9 ~~1~~ 0~~
 B₁: ~~22 ~~15~~ 14 0~~

Σ	O	-R	-S	+R	+S
44		17	10	9	0
	B ₁	16	12	13	0
51		12 11 1			

On one O plate, streak out 11-12 on surface. These colonies appeared at same time as prototrophs (24 hours) and were of comparable size.

Lac - V conjugation.

March 6, 1947.

Use colonies from Expt. 437 (grow together - chance for selection) [140 x 153 (T) by plate number].

Plate # lac - R lac + R lac - S lac + S. \bar{x} * V scoring not

9	10	2	5	1	0	8	dependable -
20		4	4	2	0	10	phage apparently
* 27	40-50	9	6	0	0	15 } very low titer.	not 8 by itself
* 22	40-50	8	10	0	0		
* 3		8	3	0	0		
8		3	1	0	0	4	
3		7	3	3	0	13	
42		1	2	1	0	4	
6		6	2	2	0	10	
4		4	2	0	0	6	lac - = 135/214
14	10	6	1	0	1	8	= 63%
40		1	1	2	0	4	compare \bar{x} 70% prev.
21		2	1	1	0	4	$V_R = 190/214$
7	10	6	3	2	1	12	= 85%
23		3	2	0	0	5	comp. \bar{x} .74 prev.
2		13	7	1	0	21	
25		5	14	2	0	21	
11		1	3	0	0	4	
10		11	1	3	0	15	
19		1	1	1	0	3	
17		5	1	0	0	6	
29		7	4	1	0	12	
		113	77	22	2	214	

The agreement of the lac + R; lac - S classes with exp from previous expts is very poor. Reexamine crosses of aberrant cultures.

There is a shift from lac - S to lac + R

in map basis

β_i	-	+		R			+
	1	1		1			1
	OM	Lac		V			TL
	+ a	-	b	S	c		-

$\langle lac - \rangle$ bvc $\langle \rangle$ a

$\langle v^R \rangle$ bva $\langle \rangle$ c

(Lc. more than b/c)

a = +R
b = -R
c = -S.

[-R would not be augmented by the
diminution of c?]

or, another interpretation, is that ~~the~~

the previous studies were revisited for T or for L, \bar{c} a larger distance
in the interval v-(T or L)

\therefore compare types \bar{c} a short or long interval
(i.e. low or high - S) biochemically.

total. 146.

#	lac-R	lac+R	lac-S	lac+S.	\bar{E}
15.	3	9	1	1	14
33	2	1	2	0	5
34	5	0	0	0	5
30	6	2	2	0	10
31	7	7	4	0	18
24	3	3	1	0	7
32	7	3	1	0	11
35	7	2	1	0	10
1	6	4	4	0	14
36	4	2	2	0	8
16	12	10	3	0	25
26	14	9	3	0.	26

$$83-76. \quad 52^{53} \quad 22^{20} \quad 1 \quad | \quad 151.$$

$$106-113 \quad 77^{76} \quad 22^{21} \quad 2 \quad | \quad 214$$

$$189 \quad 129 \quad 44 \quad 3 \quad | \quad 365$$

$$\chi^2_2 = \frac{49}{83} + \frac{49}{106} + \frac{1}{53} + \frac{1}{76} + \frac{9}{20} + \frac{9}{27} = 1.86 \quad p = .14.$$

- .59
- .46
- .02
- .01
- .45
- .33

cf. 445.
compare \bar{E} remainders
freqt.

These samples agree.

Homogeneity??

analysis of
446 vs 359 summarized.

-R	+R	-S	+S	Σ
189 ¹⁴	129 ¹¹⁷	44 ⁶³	3	365
100 ¹⁰⁵	55 ⁴⁷	50 ³⁶	4	209
289	184	94	7	574

$$\chi^2 = \frac{25}{105} + \frac{25}{184} + \frac{144}{67} + \frac{144}{117} + \frac{196}{36} + \frac{126}{63} + \frac{1}{4} + \frac{1}{3} \dots$$

= .2

.1

2.2

1.2

5.4

3.1

8.5 by this component. $p = .005$.

$$\begin{array}{r} 6.2 \\ 6.3 \\ \hline 12.7 \end{array}$$

12.7 = χ^2_3

8.5

4.2 = χ^2_2 $p = .04$

it is the difference in the frequency of -s which differentiates the distributions.

compare

194 - 189	129 ¹²⁴	318
95 - 100	55 ⁶⁰	155
289	184	473

should be 194:124
95:60

$$\chi^2_1 = \frac{25}{189} + .25 + \frac{25}{124} + \frac{25}{60} = 4.2$$

Hurdity of segregation types.

447.

March 7, 1947

Recover 446 - 22, 25 + 27. in order to ascertain suitability of dispropotionis in ratios. Compare segregation of lac + V = 440, 453 stand. in previous test:

A 22:- 8:10:0:0
 B 28:-25 9:6:0:0
 C 25:-27 5:14:2:0
 D. 440 x 453. lac, V.

	-R	+R	-S	+S
A	47	19	9	1
	44	20	7	1
B	26	12	16	0
		13	16	0
C	7	5	2	1
	80	36	32	2
D.	30 ✓	21 ✓	19	0 ✓
O ₂	10	7	5	0 ✓
N ₂	6	5	6	0 ✓
M.B.	4	4	6	0 ✓
S.H.	2	1	2	0 ✓
D. total:	52.	38	39	0

This is homogeneous with the previous tabulation: 100:55:50:4.

$$\chi^2 = \frac{100}{37} + \frac{100}{62} = 4.3$$

$$+ \frac{4}{21} + \frac{4}{36} = .28$$

$$+ \frac{64}{18} + \frac{64}{31} = 5.6$$

$$\chi^2 = 10.1$$

$$p = .007$$

Compare A E D..

47	19	9 + 1	0	76
52	38	39	0	129
99	57	49		205

See over.

Total 132 74 71 2 279.

Compare $\bar{A} \bar{c} S.$

58	66	10 ¹⁸	76
98	90	39 ³¹	129
156		49	205

$$\chi^2 = 64 \left(\frac{1}{58} + \frac{1}{98} + \frac{1}{18} + \frac{1}{31} \right)$$

$$= 64 (.017 \quad .010 \quad .055 \quad .032)$$

$$= 64 (.114)$$

$$= 7.3 \quad p = .007.$$

Therefore the total discrepancy is due to a difference in the proportion of V^R cultures. In practice, this means a deficiency of 5 cultures in the new group.

-R +R -S +S.

A. 22

~~37~~
~~44~~ 20 7 1

This is the "abnormal" distribution

B. 25

25 13 16 0

1 plate showed 13:14:0:0.

C. 27

7 5 2 1

(perhaps scoring of V^R was not proper.) Recheck! ✓

~~76~~ 38 25 2

D.

-	30	22	19	0
O ₂	10	7	5	0
N ₂	2	1	2	0
SH	6	5	6	1
MB	4	4	5	1
Σ .	52	39	37	2
	69	38	31	2

447

Plate by plate

A	6	2	0	0
	6	14	6	0
	8	1	3	0
B	17	3	4	1
	4	1	6	0
	12	10	8	1
	9	2	2	0
C	7	5	2	1

compare +, -

89	41	130
100	40	140
189	81	270

Caution! Be careful of scoring V^S .

Interpretation of 447.

448

Compare $T^- \rightarrow T^+$ in 447 lines + standard.

March 7, 1947

Using cells from exp. 447, plate the Y53 components at a 10^{-2} dilution into LB₁ agar.

Y53	turbid	10 cols
(Y53-427)	less	30 cols
25		10 "
22		1-2

There is no stable differentiation, not apparently due to scoring difficulties.

Plating medium for recombination

March 7, 1947

Factors: buffer: phosphate, citrate, acetate, phosphate. More alkaline.

RH: thioglycollate; ascorbic ac.; pyruvate-lactate; O₂; Methylene Blue. hydrolyzed??

B₁ ? - Try on BM x TP.

T(0) - V40 x Y53 mix in agar + pour on pre-poured plates.

1. Controls

2. "O₂ atmosphere" least turbid plates

3. N₂ atmosphere turbid plates; colonies v. small, countable merese me.

4. Na Thioglycollate 2mg -
.2mg -

5. Ascorbic ac. 1% no colonies (pH 4-5) No turbidity

6. Methylene blue 100v Turb. decreased
10v " "

The number of apparent colonies is correlated inversely with the gross turbidity, and was least in O₂, most in N₂. SH had no apparent effect, however. There is, however, no marked increase of colonies in the O₂ atmosphere plates, but those which do appear are larger.

see 441. Test Y78. mEMB; Lactose. cf. 453.

- also 441A - lactose - on lac → glycerol white + black colonies +
 Y78 (58-161 Sm^R) ^{lac} +++ gds. ^{gly.} +++ - ^{some blue halos} - gum sheen!
 58-161 - (±??)
 Y80 (58-161-V₁^R-gly-) -

Note, 480 on v. long incubation times a faint violet color, à la sucrose
 from previous expts + now

Y53	++	Y80	-
Y81	++	58-161	±
Y40	++		
Y78	++		

? relative amts.

58-161 is slow, but will score vs. Y80. see 453.

Y80 x Y81.
Sly. segregation.

March 9, 1947.



See 451A.

A. Repeat 440
Y80 x Y81.

T(O).	16/16 Sly+	12/14 Lac-	86%
T(B ₁)	13/13 Sly+	58/72 Lac-	78%

~~SE~~ Note, this denotes absolute linkage of Sly to B₁. This is predictable either on the mapping:

written Y80 x Y53
not Y81 x Y40
seems to be disturbed!
request!



Note (also) the segregation for Lac may be disturbed.
data for Y40 x Y53: (71% Lac-) $p = .06$

Comparing \bar{e} occur.

\bar{e} 440 $p < .025$

(This + 440) \bar{e} 71% $p = < .001$

440 \bar{e} 71% $p = .008$

B₁⁺/B₁⁻ $p = .3$

Mauch, 9, 1947.

These plates were allowed to stand 3 days before testing

B. Y80 x Y53. Test B_1^+ & ($B_1^- + B_1^+$) on Gly; Lac.

a. B_1^+ 27 Lac- : 9 Lac+ 35/35 Gly+

b. B_1^- 50 Lac- : 16 Lac+ *scouring uncutani*. 1++ ; 30+ / 31

77 Lac- : 25 Lac+ ca 75% Lac-

C. ~~Y80~~ Y81 x Y40.

Test on Lac

Gly 16/16 +
(not work)

B_1^+ 104 Lac- : 43 Lac+

B_1^- 28 : 13

132 : 56 188

70% Lac-

This experiment is not homogeneous with the carrier Y40 x Y53. mes.

D. Y53 x Y40.

-R -S +R +S
33 32 28 25 14 18 0

B_1^+
 B_1^-
18 19 12 15 13 10 1

75 → $\chi^2 = 3.2$
 $p = .2$ data not adequate
44 → $\chi^2 = 3.5$ $p = .2$

51. 40 27 1 119 of standard. $p = .06$

91 : 28 76% Lac-

Segregation of B-

452

March 9, 1947.

See 445

440 x 453 in B₂ medium. moi. helpful in T(10) + T(B).

Pick colonies to H₂O, + 50 isolates.

50 isolates. At 12 hours, 5 def. B⁻

β did not grow in either. 5 at first.

lac, V eye:

	lac	V
1	+	R
2	+	R
3	-	R
4	-	R
5	-	R.

P13 - 5 grew on B₂ not O.

Ancest, 1-4 contained primarily -S, E + R as well. 5 was uniformly -R. see 456.

To summarize B⁻

~~lac - R - S + R + S.~~

-R +R -S +S.

HHH | HHH 0 1

∴ 0-S/12.

	6	5	0	1	12
A (-)	6	3	3	.36	
B (M-B-)	1.4	9	.8	1.4	


$$\chi^2 = \frac{4}{3} + \frac{9}{3} = 4.3$$

p = .14

$$\chi^2 = 5 + \frac{16}{9} +$$

$$= 7 + .03$$

mapis



Glycol reactions, various strains.

453.

March 11, 1947.

P11. Strains on a single glycol plate (EMB-2%) the following:

		A12	P12	A13	4P13	A14
1	K-12	-	+	++	+++	+++
2	58-161	-	-	-	- + strands.	
3	Y40	-	-	±	± + strands.	
4	Y10	-	-	±	+	
5	Y53	-	-	+	++	
6	Y46	-	-	+	++	
7	Y64	-	-	±	++	
8	Y78	-	±	++	++	
9	Y77	-	-	±	+++	
! 10	Y80	-	±	±	± translucent, creamy shade, not opaque purple.	
11	Y81	-	-	+	++ ++	
12	Y82	-	-	±	±	
13	Y73	-	-	+	+	
14	Y74	-	-	+	+	
15	Y82	-	-	+	++	
16	Y83	+	++	+++	+++	
17	Y84	-	-	+	++	
18	Y79	-	-	+	++.	

- resistance of bacteria

4/11/47

On various plates, streaks: *Serratia marcescens* 1
Salmonella 20 2
 " 21 3
Phytomyces tumefaciens 4
Staph aureus 5

incubate

vor u. f. ml.

Malachite Green	M.G.	1	++	✓	++	✓	-	++	✓	-	±	++	-	✓
	10	++	✓	++	✓	++	✓	-	-	✓	-	-	-	✓
	50	-	-	-	-	-	±	-	-	-	-	-	-	-
	100	-	-	(- cols)	±	±	++	✓	-	-	-	-	-	truy cols.
Penicillin Perm.	P.G.	100	-	-	✓	-	✓	-	✓	-	✓	-	✓	✓

Streptomycin Sth	Sth	1	++ ^R	✓	+	✓	+	✓	+	++	✓	++	✓	✓
	5	++	✓	cols.	-	-	±	±	pepilles	±	++	±	✓	OK.
	10	-	✓	good cols.	-	-	-	-	✓	±	+	✓	-	-

Streptomycin Sm	Sm	1	-	✓	+	+	+	+	±	+	✓	-	✓	grow large.
	5	-	-	-	-	-	-	-	±	-	-	-	✓	✓

Penicillin si ¹	10	++	✓	-	✓	-	✓	-	✓	-	✓	-	✓	✓
Penicillin si ²	100	++	✓ ^R	++	✓	++	-	±	+	++	✓ ^Y	✓	✓	✓

Control N.A.	++	++ ^R	++	✓	++	-	±	++	-	++	++ ^Y	-	✓	✓
--------------	----	-----------------	----	---	----	---	---	----	---	----	-----------------	---	---	---

Nov 10 P11. Readings 1st. 1130A12.
 6 P12
 9 A13

Concl. - [kth. of selective conc.]: M.G. B.G. Sth Sm Perm

<i>Serratia</i>	10-50	-	<100	10	10	<1.	-	100-101
<i>Sal 20</i>	50	50	<100	5	5	1-5	>1	
S21	>100	<100	<100	>5	>5	1-5	>1	
<i>Phyto</i>	1-10	<100	<100	>10	>5	>5	>1	
<i>Staph</i>	<1	<100	<100	5	5	1	1	

Resistance mutants

454a

Staph.	M.G.	1	turbid.
		5	ca 10 Mg^R /cc; gd. inhibition of residue
		10	perfectly clear!
	Sm	1	ca 10 ⁴ Sm ^R .
		5	ca 10 Sm ^R
	Profamine	100.	Turbid.
		10	clear at 48h.
— Seratia	M.G.	20	clear at 24 hrs.
		50	a. ca 20 Mg^R (old?)
	Sm	1	clear. no Sm ^R seen.
		5	ca. 10 app. colonies . some v. small Sm ^R . 1 Sm ^R ? white!
	StH	100	ca 5 StH ^R /ml
		20	ca 100 StH ^R /0.1 ml
			col. v. small wide range of color.
	Prof.	100	turbid
— S20	Bg	100	clear, no Bg ^R !
	Sm	1	turbid
	Sm	5	turbid; but possibly Sm ^R , v. small
	Prof	100	turbid.
— S21	Bg	100	clear, no Bg ^R .
	Sm	1	turbid.
	Sm	5	as S20. inhibition incomplete.
	Prof	100	turbid.
Phyto.	M.G.	10	inhibition incomplete
	StH	50	inhibition incomplete; some selections
	"	100	clear! ca resist.
	Sm	10	incomplete inhibition
		50	clear. ca 1000 resist?
	Prof	100	clear! — ca 300 resistants

Available:

~~Staph:~~
Phyto
Sth 100
Sm 50
Prof 100

St: Sm 5
Sth 10
Zhg 5

820 B.G.
Sth 10
Sm 5

S21 Bg
Sth 10
Sm 1

Seneteq Sth 20 } color variation
50 }
100 }

Sm 5

Mg 50

Reversion detection

455

4/11/47.

As above on N.A. and T(0).

petri dishes show up as papillae in T(0) streaks, justifying this technique. Throw out NA plates.

T(0) colonies in center zone

1	++
2	++
3	+
4	++
5	++
6	++
7	++

streak out further 48 hrs. colonies of 440T + 453T.

n.g. too irregular. NA mix + plate is more reliable.

14 MAR 1947

See 452 for sources.

Colonies from 440 x 453 on Biotin were picked to water + T(O), T(B) inoculated. Here reported are 5/50 which grew on T(O) only after 2-3 days. The T(O) and T(B) tubes were both streaked on vac-v agar: (↓ not turbidity)

	T(O)	T(B)
1	-S; +R	-S; +R
2	-S; +R	-S; +R
3	-S; +R	-S; +R
4	-S; +R	-S; +R
5	+R	+R.

Since -S and +R are the parental configurations, the delayed growth [and the original colony formation] might be due to symbiosis.

∴ streak out T(O) tubes on EMPB vac to purify.

Test - (a) and (b) colony of each on B, O medium.

	a	b
1a	-	-
1b	-	-
2a	++	++
2b	-	-
3a	-	-
3b	-	-
4a	-	-
4b	-	-
5a	-	-
5b	-	-


Expl. 1. Symbioplus colony
2. Colony not picked; only interstitial growth in agar. Requires repeating.

Mucoid variations.

19 MAR 1947

Streak Y53 across T1 on EMB agar.
Suspend slimy growth from ~~Y53~~ intersection of bacteria & phage
with 0.5 + streak out.

Note, at intersections of bacteria & phage, a zone of coloration of
the bacteria as if there were there some enzymatic activity.

 mucoid colonies. Y53M.

N21. Pick from red region + from mucoid colonies to water
and streak out on EMB ke. (Y53/1).

Y53/1 same growth (probably resistant; ~~both~~ lac- + lac+).
(standard type)

1. Y53M₁ - all mucoid. pick one colony + test on T1 VR
also streaks out →

2. Y53M₂ - all mucoid. P23. Streaks out →

3. Y53M₃ - all mucoid. Pick to streak and label for subsequent
analysis. Y56

Plate Y40 x Y53 unusual. At 48 hours pick 5 largest (>) and smallest (<) colonies from each of 7 plates + compare the distributions

-R -S +R +S.

4 1 0 0

4 0 1 0

3 0 2 0

2 2 1 0

2 1 2 0

2 0 3 0

1 0 4 0

In large, +R > -S.

In small +R > -R.

18 4 13 0 }
1 1 1 1 }

$\chi^2_3 = 5.14$ $p = .16.$

27 12 30 1

But, compare all 3 groups,


$\chi^2_4 = 12.61$

$p = .013$

Random selection from these plates gave:

27 13 13 1.

selection may play a role.



221 136 142 6

is cumulative data.

Note that both in in large + small selection types, there is a marked deficiency in -S as compared to random selection & cumulative data!