

Fla⁻ H₁^{non b} serotypes.

1258

DATE: APR 28 1955

REF:

1 2 3 4 5 6 7 8 9 10

1 1252A2 } a-x b
 2 " }
 3 1252B2 } trails? in b serum.

Test by two agglutins. Isolate Fla⁻ H₁^x

B) x - FA10 (H₁^b) + b serum → ? → Fla⁺ H₁^b

A) x - FA93 H₁^a + a serum. If x = a, no swarms. If x = b, H₁^x = b.

Results:

- 1 - Not pure motile. Gave to DCG
- 2 - Pure motile. } S B no swarms
- 3 - " " } S A swarms →

20

30

40

50

DATE: MAY 3 1955

REF:

A. Staining in situ W1177.

a. in mixture c. 0.005% T2 under oil - No
stg. overnight Some inhibition?

b. Dimethylol 4000 20% (vis Penassay) + c. .1% T2,
isolated colonies only (resistant?) - sterilized in center.

? are these conditions too aerobic? Otherwise polynitrate toxicity
of T2.

Should re-isolate colonies; compare growth c, s T2 under oil.

B. Chemis from isolated cells. (see c. 4-6-57 then refer)

1. In scattered region many chemis showed intestinal lethals.
(effect of cold?)

2. Few & granules now seen.

3. Isolates immediately spread growth under them (film of
mortality from deep!) A-B-C

D 2 is tangle
D 4 c 20 cells, 7 term.

5 4 g.
6 4 g.
7 4 g.
10

X

E 1. tangle, 2 term.

2-5 n.s.

6 no T

7 no T

8 no T

9 4 g.

A4

40

50

Motility in Methocel solns.

DATE: MAY 4 1955

REF:

- | | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|----|--|---|---|---|---|---|---|---|---|----|
| A. | attempts at leaving cells in situ, according in relation to 28u MAM beads, in Methocel 4000 cps 2% in linessay. | | | | | | | | | |
| | General conclusions as stated attached, see 1258 p. 100. | | | | | | | | | |
| | Some Z^+ chains 2-100 cells terminal Z cell still intact. Infectious probably more frequent after refrigeration but this is not settled by direct observation. | | | | | | | | | |
| B. | 5/6/55. Set up to repeat 1254. began OK but slow to divide at RT (though warm) & later lost c.g. | | | | | | | | | |
| C. | " Methocel 400 cps seems to slow up motile cells (129A1+). Put. trials as selector for E cells. Part must. used 4000 in trials. | | | | | | | | | |
| D. | See 5/8/55. E - a slump: did not completely inhibit initial med. replication. | | | | | | | | | |
| | 1) Possible use of Scotchbrite MAM beads as reference markers - there is a slow drift; May be better to use 3% Methocel 4000 rather than 2%. | | | | | | | | | |
| | 2) Wasted exp. in counting E, Z chains | | | | | | | | | |
| | 3) Pulver's exp. on screening E, S cells by vesicis medium strongly successful. See ff. | | | | | | | | | |
| | 4) But most time last week or 10 days was wasted exp. to improve general ingression & technique. | | | | | | | | | |

5/6/00 1258

Lab plans: what to do? Things are a mess.

1. Currently enmeshed in the fate of Z granules. Can these really give any important information? By following a granule during the growth of a single cell, one might get a clue as to whether growth is interstitial or bipolar (in a few cases). To distinguish, one might have to show increasing separation between two granules, before fission in a single cell and this may be difficult.

It is already clear that 1) terminal granules usually remain terminal, and that this is the most common type, already suggesting a polarity in the cell. Occasionally, bi-antipolar cells are seen (more commonly than bi-synpolar), suggesting that the two poles share something distinct from the fissile center. However, the basic interest in the Z granule for the current problem is the possible correlation with E, and this, if anything is what should be pursued for now. Later it may be convenient to try to repeat experiments with a polar-flagellated organism.

Another sideline is to use the chains in stiff medium to study other problems, chiefly lethality both spontaneous and UV. Also look for data on growth of branched cells. (Twort)

2. More pertinent: 1) look for divided E further. 2) diagnose E,S cells by viscous media. 3) transfer intermediate chains for electron microscopy 4) clean up serotypes of co-segregants-- collect more? 5) For 4 and others need to complete review of data and write up.

3. TODAY: Clean up what is accumulated to look at and photograph.

Start new preps. of 93--x w/wo TZ. Use for divided clones and for Z correlation.

(Sat 5/7/55- Sun 5/8/55---)

Use T2 stained prepn. 5/6. 12n7 Checked first with 1237A1+ for swarm motility. In this series, used 2% methocel 400, diluted c. 1/10 with penassay.

a) use methocel for trap; b) isolate initials in broth trap, then TRANSFER to mcl.

The latter was found ineffective (probably still too stiff); By 4 PM, had isolated 13 cells still sluggishly motile in mcl trap, and 7 addl. which were at a distance from reservoir but not now motile. swarm cells were sluggishly motile in this methocel conc., about 50-70% were directly inhibited. This soln. probably wets glass more effectively, at any rate it tends to spread, and a few of the motiles below may be contaminants from 1237A1+.

The motile residuals above were ~~planted~~ ^{planted} in individual drops of broth for class. as Sw. or E cells.

NRs found, in first group: 6 swarms, 3 E, 2 ng, 2 E.
second 4 E 1 ng 2 E

Total 6 S 7E 3ng 4 E

which demonstrates strong selection against E cells

Detailed counts:

	growth	motiles
1.	4+	9
2	3 = sw	swarm, 50%?
3	4+	2
4	ng	
5	4+	12 (from Z cell, but Z nf)
6	trap 4+	0,1
7	sn, 1 mot cell	
8	20 sw	16, sev. shakes, prob. sw
9	sw sw	
10	sw sw	
11	like 8 sw	
12	500	12
13	200 sw sw	
21	4+	4
22	ng	
23	4+	18
24	4+	24
25	4+	2
26	4+	35
27	4+	16

(10⁴)

The occasion was also used to plant about 25 single motiles (removed before test below—perhaps should have been left in it) for ~~opportunity~~ ^{opportunity} on immediate and later motility of dividing chain cell. About 12 usable cases no discrepancies, some to one or two later divisions. As none gave two motiles, pres, none of these were E. Of remainder, most gave two app. nm at this division— it may be possible to reexamine these drops tomorrow. What is significance of this crisis in termination? Is is growth in fresh medium? (May still need a good exhausted medium to keep cell size small.)

P8 These were then used in tests for residual motility in mcl. Unf., 1,5 were washed in 5% mcl 15 (calc. visc 200) which proved also to inh. swarms. Further tests were then made with mcl 400, 1.8% and 1% (1:1 penassay), the latter being adopted as it permits almost full motility of motile swarms (from above). (This may be too fluid for accurate discrimination against E, as will be seen). From E: 12, 23, 24, 26, 27, altogether cells reisolated which remained motile were planted for further classification. → none proved definite E cells. See further below.

Until this is worked out against in further pedigree!

b a serum.

DATE: May 9, 1955

REF:

1 2 3 4 5 6 7 8 9 10
Prestained prepn. used. (st Z 9:45-12N; phage to 1:20, centr. and refr. 1:40)
(also another preph. unstained - A.

Note immoderate spreading out of methocel droplets. isolate initials in 1% methocel 400
(1:1 2% penassay).

A. isol. from unstained, plant out in droplets individually.

B. isol. Z-stained initials. to c. 3PM, some fresh isol. C to 4:30
set in single drops on initial cg. first, transfer latter as families to
isolation cg. Ditto for A-- plant out descendants.

D. 5PM B above, in broth traps: pick c. 4000 initials (somewhat late now for tests)
in serums.

(Klein visited 5/10-11.)

D: Almost all initials are inhibited in a or b serum, though cells may continue to spin
for a few minutes. 7 cells did persist in b, planted out. 3 proved viable swarms.
isolate as 1259 D1-3. See DCG for results of platings (after picking to broth) in
MGA. 2 persists in a, but neither viable.


E. same as B-D but not sublined

A: Held to 5/11 for examn, and may have partly diminished therefore.

13 clones 2? E clones. 1% Methocel probably too thin.

B: Most isolates grew out; had been separated once or twice at n₂-n₄. However, of
38 isolations, 3 ng; 5 swarms; only 4E, none interesting except:
E-clones were reexamined for content. In B2, sib to swarms had 22 motiles,
transferred to /238/ E2,3; 4; 5 (sw). The motiles in E2 tested, all gave rise to
inviable or E clones & therefore certainly not sw. cells/

6% originally looked as if only c. 100+/10⁴ but these later proved to be swarms.
The clone was not recovered (owing to drying out) to verify original low assay.



Otherwise, detailed numbers of intermediates were not recorded.
F+ 5/10 interm motiles tested in a serum: at least (28) from 5 clones were imbo.
but 2/4 from B15 were not! However, two tested swarms were inhibited; specificity
of serum should be rechecked.

Also saved 1259B1 (= b8). Swarm- test purity by plating
B2a, B (= c5 z cell removed at n₅ = nonmot, b) (not not certain record)

DCG found D1-3 all motile but with confusing clusters. B1: no definite swarms B2b
"all clusters"; a pure non-motile. Will have to be rechecked on return

no motile

5/17

(over)

E: 34 isolates planted w/o lineage afterward.

AE (9,11,15,16)
15 (1, 4, 3, 1, 1, 6, 7, 3, 3, 5, 5, 2, 1, 3, 1, 4,)
Sw
5 ng.

Only conclusion: medium not adequately selective. Try 1½% methocel 400
(v.i.: 1260)

1259 summary to 5/16

5/12 Plated in MGA

5/13 Picked possible singles. Plates were incubated too short a time at 37°, D1 & D2 had singles, swarms, & clusters; D3, B1, & B2 had singles & clusters only. Counts:

	<u>Clusters & swarms</u>	<u>Singles</u>	<u>Singles picked</u>
D1	51	3	2
D2	91	1	1
D3	59	5	4
B1	32	21	8
B2b	90	17	8

} these spotted on MGA


5/14 All "singles" picked 5/13 & spotted on MGA were motile (Spots had appearance of "clusters" rather than swarms).

Plated again: All original broths, + ^{some of} singles picked 5/13 (D1, 2; D2, 1; D3, 2; B1, 2; B2b, 2.)

Incubated 3 hrs at 37°, overnight at 22°, then refrigerated until examined 5/16.

5/16 Results of 5/14 platings:

Original broths:

- D1 Swarms, centered swarms, & col. c "satellites" 
- D2 ~ D1, higher proportion of swarms.
- D3 ~ D1.
- B1 trails, clusters, apparent singles; no swarms
- B2a pure non-motile
- B2b All clusters

Presumed Fla - :

- D1(1) all clusters
- D1(2) Clusters, swarms } no singles
- D2 Clusters, swarms, no singles
- D3(1) Clusters, swarms } no singles
- D3(2) " " " " }
- B1(1) } Clusters, trails or satellites; possibly some singles;
- B1(2) } no swarms.

N⁴ - 62



MAY 10 1955

Best resume page



- 1259B. ABCDE

In abc 3 swarms, 5E / 34 isolates

disappointing. No E cond. 2. But write out detail any how



B15 Rec. confused.



C4 (+) ^{many} to [238] H2 }
 Residue to F2, F3 }


Hold for full analyses if needed.

Cond B were evidently confused yesterday !!

MAY 1 0 1955

Do not save these swarms among top possible confusions.

But study closely B2 and C4.



C4 is $\textcircled{1} \rightarrow \frac{57100}{10^4}$.

$\textcircled{238}$ kind of pre-picking

Partitions

May 13. New prepn., unstained. (probably usual, about 90-120mins.)

Fuse drops 2:30 Collect to 3:30. Cf 1259D motile.

No initial was nearly as active as 59D. Pick those that have moved the furthest, not necessarily v. active now. Estimated yield, 10% of broth yield.

Note : to compensate for spreading of methocel solution, use cg. that has been greased (human), then flamed. This works well, especially with larger drops, but smaller drops are too convex for best visualization. Intention was partly to look for early gains (E) in the methocel, but time did not allow and most isolates were made to broth directly/(A, B resp.) Lineages were separated at n_2-3 .

A: 1,2,3,6 ok. Partitions at n_1 :

14+ :1 6:5 ng snakes. Later transferred entire clones to get fullest estimate of motiles.

A1 came out +(14):6 Sepn at $n_1 =$

B1-14,21-36. 4 ng. Mostly non E. Records show at first scanning:

2:4;1 14+ 2:1 3 7:20 0;4;1 1;3;d 3 2++ 2:1 1:0 5

sw;sw;sw;sw (1260B33 later DCG verified purity of each). 6:5 7:+

Underscores were rechecked (on ungreased slide!) and following definite values for splits on these:

1:20 8:20 2:2 4:12 3:2 7: 26 Therefore no equal splits.

General totals:

E 5
ng 4
sw 1
E

33

Little if any selection for E in 1 1/2% methocel.400. Need 2% which probably totally stops many motile cells.

No new experiments after 5/14
Trip to NY 5/18 - 5/29. Reserve lab with 1/3!

→ x 92066
Method

1261

JUN 1 1955
MAY 31 1955

1:1
Pupae 93 x 92066, 10⁵ - 11⁴⁵ (12³⁰)
c 430-545 isol. residual motiles. ^{SIC} ^{in antipyrin} Ref. to c. 4 PM.
Est discrimination factor

A) Note: to prevent spread of method, ~~plates~~ are lightly gassed with fungus (on one side); flamed; oil added. However, motile selection seemed most effective when there was appreciable wetting and spreading of the drops on the coverglass.

Notes transferred to fresh petri dishes c 6 PM, Dec 30^o
Counts of \oplus / 10² - 10⁴ : 2, 25, 20, 53, 46, 4, 6, 2, 50 ; 3, 20, 18, 3, 20, 7, 11,
13, 10, swarms.
 Σ : (6E : 3F : 1 swarm. ↓ 2 Inusable

JUN 2 1955

∴ with 2% methanol 400 there is effective discrimination. at this case, Fla⁺ (1237A1⁺) was greatly slowed down (10x ?) but most cells did continue to move.

[Note - to this point considerable interruption in continuity of work was occasioned by ① trip to NY for ascites meeting ② breakdown of manipulator - valve in diaphragm, temporarily repaired.]

∴ continue pedigree studies on preselcted initials.

swarm: manual plating of clone, in 1 ml, .01 ml gave 44 swarms
again note low ratio. Bechler's completion! 265 singles
(see photo - plate had been held at RT overnight, inc 2 1/2 hours³⁷, then RT 4 hours.)


Pedigree.

JUN 2 1955

A) grow in penassay B. grow in Mcl.
Plants pure dips c. 12¹⁰ AM.


1261 pups.

No cells continuously motile like F^{a+} clone were seen. Pick the most active. If continued, probably used to regulate the degree of wetting. Did ~~pedigree~~ to 43-45 on 34 units transferred to broth. ^{chain pedigree}
P3, scan for E, ϕ , swarm. Found only 3 E clones.



\therefore splits are 1:42 3:34 and 28?:0 (known 1:27)
2:25

One swarm clone C4, already pure. DCC checked purity of each clone by plating.



same! as 1262 C1-5
for H₁ check.

swarm totals were only 3E, 1S : 22 ϕ and 10 ϕ (= lethal)
[11-0; 5-1; 2-2, 3, 4, 6, 7],
43, 37, 28

the experiment was quite unsuccessful.

Again review salmonella data
to get paper out of the way.

July 13
1955

A) → Should first get general picture of experiments + what they were!

Write out 1138 Bf

? → X SW666 $4p^+$

Note diminished motility of large cells. Occ. early isol 1. → $1/10^3$...
(remarks on growth cycle) e.g. 1141 A4

1141 A4 v.p.
A5

B1
B3
B4 vp

A1, A3, B5, C1 n.g. (stayed motile)
C2

Σ	a	b	c	d
3	1	1	5	-
	2	6	6	11
1	0	1	1	1
	0	1	1	1

2 1 2 2 av3.

A4 (v.p.)

$\Sigma = 33$.

* ① first summe!

note partition: 19:11
or 1:(19:11):1:1

↑
But cannot use as G2 may be listed as from B4!

B4 if the latter, then

Σ				
A ₂	1	2	2	19*
B ₂	D _{1/5} 18	6	16	16
	B₂ 15	20	27	27-29
	n.g.			40
C ₂	1	2	2	2
B	D _{2/5} A 11	9	15	15
D ₂	1	2	2	7
G2		16	23	33
			23	24-26
				34
				47.

This datum is unreliable.

subscript = point of this branch in the pedigree.

7/13/55

1142. (9)

C3

Σ
750
5 tested

6 10 31 36

D1

7100
10 tested

7 24 41 45

C4

10
2,1,3,1,0,0

3 10 — —
no word

1143

E3

3

}	2	2	12	14
	3	12	—	—

E2

11

}	s_1	2	2	2	7
	s_1	2	2	3	3d
}	s_2	3	10	16	23
		3	10	42	48
}	s_7	5	11	27	31
		5	11	31	38
		5	11	14	—
		5	11	—	15

1144

Leifson cultures.

1272
SEP 9 1955

all 6 cultures grow as well on both at 30° as at 37, except 205.

For preliminary comparisons, re-inoculate H1, H300, H32, H37

1:5 in broth + re-inoculate 9AM -


SEP 8 1955

Leifson's slides Acetabularia

figure pairs primarily

H1 A. 1700x apoch.


P. acynozoa type.



no clear
antipolar
pairs.
speaking?

✓ polar mother
unip.

H 300



ditto
cells larger than above.


H 32



v. bio?
united, antipolar


"alcaligenis"
(Lyphobacter)

→ H 37



large cells. ✓

H 205.



more usual

rare


prob. intermediate

polar multitrunk
same bipolar!

H 430



usually unipolar




(over)

see [242] protocols.

Conclusions

a) 1 pedigree possibly



b) No great regularity; some + < -

Some + < +

Could be studied further
H1 is best culture