

7/8/49

16 cultures tested, streaked on Xyl EMB.

All streaks are Xyl +, - mixture. Occasional v seen. Pick for propagation on Lac EMB.

Xyl v recovered. Streak out on MREMB for material to test Xyl loc.

(320) MH v colonies brushed on Xyl EMB; Lac EMS.

3 Xyl - colonies noted. Grow very poorly on EMS Lac, probably segregated, but streak out on MREMB.

7/17/49

Does buffer activation increase enzyme?

cells harvested from 100 ml Y2 Lac (36 h.) to 5 ml water.

A) 2.5 ml cells + 2.5 ml water

B) " " NaP 4/5. Incubate 12²⁰ PM - 2:15.Assay — samples. Wash treated cells with water, save supernatants, and dry sediments over P₂O₅.

Di

Super
.5 mlA
B016
420 (5 minis.)

No activity is extracted by water. Ca 8-10 fold activation by buffer, and ca. 1/3 activity is extracted.

cells
0.5 mlA
B080
056111 271 (20 minis)
490 (5 minis)∴ Total sediment of B) should have activity of $4.5 \times \frac{1}{.05} \times \frac{450 \times \frac{20}{5}}{100}$
= 1,600 u.and A) should have $4.5 \times \frac{1}{.05} \times \frac{200}{100} = 180$ u

Recover dry cells.

A 7.5 mg.

B 11.0 mg.

B formed a much more compact pellet than A, of which ca 1/3 was estimated lost in preparing for drying and scraping from slides.
Calculate on predicted basis of 12 mg. cells in each.Triturate each in 5 ml H₂O to prepare for assay.

Predicted activity, ca. 150 u/mg for A and 13,000 u/mg for B.

Note: These Lac cells also give brown purple color + iodine.

Two kinds of phage apparent:

ϕ_A carried by 7, 13, 15, 18, 24, 32, 34, 36. type: W1029.

ϕ_B 17, 19, 23, 31

	ϕA^+	ϕB^+	ϕO^+
A^S	—	—	6, 14, 27, 29,
B^S	7, 15, 18, 24, 32, 34, 36	—	10, 16, 21
O^S	13	17, 19, 23, 31.	++

40 cultures, 8 carried A, 4 B = 12/40 = 30%

Have A^S ; 7, B^S = 11/40.

The strongest phage action was that of #17.

#24 responded moderately to λ —

11/29/30: Throw out all except 10, 13, 14, 17, 24

Lysozymicity tests on Shapiro's
E. coli cultures.

7/16/49.

Test by seeding λ each strain as indicator, then spotting each strain on surface. Active combinations indicated:

Host

1			
2			
3			
4			
5			
6		#7, 13, 15, 18	
7			
8		17± 19±	
9	s.g.		
10			
11	s.g.	17± 19±	
12			
13	s.g.		
14			
15		7, 13, 15, 18	
16		17, 19 ++	
17		17, 19	
18			
19		17, 19	
20			

Following groups noted.

6, 14 sensitive to 7, 13, 15, 18

7, 10, 15, 16, 18 sensitive to 17, 19.

Note: 19 → 18 → 14.
6

(7, 15, 18) $P_a^+ P_b^s$
 (6, 14) $P_a^s P_b^+$
 (17, 19) $P_a^- P_b^s$
 (10, 16) $P_a P_b$

Shapiro	Key:		
835	w 1028	# 6	sensitive to p 7
866	w 1029	# 7	" p 17
848	w 1030	# 10	" " ; canis p 7
116	w 1031	# 17	canis p 17

Should include $\lambda - \lambda^s$; $\lambda - \lambda^R$ and λ^+ in these tests.

7/19/49.

Checks, intense, # 21-40.

21	v. str.	23, 31	
22		—	
23		—	
24		23, 31	
25		—	
26		—	
27		24, 32, 34 (all weak), 36.	
28		—	
29		24, 32, 34 (v. weak), 36.	
30		—	
31		—	
32		23, 31 v. strong.	
33		—	
34		23, 31 wh.	
35		—	
36		23, 31 ++	
37		—	
38		—	
39		—	
40		—	

21, 24, 32, 34, 36

← 23, 31

27, 29

← 24, 32, 34, 36

N.B. 24, 36 → 29
 ↑
 31 →

2 or 3 λ's indicated.

Test 21-40, λ, W1029 and W1031 m.

27
 32
 1028
 1029
 1030
 818

24, 32, 34, 36, 1029
 no λ's.
 24, 32, 34, 36, 1029
 no λ's.
 no λ's.
 no λ's

λ?? exceedingly weak if at all.

(even λ did not show up).

7/15, 17/49.

- c) 1. W478 x W1022. 635 Lact+ : / 14 Lact- .
 200 Lact+ tested for lac_v. None found.
 100 additional tested. Select those that appear weakest Lact+
 5 suspicious colonies reincubated. 3 are lac_v.
 Pick A) weak lac_v and B) broad streak to LacEMS, LacEMB, MalETIB.
 ulm are data?
- A) (2)⁽¹⁾ W67 x W1022 1 Lact+ / 3000 prototrophs.
lac_v Mal+ Xyl+ MH_v?
- B) 3. 440/6 x W1022. 10 plates LacEMS. To determine whether
 the prototrophs from this cross were diploid. Test 20 Lact+,
 12 Lact- as T(0) with T6.
 Lact+ : 19 R 1 S
 Lact- : 6 S 4 unucoid, unscorable. 2 too dilute.
 Indicates that Lact+ are not Lact+ v_6^R/v_6^S , and therefore
 that they are haploid.

W1033, 1034 received from Cavalli 7/21/49.

Both are ~~F⁺~~ lact. 1033 is (B)M 1034 is BMB, ? as labelled.

Cross \bar{c} w677, cf. 58-161

Harvest and concentrate ca 5:1 as usual. Cell concentrated cells 1ml = .1, and plate various dilutions of 1033 on EMS lac or T(0) plates smeared with 677 (= 1ml).

Count: 10^{-6} ml 1033 on EMS lac gave 580 colonies.

Count = 5.8×10^8 .

No prototrophs appeared at 10^{-7} or 10^{-6} .

10^{-5} (on T(0)) gave 1 10^{-4} gave 30.

False rate as $\frac{30 \times 10^4}{580 \times 10^6} = \text{ca. } .05\% = 5/10,000$.

but 58-161 at this dilution gave 34 colonies !! (mixup?)

On EMS, No colonies were seen at 10^{-4} from 1033 x 677.

10^{-3} gave 18 (5+13-)

However streak of 1033 on 677 background on EMS lac B, gave very much higher yields than did 58-161. The effect must be largely based upon microcolony formation. Efficiency of 58-161 should be checked!

7/24/49

Inoculate 677 and 1033 heavily together in Y2 broth (1:5 ca.)
 After 24 h. streak out on lac EM5 and test single +, - colonies
 on Xyl.

21 lac- were Xyl-
 of 8 lac+, 6 were Xyl+, 2 were Xyl-.

Recheck these 2 (585 B1, B2) on lac, Xyl, and mutation.

JUL 25 1949

B1 and B2 are both lac+ Xyl- Mal-. Store as W1048, 1049
 LB₁-

Additional tests of same plating:

13 Xyl- → 10 lac-, 2 lac+, 1 mixed.

36 Xyl- → 29 lac- 5 lac+, 2 mixed

49 Xyl- : 39 lac- 7 lac+ 3 mixed.

5 Xyl+ : 5 lac+

14 lac- : 14 Xyl-

24 lac+ : 18 Xyl+ 5 Xyl- 1 mixed.

There are exceptional lac+ Xyl- present. 1033 is pure,
 but 677 needs re-verification. Streak out on lac EM5 to
 check! OK!

7/25.

Cross check plated mixture of super. Xyl and Mal.

39 Xyl+	36 Mal+	3 Mal-	} 1-5] 6	6 Xyl-Mal recomb.
5 Xyl+	3 +	2 -			79 keto
35 Xyl-	34 Mal-	1 Mal+			6.

	lac	Mal	Xyl	Gal	MHE	U R S R S R R R R S	Nutr.	W
1	+	-	+	-?	-		L	1060
2	+	-	+	-?	+		MT	1061
3	-	-	+	-	+		TLB ₁	1062
4	-	-	+	-	-		B ₁	1063
5	-	-	+	+	+		MTL	1064
6	-	+	-	-	-		TLB ₁	1065
677	-	-	-	-	-		TLB ₁	1066
"H6" = 1033	+	+	+	+	+		M	1067
25 Xyl-		1 Mal+						
15 Mal+		1 Xyl-						
20 Mal-		All Xyl-						
7	-	+	-	+	-	R	TLB ₁	1066
8	+	+	-	-	-	R	ML	1067
							#7	1066
							#8	1067

JUL 23 1949

A	1014 x 1015	7(0)	
B	1014 x 588	233+ : 110-	Lac EMS.
C	1033 x 477		

A. Yield quite low, ca 5-10/plate instead of > 100.
 Streakout on EM3 Mal. 31- : 7+ No Mal v.
 Note: BM Mal- x TLB, Mal+ should be excess of Mal+.
 cf. W1022.

B.
 100 tests
 Emphasis
 on local +
 prototrophs.

4 distinct, 5 uncutaminocuv. → (7) heavily Lac v.
 Streakout a) isolated Lac v and b) broad brush on
 Lac EMS, Lac EMS, Mal EMS.

52 addnl.

C.
 100 tests

1 distinct Lac v. ~~2,3~~ ?? Lac v. all Lac v.

B)

	lacEMB	MalEMB.	
1	✓	+	
2	✓	+	
3	✓	+	
4	✓	+	
5	✓	+	
6	✓	+	
7			
8	✓	+	1 v ⁺ colony?
11	✓	+	
12	✓	+	
13	✓	+	
14	✓	+	
15	✓	+	
16	✓	+	

None Mal ✓. All Mal+

C)

1	✓	
2	✓ ⁺	
3	✓ ⁺ ?	v ⁺ ?

Crosses on double-sugar
Hfr.

585.

JUL 25 1949

(A) (B) (C)
W1048, or W1049 mixed with Y87/6 and inoculated
1:10 into Y2 3^{PM}

3^{PM}. Streak out ~~A, B, C~~ A, B, C on Lac, Mal and ~~Lac+Mal~~ Malac

Plate out AC and BC on EMB Malac

Repeat P25.

A, B were pure Lac+Mal-; C was Lac-Mal+ All (A, B, C) were
pure+ on Malac.

BC. 4 plates ca 150 scorable. 1 Malac - colony noted.

AC. 5 plates ca 250 (=1250) 4 Malac - colonies? [might be a
contaminant.] Isolate to analysis.

Repeat:

AC: (2 x 100) No -.

BC: (3 x 150) No -.

These might ~~be~~ not be Hfr.

See 589

24 hr heavy mixed cultures plated on FMB Malac

A	487 + W814	0	487 + W814 pure lact, lact resp.
B	1056	"	✓
C	1088	"	(✓)
D	1059	"	✓✓

4 plates each.

- A). 1600 cols. No Malac -
- B). 4 Malac - 2 sectorial. 1 sectorial not isolable.
- C). 3 - and 2 sect??
- D). 7 - and 6 sectorial, not all isolable

see FMB. (2 plates).

- D). 2 sectorial
- B). 1 " + 2 unisolable
- C). 1
- A). 0 (1 plate).

JUL 2X 1949

Sh	Yr	Origin	Media	Notes	Media	Notes
Sh 1		(5)	HC, YE, (AS)	YNA +	A6 ±	W1045. <u>Proline</u>
Sh 2		(5)	"			"
Sh 3		(5)	"			"
Sh 4	72	K-12	+ all AA			
Sh 5	75	"	± all AA A12, A4++			
Sh 6	76	"	+ in 48 hrs. + all AA.			
Sh 7	77	"	± all amino acids. A4, A12 ++			
Sh 8	50	"	like II			
Sh 9	51	"	+ or ± all amino acids			
Sh 10	53	"	± minimal 36h. do. II			
Sh 11	54	"	24h: HC, YE, (YNA), A4, (A12) ± others			

→ All grow promptly on tyrosine, later adapt.

Sh 1-3 were 3 units/20 tests.

47 additional colonies tested. 6 mutants found: # 4, 22, 26, 35, 40, 43.

Each also showed up very well on minimal agar.

4, 22, 35 were prolineless. T.O.

Random remaining 3: A: A12
B: cystine
C: histidine

W
1050
1051
1052

of Sh 4-11.

Summary. 9/67 from treatment and penicillinism on the wild type, W1045, were mutants. of these 9, 6 were prolineless. Only one was preserved as W1046. The others were W1050-1052, respectively, :

1050
1051
1052

A series of 8 "mutants" were isolated from a K-12 strain. All proved to be rather revertible tyrosineless. Keep # Sh 11 as W1053

JUL 25 1949

W1050 = A

1051 = B

1052 = C.

Plate separately and in combination on T(0) plates.

(Also inoculate into Y2.

At 36 hours, A-C were blank, AB and AC had numerous prototrophs. BC had a few.

A27. (48h.) A reversal noted. Not so numerous as in AB and AC. ?? Reversions or recombinants??

check 1050 with various ψ stocks. Cf. 1033. In Y agar.

	T1	2 λ	3 λ	4	5	6	7
1050	R	R	R	R	R	R	R.
1033	S	S	R	S	S	S	S
			↑				
			N.G.				

Need double mutants for conclusive result

New penicillin^r uses gene low yields of double mutants

#1 from 1050

= W1069

lys + methionine

tryptophane

#3 from 1051

W1070

Cysteine.

Have growth responses to A12
also on Vits.

16 hrs: +++ on methionine or lysine

Also give smaller responses on:

Vitamin mix

B₁

Put

Inositol

K

B₁₂

Probably recessive
or tube contamination

24 hours - growth on T(0)

just a "slow" mutant.

Crosses of Lac- mutants.

7/24/49

radiate 1033 on 18 Lac⁻EMB plates. Survived high (N_g^R!)
 6 mutants recovered. Ca 6000 colonies.

	Lac	Mal	bla	gal	
1	-	-	-	+	} W1054 1055-1059. "5-9"
2	-	+	+	+	
3	-	+	+	+	
4	-	+	+	+	
5	-	+	+	+	
6	-	+	+	+	

Cross 5,7,8,9 $\bar{\times}$ W677 for Lac alleles, on EMB^{lac} and
 in Y2, then to EMB^{lac}.

5x	All Lac-
6x	2-3% Lac+
7x	All Lac-
8x	20% Lac+
9x	All Lac-
Parents	All -

} None very suitable as
 recombination nucleus.

EMB^{lac}. (Mixtures plated out after 24h.)

5:	0/400 +.
6:	0/500
7:	0/150
8:	0/800
9:	0/25.

Lac, \times Lac, recombination unfeasible.

7/27/49.

As supra. 677 background, m T(B₁).

1 ml = 10⁹ cells.

Dilution

58-161

1033

0

>1000

3

11

>1000

4

2

200

5

4

20+

6

4

[Photographs]

3/21/50: Conclusion is ca 100-fold augmentation in H₂O.

7/29/49

Stratout sectional colonies on FMB Lac. (see 585a)

1. D Lac
2. D Lac
3. B Lac
4. C Lac
5. B Malac
6. B Malac
7. C? Malac
8. C? Malac X
9. D Malac
10. D Malac
11. D Malac
12. D? Malac
13. D Malac

Also, 4 - from B, 4 - from C and

6 - from D

	Lac	Mal	Gal	Xyl	MHL	V ₁
B 1	-	-	+	Xyl	-	V ₁
2	-	-	+	-	-	R
3	-	-	+	-	-	R
4	-	-	+	-	-	S
C 1	-	-	-	-	-	S
2	-	-	-	-	-	R
3	-	-	-	-	-	R
4	-	-	-	-	-	R
D 1	-	-	-	-	-	R
2	-	-	-	-	-	S
3	-	-	-	-	-	R
4	-	-	-	-	-	S
5	-	-	-	-	-	S
6	-	-	-	-	-	S
	+	-	-	-	-	R
	-	+	+	+	+	S

10 - from 589a D.

- 3 Lac - Mal - Gal - Xyl - MHL - V₁^R #3,4,5
- 6 " " " V₁^S
- 1 Lac - Mal - Gal - Xyl + MHL - V₁^S #9

launt.

A-D plated at 12h to EMB Lac. Pick +, r to
 then sugar for random test.

A) 36 Lac+ : 34 Mal-Xyl- " 2 Mal+Xyl+" Recheck!
 1 Lac- " Both were Lac-

40 Lac- : 40 Mal+Xyl+

B) 10 Lac- : 10 Mal+Xyl+
 40 Lac+ : 40 Mal-Xyl-

D) 40 Lac+ : Mal-Xyl-
 20 Lac+ : Mal-Xyl-

60

16 Lac- : Mal+Xyl+

C) 36 Lac+ Mal-Xyl-
 20 Lac- Mal+Xyl+

∴ no "recombinations" except in A. 12 hours too short?

Recheck:

1. Lac-	13	All V_1^S	5 Mal- Xyl- 8 Mal+ Xyl+				
			1, 4, 5, 6, 7,				
			9, 10,				
			2, 3, 8, 11, 13				
Lac+	22	All V_1^R	All Xyl- Mal-			!!	No Recomb. Mal+
2. Lac-	16	V_1^S # 25 V_1^R	All Xyl- Mal-				
Lac+	14	All V_1^R	All Xyl- Mal-				No Mal+!
3. #10 adfect							
Lac+	14	V_1^R	Mal* - Xyl-			(No recombination here.)	
Lac-	6	V_1^S	Mal+ Xyl+				
4. Lac+	29	V_1^R	Mal- Xyl-	} 37			
	8	V_1^R	Mal- Xyl-				No Mal+
Lac-	4	V_1^R	Mal- Xyl-				
5. Lact	8	V_1^S	Mal- Xyl-			<u>2 Recombinants</u>	
Lac-	2	V_1^R	Mal- Xyl-				

6.					
lact+	13.	V_1^R	Mal-	Xyl-	
lact-	8	V_1^S	Mal-	Xyl-	
8.					
Lact+	25	V_1^R	Mal-	Xyl-	
lact-	11	indistinct	} V_1^S	Mal+	Xyl+?
9.					
lact-	10	V_1^R	Mal-	Xyl-	
lact+	11	V_1^R	Mal-	Xyl-	
10.					
lact-	12	V_1^R	Mal-	Xyl-	
Lact+	23	V_1^R	Mal-	Xyl-	

No - colonies on
Malac! - Not recombining

Summary: Parents are V_1^R Lac, Mal, V_1^R I:

+ - R
- + S

①	--S	--+S	+ - R
②	--S		+ - R
③	--R		+ - R
④	--R		+ - R
⑤	+ - S		
⑥	--S		+ - R
⑨	--R		+ - R
⑩	--R		+ - R

Selected as Lac^V .

7/30/49

Replate 589D, after 2 hrs. additional at room temperature, on EM5 lac and Malac.

10 Lac plates Lac+ > Lac- ca 500 colonies.

2 ? Lac

20 Malac plates. ca 1000 colonies.

###	###	10	Malac -
		2	Malac v (+/-)
		2	? Malac v (maybe +/-)

14

1.4%

- 14 D Lac → Pure Lac+! Hazy margins.
- 15 D Lac → Test +, - on Mal. Pure Mal-! 2 L-M- 11 L-M+; 10 L+M-
- 16 D Malac
- 17 D Malac
- 18 D Malac? (+, -)
- 19 D Malac? (+, -)

maybe +/-

[40 Lac tested. All Mal-]

Lac segregation seems to be a good criterion of recombination.

17. 5 Lac+ : Mal-

35 Lac- : 22 Mal+; 17 Mal-

18. 20 Lac+ Mal-

20 Mal+ Lac-

19. 30 Lac+ Mal-

3 Lac- Mal-

7 Lac- Mal+


14. 8 Lac- : 3 Mal-

5 Mal+

33 Lac+ : all Mal-

16: 40 Lac- : Mal-

40 Lac+ : Mal-



No Lac+ Mal+!

looks for recombination of other factors — all parental reperm + U, R

7/29/49

W1034x677

100 Lac+ tested for lac^v. All Lac⁺

Many prototrophs were selected on EMS Lac!

January 31, 1949.

A) N1058 + 477 (for persistent heterozygotes: lac $\frac{1}{2}$ v.)

B) N1058 + 814 24hmi. 72h. Room temp.

C) N1056 + " " " "

D) N1059 + " " " "


E) N1022 + 1033 (Reverse symmetry)

8/1 F) 487 + 814. [like B-D + 24h.]

A: w 10x200. No lac+ noted.

B-D: high proportion of lac-Mal-

Select segregating colonies from lac EMB:
But frequency of Mal-lac- makes it difficult to decide what parents are!



F. 8/2. ca 300 colonies. No Mal-lac-. Antlac, +, - seen =

E' p. 8/2 1 lac v colony. Also Xyl, Mal v. Cross test.

G. 8/2. 1033 + 595 24h.

E'' p2. 8x100. lac EMB. Almost all+. No v noted.

D: ca 200/plate. 10 lac: - exceed. In termination unavoidable.
Several weak + colonies noted (9) may be Mal-lac+. 1-6 lac v.

2 Xyl	1 v		#7
2 Mal	1 v?		#8
5 Mal	2 v?		9-10
1 Xyl Mal	1 v	⊕	11

E: ~~15~~ 300 lac+ Mal+ 18 lac-Mal-. No recombinants! (Do 1022 anti Mfr?)

August 4-6, 1949

P. (W1033+ W595).

selection	lac	Mal	Bal	Xyl	M+L	(lac) T1	
1	+-	++	++	++	+	SR	4L+ : MG+
2	+-	+-	++	++	+	SR	8L+ : 6MG+ (2MG-)
3	+-	+-	++	++	+	SR	10L+ : MG+ 8L- : MG-
4	+-	+-	++	++	+	SR	10L+ : M+G+ 10L- : M-G- (1) M±
5	+-	+-	++	++	+	SR	10 " 7 "
6	+-	+-	++	++	+	SR	10L+ → M+G+ 6L- → M-G-
7	+-	++	++	+	+(-)	+S-R	10M+ → L- 10M- → L-
8	-	++	-	-	-	-R	10L+ : M+G+ 10L- : M-G-
9	+-	+-	-	+	+(-)	SR	10L+ : M+G+ 10L- : M-G-
10	+-	+-	+	+	"	SR	10L+ : M+G+ 10L- : M-G-; SMG-
11	+-	++	+	+	"	SR	

lac
xyl v
Mal v
Bal v
" "
xyl Mal

NoL-!

#11 → 10 Xyl+ → LMG+ 10 Xyl- → 9 LMG- 1 LG-M±?

#8 is Mal v all else -R!

#7 → 10 Xyl+ → LMG+ 10 Xyl- → 10 L+G-M- sic?

Summary:

- #1 was "pure" lac+M+G+ (parental). - not seen on plate but were there!
- #3, 5, 6, 9, 11 parentals only (re lac Mal Bal)
- 2 : lac+M+G+ (parental) and (lac+M-G-) (recomb.)
- 4 : Both parents and (?) L-M+G-?
- 7 : L+M+G+ X+ (par.) and (L+G-M-X-) (recom.)
- 8 : LMGX- (par) and (M+LGX-)
- 10 : LMG+ (par) and; LMG- (par). and (M+L-G-)
- 11 : ~~LMGX+ (par) and LMGX- (par)~~

only 4 likely recombinants. Exhaustively check all of these. (2, 7, 8, 10)

8/7/49

M attenuated!

#7. 10 lac+ ^{Mal+}Mal- Xyl+ Gal+ MHL+ V₁^S } ^{M₃} Complementary Recombinants
 10 lac- Mal+ Xyl- Gal- MHL- V₁^R } TLB

Mal scoring questionable. Also some inconsistency. Colonies which originally scored (Xyl+)lac+ and (Xyl-)(lac+) → lac-! Recheck! and mutation

#10 (A) 10 Lac+ Gal+ Xyl+ MHL+ ^{a?}Mal- V₁^S Recomb?
 (B) 5 Lac- Gal- Xyl- MHL- Mal+ V₁^R Recomb. Recomb.
 (C) 5 Lac- Gal- Xyl- MHL- ^{a?}Mal- V₁^R Parental

These appear quite difficult. a may be Mal+. Do there are interactions between lac or other factors on Mal expression?

#8 10 Mal+ lac Gal Xyl MHL- V₁^R TLB, Recomb.
 10 Mal- lac Gal Xyl MHL- V₁^R TLB, Par.

#2. 6 lac+ Mal+(?) Xyl+ MHL+ Gal+ V₁^S
 2 lac+ Mal- Xyl- MHL- Gal- V₁^S

Mal+ was mucoid when examined.

8/10 Confirmation: 1) lac+ is weak Mal+
 lac- is strong Mal+ } Both Mal+!
 2) Gal+ is Mal+(slow)
 Gal- " "

10) A Mal slow
 B Mal+
 C Mal-

Interpretation of Mal slow??

PH. Replate G.

20 Lac. ca 150/

Lac - recomb. 2 v (difficult to ascertain)

15 Mal

3 v

Many not stable or certain

5 Xyl

4 v

Cleanest plates.

5 Gal

No v

Note - : + scored higher on Xyl and Lac than on Mal! (i.e., exc Mal + Lac-?)

Selection	Lac	T ¹	Mal	Xyl	Gal	Strains
1 Lac	+S	-R	+ -	+	+	Lac + -
2 Lac	+S	-R	"	+	+	Lac + (-)
3 Mal	+	-R	"	-	-	Mal + -
4 Mal	+S	-R	"	+ -	+	Mal + -
5 Mal	+S	-R	"	"	"	Mal + -
6 Xyl	+S	-A	"	"	"	Xyl + -
7 Xyl	+S	-A	"	"	"	Xyl + -
8 Xyl	+S	-R	"	"	"	Xyl + -
11 5926X	Xyl	-S	"	+	"	Xyl +, -
12 5926M1		-S	"	+	"	Mal ++
13 5926M2		++SR	"	+	"	Mal ++

Strains on selection sugar. Recomb on Lac, Mal, Gal, Xyl, T¹

see lac-S; 3 Xyl-Mal+; 7 Xyl+Gal+
Not recomb. (T.O.)

Patent Recomb: #3: Mal + - : Lac - Xyl - Gal - (R?)

Lac +

<p>11 (592B) Xyl±</p>	<p>Xyl+ Lac- Mal+ V_1^S</p>	<p>Parental</p>							
	<p>Xyl- Lac- Mal+ V_1^S</p>	<p>recomb.</p>							
<p>13 Mal±</p>	<p>10 Mal+ Lac- V_1^S</p>	<p>Xyl+</p>	<p>} Parentals only.</p>						
	<p>10 Mal- Lac+ V_1^R</p>	<p>Xyl-</p>							

P4. Plate again from "B" in various: Ca 100 / plate

20 Lac 16 v. [2 BL]

15 Mal (lytic reaction) 2 v MI-2

5 Xyl 1 v *

5 Malac 4 - No v.

} See 591b for analysis

of 1-16, # 4, 7, 8, 11 were pure Mal- but Lac+, -
Analyze these for recombinants. (v?)

a) Verify 4, 7, 8, 11 as having Lac+ Mal- and Lac- Mal-

b) Sample ca 10 crashed cells
Lac+ Mal- and Lac- Mal+

- 1 Lac+ Mal- and Lac- Mal+
- 2 " " "
- 3 " " "
- 5 " " "
- 6 " " "
- 9 " " "
- 10 " " "
- 12 " " "
- 13 " " "
- 14 " " "
- 15 " " "
- 16 " " "

Parentals only.
re Lac; Mal

c) Reanalyze 4, 7, 8, 11.

	Lact	Mal	Sal	Xyl	MR	V _i	R	TLB _i	Rec
4	Lact+	Mal-	Sal+	Xyl-	MR-	V _i	R	TLB _i	Rec
	Lact-	Mal-	Sal-	Xyl-	MR-	V _i	R	TLB _i	Rec
7	Lact+	Mal-	Sal+	Xyl-	MR-	V _i	R	TLB _i	Rec
	Lact-	Mal-	Sal-	Xyl-	MR-	V _i	R	TLB _i	Rec
8	Lact+	Mal-	Sal+	Xyl-	MR-	V _i	R	TLB _i	Rec
	Lact-	Mal-	Sal-	Xyl-	MR-	V _i	R	TLB _i	Rec
11	Lact+	Mal-	Sal+	Xyl-	MR-	V _i	S	TLB _i	Rec
	Lact-	Mal-	Sal-	Xyl-	MR-	V _i	R	TLB _i	Rec

Parents + - - - R TLB_i
- + (-) + + S M

Recheck Sal char!

August 3, 1949

- A) Mix 1/2 ml ca. grown cultures of 814 and 1059
 B) Mix 1 ml ca. in 10 ml Y2.

Plate after 12h.

A) Malac 2x50. No -
 Lac ca 150. No Lac v.

B) Malac 1x100 No -
 Lac 2x100. Several Lac v.

Streak out and brush on Lac, Mal, Xyl etc.

	Lac	Mal	Xyl	Mtl	TI (mal)	
1	+	-	-	-	R	Lac-Mal+ 27; 16+- All Mal-
2	+	-	-	-	+S -R	
3	+	-	-	-	R	10L-M+ 25L+M- 15L-M+ 13L+M-
4	+	-	-	-	R	
5	+	-	-	-	+S -R	
6	+	-	-	-	+S -R	

∴ 1, 3, 4 are pure Mal-Xyl-Mtl-Lac v. R.

2, 5, 6 are mixed, but show parental combinations only (No Mal+Lac)

Plate B after 36 hours for further segregation material.

14 Lac EMB. 3 Mal EMB. 2 Mal Xyl EMB. ca 150 ca. plate

3 Mal v. Only 2 isolable (7, 8) No Mal Xyl v

All Lac v, some were distinctive v, others were 9,10, and some were either v or conjunctions.

11-19 were distinctive Lac v. 20-24 singly assessed + and - is a common ratio. #12 may have a fortuitous Lac-; #13 Lac

P2.

Analyses of 7-24.

592a.

August 4-6, 1949
streaks

	lac	Mal	Xyl	MHC	TI		
<i>Mal</i> + -	7 = +	+ (-)	++	+ (-)	+S -R	(MR)	10L- : M+ X+ ; 2L+ M-X-
	8 = +	"	+	"	"	"	10L- " 8L+ M-X-
	9 +-	"	+ (-)	"	"	"	10L- M+ 10L+ M-
	10 +-	"	"	"	"	"	" "
	11 +-	"	"	"	"	"	L-M+ L+M-
	12 +-	"	"	"	"	"	L-M+ L+M-
	13 ++, -	"	"	"	"	"	L-M+ L+M-
	14 +-	"	"	"	"	"	L-M+ L+M-
	15 +-	"	"	"	"	"	L-M+ L+M-
	16 +-	"	"	"	"	"	L-M+ L+M-
	17 +-	"	"	"	"	"	L-M+ ; L+M-
	18 ++ <i>too wild</i>	"	"	"	"	"	10L+M- 6L+M+
	19 +-	"	"	"	"	"	L-M+ L+M-
	20 +-	"	"	"	"	"	L-M+ L+M-
	21 +-	"	"	"	"	"	" "
	22 +-	"	"	"	"	"	" "
	23 +-	"	"	"	"	"	L-M+ L+M-
	24 +-	"	"	"	"	"	" "

None of these (7-22) were recombinants of lac with Mal!

X-	Lac+Mal-	Lac-Mal-X+
3A	W814 + W1062	
3B	" W1063	
3C	" W1064	

No recombination

Plate on EMB ~~with~~ Lac and Xyl. 24h. 37°
 B, C had one Lac⁺ each. (ca 5 x 100 cols)

B had several Lac⁺ colonies. One of these streaked out and cross-killed Lac... Xyl. All parentals: P1: 14 Lac-Xyl+
 12 Lac+Xyl-
 20; 23 resp.

A-C' subplated after 48 hours.

1 ^{2,3} from Lac EMB 4 ⊙ from MalXyl EMB.

3A-C. Ca 300 cols. ca. No Lac-Xyl-. Probably not Hfr!
 unless Xyl+ is favored in these combinations.
 1 Lac⁺ seen on 3A streak and analyze as 593-5.

593:1-5.

Cross test.

	Lac	Xyl
1	+-	+-
2	++-	+-
3	++-	+-
4	++-	=, +

} All Lac- were Xyl+
 Lac+ were Xyl-

No Recombination's apparent

W1062-1064 apparently not Hfr

Conditions of Hfr recomb.

594

August 4, 1949

GPM. Mix overheavily 814 + 1059 (20x)

- A) In water (saline)
- B) in = vol. 1/10
- C) in = vol 1/2.

Final suspensions ca 5x.

A6: dilute and plate out on Malac and lac EMB
(36 hours). (3 plates ea.)

A. Malac: ca 100 cols. total No -
lac +, - = No.

B: ~~##~~ No colonies

C: Numerous Malac - ! 15- / 40+
Numerous lac u. ca 1/20

∴ Recombination occurs in heavy 1/2 suspensions
but not in saline

August 8, 1949.

- A = 1069 (LyMeth + fyp)
- B = 1077 (hist + iso-val)
- C = 1078 (cyst + iso-val)
- D = 1081 (cyst + LyMeth)
- K = 1033 (BM Hfr)

inc. separately P7, and also AB, AC, AD, and BK
~~th.~~ 48h.

- A —
- B —
- C —
- D —
- K —

- AB B-4 diffuse colonies *
- AC —
- CD —
- BK —

- A+B —
- A+C —
- A+K —
- B+K —
- B+D —
- C+K —
- D+K —

Pick + streak on T(10)

~~W1073~~

W1073 x W909

August 8, 1949

W1073 BM Hfr lac-Mal- x W909 Y10 Gal- plated after 36 hrs. in Y2 lac
ca 100/plate






16 lac ~~5~~ possible lac⁻. 1-19 fairly distinctive. 20-32 ^{10 10} \ominus
ca 10 tests each (L⁻, L⁺)

5 Mal 2 Mal⁻? 33-34

5 Gal ^{1??} Gal⁻. 35 | Amicubate.

	Lac	Mal	Gal	L+M+G-		
1	All L+	+	-			
2			-	- + -		++ -
3			-	- + -		++ -
4			-	- + -		++ -
5			-	- + -		++ -
6			-	- + -		++ -
7			-	- + -		++ -
8			-	- + -		++ -
9			-	- + -		++ -
10			-	- + -		++ -
11			-	- + -		++ -
12			-	- + -		++ -
13			-	- + -		++ -
14			-	- + -		++ -
15			-	- + -		++ -
16			-	- + -		++ -
17			-	- + -		++ -
18			-	- + -		++ -
19			+	- + -		++ -
20			-	- + -		++ -
21	ogly+ me papillae	- +	+	- + -	(++ -)	++ -
22		+	-	- + -		++ -
23			-	- + -		++ -
24			-	- + -		++ -
25			-	- + -		++ -
26			-	- + -		++ -
27			-	- + -		++ -
28			-	- + -		++ -
29			-	- + -		++ -
30			-	- + -		++ -
31			-	- + -		++ -
32			-	- + -		++ -
33			+	- + -		++ -
34			+	- + -		++ -
35			+	- + -		++ -

P9: ca 5% Galac- or Galac⁻



But the elimination must be due to deficiency, as deficient chromosomes can be kept in heterozygotes!

8/17/49

Recheck of results of 596. One + and one - picked from each of 1-34 and rechecked on vac, mal, Gal.

	LMG	LMG	
1	++-	++-	(no 1-)
2	-+-	++-	
3	-+-	++-	
4	-+-	++-	
5	-+-	++-	
6	-+-	++-	
7	-+-	++-	
8	-+-	++-	
9	-+-	++-	
10	-+-	++-	
11	-+-	++-	
12	-+-	++-	
13	-+-	++-	
14	-+-	++-	
15	-+-	++-	
16	-+-	++-	
17	-+-	++-	
18	-+-	++-	
19	--+	++-	
20	-+-	++-	
21	--+	++-	++- (no 21+)
22	-+-	++-	
23	-+-	++-	
24	-+-	++-	
25	-+-	++-	
26	-+-	++-	
27	-+-	++-	
28	-+-	++-	
29	-+-	++-	
30	-+-	++-	
31	-+-	++-	
32	-+-	++-	
33	--	++-	
34	--+	++-	-+-

w 909
w 1073

++-
--+

vector characteristic of w 909 weak.

August 9, 1949.

Plate 24 h. cultures on Lac, Mal, Malac. (M-L+ x M-L-).

No: ca 150/

6 Malac: No - found.

8 Lac: No - found.

8 Mal: 2 u, unusable.

Replate 3P/0.

ca 200/

5 Malac No -

10 Lac

10 Mal

Replate 1A14:

No Malac -

No Lac u

Replate from 8/14 on 8/20. (Malac)

August 15th 1949

Mutant yields not recorded.

W	sh 3						
1097	A.	histidine					
1098	C.	aromatic?	mutant!	Any 2 of	tyro; Val;	tryp	esp.
1104	D.	threonine					
1102	E.						
1099	F.	histidine					
1101	G.						
	H.	Valine					
	I.						
1105	L	Threonine	or	any pair from	Arg Lys Meth Cyst.		
1100	M	histidine					

Arg+Lys Cyst+Meth 1st 2nd A L C M

	sh 4		
1106	B	histidine	
1107	D	histidine	
1109	E	Val	2 nd
1108	F	histidine	
1110	K	Val	2 nd

8/23/49

2/10/49

Nutritional tests of 576 B

1
2
3
4
5
6

-			+		
M	T	L	B	i	t
+	-	-	-	+	
+	-	-	-	+	
-	-	+	-	+	
+	-	+	-	+	
+	+	+	-	+	

M	T	L	B	i	t
+	-	-	-	+	
+	-	-	-	+	
-	-	-	-	±	
-	-	-	-	+	
+	-	-	-	+	

August 20, 1949.

9P20. Inoc heavily in Pennessay:

	L+M-	L-M+	
D. #	W 1084	* W 1059	(Mfr x Mfr)
E.	W 1084	+ W 842 W 842	(Mfr x +)

1) plated P22. (ca 40 hours).

D Malac 10 plates ¹⁰⁰ 10- (all ^{two} ~~one~~ plates!)

E 30/ No -

2) Plated 123.

Yield very low 0-2-16 / plate
No -

3

Plated P24 10⁻⁶ ml 5 plates Malac each.

D No Malac - // 400 cols.
E 1? or No Malac - / 600 cols.

(Crater Lat. : Mal-??)

August 16th 1949

	Mal:		Plates (sample)	lac:		
	+	-		+	-	
W1073 x	+	-		+	-	
A. W1033	29	24	0	15	25	1
B. 58-161	28	61	4	45	77	9
C. K12.	22	26	1	12	27	3

Incubated together 48 hours in 1/2 of Punnassay. Plated on 10 plates each of lac and Mal EMB. Possible sectored colonies streaked out on homologous medium.

[sectored colonies in 10 plates in table above ↑] ca 10+ests each
-, +.

A: lac^v. L+M+; L-M-

B. 9 lac^v All: L+M+; L-M-

4 Mal^v. 4/5: L+M+ L-M-

C. 3 lac^v All L+M+ L-M-

1 Mal^v ~~L+M+ L-M-~~
L+M- L-M-

No Recombination!

Plate 122. 10 lac each. Cultures are badly changed.

A) Ca 25 ea. + > - 3 lac^v

B) Ca 2 ea.

C) Ca 150 ea.

10 colonies on EMS: all +.

5 lac^v.

→ #3 all lac+Mal+

Plated P23. Yields v. low

10 Lac, 5 Mal ca.

2. A) ca 20/
3 Lac v.

High variance of +/- points to clumping
but lac+ >> -

B) ca 5/
No v.

C) ca 2/

Edary yield too low for any conclusions

3 Plate P24 ca 10^{-6} ml./plate

10 Lac 10 Mal ser. ca 200/ for A, B ca 100/ C.

A 3 Lac v 4 Mal v

B 3 Lac v 11 Mal v

C 3 Lac v

} All parentals only

August 25, 1949.

In NB mix hairy as follows:

Plate 10^{-7} ml: 8/27.

A	1033 x 677	Lac 15	Mal 10	
B	1059 x 814	Lac 10	Malac 5	
C	1059 x 1084	Lac 10	Malac 15	Mal Mal 10
D	1073 x 1033	Lac 15	Malac 1	
E	1073 x 58161	Lac 15		
F	1073 x 10 K12	Lac 15	EMS Lac 5 (2 drops)	

70

A ca 80/ lact >> - (20:1)

B ca 30/ No Malac -

C ca 20-50/ No Malac - Mal - = Lac -

D ca 200/ lact >> Lac -

E ca 150/

F ca 100/

A.	1 Malv	0 Lacv	/ 800; 1200 resp. [Note excess of + > -].
B.	4 Lacv		/ 500. 0 Malac / 150
C.	2 Lacv		/ 400. 0 Malac - / 600.
D.	3 Lacv?		/ 2000
E.	8 Lacv		/ 2200
F.	2 Lacv		/ 1500 EMS: No -

		L+	L-							
D	3.	4M+	10L-?							
	1	10M+								
	2									
E	1	10M+	10M-							
	2	7M+	5M-							
	3	10M+	10M-							
	4	10M+	10M-							
	5	10M+	10M-							
	6	9M+	6M-							
	7	5M+	10M-							
	8	10M+	10M-							
F	1	7M+	10M-							
	2	10M+	5M-							

~~No Recombination~~