

Strains taken to L. S. H.

LT 2 = Lilleengen 85  
LT 7 = " 578  
LT22 = " 409  
SY79 = (S. typhi Watson, from Kauffmann)  
S 57 = (S. typhi H901 W. Edwards, circular 54.)  
SW 351 = LT22, (D al. + tyr.)-, tryp.-, gal-, xyl-  
SW 414 = LT2 meth.-, hist.-.  
SW 435 = SW351, SR (500 units)  
SW 520 = SY79 + FA. LT7. (H-1)  
SW 532 = SY79 + FA. LT2 (Rhamnose)  
SW 534 = O from SW 533 (phage)  
SW 536 = O from SW 535 (phage)  
SW 537 = S. typhi H901 Boulgakoff  
SW 538 = O from SW 537 (phage)  
SW 539 = S. typhi 383 Boulgakoff  
SW 540 = S. typhi rough Boulgakoff  
SW 541 = S. t-m copenhagen O (F.K.)  
SW 542 = S. typhi O.901 (F.K. 58)  
SW 543 = S. para B O (F.K. 248)  
SW 544 = S. t-m O (F.K. 13)  
SW 545 = S. t-m O 4937/50 Edwards  
SW 546 = S. para B phase II N-25 Edwards  
SW 547 = S. t-m phase II 191 Edwards  
SW 548 = S. group B, O 2294/49 "  
SW 549 = S. group B, O 117/51 "  
SW 552 = S. group D, O 1524/51 "  
SW 553 = S. group D, O 1520/51 "  
SW 558 = S. gallinarum 30953 "  
SW 563 = SW553 + FA. LT2, motile (s.v.)  
SW 569 = SW537 (H.901) + FA. LT2, motile (1)  
SW 570 = SW537 (H.901) + FA. LT2, " "  
SW 572 = Motile mutant of SW 543 (1)  
and SW 573 to SW 656.