

Mental Health: A Report of the Surgeon General

services, and, in some cases, medical care. In short, more resources must be devoted to programs that integrate mental health rehabilitative services into long-term care in both community and institutional settings.

Financing Services for Older Adults

Financing policies furnish incentives that favor utilization of some services over others (e.g., nursing homes rather than state mental hospitals) or preclude the provision of needed services (e.g., mental health services in nursing homes). Details on financing and organizing mental health services, with a special focus on access, are presented in Chapter 6. Selected issues germane to older adults are addressed here.

Historically, Federal financing policy has imposed special limits on reimbursement for mental health services. Medicaid precluded payment for care in so-called "institutions for mental diseases," Medicaid's term for mental hospitals and the small percentage of nursing homes with specialized mental health services. This Medicaid policy provided a disincentive for the majority of nursing homes to specialize in delivering mental health services for fear of losing Medicaid payments (Taube et al., 1990). Under Medicare, the most salient limits were higher copayments for outpatient mental health services and a limited number of days for hospital care. Medicare's special limits on outpatient mental health services were changed over the past decade, resulting in significantly increased access to and utilization of such services (Goldman et al., 1985; Rosenbach & Ammering, 1997). The concern, however, is that the gains made as a result of policy changes easily could be eroded by the shift to managed care (Rosenbach & Ammering, 1997).

Increased Role of Managed Care

Projections are that 35 percent of all Medicare beneficiaries will be in managed care plans by the year 2007, amounting to approximately 15.3 million people (Komisar et al., 1997). Although the managed care industry has the potential to provide a range of integrated services for people with long-term care needs, managed care's awareness of and response to

chronic care are rudimentary (Institute for Health and Aging, 1996). Despite the potential of systems of managed health care, such as HMOs, to provide comprehensive preventive, acute, and chronic care services, their current specialized geriatric programs and clinical case management for older persons tend to be inadequate or poorly implemented (Friedman & Kane, 1993; Pacala et al., 1995; Kane et al., 1997). In addition, older patients are likely to be poorly served in primary care settings (including primary care HMOs) because of minimal use of specialty providers and suboptimal pharmacological management (Bartels et al., 1997). Further, current systems lack the array of community support, residential, and rehabilitative services necessary to meet the needs of older persons with more severe mental disorders (Knight et al., 1995). These shortcomings are unlikely to be remedied until more research becomes available demonstrating cost-effective models for treating older people with mental illness.

Carved-In Mental Health Services for Older Adults

The types of mental health services available within managed care organizations vary greatly with respect to how services are provided. In some organizations, mental health care is directly integrated into the package of general health care services ("carved-in" mental health services), while it is provided in others through a contract with a separate specialty mental health organization that provides only these services and accepts the financial risk ("carved-out" mental health services).

Proponents of carved-in mental health services argue that this model better integrates physical and mental health care, decreases barriers to mental health care due to stigma, and is more likely to produce cost-offsets and overall savings in general health care expenditures. These features are particularly relevant to older persons, as they commonly have comorbid somatic disorders for which they take multiple medications that may affect mental disorders, often avoid specialty mental health settings, and incur significant health care expenses related to psychiatric

symptoms (George, 1992; Paveza & Cohen, 1996; Moak, 1996; Riley et al., 1997). Unfortunately, mental health specialty services for older persons tend to be a low priority in managed health care organizations, by comparison with medical or surgical specialty services (Bartels et al., 1997). More importantly, carved-in mental health care may have superior potential for individuals with diagnoses such as minor depression and anxiety disorders but tends to shortchange older patients with SPMD who require intensive and long-term mental health care (Mechanic, 1998). The range of outreach, rehabilitative, residential, and intensive services needed for patients with SPMD is likely to exceed the capacity, expertise, and investment of most general health care providers.

Economic factors also may limit the usefulness of mental health carve-ins in serving the needs of older individuals with SPMD. First, evidence from private sector health plans suggests that without mandated parity, insurers offer inferior coverage of mental health care (Frank et al., 1997b, 1997c). Furthermore, if providers or payers compete for enrollees, there is strong incentive to avoid enrollees expected to have higher costs from mental health problems (e.g., older persons with SPMD). To avoid such discrimination, equal coverage of mental health care would have to be mandated through legislation on mental health parity or through specialized contract requirements with managed care organizations.

Carved-Out Mental Health Services for Older Adults

Proponents of mental health service carve-outs for older persons argue that separate systems of financing and services are likely to be superior for individuals needing specialty mental health services, especially those with SPMD. In particular, advocates suggest that carved-out mental health organizations have superior technical knowledge, specialized skills, a broader array of services, greater numbers and varieties of mental health providers with experience treating severe mental disorders, and a willingness and commitment to service high-risk populations (Riley et al., 1997). From an economic perspective, since competition is largely over

the carve-out contract with the payer (generally a public organization or an employer), there is less incentive to compete on risk selection, and risk adjustment becomes unnecessary. In addition, mental health carve-out organizations may be better equipped to provide rehabilitative and community support mental health services necessary to care for older persons with SPMD. Finally, growth of innovative outpatient alternatives could be stimulated by reinvestment of savings by the payer from any decrease in inpatient service use.

Unfortunately, research is lacking on outcomes and costs for older persons with SPMD in mental health carve-outs. A carve-out arrangement could lead to adverse clinical outcomes in older patients due to fragmentation of medical and mental health care services in a population with high risk of complications of comorbidity and polypharmacy. Also, from a financial perspective, the combination of physical and mental comorbidities seen in older adults, especially those with SPMD, may reduce the economic advantages of carved-out services (Bazemore, 1996; Felker et al., 1996; Tsuang & Woolson, 1997). If the provider cannot appropriately manage services and costs associated with the combination of somatic and mental health disorders, anticipated savings may not materialize. Furthermore, fragmentation of reimbursement streams would likely complicate the assessment of cost-effectiveness or cost-offsets. For example, apparent savings of mental health carve-outs under Medicare actually may be due to shifting costs when an individual is also covered under Medicaid. In this situation, Medicaid may cover prescription drugs, long-term care, and other services that are not paid for by Medicare. In order to offer true efficiencies, Medicare mental health carve-outs need to find a way to bridge the fragmentation of financing care for older persons.

Outcomes Under Managed Care

There do not appear to be any studies of mental health outcomes for older adults under managed care. In general, the available research on mental health outcomes for other adults consistently finds that

Mental Health: A Report of the Surgeon General

managed care is successful at reducing mental health care costs (Busch, 1997; Sturm, 1997), yet clinical outcomes (especially for the most severely and chronically ill) are mixed and difficult to interpret due to differences in plans and populations served. Several studies suggest that outcomes under managed care for younger adults are as favorable as, or better than, those under fee-for-service (Lurie et al., 1992; Cole et al., 1994). In contrast, others report that the greater use of nonspecialty services for mental health care under managed care is associated with less cost-effective care (Sturm & Wells, 1995), and that older and poor chronically ill patients may have worse health outcomes or outcomes that vary substantially by site and patient characteristics (Ware et al., 1996). A recent review of health outcomes for both older and younger adults in the managed care literature (Miller & Luft, 1997) concluded that there were no consistent patterns that suggested worse outcomes. However, negative outcomes were more common in patients with chronic conditions, those with diseases requiring more intensive services, low-income enrollees in worse health, impaired or frail elderly, or home health patients with chronic conditions and diseases. These risk factors apply to older adults with SPMD, suggesting that this group is at high risk for poor outcomes under managed care programs that lack specialized long-term mental health and support services. To definitively address the question of mental health outcomes for older persons under managed care, appropriate outcome measures for older adults with mental illness will need to be developed and implemented in the evolving health care delivery systems (Bartels et al., in press).

Other Services and Supports

Older adults and their families depend on a multiplicity of supports that extend beyond the health and mental health care systems. Patients and caregivers need access to education, support networks, support and self-help groups, respite care, and human services, among other supports (Scott-Lennox & George, 1996). These services assume heightened importance for older people who are living alone, who are uncomfortable with formal mental health services, or who are

inadequately treated in primary care. Services and supports appear to be instrumental not only for the patient but also for the family caregiver, as this section explains, but research on their efficacy is sparse. The strongest evidence surrounds the efficacy of services for family caregivers. Support for family caregivers is crucial for their own health and mental health, as well as for controlling the high costs of institutionalization of the family member in their care. The longer the patient remains home, the lower the total cost of institutional care for those who eventually need it.

Support and Self-Help Groups

Support groups, which are an adjunct to formal treatment, are designed to provide mutual support, information, and a broader social network. They can be professionally led by counselors or psychologists, but when they are run by consumers⁸ or family members, they are known as self-help groups. The distinction is somewhat clouded by the fact that mental health professionals and community organizations often aid self-help groups with logistical support, start-up assistance, consultation, referrals, and education (Waters, 1995). For example, self-help support groups sponsored by the Alzheimer's Association use professionals to provide consultation to groups orchestrated by lay leaders.

Support groups for people with mental disorders and their families have been found helpful for adults (see Chapter 4). Participation in support groups, including self-help groups, reduces feelings of isolation, increases knowledge, and promotes coping efforts. What little research has been conducted on older people is generally positive but has been limited mostly to caregivers (see later section) and widows (see below), rather than to older people with mental disorders.

Despite the scant body of research, there is reason to believe that support and self-help group participation is as beneficial, if not more beneficial, for older people with mental disorders. Older people tend to live alone

⁸ Consumers are people engaged in and served by mental health services.

and to be more socially isolated than are other people. They also are less comfortable with formal mental health services. Therefore, social networks established through support and self-help groups are thought to be especially vital in preventing isolation and promoting health. Support programs also can help reduce the stigma associated with mental illness, to foster early detection of illnesses, and to improve compliance with formal interventions.

Earlier sections of this chapter documented the untoward consequences of prolonged bereavement: severe emotional distress, adjustment disorders, depression, and suicide. Outcomes have been studied for two programs of self-help for bereavement. One program, They Help Each Other Spiritually (THEOS), had robust effects on those who were more active in the program. Those widows and widowers displayed the improvements on health measures such as depression, anxiety, somatic symptoms, and self-esteem (Lieberman & Videka-Sherman, 1986). The other program, Widow to Widow: A Mutual Health Program for the Widowed, was developed by Silverman (1988). The evaluation in a controlled study found program participants experienced fewer depressive symptoms and recovered their activities and developed new relationships more quickly (Vachon, 1979; Vachon et al., 1980, 1982).

Education and Health Promotion

There is a need for improved consumer-oriented public information to educate older persons about health promotion and the nature of mental health problems in aging. Understanding that mental health problems are not inevitable and immutable concomitants of the aging process, but problems that can be diagnosed, treated, and prevented, empowers older persons to seek treatment and contributes to more rapid diagnosis and better treatment outcomes.

With respect to health promotion, older persons also need information about strategies that they can follow to maintain their mental health. Avoiding disease and disability, sustaining high cognitive and physical function, and engaging with life appear to be

important ways to promote mental and physical health (Rowe & Kahn, 1997). The two are interdependent.

Established programs for health promotion in older people include wellness programs, life review, retirement, and bereavement groups (see review by Waters, 1995). Although controlled evaluations of these programs are infrequent, bereavement and life review appear to be the best studied. Bereavement groups produce beneficial results, as noted above, and life review has been found to produce positive outcomes in terms of stronger life satisfaction, psychological well-being, self-esteem, and less depression (Haight et al., 1998). Life review also was investigated through individualized home visits to homebound older people in the community who were not depressed but suffered chronic health conditions. Life review for these older people was found to improve life satisfaction and psychological well-being (Haight et al., 1998).

Another approach to promoting mental health is to develop a “social portfolio,” a program of sound activities and interpersonal relationships that usher individuals into old age (Cohen, 1995b). While people in the modern work force are advised to plan for future economic security—to strive for a balanced financial portfolio—too little attention is paid to developing a balanced social portfolio to help to plan for the future. Ideally, such a program will balance *individual* with *group* activities and *high mobility/energy* activities requiring significant physical exertion with *low mobility/energy* ones. The social portfolio is a mental health promotion strategy for helping people develop new strengths and satisfactions.

Families and Caregivers

Among the many myths about aging is that American families do not care for their older members. Such myths are based on isolated anecdotes as opposed to aggregate data. Approximately 13 million caregivers, most of whom are women, provide unpaid care to older relatives (Biegel et al., 1991). Families are committed to their older members and provide a spectrum of assistance, from hands-on to monetary help (Bengston et al., 1985; Sussman, 1985; Gatz et al., 1990; Cohen,

Mental Health: A Report of the Surgeon General

1995a). Problems occur with older individuals who have no children or spouse, thereby reducing the opportunity to receive family aid. Problems also occur with the "old-old," those over 85 whose children are themselves old and, therefore, unable to provide the same intensity of hands-on help that younger adult children can provide. These special circumstances highlight the need for careful attention to planning for mental health service delivery to older individuals with less access to family or informal support systems.

Conversely, a large and growing number of older family members care for chronically mentally ill and mentally retarded younger adults (Bengston et al., 1985; Gatz et al., 1990; Eggebeen & Wilhelm, 1995). Too little is known about ways to help the afflicted younger individuals and their caregiving parents. Families are eager to help themselves, and society needs to find ways to better enable them to do so.

There is a great need to better educate families about what they can do to help promote mental health and to prevent and treat mental health problems in their older family members. Families fall prey to negative stereotypes that little can be done for late-life mental health problems. They need to know that mental health problems in later life, like physical health problems, can be treated. They need to understand how to better recognize symptoms or signals of impending mental health problems among older adults so that they can help their loved ones receive early interventions. They need to know what services are available, where they can be found, and how to help their older relatives access such help when necessary.

The plight of family caregivers is pivotal. As noted earlier, the burden of caring for an older family member places caregivers at risk for mental and physical disorders. Virtually all studies find elevated levels of depressive symptomatology among caregivers, and those using diagnostic interviews report high rates of clinical depression and anxiety (Schultz et al., 1995). Ensuring their mental and physical health is not only vital for their well-being but also is vital for the older people in their care. Support groups and services aimed at caregivers can improve their health and quality of

life, can improve management of patients in their care, and can delay their institutionalization.

Communities and Social Services

Family support is often supplemented by enduring long-term relationships between older people and their neighbors and community, including religious, civic, and public organizations (Scott-Lennox & George, 1996). Linkages to these organizations instill a sense of belonging and companionship. Such linkages also provide a safety net, enabling some older people to live independently in spite of functional decline.

While the vast majority of frail and homebound older people receive quality care at home, abuse does occur. Estimates vary, but most studies find rates of abuse by caregivers (either family or nonfamily members) to range up to 5 percent (Coyne et al., 1993; Scott-Lennox & George, 1996). Abuse is generally defined in terms of being either physical, psychological, legal, or financial. The abuse is most likely to occur when the patient has dementia or late-life depression, conditions that impart relatively high psychological and physical burdens on caregivers (Coyne et al., 1993). A recent report by the Institute of Medicine describes the range of interventions for protection against abuse of older people, including caregiver participation in support groups and training programs for behavioral management (especially for Alzheimer's disease) and social services programs (e.g., adult protective services, casework, advocacy services, and out-of-home placements). While there are very few controlled evaluations of these services (IOM, 1998), communities need to ensure that there are programs in place to prevent abuse of older people. Programs can incorporate any of a number of effective psychosocial and support interventions for patients with Alzheimer's disease and their caregivers—interventions that were presented earlier in this section and the section on Alzheimer's disease.

Communities need to ensure the availability of adult day care and other forms of respite services to aid caregivers striving to care for family members at home. They also can provide assistance to self-help and other support programs for patients and caregivers. In the

process of facilitating or providing services, communities need to consider the diversity of their older residents—racial and ethnic diversity, socioeconomic diversity, diversity in settings where they live, and diversity in levels of general functioning. Such diversity demands comprehensive program planning, information and referral services (including directories of what is available in the community), strong outreach initiatives, and concerted ways to promote accessibility. Moreover, each component of the community-based delivery system targeting older adults should incorporate a clear focus on mental health. Too often, attention to mental health services for older people and their caregivers is negligible or absent, despite the fact, as noted earlier, that mental health problems and caregiver distress are among the leading reasons for institutionalization (Lombardo, 1994). Important life tasks remain for individuals as they age. Older individuals continue to learn and contribute to society, in spite of physiologic changes due to aging and increasing health problems.

Conclusions

1. Important life tasks remain for individuals as they age. Older individuals continue to learn and contribute to the society, in spite of physiologic changes due to aging and increasing health problems.
2. Continued intellectual, social, and physical activity throughout the life cycle are important for the maintenance of mental health in late life.
3. Stressful life events, such as declining health and/or the loss of mates, family members, or friends often increase with age. However, persistent bereavement or serious depression is not “normal” and should be treated.
4. Normal aging is not characterized by mental or cognitive disorders. Mental or substance use disorders that present alone or co-occur should be recognized and treated as illnesses.
5. Disability due to mental illness in individuals over 65 years old will become a major public health problem in the near future because of demographic changes. In particular, dementia, depression, and

schizophrenia, among other conditions, will all present special problems in this age group:

- a. Dementia produces significant dependency and is a leading contributor to the need for costly long-term care in the last years of life;
 - b. Depression contributes to the high rates of suicide among males in this population; and
 - c. Schizophrenia continues to be disabling in spite of recovery of function by some individuals in mid to late life.
6. There are effective interventions for most mental disorders experienced by older persons (for example, depression and anxiety), and many mental health problems, such as bereavement.
 7. Older individuals can benefit from the advances in psychotherapy, medication, and other treatment interventions for mental disorders enjoyed by younger adults, when these interventions are modified for age and health status.
 8. Treating older adults with mental disorders accrues other benefits to overall health by improving the interest and ability of individuals to care for themselves and follow their primary care provider’s directions and advice, particularly about taking medications.
 9. Primary care practitioners are a critical link in identifying and addressing mental disorders in older adults. Opportunities are missed to improve mental health and general medical outcomes when mental illness is underrecognized and undertreated in primary care settings.
 10. Barriers to access exist in the organization and financing of services for aging citizens. There are specific problems with Medicare, Medicaid, nursing homes, and managed care.

References

- Aarts, P., & Op den Velde, W. (1996). Prior traumatization and the process of aging. In B. A. van der Kolk, A. C. McFarlane, & L. Weisath (Eds.), *Traumatic stress: The effects of overwhelming experience on mind, body and society* (pp. 359–377). New York: Guilford Press.

Mental Health: A Report of the Surgeon General

- Abrams, R. C., Rosendahl, E., Card, C., & Alexopoulos, G. S. (1994). Personality disorder correlates of late and early onset depression. *Journal of the American Geriatrics Society*, *42*, 727-731.
- Adams, W. L., Garry, P. J., Rhyne, R., Hunt, W. C., & Goodwin, J. S. (1990). Alcohol intake in the healthy elderly. Changes with age in a cross-sectional and longitudinal study. *Journal of the American Geriatrics Society*, *38*, 211-216.
- Aevarsson, O., & Skoog, I. (1996). A population-based study on the incidence of dementia disorders between 85 and 88 years of age. *Journal of the American Geriatrics Society*, *44*, 1455-1460.
- Aiken, L. (1990). Chronic mental illness. In B. S. Fogel, A. Furino, & G. Gottlieb (Eds.), *Mental health policy for older Americans: Protecting minds at risk*. Washington, DC: American Psychiatric Press.
- Albert, M. S., Jones, K., Savage, C. R., Berkman, L., Seeman, T., Blazer, D., & Rowe, J. W. (1995). Predictors of cognitive change in older persons: MacArthur studies of successful aging. *Psychology and Aging*, *10*, 578-589.
- Alexopoulos, G. S. (1997, November 6). *Epidemiology, nosology and treatment of geriatric depression*. Paper presented at Exploring Opportunities to Advance Mental Health Care for an Aging Population, meeting sponsored by the John A. Hartford Foundation, Rockville, MD.
- Alexopoulos, G. S., & Abrams, R. C. (1991). Depression in Alzheimer's disease. *Psychiatric Clinics of North America*, *14*, 327-340.
- Alexopoulos, G. S., & Chester, J. G. (1992). Outcomes of geriatric depression. *Clinics in Geriatric Medicine*, *8*, 363-376.
- Alexopoulos, G. S., Meyers, B. S., Young, R. C., Campbell, S., Silbersweig, D., & Charlson, M. (1997). "Vascular depression" hypothesis. *Archives of General Psychiatry*, *54*, 915-922.
- Alexopoulos, G. S., Meyers, B. S., Young, R. C., Kakuma, T., Feder, M., Einhorn, A., & Rosendahl, E. (1996). Recovery in geriatric depression. *Archives of General Psychiatry*, *53*, 305-312.
- Alexopoulos, G. S., & Salzman, C. (1998). Treatment of depression with heterocyclic antidepressants, monoamine oxidase inhibitors, and psychomotor stimulants. In C. Salzman (Ed.), *Clinical geriatric psychopharmacology* (pp. 184-244). Baltimore: Williams & Wilkins.
- American Association for Geriatric Psychiatry. (1997). *Recommendations from primary care physicians: When to refer depressed elderly patients to a geriatric psychiatrist*. Bethesda, MD: American Association for Geriatric Psychiatry.
- American Psychiatric Association. (1980). *Diagnostic and statistical manual of mental disorders* (3rd ed.). Washington, DC: American Psychiatric Press.
- American Psychiatric Association. (1993). Practice guidelines for major depressive disorders in adults. *American Journal of Psychiatry*, *150*, 1-26.
- American Psychiatric Association. (1994). *Diagnostic and statistical manual of mental disorders* (4th ed.). Washington, DC: Author.
- Andersen, K., Launer, L. J., Ott, A., Hoes, A. W., Breteler, M. M., & Hofman, A. (1995). Do nonsteroidal anti-inflammatory drugs decrease the risk for Alzheimer's disease? The Rotterdam Study. *Neurology*, *45*, 1441-1445.
- Anderson, G. M., Kerluke, K. J., Pulcins, I. R., Hertzman, C., & Barer, M. L. (1993). Trends and determinants of prescription drug expenditures in the elderly: Data from the British Columbia Pharmacare Program. *Inquiry*, *30*, 199-207.
- Anthony, J. C., & Helzer, J. E. (1991). Syndromes of drug abuse and dependence. In L. N. Robins & D. A. Regier (Eds.), *Psychiatric disorders in America: The Epidemiologic Catchment Area study* (pp. 116-154). New York: Free Press.
- Arean, P. A., Perri, M. G., Nezu, A. M., Schein, R. L., Christopher, F., & Joseph, T. X. (1993). Comparative effectiveness of social problem-solving therapy and reminiscence therapy as treatments for depression in older adults. *Journal of Consulting and Clinical Psychology*, *61*, 1003-1010.
- Atkinson, R. M., Ganzini, L., & Bernstein, M. J. (1992). Alcohol and substance-use disorders in the elderly. In J. E. Birren, B. Sloane, & G. D. Cohen (Eds.), *Handbook of mental health and aging* (2nd ed., pp. 515-555). San Diego: Academic Press.
- Bachman, D. L., Wolf, P. A., Linn, R., Knoefel, J. E., Cobb, J., Belanger, A., D'Agostino, R. B., & White, L. R. (1992). Prevalence of dementia and probable senile dementia of the Alzheimer type in the Framingham Study. *Neurology*, *42*, 115-119.
- Bailey, M. B., Haberman, P. W., & Alksne, H. (1965). The epidemiology of alcoholism in an urban residential area. *Quarterly Journal of Studies on Alcohol*, *26*, 19-40.

- Baldwin, R. C., & Tomenson, B. (1995). Depression in later life. A comparison of symptoms and risk factors in early and late onset cases. *British Journal of Psychiatry*, *167*, 649–652.
- Baltes, P. B., Sowarka, D., & Kliegl, R. (1989). Cognitive training research on fluid intelligence in old age: What can older adults achieve by themselves? *Psychology and Aging*, *4*, 217–221.
- Barbone, F., McMahon, A. D., Davey, P. G., Morris, A. D., Reid, I. C., McDevitt, D. G., & MacDonald, T. M. (1998). Association of road-traffic accidents with benzodiazepine use. *Lancet*, *352*, 1331–1336.
- Barinaga, M. (1998). Alzheimer's treatments that work now. *Science*, *282*, 1030–1032.
- Bartels, S. J., & Colenda, C. C. (1998). Mental health services for Alzheimer's disease. Current trends in reimbursement and public policy, and the future under managed care. *American Journal of Geriatric Psychiatry*, *6*, S85–S100.
- Bartels, S. J., Horn, S., Sharkey, P., & Levine, K. (1997). Treatment of depression in older primary care patients in health maintenance organizations. *International Journal of Psychiatry in Medicine*, *27*, 215–231.
- Bartels, S. J., & Levine, K. J. (1998). Meeting the needs of older adults with severe and persistent mental illness: Public policy in an era of managed and long-term reform. *Public Policy and Aging Report*, *9*, 1–6.
- Bartels, S. J., Levine, K. J., & Shea, D. (in press). Mental health long-term care for older persons with severe and persistent mental disorders in an era of managed care. *Psychiatric Services*.
- Bass, S. A. (Ed.). (1995). *Older and active: How Americans over 55 are contributing to society*. New Haven: Yale University Press.
- Baum, C., Kennedy, D. L., Forbes, M. B., & Jones, J. K. (1984). Drug use in the United States in 1981. *Journal of the American Medical Association*, *251*, 1293–1297.
- Bazemore, P. H. (1996). *Medical problems of the seriously and persistently mentally ill*. Seattle: Hogrefe and Huber.
- Beck, C., Heacock, P., Mercer, S. O., Walls, R. C., Rapp, C. G., & Vogelpohl, T. S. (1997). Improving dressing behavior in cognitively impaired nursing home residents. *Nursing Research*, *46*, 126–132.
- Beck, J. G., & Stanley, M. A. (1997). Anxiety disorder in the elderly: The emerging role of behavior therapy. *Behavior Therapy*, *28*, 83–100.
- Beekman, A. T., Deeg, D. J., van Tilburg, T., Smit, J. H., Hooijer, C., & van Tilburg, W. (1995). Major and minor depression in later life: A study of prevalence and risk factors. *Journal of Affective Disorders*, *36*, 65–75.
- Behl, C., Widmann, M., Trapp, T., & Holsboer, F. (1995). 17-beta estradiol protects neurons from oxidative stress-induced cell death in vitro. *Biochemical and Biophysical Research Communications*, *216*, 473–482.
- Bengston, V. L., Cutler, N. E., & Mangen, D. J., & Marshall, V. W. (1985). Generations, cohorts and relations between age groups. In R. H. Binstock & E. Shanas (Eds.), *Handbook of aging and social sciences* (2nd ed., pp. 415–449). New York: Van Nostrand Reinhold.
- Bernstein, M. A., & Hensley, R. (1993). Developing community-based program alternatives for the seriously and persistently mentally ill elderly. *Journal of Mental Health Administration*, *20*, 201–207.
- Beutler, L. E., Scogin, F., Kirkish P., Schretlen, D., Corbishley, A., Hamblin, D., Meredith, K., Potter, R., Bamford, C. R., & Levenson, A. I. (1987). Group cognitive therapy and alprazolam in the treatment of depression in older adults. *Journal of Consulting and Clinical Psychology*, *55*, 550–556.
- Biegel, D. E., Sales, E., & Schulz, R. (1991). *Family caregiving in chronic illness*. Newbury Park, CA: Sage Publications.
- Bland, R. C., Newman, S. C., & Orn, H. (1988). Prevalence of psychiatric disorders in the elderly in Edmonton. *Acta Psychiatrica Scandinavica Supplement*, *338*, 57–63.
- Blazer, D. (1989). Depression in the elderly. *New England Journal of Medicine*, *320*, 164–166.
- Blazer, D. G. (1996a). Alcohol and drug problems. In E. W. Busse & D. G. Blazer (Eds.), *The American Psychiatric Press textbook of geriatric psychiatry* (2nd ed., pp. 341–359). Washington, DC: American Psychiatric Press.
- Blazer, D. G. (1996b). Epidemiology of psychiatric disorders in late life. In E. W. Busse & D. G. Blazer (Eds.), *The American Psychiatric Press textbook of geriatric psychiatry* (2nd ed., pp. 155–171). Washington, DC: American Psychiatric Press.
- Bodick, N. C., Offen, W. W., Levey, A. I., Cutler, N. R., Gauthier, S. G., Satlin, A., Shannon, H. E., Tollefson, G. D., Rasmussen, K., Bymaster, F. P., Hurley, D. J., Potter, W. Z., & Paul, S. M. (1997). Effects of xanomeline, a selective muscarinic receptor agonist, on cognitive function and behavioral symptoms in Alzheimer disease. *Archives of Neurology*, *54*, 465–473.

Mental Health: A Report of the Surgeon General

- Bohm, C., Robinson, D. S., Gammans, R. E., Shrotriya, R. C., Alms, D. R., Leroy, A., & Placchi, M. (1990). Buspirone therapy in anxious elderly patients: A controlled clinical trial. *Journal of Clinical Psychopharmacology*, *10*, 47S–51S.
- Borkan, G. A., Hults, D. E., Gerzof, S. G., Robbins, A. H., & Silbert, C. K. (1983). Age changes in body composition revealed by computed tomography. *Journal of Gerontology*, *38*, 673–677.
- Borson, S. (1995). *Comprehensive textbook of psychiatry*. Baltimore: Williams & Wilkins.
- Borson, S., Barnes, R. A., Kukull, W. A., Okimoto, J. T., Veith, R. C., Inui, T. S., Carter, W., & Raskind, M. A. (1986). Symptomatic depression in elderly medical outpatients. I. Prevalence, demography, and health service utilization. *Journal of American Geriatric Society*, *34*, 341–347.
- Borson, S., & Raskind, M. A. (1997). Clinical features and pharmacologic treatment of behavioral symptoms of Alzheimer's disease. *Neurology*, *48*, S17–S24.
- Branconner, R. J., Cole, J. O., Ghazvinian, S., Spera, K. F., Oxenkrug, G. F., & Bass, J. L. (1983). Clinical pharmacology of bupropion and imipramine in elderly depressives. *Journal of Clinical Psychiatry*, *44*, 130–133.
- Branconner, R. J., DeVitt, D. R., Cole, J. O., & Spera, K. F. (1982). Amitriptyline selectively disrupts verbal recall from secondary memory of the normal aged. *Neurobiology of Aging*, *3*, 55–59.
- Breitner, J. C. (1991). Clinical genetics and genetic counseling in Alzheimer disease. *Annals of Internal Medicine*, *115*, 601–606.
- Breteler, M. M., Claus, J. J., van Duijn, C. M., Launer, L. J., & Hofman, A. (1992). Epidemiology of Alzheimer's disease. *Epidemiology Review*, *14*, 59–82.
- Broadhead, W. E., Blazer, D. G., George, L. K., & Tse, C. K. (1990). Depression, disability days, and days lost from work in a prospective epidemiologic survey. *Journal of the American Medical Association*, *264*, 2524–2528.
- Brody, D. S., Khaliq, A. A., & Thompson, T. L., Jr. (1997). Patients' perspectives on the management of emotional distress in primary care settings. *Journal of General Internal Medicine*, *12*, 403–406.
- Bruce, M. L., & Hoff, R. A. (1994). Social and physical health risk factors for first-onset major depressive disorder in a community sample. *Social Psychiatry and Psychiatric Epidemiology*, *29*, 165–171.
- Bruce, M. L., Kim, K., Leaf, P. J., & Jacobs, S. (1990). Depressive episodes and dysphoria resulting from conjugal bereavement in a prospective community sample. *American Journal of Psychiatry*, *147*, 608–611.
- Burgio, L. D., Engel, B. T., Hawkins, A., McCormick, K., Scheve, A., & Jones, L. T. (1990). A staff management system for maintaining improvements in continence with elderly nursing home residents. *Journal of Applied Behavior Analysis*, *23*, 111–118.
- Burgio, L., Scilley, K., Hardin, J. M., Hsu, C., & Yancey, J. (1996). Environmental "white noise": An intervention for verbally agitated nursing home residents. *Journals of Gerontology. Series B, Psychological Sciences and Social Sciences*, *51*, P364–P373.
- Burns, B. J. (1991). Mental health services research on the hospitalized and institutionalized CMI elderly. In E. Light & B. D. Lebowitz (Eds.), *The elderly with chronic mental illness* (pp. 207–215). New York: Springer.
- Busch, S. (1997). Carving-out mental health benefits to Medicaid beneficiaries: A shift toward managed care. *Administration and Policy in Mental Health*, *24*, 301–321.
- Butler, R. N. (1974). Successful aging and the role of the life review. *Journal of the American Geriatrics Society*, *22*, 529–535.
- Butler, R. N., Lewis, M. I., & Sunderland, T. (1991). *Aging and mental health: Positive psychosocial and biomedical approaches*. Columbus, OH: Charles E. Merrill.
- Buysse, D. J., Reynolds, C. F., III, Hoch, C. C., Houck, P. R., Kupfer, D. J., Mazumdar, S., & Frank, E. (1996). Longitudinal effects of nortriptyline on EEG sleep and the likelihood of recurrence in elderly depressed patients. *Neuropsychopharmacology*, *14*, 243–252.
- Cadoret, R. J., & Widmer, R. B. (1988). The development of depressive symptoms in elderly following onset of severe physical illness. *Journal of Family Practice*, *27*, 71–76.
- Cafferata, G. L., Kasper, J., & Bernstein, A. (1983). Family roles, structure, and stressors in relation to sex differences in obtaining psychotropic drugs. *Journal of Health and Social Behavior*, *24*, 132–143.
- Caine, E. D., Lyness, J. M., & Conwell, Y. (1996). Diagnosis of late-life depression: Preliminary studies in primary care settings. *American Journal of Geriatric Psychiatry*, *4*, S45–S50.

- Callahan, C. M., Hall, K. S., Hui, S. L., Musick, B. S., Unverzagt, F. W., & Hendrie, H. C. (1996a). Relationship of age, education, and occupation with dementia among a community-based sample of African Americans. *Archives of Neurology*, *53*, 134–140.
- Callahan, C. M., Hendrie, H. C., Nienaber, N. A., & Tierney, W. M. (1996b). Suicidal ideation among older primary care patients. *Journal of the American Geriatrics Society*, *44*, 1205–1209.
- Callahan, C. M., Hendrie, H. C., & Tierney, W. M. (1995). Documentation and evaluation of cognitive impairment in elderly primary care patients. *Annals of Internal Medicine*, *122*, 422–429.
- Callahan, C. M., Hui, S. L., Nienaber, N. A., Musick, B. S., & Tierney, W. M. (1994). Longitudinal study of depression and health services use among elderly primary care patients. *Journal of the American Geriatrics Society*, *42*, 833–838.
- Callahan, C. M., Nienaber, N. A., Hendrie, H. C., & Tierney, W. M. (1992). Depression of elderly outpatients: Primary care physicians' attitudes and practice patterns. *Journal of General Internal Medicine*, *7*, 26–31.
- Callahan, C. M., & Wolinsky, F. D. (1995). Hospitalization for major depression among older Americans. *Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, *50*, M196–M202.
- Caramel, V. M., Remarque, E. J., Knook, D. L., Lagaay, A. M., & Van den Brande, K. J. (1998). Benzodiazepine users aged 85 and older fall more often. *Journal of the American Geriatrics Society*, *46*, 1178–1179.
- Carman, M. B. (1997). The psychology of normal aging. *Psychiatric Clinics of North America*, *20*, 15–24.
- Castle, D. J., & Howard, R. (1992). What do we know about the aetiology of late-onset schizophrenia. *European Psychiatry*, *7*, 99–108.
- Castle, D. J., Wessely, S., Howard, R., & Murray, R. M. (1997). Schizophrenia with onset at the extremes of adult life. *International Journal of Geriatric Psychiatry*, *12*, 712–717.
- Centers for Disease Control and Prevention. (1999). *Suicide deaths and rates per 100,000* [On-line]. Available: <http://www.cdc.gov/ncipc/data/us9794/suic.htm>.
- Chrischilles, E. A., Foley, D. J., Wallace, R. B., Lemke, J. H., Semla, T. P., Hanlon, J. T., Glynn, R. J., Ostfeld, A. M., & Guralnik, J. M. (1992). Use of medications by persons 65 and over: Data from the established populations for epidemiologic studies of the elderly. *Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, *47*, M137–M144.
- Cohen D., & Cairl R. (1996). Mental health care policy in an aging society. In B. L. Levin & J. Petrila (Eds.), *Mental health services: A public health perspective* (pp. 301–319). New York: Oxford University Press.
- Cohen, G. D. (1988). *The brain in human aging*. New York: Springer.
- Cohen, G. D. (1990). Lessons from longitudinal studies of mentally ill and mentally healthy elderly: A 17-year perspective. In M. Bergener and S. I. Finkel (Eds.), *Clinical and scientific psychogeriatrics* (Vol. 1, pp. 135–148). New York: Springer.
- Cohen, G. D. (1994). The geriatric landscape: Toward a health and humanities research agenda in aging. *American Journal of Geriatric Psychiatry*, *2*, 185–187.
- Cohen, G. D. (1995a). Intergenerationalism: A new “ism” with positive mental health and social policy potential. *American Journal of Geriatric Psychiatry*, *3*, 1–5.
- Cohen, G. D. (1995b). Mental health promotion in later life: The case for social portfolio. *American Journal of Geriatric Psychiatry*, *3*, 277–279.
- Cohen, G. D. (1998a). Creativity and aging: Ramifications for research, practice, and policy. *Geriatrics*, *53*(Suppl. 1), S4–S8.
- Cohen, G. D. (1998b). Do health science concepts influence care? The case for a new landscape of aging. *American Journal of Geriatric Psychiatry*, *6*, 273–276.
- Cohen, G. D. (in press). Human potential phases in the second half of life: Mental health theory development. *American Journal of Geriatric Psychiatry*.
- Cohen-Mansfield, J., & Werner, P. (1997). Management of verbally disruptive behaviors in nursing home residents. *Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, *52*, M369–M377.
- Cole, M. G., & Bellavance, F. (1997). The prognosis of depression in old age. *American Journal of Geriatric Psychiatry*, *5*, 4–14.

Mental Health: A Report of the Surgeon General

- Cole, R. E., Reed, S. K., Babigian, H. M., Brown, S. W., & Fray, J. (1994). A mental health capitation program: I. Patient outcomes. *Hospital and Community Psychiatry, 45*, 1090–1096.
- Collins, M. W., & Abeles, N. (1996). Subjective memory complaints in the able elderly. *Clinical Gerontologist, 16*, 29–54.
- Constantino, R. E. (1988). Comparison of two group interventions for the bereaved. *Image—The Journal of Nursing Scholarship, 20*, 83–87.
- Conwell, Y. (1996). *Diagnosis and treatment of depression in late life*. Washington, DC: American Psychiatric Press.
- Conwell, Y., Duberstein, P. R., Cox, C., Hermann, J. H., Forbes, N. T., & Caine, E. D. (1996). Relationships of age and axis I diagnoses in victims of completed suicide: A psychological autopsy study. *American Journal of Psychiatry, 153*, 1001–1008.
- Cooper, J. K., Love, D. W., & Raffoul, P. R. (1982). Intentional prescription nonadherence (noncompliance) by the elderly. *Journal of the American Geriatrics Society, 30*, 329–333.
- Cooper-Patrick, L., Crum, R. M., & Ford, D. E. (1994). Characteristics of patients with major depression who received care in general medical and specialty mental health settings. *Medical Care, 32*, 15–24.
- Copeland, J. R., Dewey, M. E., Wood, N., Searle, R., Davidson, I. A., & McWilliam, C. (1987a). Range of mental illness among the elderly in the community. Prevalence in Liverpool using the GMS-AGECAT package. *British Journal of Psychiatry, 150*, 815–823.
- Copeland, J. R., Gurland, B. J., Dewey, M. E., Kelleher, M. J., Smith, A. M., & Davidson, I. A. (1987b). Is there more dementia, depression and neurosis in New York? A comparative study of the elderly in New York and London using the computer diagnosis AGECAT. *British Journal of Psychiatry, 151*, 466–473.
- Corder, E. H., Saunders, A. M., Strittmatter, W. J., Schmechel, D. E., Gaskell, P. C., Small, G. W., Roses, A. D., Haines, J. L., & Pericak-Vance, M. A. (1993). Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science, 261*, 921–923.
- Coyne, A. C., Reichman, W. E., & Berbig, L. J. (1993). The relationship between dementia and elder abuse. *American Journal of Psychiatry, 150*, 643–646.
- Crook, T. (1993). Diagnosis and treatment of memory loss and older patients who are not demented. In Levy, R., Howard R., & Burns A. (Eds.), *Treatment and care in old age psychiatry* (pp. 95–111). New York: Wrightson Biomedical Publishing.
- Cuffel, B. J., Jeste, D. V., Halpain, M., Pratt, C., Tarke, H., & Patterson, T. L. (1996). Treatment costs and use of community mental health services for schizophrenia by age cohorts. *American Journal of Psychiatry, 153*, 870–876.
- Cummings, J. L., Cyrus, P. A., Bieber, F., Mas, J., Orazem, J., & Gulanski, B. (1998a). Metrifonate treatment of the cognitive deficits of Alzheimer's disease. Metrifonate Study Group. *Neurology, 50*, 1214–1221.
- Cummings, J., & Jeste, D. (1999). Alzheimer's disease and its management in the year 2010. *Psychiatric Services, 50*, 1173–1177.
- Cummings, J. L., Vinters, H. V., Cole, G. M., & Khachaturian, Z. S. (1998b). Alzheimer's disease: Etiologies, pathophysiology, cognitive reserve, and treatment opportunities. *Neurology, 51*, S2–S17.
- DeLisi, L. E. (1992). The significance of age of onset for schizophrenia. *Schizophrenia Bulletin, 18*, 209–215.
- Depression Guideline Panel. (1993). *Clinical practice guideline Number 5: Depression in primary care II* (AHCPR Publication No. 93-0551). Rockville, MD: U.S. Department of Health and Human Services.
- Devanand, D. P., Jacobs, D. M., Tang, M. X., Del Castillo-Castaneda, C., Sano, M., Marder, K., Bell, K., Bylsma, F. W., Brandt, J., Albert, M., & Stern, Y. (1997). The course of psychopathologic features in mild to moderate Alzheimer disease. *Archives of General Psychiatry, 54*, 257–263.
- Devor, M., Wang, A., Renvall, M., Feigal, D., & Ramsdell, J. (1994). Compliance with social and safety recommendations in an outpatient comprehensive geriatric assessment program. *Journal of Gerontology, 49*, M168–M173.
- Dew, M. A., Reynolds, C. F., III, Houck, P. R., Hall, M., Buysse, D. J., Frank, E., & Kupfer, D. J. (1997). Temporal profiles of the course of depression during treatment. Predictors of pathways toward recovery in the elderly. *Archives of General Psychiatry, 54*, 1016–1024.
- DSM-IV. See American Psychiatric Association (1994).

- Duara, R., Barker, W. W., Lopez-Alberola, R., Loewenstein, D. A., Grau, L. B., Gilchrist, D., Sevush, S., & St. George-Hyslop, S. (1996). Alzheimer's disease: Interaction of apolipoprotein E genotype, family history of dementia, gender, education, ethnicity, and age of onset. *Neurology*, *46*, 1575–1579.
- Eaton, W. W., Anthony, J. C., Gallo, J., Cai, G., Tien, A., Romanoski, A., Lyketsos, C., & Chen, L. S. (1997). Natural history of Diagnostic Interview Schedule/DSM-IV major depression. The Baltimore Epidemiologic Catchment Area follow-up. *Archives of General Psychiatry*, *54*, 993–999.
- Eaton, W. W., Kramer, M., Anthony, J. C., Dryman, A., Shapiro, S., & Locke, B. Z. (1989). The incidence of specific DIS/DSM-III mental disorders: Data from the NIMH Epidemiologic Catchment Area Program. *Acta Psychiatrica Scandinavica*, *79*, 163–178.
- Esgebeen, D. J., & Wilhelm, M. O. (1995). Patterns of support given by older Americans to their children. In S. A. Bass (Ed.), *Older and active: How Americans over 55 are contributing to society* (pp. 122–168). New Haven: Yale University Press.
- Ekerdt, D. J., De Labry, L. O., Glynn, R. J., & Davis, R. W. (1989). Change in drinking behaviors with retirement: Findings from the normative aging study. *Journal of Studies on Alcohol*, *50*, 347–353.
- Ellor, J. R., & Kurz, D. J. (1982). Misuse and abuse of prescription and nonprescription drugs by the elderly. *Nursing Clinics of North America*, *17*, 319–330.
- Elseviers, M. M., & De Broe, M. E. (1998). Analgesic abuse in the elderly. Renal sequelae and management. *Drugs & Aging*, *12*, 391–400.
- Erikson, E. (1950). *Childhood and society*. New York: W. W. Norton.
- Ernst, R. L., & Hay, J. W. (1994). The US economic and social costs of Alzheimer's disease revisited. *American Journal of Public Health*, *84*, 1261–1264.
- Evans, D. A., Funkenstein, H. H., Albert, M. S., Scherr, P. A., Cook, N. R., Chown, M. J., Hebert, L. E., Hennekens, C. H., & Taylor, J. O. (1989). Prevalence of Alzheimer's disease in a community population of older persons. Higher than previously reported. *Journal of the American Medical Association*, *262*, 2551–2556.
- Eyler-Zorilla, L., Simjee-McClure, F., Patterson, T. L., Palmer, B. W., Harris, M. J., & Jeste, D. V. (1999). *Clinical, functional, and cognitive characteristics of elderly, community-dwelling schizophrenia patients*. New Orleans, LA: American Association of Geriatric Psychiatry.
- Feinberg, I. (1983). Schizophrenia: Caused by a fault in programmed synaptic elimination during adolescence? *Journal of Psychiatric Research*, *17*, 319–334.
- Felker, B., Yazel, J. J., & Short, D. (1996). Mortality and medical comorbidity among psychiatric patients: A review. *Psychiatric Services*, *47*, 1356–1363.
- Ferris, S. H., Steinberg, G., Shulman, E., Kahn, R., & Reisberg, B. (1987). Institutionalization of Alzheimer's patients: Reducing precipitating factors through family counseling. *Home Health Care Services Quarterly*, *8*, 23–61.
- Finch, C. E., & Tanzi, R. E. (1997). Genetics of aging. *Science*, *278*, 407–411.
- Finkel, S. I., Costa e Silva, J., Cohen, G., Miller, S., & Sartorius, N. (1996). Behavioral and psychological signs and symptoms of dementia: A consensus statement on current knowledge and implications for research and treatment. *International Psychogeriatrics*, *8*(Suppl. 3), 497–500.
- Finkelstein, S. N., Berndt, E. R., & Greenberg, P. E. (1996, January 17–18). *Economics of depression: A summary and review*. National Depressive and Manic-Depressive Association sponsored Consensus Conference on the Undertreatment of Depression, Washington, DC.
- Finlayson, R. E., & Davis, L. J., Jr. (1994). Prescription drug dependence in the elderly population: Demographic and clinical features of 100 inpatients. *Mayo Clinic Proceedings*, *69*, 1137–1145.
- Finney, J. W., & Moos, R. H. (1991). The long-term course of treated alcoholism: I. Mortality, relapse and remission rates and comparisons with community controls. *Journal of Studies on Alcohol*, *52*, 44–54.
- Flint, A. J. (1994). Epidemiology and comorbidity of anxiety disorders in the elderly. *American Journal of Psychiatry*, *151*, 640–649.
- Ford, D. E., & Kamerow, D. B. (1989). Epidemiologic study of sleep disturbances and psychiatric disorders. An opportunity for prevention? *Journal of the American Medical Association*, *262*, 1479–1484.

Mental Health: A Report of the Surgeon General

- Frank, E., Prigerson, H. G., Shear, M. K., & Reynolds, C. F., III. (1997a). Phenomenology and treatment of bereavement-related distress in the elderly. *International Clinical Psychopharmacology*, *12*(Suppl. 7), S25–S29.
- Frank, R. G., Koyanagi, C., & McGuire, T. G. (1997b). The politics and economics of mental health “parity” laws. *Health Affairs*, *16*, 108–119.
- Frank, R., McGuire, T., & Newhouse, J. (1997c). Risk contracts in managed mental health care. *Health Affairs*, *14*, 50–64.
- Frasure-Smith, N., Lesperance, F., & Talajic, M. (1993). Depression following myocardial infarction. Impact on 6-month survival. *Journal of the American Medical Association*, *270*, 1819–1825.
- Frasure-Smith, N., Lesperance, F., & Talajic, M. (1995). Depression and 18-month prognosis after myocardial infarction. *Circulation*, *91*, 999–1005.
- Friedman, B., & Kane, R. L. (1993). HMO medical directors’ perceptions of geriatric practice in Medicare HMOs. *Journal of the American Geriatrics Society*, *41*, 1144–1149.
- Galasko, D., Hansen, L. A., Katzman, R., Wiederholt, W., Masliah, E., Terry, R., Hill, L. R., Lessin, P., & Thal, L. J. (1994). Clinical-neuropathological correlations in Alzheimer’s disease and related dementias. *Archives of Neurology*, *51*, 888–895.
- Gallagher, D., & Thompson, L. (1982). Treatment of major depressive disorder in older adult outpatients with brief psychotherapies. *Psychotherapy: Theory, Research and Practice*, *19*, 482–490.
- Gallagher-Thompson, D., Hanley-Peterson, P., & Thompson, L. W. (1990). Maintenance of gains versus relapse following brief psychotherapy for depression. *Journal of Consulting and Clinical Psychology*, *58*, 371–374.
- Gallo, J. J., Anthony, J. C., & Muthen, B. O. (1994). Age differences in the symptoms of depression: A latent trait analysis. *Journal of Gerontology*, *49*, P251–P264.
- Gallo, J. J., & Lebowitz, B. D. (1999). The epidemiology of common late-life mental disorders in the community: Themes for the new century. *Psychiatric Services*, *50*, 1158–1166.
- Gallo, J. J., Rabins, P. V., Lyketsos, C. G., Tien, A. Y., & Anthony, J. C. (1997). Depression without sadness: Functional outcomes of nondysphoric depression in later life. *Journal of the American Geriatrics Society*, *45*, 570–578.
- Gallo, J. J., Royall, D. R., & Anthony, J. C. (1993). Risk factors for the onset of depression in middle age and later life. *Social Psychiatry and Psychiatric Epidemiology*, *28*, 101–108.
- Gallo, J. J., Ryan, S. D., & Ford, D. (in press). Attitudes, knowledge, and behavior of family physicians regarding depression in late life. *Archives of Family Medicine*.
- Gask, L., Sibbald, B., & Creed, F. (1997). Evaluating models of working at the interface between mental health services and primary care. *British Journal of Psychiatry*, *170*, 6–11.
- Gatz, M., Bengtson, V. L., & Blum, M. J. (1990). Caregiving families. In J. E. Birren & K. W. Schaie (Eds.), *Handbook of the psychology of aging* (pp. 404–426). San Diego: Academic Press.
- Gatz, M., & Smyer, M. A. (1992). The mental health system and older adults in the 1990s. *American Psychologist*, *47*, 741–751.
- George, L. K. (1992). Community and home care for mentally ill older adults. In J. E. Birren, R. B. Sloane, G. D. Cohen, N. R. Hooyman, B. D. Lebowitz, & M. I. Wykle (Eds.), *Handbook of mental health and aging* (2nd ed., pp. 793–813). San Diego: Academic Press.
- Glasser, M., & Gravdal, J. A. (1997). Assessment and treatment of geriatric depression in primary care settings. *Archives of Family Medicine*, *6*, 433–438.
- Glassman, A. H., Roose, S. P., & Bigger, J. T., Jr. (1993). The safety of tricyclic antidepressants in cardiac patients. Risk-benefit reconsidered. *Journal of the American Medical Association*, *269*, 2673–2675.
- Goldman, H. H., Cohen, G. D., & Davis, M. (1985). Expanded Medicare outpatient coverage for Alzheimer’s disease and related disorders. *Hospital and Community Psychiatry*, *36*, 939–942.
- Gordon, T., & Kannel, W. B. (1983). Drinking and its relation to smoking, BP, blood lipids, and uric acid. The Framingham study. *Archives of Internal Medicine*, *143*, 1366–1374.
- Gottesman, L. E., Peskin, E., Kennedy, K., & Mossey, J. (1991). Implications of a mental health intervention for elderly mentally ill residents of residential care facilities. *International Journal of Aging and Human Development*, *32*, 229–245.
- Gottlieb, G. L. (1995). Geriatric psychiatry. In H. H. Goldman (Ed.), *Review of general psychiatry* (4th ed., pp. 483–491). Norwalk, CT: Appleton and Lange.

- Grant, B. F., Harford, T. C., Dawson, D. A., Chou, P. S., & Pickering, R. P. (1994). Prevalence of DSM-IV alcohol abuse and dependence: United States, 1992. *Alcohol Health and Research World*, 18, 243.
- Green, M. F., Marshall, B. D., Jr., Wirshing, W. C., Ames, D., Marder, S. R., McGurk, S., Kern, R. S., & Mintz, J. (1997). Does risperidone improve verbal working memory in treatment-resistant schizophrenia? *American Journal of Psychiatry*, 154, 799–804.
- Greenberg, J., Leutz, W., Greenlick, M., Malone, J., Ervin, S., & Kodner, D. (1988). The social HMO demonstration: Early experience. *Health Affairs*, 7, 66–79.
- Greenblatt, D. J., Sellers, E. M., & Shader, R. I. (1982). Drug disposition in old age. *New England Journal of Medicine*, 306, 1081–1088.
- Greenwald, B. S., Kramer-Ginsberg, E., Bogerts, B., Ashtari, M., Aupperle, P., Wu, H., Allen, L., Zeman, D., & Patel, M. (1997). Qualitative magnetic resonance imaging findings in geriatric depression. Possible link between later-onset depression and Alzheimer's disease? *Psychological Medicine*, 27, 421–431.
- Gurland, B. J., Cross, P. S., & Katz, S. (1996). Epidemiological perspectives on opportunities for treatment of depression. *American Journal of Geriatric Psychiatry*, 4(Suppl. 1), S7–S13.
- Haastруп, S., & Jepsen, P. W. (1988). Eleven year follow-up of 300 young opioid addicts. *Acta Psychiatrica Scandinavica*, 77, 22–26.
- Hagnell, O., Lanke, J., Rorsman, B., & Ojesjo, L. (1982). Are we entering an age of melancholy? Depressive illnesses in a prospective epidemiological study over 25 years: The Lundby Study, Sweden. *Psychological Medicine*, 12, 279–289.
- Haight, B. K., Michel, Y., & Hendrix, S. (1998). Life review: Preventing despair in newly relocated nursing home residents' short- and long-term effects. *International Journal of Aging and Human Development*, 47, 119–142.
- Hardy, J. A., & Higgins, G. A. (1992). Alzheimer's disease: The amyloid cascade hypothesis. *Science*, 256, 184–185.
- Harlow, S. D., Goldberg, E. L., & Comstock, G. W. (1991). A longitudinal study of risk factors for depressive symptomatology in elderly widowed and married women. *American Journal of Epidemiology*, 134, 526–538.
- Harwood, H., Fountain, D., Livermore, G., & Lewin Group. (1998). *The economic costs of alcohol and drug abuse in the United States, 1992*. Rockville, MD: U.S. Department of Health and Human Services, National Institutes of Health, National Institute on Drug Abuse, Office of Science Policy and Communications, National Institute on Alcohol Abuse and Alcoholism, Office of Policy Analysis.
- Hawton, K., & Kirk, J. (1989). Problem solving. In K. Hawton, P. M. Salkovskis, J. Kirk, & D. M. Clark (Eds.), *Cognitive behaviour therapy for psychiatric patients: A practical guide*. Oxford, England: Oxford University Press.
- Heaton, R., Paulsen, J. S., McAdams, L. A., Kuck, J., Zisook, S., Braff, D., Harris, J., & Jeste, D. V. (1994). Neuropsychological deficits in schizophrenics. Relationship to age, chronicity, and dementia. *Archives of General Psychiatry*, 51, 469–476.
- Hebert, L. E., Scherr, P. A., Beckett, L. A., Albert, M. S., Pilgrim, D. M., Chown, M. J., Funkenstein, H. H., & Evans, D. A. (1995). Age-specific incidence of Alzheimer's disease in a community population. *Journal of the American Medical Association*, 273, 1354–1359.
- Helzer, J. E., Burnam, A., & McEvoy, L. T. (1991). Alcohol abuse and dependence. In L. N. Robins & D. A. Regier (Eds.), *Psychiatric disorders in America: The Epidemiologic Catchment Area study* (pp. 81–115). New York: Free Press.
- Hill, L. R., Klauber, M. R., Salmon, D. P., Yu, E. S., Liu, W. T., Zhang, M., & Katzman, R. (1993). Functional status, education, and the diagnosis of dementia in the Shanghai survey. *Neurology*, 43, 138–145.
- Himmelfarb, S., & Murrell, S. A. (1984). The prevalence and correlates of anxiety symptoms in older adults. *Journal of Psychology*, 116, 159–167.
- Hirschfeld, R. M., Keller, M. B., Panico, S., Arons, B. S., Barlow, D., Davidoff, F., Endicott, J., Froom, J., Goldstein, M., Gorman, J. M., Marek, R. G., Maurer, T. A., Meyer, R., Phillips, K., Ross, J., Schwenk, T. L., Sharfstein, S. S., Thase, M. E., & Wyatt, R. J. (1997). The National Depressive and Manic-Depressive Association consensus statement on the undertreatment of depression. *Journal of the American Medical Association*, 277, 333–340.
- Holroyd, S., & Duryee, J. J. (1997). Differences in geriatric psychiatry outpatients with early- vs late-onset depression. *International Journal of Geriatric Psychiatry*, 12, 1100–1106.

Mental Health: A Report of the Surgeon General

- Hoyert, D. L., Kochanek, K. D., & Murphy, S. L. (1999). Deaths: Final data for 1997. *National Vital Statistics Reports, 47* (9). Hyattsville, MD: National Center for Health Statistics.
- Hser, Y. I., Anglin, D., & Powers, K. (1993). A 24-year follow-up of California narcotics addicts. *Archives of General Psychiatry, 50*, 577–584.
- Ingster, L. M., & Cartwright, W. S. (1995). Drug disorders and cardiovascular disease: The impact on annual hospital length of stay for the Medicare population. *American Journal of Drug and Alcohol Abuse, 21*, 93–110.
- Institute for Health and Aging. (1996). *Chronic care in America: A 21st century challenge*. Princeton, NJ: Robert Wood Johnson Foundation.
- Institute of Medicine. (1984). *Bereavement: Reactions, consequences and care*. Washington, DC: National Academy Press.
- Institute of Medicine. (1986). *Improving the quality of care in nursing homes*. Washington, DC: National Academy Press.
- Institute of Medicine. (1998). *Violence in families: Assessing prevention and treatment programs*. Washington, DC: National Academy Press.
- Institute of Medicine. (1999). *Reducing the burden of injury: Advancing prevention and treatment*. Washington, DC: National Academy Press.
- Jeste, D. V., Caligiuri, M. P., Paulsen, J. S., Heaton, R. K., Lacro, J. P., Harris, M. J., Bailey, A., Fell, R. L., & McAdams, L. A. (1995a). Risk of tardive dyskinesia in older patients. A prospective longitudinal study of 266 outpatients. *Archives of General Psychiatry, 52*, 756–765.
- Jeste, D. V., Harris, M. J., Krull, A., Kuck, J., McAdams, L. A., & Heaton, R. (1995b). Clinical and neuropsychological characteristics of patients with late-onset schizophrenia. *American Journal of Psychiatry, 152*, 722–730.
- Jeste, D. V., Lacro, J. P., Bailey, A., Rockwell, E., Harris, M. J., & Caligiuri, M. P. (in press). Lower incidence of tardive dyskinesia with risperidone compared with haloperidol in older patients. *Journal of the American Geriatrics Society*.
- Jeste, D. V., & Palmer, B. (1998). Secondary psychoses: An overview. *Seminars in Clinical Neuro Psychiatry, 3*, 2–3.
- Jeste, D. V., Symonds, L. L., Harris, M. J., Paulsen, J. S., Palmer, B. W., & Heaton, R. K. (1997). Nondementia nonpraecox dementia praecox? Late-onset schizophrenia. *American Journal of Geriatric Psychiatry, 5*, 302–317.
- Jeste, D. V., Wragg, R. E., Salmon, D. P., Harris, M. J., & Thal, L. J. (1992). Cognitive deficits of patients with Alzheimer's disease with and without delusions. *American Journal of Psychiatry, 149*, 184–189.
- Johansson, B., & Zarit, S. H. (1995). Prevalence and incidence of dementia in the oldest-old: A longitudinal study of a population-based sample of 84–90 year olds in Sweden. *International Journal of Geriatric Psychiatry, 10*, 359–366.
- Jorm, A. F., Korten, A. E., & Henderson, A. S. (1987). The prevalence of dementia: A quantitative integration of the literature. *Acta Psychiatrica Scandinavica, 76*, 465–479.
- Judd, L. L., Rapaport, M. H., Paulus, M. P., & Brown, J. L. (1994). Subsyndromal symptomatic depression: A new mood disorder? *Journal of Clinical Psychiatry, 55*(Suppl.), 18–28.
- Kachur, S. P., Potter, L. B., James, S. P., & Powell, K. E. (1995). *Suicide in United States 1980–1992* (Violence Surveillance Summary Series, No. 1). Atlanta, GA: National Center for Injury Prevention and Control.
- Kane, J. M., Cole, K., Sarantakos, S., Howard, A., & Borenstein, M. (1983). Safety and efficacy of bupropion in elderly patients: Preliminary observations. *Journal of Clinical Psychiatry, 44*, 134–136.
- Kane, J. M., Woerner, M. G., Pollack, S., Safferman, A. Z., & Lieberman, J. A. (1993). Does clozapine cause tardive dyskinesia? *Journal of Clinical Psychiatry, 54*, 327–330.
- Kane, R. A. (1995). Expanding the home care concept: Blurring distinctions among home care, institutional care, and other long-term-care services. *Milbank Quarterly, 73*, 161–186.
- Kane, R. L., Kane, R. A., Finch, M., Harrington, C., Newcomer, R., Miller, N., & Hulbert, M. (1997). S/HMOs, the second generation: Building on the experience of the first Social Health Maintenance Organization demonstrations. *Journal of the American Geriatrics Society, 45*, 101–107.
- Kang, D. E., Saitoh, T., Chen, X., Xia, Y., Masliah, E., Hansen, L. A., Thomas, R. G., Thal, L. J., & Katzman, R. (1997). Genetic association of the low-density lipoprotein receptor-related protein gene (LRP), an apolipoprotein E receptor, with late-onset Alzheimer's disease. *Neurology, 49*, 56–61.

- Kastenbaum, R. (1985). Dying and death: A life-span approach. In J. E. Birren & K. W. Schaie (Eds.), *Handbook of the psychology of aging* (2nd ed., pp. 619–643). New York: Van Nostrand Reinhold.
- Katon, W. J., Robinson, P., Von Korff, M., Lin, E., Bush, T., Ludman, E., Simon, G., & Walker, E. (1996). A multifaceted intervention to improve treatment of depression in primary care. *Archives of General Psychiatry*, *53*, 924–932.
- Katon, W. J., & Gonzales, J. (1994). A review of randomized trials of psychiatric consultation-liaison studies in primary care. *Psychosomatics*, *35*, 268–278.
- Katon, W., Von Korff, M., Lin, E., Bush, T., & Ormel, J. (1992). Adequacy and duration of antidepressant treatment in primary care. *Medical Care*, *30*, 67–76.
- Katon, W., Von Korff, M., Lin, E., Unutzer, J., Simon, G., Walker, E., Ludman, E., & Bush, T. (1997). Population-based care of depression: Effective disease management strategies to decrease prevalence. *General Hospital Psychiatry*, *19*, 169–178.
- Katona, C. L., Hunter, B. N., & Bray, J. (1998). A double-blind comparison of the efficacy and safety of paroxetine and imipramine in the treatment of depression with dementia. *International Journal of Geriatric Psychiatry*, *13*, 100–108.
- Katz, I. R. & Parmelee, P. A. (1997). Overview. In R. L. Rubinstein & M. P. Lawton (Eds.), *Depression in long term and residential care* (pp. 1–28). New York: Springer.
- Katzman, R. (1993). Education and the prevalence of dementia and Alzheimer's disease. *Neurology*, *43*, 13–20.
- Kaufers, D. I., Cummings, J. L., & Christine, D. (1996). Effect of tacrine on behavioral symptoms in Alzheimer's disease: An open-label study. *Journal of Geriatric Psychiatry and Neurology*, *9*, 1–6.
- Kaufers, D. I., Cummings, J. L., Christine, D., Bray, T., Castellon, S., Masterman, D., MacMillan, A., Ketchel, P., & DeKosky, S. T. (1998). Assessing the impact of neuropsychiatric symptoms in Alzheimer's disease: The Neuropsychiatric Inventory Caregiver Distress Scale. *Journal of the American Geriatrics Society*, *46*, 210–215.
- Kendell, R. E., Brockington, I. F., & Leff, J. P. (1981). Prognostic implications of six alternative definitions of schizophrenia. *Archives of General Psychiatry*, *36*, 25–31.
- Kessler, R. C., Berglund, P. A., Zhao, S., Leaf, P. J., Kouzis, A. C., Bruce, M. L., Friedman, R. M., Grosser, R. C., Kennedy, C., Narrow, W. E., Kuehnel, T. G., Laska, E. M., Manderscheid, R. W., Rosenheck, R. A., Santoni, T. W., & Schneier, M. (1996). The 12-month prevalence and correlates of serious mental illness (SMI). In R. W. Manderscheid & M. A. Sonnenschein (Eds.), *Mental health United States, 1996* (DHHS Publication No. SMA 96-3098, pp. 59–70). Washington, DC: Superintendent of Documents, U.S. Government Printing Office.
- King, C. J., Van Hasselt, V. B., Segal, D. L., & Hersen, M. (1994). Diagnosis and assessment of substance abuse in older adults: Current strategies and issues. *Addictive Behaviors*, *19*, 41–55.
- Klapow, J. C., Evans, J., Patterson, T. L., Heaton, R. K., Koch, W. L., & Jeste, D. V. (1997). Direct assessment of functional status in older patients with schizophrenia. *American Journal of Psychiatry*, *154*, 1022–1024.
- Klausner, E. J., & Alexopolous, G. S. (in press). The future of psychosocial treatments for the elderly. *Psychiatric Research*.
- Klerman, G. L., Weissman, M. M., & Rounsaville, B. J., Sherron, E. S. (1984). *Interpersonal psychotherapy of depression*. New York: Basic Books.
- Knight, B. G., Lutzky, S. M., & Macofsky-Urban, F. (1993). A meta-analytic review of interventions for caregiver distress: Recommendations for future research. *Gerontologist*, *33*, 240–248.
- Knight, B. G., Rickards, L., Rabins, P., Buckwalter K., & Smith M. (1995). Community-based services for older adults: A role for psychologists? In B. Knight, L. Teri, P. Wohlford, & J. Santos (Eds.), *Mental health services for older adults: Implications for training and practice in geropsychology* (pp. 21–29). Washington, DC: American Psychological Association.
- Knight, B. G., Woods, E., & Kaskie, B. (1998). *Community mental health services in the United States and the United Kingdom: A comparative systems approach*. Oxford, England: Elsevier.
- Knopman, D., Schneider, L., Davis, K., Talwalker, S., Smith, F., Hoover, T., & Gracon, S. (1996). Long-term tacrine (Cognex) treatment: Effects on nursing home placement and mortality, Tacrine Study Group. *Neurology*, *47*, 166–177.

Mental Health: A Report of the Surgeon General

- Kofoed, L. L. (1984). Abuse and misuse of over-the-counter drugs by the elderly. In R. M. Atkinson (Ed.), *Alcohol and drug abuse in old age* (pp. 49–59). Washington, DC: American Psychiatric Press.
- Komisar, H., Reuter, J., & Feder, J. (1997). *Medicare chart book*. Menlo Park, CA: The Henry J. Kaiser Family Foundation.
- Kraepelin, E. (1971). *Dementia praecox and paraphrenia*. Huntington, NY: R. E. Krieger Publishing Company.
- Kramer, M., Simonsick, E., Lima, B., & Levav, I. (1992). *The epidemiological basis for mental health care in primary health care: A case for action*. New York: Tavistock/Routledge.
- Kramer, M., Taube, C. A., & Redick, R. W. (1973). Patterns of use of psychiatric facilities by the aged: Past, present and future. In C. Eisdorfer & M. P. Lawton (Eds.), *The psychology of adult development and aging* (pp. 428–528). Washington, DC: American Psychological Association.
- Krishnan, K. R., Hays, J. C., Tupler, L. A., George, L. K., & Blazer, D. G. (1995). Clinical and phenomenological comparisons of late-onset and early-onset depression. *American Journal of Psychiatry*, *152*, 785–788.
- Laghrissi-Thode, F., Pollock, B. G., Miller, M. C., Mulsant, B. H., Altieri, L., & Finkel, M. S. (1995). Double-blind comparison of paroxetine and nortriptyline on the postural stability of late-life depressed patients. *Psychopharmacology Bulletin*, *31*, 659–663.
- Lamy, P. P., Salzman, C., & Nevis-Olesen, J. (1992). Drug prescribing patterns, risks, and compliance guidelines. In C. Salzman (Ed.), *Clinical geriatric psychopharmacology* (2nd ed., pp. 15–37). Baltimore: William & Wilkins.
- Lawrence, A. D., & Sahakian, B. J. (1998). The cognitive psychopharmacology of Alzheimer's disease: Focus on cholinergic systems. *Neurochemical Research*, *23*, 787–794.
- Lazarus, L. W., & Sadavoy, J. (1996). Individual psychotherapy. In J. Sadavoy (Ed.), *Comprehensive review of geriatric psychiatry* (2nd ed., pp. 819–826). Washington, DC: American Psychiatric Press.
- Lebowitz, B. D., & Pearson, J. L. (in press). Intervention research in psychosis: Prevention trials. *Schizophrenia Bulletin*.
- Lebowitz, B. D., Pearson, J. L., & Cohen, G. D. (1998). *Clinical geriatric psychopharmacology*. Baltimore: Williams & Wilkins.
- Lebowitz, B. D., Pearson, J. L., Schneider, L. S., Reynolds, C. F., III, Alexopoulos, G. S., Bruce, M. L., Conwell, Y., Katz, I. R., Meyers, B. S., Morrison, M. F., Mossey, J., Niederehe, G., & Parmelee, P. (1997). Diagnosis and treatment of depression in late life. Consensus statement update. *Journal of the American Medical Association*, *278*, 1186–1190.
- Leff, J. (1993). The TAPS project: Evaluating community placement of long-stay psychiatric patients. *British Journal of Psychiatry*, *162*, 3–56.
- Levy, L. H., Derby, J. F., & Martinkowski, K. S. (1993). Effects of membership in bereavement support groups on adaptation to conjugal bereavement. *American Journal of Community Psychology*, *21*, 361–381.
- Levy, B., Cushman, J., & Abeles, N. (in press). Memory complaints in the able elderly. *Clinical Gerontologist*.
- Liberto, J. G., Oslin, D. W., & Ruskin, P. E. (1992). Alcoholism in older persons: A review of the literature. *Hospital and Community Psychiatry*, *43*, 975–984.
- Lieberman, M. A., & Videka-Sherman, L. (1986). The impact of self-help groups on the mental health of widows and widowers. *American Journal of Orthopsychiatry*, *56*, 435–449.
- Light, E., & Lebowitz, B. D. (Eds.). (1991). *The elderly with chronic mental illness*. New York: Springer.
- Light, E., Lebowitz, B. D., & Bailey, F. (1986). CMHC's and elderly services: An analysis of direct and indirect services and service delivery sites. *Community Mental Health Journal*, *22*, 294–302.
- Lindesay, J., Briggs, K., & Murphy, E. (1989). The Guy's/Age Concern Survey. Prevalence rates of cognitive impairment, depression and anxiety in an urban elderly community. *British Journal of Psychiatry*, *155*, 317–329.
- Lipsman, R. (1996). Services and supports to the homebound elderly with mental health needs. *Journal of Long-Term Home Health Care*, *15*, 24–38.
- Little, J. T., Reynolds, C. F., III, Dew, M. A., Frank, E., Begley, A. E., Miller, M. D., Cornes, C., Mazumdar, S., Perel, J. M., & Kupfer, D. J. (1998). How common is resistance to treatment in recurrent, nonpsychotic geriatric depression? *American Journal of Psychiatry*, *155*, 1035–1038.
- Locascio, J. J., Growdon, J. H., & Corkin, S. (1995). Cognitive test performance in detecting, staging, and tracking Alzheimer's disease. *Archives of Neurology*, *52*, 1087–1099.

- Loewenstein, D. A., Amigo, E., Duara, R., Guterman, A., Hurwitz, D., Berkowitz, N., Wilkie, F., Weinberg, G., Black, B., Gittelman, B., & Eisdorfer, C. (1989). A new scale for the assessment of functional status in Alzheimer's disease and related disorders. *Journal of Gerontology*, *44*, P114–P121.
- Lohr, J. B., Alder, M., Flynn, K., Harris, M. J., & McAdams, L. A. (1997). Minor physical anomalies in older patients with late-onset schizophrenia, early-onset schizophrenia, depression, and Alzheimer's disease. *American Journal of Geriatric Psychiatry*, *5*, 318–323.
- Lombardo, N. E. (1994). *Barriers to mental health services for nursing home residents*. Washington, DC: American Association of Retired Persons.
- Lurie, N., Moscovice, I. S., Finch, M., Christianson, J. B., & Popkin, M. K. (1992). Does capitation affect the health of the chronically mentally ill? Results from a randomized trial. *Journal of the American Medical Association*, *267*, 3300–3304.
- Maddox, G. L., Steinhauer, S., & Bolda, E. (1996). Continuum of care: Movement toward the community. In E. W. Busse & D. G. Blazer (Eds.), *The American Psychiatric Press textbook of geriatric psychiatry* (2nd ed., pp. 433–459). Washington, DC: American Psychiatric Press.
- Malmgren, R. (1994). Epidemiology of aging. In C. E. Coffey & J. L. Cummings (Eds.), *The American Psychiatric Press textbook of geriatric neuropsychiatry* (pp. 17–33). Washington, DC: American Psychiatric Press.
- Manton, K. G., Corder, L., & Stallard, E. (1997). Chronic disability trends in elderly United States populations: 1982–1994. *Proceedings of the National Academy of Sciences, USA*, *94*, 2593–2598.
- Markowitz, J., Brown, R., Sweeney, J., & Mann, J. J. (1987). Reduced length and cost of hospital stay for major depression in patients treated with ECT. *American Journal of Psychiatry*, *144*, 1025–1029.
- Marmar, C. R., Horowitz, M. J., Weiss, D. S., Wilner, N. R., & Kaltreider, N. B. (1988). A controlled trial of brief psychotherapy and mutual-help group treatment of conjugal bereavement. *American Journal of Psychiatry*, *145*, 203–209.
- McCurry, S. M., Logsdon, R. G., & Teri, L. (1996). Behavioral treatment of sleep disturbance in elderly dementia caregivers. *Clinical Gerontologist*, *17*, 35–50.
- McEwen, B. S. (1998). Protective and damaging effects of stress mediators. *New England Journal of Medicine*, *338*, 171–179.
- McFarlane, A. C., & Yehuda, R. (1996). Resilience, vulnerability, and the course of post-traumatic reactions. In B. A. van der Kolk, A. C. McFarlane, & L. Weisath (Eds.), *Traumatic stress: The effects of overwhelming experience on mind, body and society* (pp. 155–181). New York: Guilford Press.
- McGeer, P. L., Schulzer, M., & McGeer, E. G. (1996). Arthritis and anti-inflammatory agents as possible protective factors for Alzheimer's disease: A review of 17 epidemiologic studies. *Neurology*, *47*, 425–432.
- McKhann, G., Drachman, D., Folstein, M., Katzman, R., Price, D., & Stadlan, E. M. (1984). Clinical diagnosis of Alzheimer's disease: Report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology*, *34*, 939–944.
- Mechanic, D. (1998). Emerging trends in mental health policy and practice. *Health Affairs*, *17*, 82–98.
- Meeks, S., Carstensen, L. L., Stafford, P. B., Brenner, L. L., Weathers, F., Welch, R., & Oltmanns, T. F. (1997). Mental health needs of the chronically mentally ill elderly. *Psychology and Aging*, *5*, 163–171.
- Meeks, S., & Murrell, S. A. (1997). Mental illness in late life: Socioeconomic conditions, psychiatric symptoms, and adjustment of long-term sufferers. *Psychology and Aging*, *12*, 296–308.
- Mega, M. S., Cummings, J. L., Fiorello, T., & Gornbein, J. (1996). The spectrum of behavioral changes in Alzheimer's disease. *Neurology*, *46*, 130–135.
- Mellinger, G. D., Balter, M. B., & Uhlenhuth, E. H. (1984). Prevalence and correlates of the long-term regular use of anxiolytics. *Journal of the American Medical Association*, *251*, 375–379.
- Mendes de Leon, C. F., Kasl, S. V., & Jacobs, S. (1994). A prospective study of widowhood and changes in symptoms of depression in a community sample of the elderly. *Psychological Medicine*, *24*, 613–624.
- Mesulam, M. M. (1996). The system-level organization of cholinergic innervation in the human cerebral cortex and its alterations in Alzheimer's disease. *Progress in Brain Research*, *109*, 285–297.
- Meyers, B. (1996). Psychiatric interventions to improve primary care diagnosis and treatment of depression. *American Journal of Geriatric Psychiatry*, *4*, S91–S95.

Mental Health: A Report of the Surgeon General

- Miller, N. S., Belkin, B. M., & Gold, M. S. (1991). Alcohol and drug dependence among the elderly: Epidemiology, diagnosis, and treatment. *Comprehensive Psychiatry*, 32, 153-165.
- Miller, R. A. (1996). The aging immune system: Primer and prospectus. *Science*, 273, 70-74.
- Miller, R. H., & Luft, H. S. (1997). Does managed care lead to better or worse quality of care? *Health Affairs*, 16, 7-25.
- Mittelman, M. S., Ferris, S. H., Shulman, E., Steinberg, G., & Levin, B. (1996). A family intervention to delay nursing home placement of patients with Alzheimer disease. A randomized controlled trial. *Journal of the American Medical Association*, 276, 1725-1731.
- Mittelman, M. S., Ferris, S. H., Steinberg, G., Shulman, E., Mackell, J. A., Ambinder, A., & Cohen, J. (1993). An intervention that delays institutionalization of Alzheimer's disease patients: Treatment of spouse-caregivers. *Gerontologist*, 33, 730-740.
- Moak, G. S. (1996). When the seriously mentally ill patient grows old. In S. M. Soreff (Ed.), *Handbook for the treatment of the seriously mentally ill* (pp. 279-293). Seattle: Hogrefe and Huber.
- Moak, G. S., & Fisher, W. H. (1991). Geriatric patients and services in state hospitals: Data from a national survey. *Hospital and Community Psychiatry*, 42, 273-276.
- Mohs, R. C., Breitner, J. C., Silverman, J. M., & Davis, K. L. (1987). Alzheimer's disease. Morbid risk among first-degree relatives approximates 50% by 90 years of age. *Archives of General Psychiatry*, 44, 405-408.
- Molchan, S. E., Martinez, R. A., Hill, J. L., Weingartner, H. J., Thompson, K., Vitiello, B., & Sunderland, T. (1992). Increased cognitive sensitivity to scopolamine with age and a perspective on the scopolamine model. *Brain Research and Brain Research Reviews*, 17, 215-226.
- Moos, R. H., Brennan, P. L., & Mertens, J. R. (1994). Mortality rates and predictors of mortality among late-middle-aged and older substance abuse patients. *Alcoholism: Clinical and Experimental Research*, 18, 187-195.
- Morris, J. C., Cyrus, P. A., Orazem, J., Mas, J., Bieber, F., Ruzicka, B. B., & Gulanski, B. (1998). Metrifonate benefits cognitive, behavioral, and global function in patients with Alzheimer's disease. *Neurology*, 50, 1222-1230.
- Mosher, B., Ashley, P. M. (1989). A community based residential care program for elderly people with chronic mental illness. *Adult Residential Care Journal*, 3, 267-278.
- Mossey, J. M., & Shapiro, E. (1985). Physician use by the elderly over an eight-year period. *American Journal of Public Health*, 75, 1333-1334.
- Mrak, R. E., Sheng, J. G., & Griffin, W. S. (1995). Glial cytokines in Alzheimer's disease: Review and pathogenic implications. *Human Pathology*, 26, 816-823.
- Mynors, B., Wallis, L. M., Gath, D. H., Lloyd-Thomas, A. R., & Tomlinson, D. (1995). Randomized controlled trial comparing problem solving treatment with amitriptyline and placebo for major depression in primary care. *British Medical Journal*, 310, 441-445.
- Napoliello, M. J. (1986). An interim multicentre report on 677 anxious geriatric out-patients treated with buspirone. *British Journal of Clinical Practice*, 40, 71-73.
- Nasar, S. (1998). *A beautiful mind: A biography of John Forbes Nash, Jr. winner of the Nobel Prize in economics, 1994*. New York: Simon and Schuster.
- National Center for Health Statistics. (1993). *Advance report of final mortality statistics, 1990*. Hyattsville, MD: Author.
- National Institute on Aging. (1996). *Progress report on Alzheimer's disease*. Washington, DC: Author.
- National Institute on Drug Abuse. (1992). *Annual medical examiner data 1991: Data from the Drug Abuse Warning Network (DAWN)*. Washington, DC: U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration.
- National Institutes of Health & National Institute of Mental Health. (1985). Consensus conference. Electroconvulsive therapy. *Journal of the American Medical Association*, 254, 2103-2108.
- National Institutes of Health Consensus Development Panel on Depression in Late Life. (1992). Diagnosis and treatment of depression in late life. *Journal of the American Medical Association*, 268, 1018-1024.
- Nemeroff, C. B., DeVane, C. L., & Pollock, B. G. (1996). Newer antidepressants and the cytochrome P450 system. *American Journal of Psychiatry*, 153, 311-320.

- Newman, A. B., Enright, P. L., Manolio, T. A., Haponik, E. F., & Wahl, P. W. (1997). Sleep disturbance, psychosocial correlates, and cardiovascular disease in 5201 older adults: The Cardiovascular Health Study. *Journal of the American Geriatrics Society*, *45*, 1-7.
- Norberg, A. (1996). Pharmacological treatment of cognitive dysfunction in dementia disorders. *Acta Neurologica Scandinavica*, *168*, 87-92.
- O'Connor, D. W., Pollitt, P. A., Hyde, J. B., Brook, C. P., Reiss, B. B., & Roth, M. (1988). Do general practitioners miss dementia in elderly patients? *British Medical Journal*, *297*, 1107-1110.
- Ohayon, M., Caulet, M., & Lemoine, P. (1996). [The elderly, sleep habits and use of psychotropic drugs by the French population]. *Encephale*, *22*, 337-350.
- Olichney, J. M., Galasko, D., Salmon, D. P., Hofstetter, C. R., Hansen, L. A., Katzman, R., & Thal, L. J. (1998). Cognitive decline is faster in Lewy body variant than in Alzheimer's disease. *Neurology*, *51*, 351-357.
- Olichney, J. M., Iragui, V. J., & Jeste, D. V. (1995). P300 and N400 abnormalities in older schizophrenic patients [abstract]. *Milbank Quarterly*, *31*, 254.
- Olichney, J. M., Iragui, V. J., Kutas, M., Nowacki, R., & Jeste, D. V. (1996). Normal P300 amplitudes in late onset schizophrenia and related psychoses. *Neurosciences Abstracts*, *22*, 239.
- Oslin, D., Liberto, J. G., O'Brien, J., Krois, S., & Norbeck, J. (1997). Naltrexone as an adjunctive treatment for older patients with alcohol dependence. *American Journal of Geriatric Psychiatry*, *5*, 324-332.
- Oxman, T. E., Barrett, J. E., Barrett, J., & Gerber, P. (1990). Symptomatology of late-life minor depression among primary care patients. *Psychosomatics*, *31*, 174-180.
- Pacala, J. T., Boulton, C., Hepburn, K. W., Kane, R. A., Kane, R. L., Malone, J. K., Morishita, L., & Reed, R. L. (1995). Case management of older adults in health maintenance organizations. *Journal of the American Geriatrics Society*, *43*, 538-542.
- Paganini, B., Hill, A., & Henderson, V. W. (1994). Estrogen deficiency and risk of Alzheimer's disease in women. *American Journal of Epidemiology*, *140*, 256-261.
- Parmelee, P. A., Katz, I. R., & Lawton, M. P. (1989). Depression among institutionalized aged: Assessment and prevalence estimation. *Journal of Gerontology*, *44*, M22-M29.
- Pasternak, R. E., Prigerson, H., Hall, M., Miller, M. D., Fasiczka, A., Mazumdar, S., & Reynolds, C. F., III. (1997). The posttreatment illness course of depression in bereaved elders. High relapse/recurrence rates. *American Journal of Geriatric Psychiatry*, *5*, 54-59.
- Patterson, M. B., & Bolger, J. P. (1994). Assessment of behavioral symptoms in Alzheimer disease. *Alzheimer Disease and Associated Disorders*, *8*(Suppl. 3), 4-20.
- Patterson, T. L., & Jeste, D. V. (1999). The potential impact of the baby-boom generation on substance abuse among elderly persons. *Psychiatric Services*, *50*, 1184-1188.
- Patterson, T. L., Kaplan, R. M., Grant, I., Semple, S. J., Moscona, S., Koch, W. L., Harris, M. J., & Jeste, D. V. (1996). Quality of well-being in late-life psychosis. *Psychiatry Research*, *63*, 169-181.
- Paulsen, J. S., Heaton, R. K., Sadek, J. R., Perry, W., Delis, D. C., Braff, D., Kuck, J., Zisook, S., & Jeste, D. V. (1995). The nature of learning and memory impairments in schizophrenia. *Journal of International Neuropsychological Society*, *1*, 88-99.
- Paulsen, J. S., Romero, R., Chan, A., Davis, A. V., Heaton, R. K., & Jeste, D. V. (1996). Impairment of the semantic network in schizophrenia. *Psychiatry Research*, *63*, 109-121.
- Paveza, G. J., & Cohen, D. (1996). Treatment of mental health problems in the elderly. In B. L. Levin & J. Petrila (Eds.), *Mental health services: A public health perspective* (pp. 290-300). New York: Oxford University Press.
- Paykel, E. S., Brayne, C., Huppert, F. A., Gill, C., Barkley, C., Gehlhaar, E., Beardsall, L., Girling, D. M., Pollitt, P., & O'Connor, D. (1994). Incidence of dementia in a population older than 75 years in the United Kingdom. *Archives of General Psychiatry*, *51*, 325-332.
- Pearson, J. L., Teri, L., Reifler, B. V., & Raskind, M. A. (1989). Functional status and cognitive impairment in Alzheimer's patients with and without depression. *Journal of the American Geriatrics Society*, *37*, 1117-1121.
- Penninx, B. W., Guralnik, J. M., Pahor, M., Ferrucci, L., Cerhan, J. R., Wallace, R. B., & Havlik, R. J. (1998). Chronically depressed mood and cancer risk in older persons. *Journal of the National Cancer Institute*, *90*, 1888-1893.

Mental Health: A Report of the Surgeon General

- Pincus, H. A., & Wakefield-Davis, W. (1997, September 5-6). *Subthreshold mental disorders: Review and synthesis of studies on minor depression and other brand names*. Eleventh Conference on Mental Health Problems in the General Health Care Sector, Washington, DC.
- Pliskin, N. H., Cunningham, J. M., Wall, J. R., & Cassisi, J. E. (1996). Cognitive rehabilitation for cerebrovascular accidents and Alzheimer's disease. In P. W. Corrigan & S. C. Yudofsky (Eds.), *Cognitive rehabilitation for neuropsychiatric disorders* (p. 459). Washington, DC: American Psychiatric Press.
- Pollock, B. G., & Mulsant, B. H. (1995). Antipsychotics in older patients. A safety perspective. *Drugs and Aging*, 6, 312-323.
- Pollock, G. H. (1987). The mourning-liberation process: Ideas on the inner life of the older adult. In J. Sadavoy & M. Leszcz (Eds.), *Treating the elderly with psychotherapy: The scope for change in later life* (pp. 3-29). Madison, CT: International Universities Press.
- Prigerson, H. G., Bierhals, A., & Maciejewsky, P. (in press). Predictors of depression among aged widows and widowers: A literature review and preliminary results. *Gerontologist*.
- Rabins, P. V. (1996). Barriers to diagnosis and treatment of depression in elderly patients. *American Journal of Geriatric Psychiatry*, 4, S79-S83.
- Rabins, P. V., Mace, N. L., & Lucas, M. J. (1982). The impact of dementia on the family. *Journal of the American Medical Association*, 248, 333-335.
- Raskind, M. A., Sadowsky, C. H., Sigmund, W. R., Beitler, P. J., & Auster, S. B. (1997). Effect of tacrine on language, praxis, and noncognitive behavioral problems in Alzheimer disease. *Archives of Neurology*, 54, 836-840.
- Redick, R. W., Kramer, M., & Taube, C. A. (1977). Epidemiology of mental illness and utilization of psychiatric facilities among older persons. In E. W. Busse & W. Pfeiffer (Eds.), *Behavior and adaptation in late life*. Boston: Little, Brown.
- Regier, D. A., Boyd, J. H., Burke, J. D. Jr, Rae, D. S., Myers, J. K., Kramer, M., Robins, L. N., George, L. K., Karno, M., & Locke, B. Z. (1988). One-month prevalence of mental disorders in the United States. Based on five Epidemiologic Catchment Area sites. *Archives of General Psychiatry*, 45, 977-986.
- Reid, M. C., & Anderson, P. A. (1997). Geriatric substance use disorders. *Medical Clinics of North America*, 81, 999-1016.
- Reifler, B. V. (1997). The practice of geriatric psychiatry in three countries: Observations of an American in the British Isles. *International Journal of Geriatric Psychiatry*, 12, 795-807.
- Reifler, B. V., Cox, N. J., Jones, B. J., Rushing, J., & Yates, K. (in press). Service use and financial performance in a replication program for adult day centers. *American Journal of Geriatric Psychiatry*.
- Reifler, B. V., Henry, R. S., Rushing, J., Yates, M. K., Cox, N. J., Bradham, D. D., & McFarlane, M. (1997). Financial performance among adult day centers: Results of a national demonstration program. *Journal of the American Geriatrics Society*, 45, 146-153.
- Reifler, B. V., Teri, L., Raskind, M., Veith, R., Barnes, R., White, E., & McLean, P. (1989). Double-blind trial of imipramine in Alzheimer's disease patients with and without depression. *American Journal of Psychiatry*, 146, 45-49.
- Reynolds, C. F., III. (1998, March 8-11). *The challenge of treatment in 70+ year olds with recurrent major depression: Excellent short-term but brittle long-term response*. Annual meeting of the American Association for Geriatric Psychiatry, San Diego, CA.
- Reynolds, C. F., III, Frank, E., Perel, J. M., Imber, S. D., Cornes, C., Morycz, R. K., Mazumdar, S., Miller, M. D., Pollock, B. G., Rifai, A. H., Stack, J. A., George, C. J., Houck, P. R., & Kupfer, D. J. (1992). Combined pharmacotherapy and psychotherapy in the acute and continuation treatment of elderly patients with recurrent major depression: A preliminary report. *American Journal of Psychiatry*, 149, 1687-1692.
- Reynolds, C. F., III, Frank, E., Kupfer, D. J., Thase, M. E., Perel, J. M., Mazumdar, S., & Houck, P. R. (1996). Treatment outcome in recurrent major depression: A post hoc comparison of elderly ("young old") and midlife patients. *American Journal of Psychiatry*, 153, 1288-1292.
- Reynolds, C. F., III, Frank, E., Perel, J. M., Imber, S. D., Cornes, C., Miller, M. D., Mazumdar, S., Houck, P. R., Dew, M. A., Stack, J. A., Pollock, B. G., & Kupfer, D. J. (1999). Nortriptyline and interpersonal psychotherapy as maintenance therapies for recurrent major depression: A randomized controlled trial in patients older than 59 years. *Journal of the American Medical Association*, 281, 39-45.

- Reynolds, C. F., III, Frank, E., Perel, J. M., Mazumdar, S., & Kupfer, D. J. (1995). Maintenance therapies for late-life recurrent major depression: Research and review circa 1995. *International Psychogeriatrics*, *7*(Suppl.), 27-39.
- Reynolds, C. F., III, Frank, E., Perel, J. M., Miller, M. D., Cornes, C., Rifai, A. H., Pollock, B. G., Mazumdar, S., George, C. J., Houck, P. R., & Kupfer, D. J. (1994). Treatment of consecutive episodes of major depression in the elderly. *American Journal of Psychiatry*, *151*, 1740-1743.
- Reynolds, C. F., & Kupfer, D. J. (1999). Depression and aging: A look to the future. *Psychiatric Services*, *50*, 1167-1172.
- Rihmer, Z., Rutz, W., & Pihlgren, H. (1995). Depression and suicide on Gotland. An intensive study of all suicides before and after a depression-training programme for general practitioners. *Journal of Affective Disorders*, *35*, 147-152.
- Riley, T., Rawling-Sekunda, J., & Pernice, C. (1997). *Transitioning to managed care: Medicaid managed care in mental health* (3rd ed., The Kaiser-HCFA State Symposia Series). Portland, ME: National Academy for State Health Policy.
- Ritchie, K., & Kildea, D. (1995). Is senile dementia "age-related" or "ageing-related"? Evidence from meta-analysis of dementia prevalence in the oldest old. *Lancet*, *346*, 931-934.
- Ritchie, K., Touchon, J., & Ledesert, B. (1998). Progressive disability in senile dementia is accelerated in the presence of depression. *International Journal of Geriatric Psychiatry*, *13*, 459-461.
- Robins, C., & Clayton, R. R. (1989). Gender-related differences in psychoactive drug use among older adults. *Journal of Drug Issues*, 207-219.
- Robins, L. N., & Regier, D. A. (1991). *Psychiatric disorders in America: The Epidemiologic Catchment Area study*. New York: Free Press.
- Robinson, D., Napoliello, M. J., & Schenk, J. (1988). The safety and usefulness of buspirone as an anxiolytic drug in elderly versus young patients. *Clinical Therapeutics*, *10*, 740-746.
- Robinson, G. K. (1990a). *Mental health policy for older Americans: Protecting minds at risk*. Washington, DC: American Psychiatric Press.
- Robinson, G. K. (1990b). The psychiatric component of long-term care models. In B. S. Fogel, G. L. Gottlieb, & A. Furino (Eds.), *Mental health policy for older Americans: Protecting minds at risk* (pp. 157-178). Washington, DC: American Psychiatric Press.
- Rockwell, E., Jackson, E., Vilke, G., & Jeste, D. V. (1994). A study of delusion in a large cohort of Alzheimer's disease patients. *American Journal of Geriatric Psychiatry*, *2*, 157-164.
- Rodin, J., McAvay, G., & Timko, C. (1988). A longitudinal study of depressed mood and sleep disturbances in elderly adults. *Journal of Gerontology*, *43*, P45-P53.
- Rogers, S. L., Doody, R. S., Mohs, R. C., & Friedhoff, L. T. (1998). Donepezil improves cognition and global function in Alzheimer disease: A 15-week, double-blind, placebo-controlled study. Donepezil Study Group. *Archives of Internal Medicine*, *158*, 1021-1031.
- Rokhlina, M. L. (1975). [A comparative clinico-genetic study of attack-like schizophrenia with late and early manifestation with regard to age]. *Zhurnal Nevropatologii Ii Psikiatrii Imeni S.S. Korsakova*, *75*, 417-424.
- Romanoski, A. J., Folstein, M. F., Nestadt, G., Chahal, R., Merchant, A., Brown, C. H., Gruenberg, E. M., & McHugh, P. R. (1992). The epidemiology of psychiatrist-ascertained depression and DSM-III depressive disorders. Results from the Eastern Baltimore Mental Health Survey Clinical Reappraisal. *Psychological Medicine*, *22*, 629-655.
- Roose, S. P., Glassman, A. H., Giardina, E. G., Walsh, B. T., Woodring, S., & Bigger, J. T. (1987). Tricyclic antidepressants in depressed patients with cardiac conduction disease. *Archives of General Psychiatry*, *44*, 273-275.
- Rosenbach, M. L., & Ammering, C. J. (1997). Trends in Medicare part B mental health utilization and expenditures: 1987-92. *Health Care Financing Review*, *18*, 19-42.
- Rovner, B. W. (1993). Depression and increased risk of mortality in the nursing home patient. *American Journal of Medicine*, *94*, 19S-22S.
- Rowe, J. W., & Kahn, R. L. (1997). Successful aging. *Gerontologist*, *37*, 433-440.
- Rudorfer, M. V., Henry, M. E., & Sackheim, H. A. (1997). Electroconvulsive therapy. In A. Tasman, J. Kay, & J. A. Lieberman (Eds.), *Psychiatry* (pp. 1535-1556). Philadelphia: W.B. Saunders.

Mental Health: A Report of the Surgeon General

- Sackeim, H. A. (1994). Use of electroconvulsive therapy in late life depression. In L. S. Schneider, C. F. Reynolds, B. D. Lebowitz, & A. J. Friedhoff (Eds.), *Diagnosis and treatment of depression in late life* (pp. 259–277). Washington, DC: American Psychiatric Press.
- Salloway, S., Malloy, P., Kohn, R., Gillard, E., Duffy, J., Rogg, J., Tung, G., Richardson, E., Thomas, C., & Westlake, R. (1996). MRI and neuropsychological differences in early- and late-life-onset geriatric depression. *Neurology*, *46*, 1567–1574.
- Saltz, B. L., Woerner, M. G., Kane, J. M., Lieberman, J. A., Alvir, J. M., Bergmann, K. J., Blank, K., Koblenzer, J., & Kahaner, K. (1991). Prospective study of tardive dyskinesia incidence in the elderly. *Journal of the American Medical Association*, *266*, 2402–2406.
- Salzman, C. (1991). The APA Task Force report on benzodiazepine dependence, toxicity, and abuse [editorial]. *American Journal of Psychiatry*, *148*, 151–152.
- Sano, M., Ernesto, C., Thomas, R. G., Klauber, M. R., Schafer, K., Grundman, M., Woodbury, P., Growdon, J., Cotman, C. W., Pfeiffer, E., Schneider, L. S., & Thal, L. J. (1997). A controlled trial of selegiline, alpha-tocopherol, or both as treatment for Alzheimer's disease. The Alzheimer's Disease Cooperative Study. *New England Journal of Medicine*, *336*, 1216–1222.
- Saunders, P. A., Copeland, J. R., Dewey, M. E., Davidson, I. A., McWilliam, C., Sharma, V., & Sullivan, C. (1991). Heavy drinking as a risk factor for depression and dementia in elderly men. Findings from the Liverpool longitudinal community study. *British Journal of Psychiatry*, *159*, 213–216.
- Schafft, G. E., & Randolph, F. L. (1994). *Innovative community-based services for older persons with mental illness*. Rockville, MD: Center for Mental Health Services, Division of Demonstration Programs, Community Support Section.
- Schneider, L. S. (1995). Efficacy of clinical treatment for mental disorders among older persons. In M. Gatz (Ed.), *Emerging issues in mental health and aging* (pp. 19–71). Washington, DC: American Psychological Association.
- Schneider, L. S. (1996). Pharmacological considerations in the treatment of late life depression. *American Journal of Geriatric Psychiatry*, *4*, S51–S65.
- Schnelle, J. F., Cruise, P. A., Rahman, A., & Ouslander, J. G. (1998). Developing rehabilitative behavioral interventions for long-term care: Technology transfer, acceptance, and maintenance issues. *Journal of the American Geriatrics Society*, *46*, 771–777.
- Schnelle, J. F., MacRae, P. G., Ouslander, J. G., Simmons, S. F., & Nitta, M. (1995). Functional incidental training, mobility performance, and incontinence care with nursing home residents. *Journal of the American Geriatrics Society*, *43*, 1356–1362.
- Schulberg, H. C., Madonia, M. J., Block, M. R., Coulehan, J. L., Scott, C. P., Rodriguez, E., & Black, A. (1995). Major depression in primary care practice. Clinical characteristics and treatment implications. *Psychosomatics*, *36*, 129–137.
- Schultz, R., O'Brien, A. T., Bookwala, J., & Fleissner, K. (1995). Psychiatric and physical morbidity effects of dementia caregiving: Prevalence, correlates, and causes. *Gerontologist*, *35*, 771–791.
- Scogin, F., & McElreath, L. (1994). Efficacy of psychosocial treatments for geriatric depression: A quantitative review. *Journal of Consulting and Clinical Psychology*, *62*, 69–74.
- Scott-Lennox, J. A., & George, L. (1996). Epidemiology of psychiatric disorders and mental health services use among older Americans. In B. L. Levin & J. Petriola (Eds.), *Mental health services: A public health perspective* (pp. 253–289). New York: Oxford University Press.
- Seidel, W. F., Cohen, S. A., Bilwise, N. G., & Dement, W. C. (1995). Buspirone: An anxiolytic without sedative effect. *Psychopharmacology*, *87*, 371–373.
- Semke, J., & Jensen, J. (1997). High utilization of inpatient psychiatric services by older adults. *Psychiatric Services*, *48*, 172–174, 176.
- Sheikh, J. I. (1994). Anxiety disorders. In C. E. Coffey & J. L. Cummings (Eds.), *The American Psychiatric Press textbook of geriatric neuropsychiatry* (pp. 280–296). Washington, DC: American Psychiatric Press.
- Sheldon, A., McEwan, P. J. M., & Ryser, C. P. (1975). *Retirement: Patterns and predictions* (DHEW Publication No. ADM 74-49). Washington, DC: U.S. Government Printing Office.
- Shorr, R. I., Bauwens, S. F., & Landefeld, C. S. (1990). Failure to limit quantities of benzodiazepine hypnotic drugs for outpatients: Placing the elderly at risk. *American Journal of Medicine*, *89*, 725–732.

- Siegler, I. C., Poon, L. W., Madden, D. J., & Welsh, K. A. (1996). Psychological aspects of normal aging. In E. W. Busse & D. G. Blazer (Eds.), *The American Psychiatric Press textbook of geriatric psychiatry* (2nd ed., pp. 105–128). Washington, DC: American Psychiatric Press.
- Silverman, P. R. (1988). Widow-to-widow: A mutual help program for the widowed. In R. Price, E. Cowen, R. P. Lorian, & J. J. Ramos-McKay (Eds.), *Fourteen ounces of prevention: A casebook for practitioners* (pp. 175–186). Washington, DC: American Psychological Association.
- Skoog, I. (1993). The prevalence of psychotic, depressive, and anxiety syndromes in demented and non-demented 85-year olds. *International Journal of Geriatric Psychiatry*, 8, 247–253.
- Skoog, I., Aevarsson, O., & Beskow, J. (1996). Suicidal feelings in a population sample of 85 year olds. *American Journal of Psychiatry*, 153, 1015–1020.
- Sloane, R. B., Staples, F. R., & Schneider, L. S. (1985). Interpersonal therapy vs. nortriptyline for depression in the elderly. In G. D. Burrows, T. R. Norman, & L. Denerstein (Eds.), *Clinical and pharmacological studies in psychiatric disorders* (pp. 344–346). London: John Libby.
- Small, G. W., Rabins, P. V., Barry, P. P., Buckholtz, N. S., DeKosky, S. T., Ferris, S. H., Finkel, S. I., Gwyther, L. P., Khachaturian, Z. S., Lebowitz, B. D., McRae, T. D., Morris, J. C., Oakley, F., Schneider, L. S., Streim, J. E., Sunderland, T., Teri, L. A., & Tune, L. E. (1997). Diagnosis and treatment of Alzheimer's disease and related disorders. Consensus statement of the American Association for Geriatric Psychiatry, the Alzheimer's Association, and the American Geriatrics Society. *Journal of the American Medical Association*, 278, 1363–1371.
- Small, G. W., & Salzman, C. (1998). Treatment of depression with new and atypical antidepressants. In C. Salzman (Ed.), *Clinical geriatric psychopharmacology* (pp. 245–261). Baltimore: Williams & Wilkins.
- Smith, S. L., Sherrill, K. A., & Colenda, C. C. (1995). Assessing and treating anxiety in elderly persons. *Psychiatric Services*, 46, 36–42.
- Special Committee on Aging. (1987). *Medicare prescription drug issues. Report to the Chairman, Special Committee on Aging*. Washington, DC: General Accounting Office.
- Stanley, M. A., Beck, J. G., & Glassco, J. D. (1996). Treatment of generalized anxiety in older adults: A preliminary comparison of cognitive-behavioral and supportive approaches. *Behavioral Therapy*, 27, 565–581.
- Steffens, D. C., & Krishnan, K. R. (1998). Structural neuroimaging and mood disorders: Recent findings, implications for classification, and future directions. *Biological Psychiatry*, 43, 705–712.
- Stern, Y., Gurland, B., Tatemichi, T. K., Tang, M. X., Wilder, D., & Mayeux, R. (1994). Influence of education and occupation on the incidence of Alzheimer's disease. *Journal of the American Medical Association*, 271, 1004–1010.
- Stevens, A. B., Burgio, L. D., Bailey, E., Burgio, K. L., Paul, P., Capilouto, E., Nicovich, P., & Hale, G. (1998). Teaching and maintaining behavior management skills with nursing assistants in a nursing home. *Gerontologist*, 38, 379–384.
- Stolee, P., Kessler, L., & Le Clair, J. K. (1996). A community development and outreach program in geriatric mental health: Four years' experience. *Journal of the American Geriatrics Society*, 44, 314–320.
- Sturm, R. (1997). How expensive is unlimited mental health care coverage under managed care? *Journal of the American Medical Association*, 278, 1533–1537.
- Sturm, R., & Wells, K. B. (1995). How can care for depression become more cost-effective? *Journal of the American Medical Association*, 273, 51–58.
- Sultzer, D. L., Mahler, M. E., Mandelkern, M. A., Cummings, J. L., Van Gorp, W. G., Hinkin, C. H., & Berisford, M. A. (1995). The relationship between psychiatric symptoms and regional cortical metabolism in Alzheimer's disease. *Journal of Neuropsychiatry and Clinical Neurosciences*, 7, 476–484.
- Sussman, M. B. (1985). The family life of old people. In R. H. Binstock & E. Shanas (Eds.), *Handbook of aging and social sciences* (pp. 415–449). New York: Van Nostrand Reinhold.
- Swartz, M., Landerman, R., George, L. K., Melville, M. L., Blazer, D., & Smith, K. (1991). Benzodiazepine anti-anxiety agents: Prevalence and correlates of use in a southern community. *American Journal of Public Health*, 81, 592–596.
- Taube, C. A., Goldman, H. H., & Salkever, D. (1990). Medicaid coverage for mental illness: Balancing access and costs. *Health Affairs*, 9, 5–18.

Mental Health: A Report of the Surgeon General

- Temple, M. T., & Leino, E. V. (1989). Long-term outcomes of drinking: A 20-year longitudinal study of men. *British Journal of Addiction*, *84*, 889–899.
- Teri, L., & Gallagher-Thompson, D. (1991). Cognitive-behavioral interventions for treatment of depression in Alzheimer's patients. *Gerontologist*, *31*, 413–416.
- Teri, L., Logsdon, R. G., Uomoto, J., & McCurry, S. M. (1997). Behavioral treatment of depression in dementia patients: A controlled clinical trial. *Journals of Gerontology. Series B, Psychological Sciences and Social Sciences*, *52*, 156–166.
- Teri, L., & Uomoto, J. (1991). Reducing excess disability in dementia patients: Training caregivers to manage patient depression. *Clinical Gerontologist*, *10*, 49–63.
- Thal, L. J. (1996). Cholinomimetic treatment of Alzheimer's disease. *Progress in Brain Research*, *109*, 299–309.
- Thomas, C. S., & Lewis, S. (1998). Which atypical antipsychotic [editorial]? *British Journal of Psychiatry*, *172*, 106–109.
- Trieman, N., Wills, W., & Leff, J. (1996). TAPS project 28: Does reprovision benefit elderly long-stay mental patients? *Schizophrenia Research*, *21*, 199–208.
- Tsuang, M. T., & Woolson, R. F. (1997). Mortality in patients with schizophrenia, mania, depression and surgical conditions: A comparison with general population mortality. *British Journal of Psychiatry*, *130*, 162–166.
- Unutzer, J., Katon, W. J., Simon, G., Walker, E. A., Grembowski, D., & Patrick, D. (1996). Depression, quality of life, and use of health services in primary care patients over 65: A 4-year prospective study. *Psychosomatics*, *37*, 35.
- Unutzer, J., Katon, W., Sullivan, M., & Miranda, J. (1997a, November 6). *The effectiveness of treatments for depressed older adults in primary care*. Paper presented at Exploring Opportunities to Advance Mental Health Care for an Aging Population, meeting sponsored by the John A. Hartford Foundation, Rockville, MD.
- Unutzer, J., Patrick, D. L., Simon, G., Grembowski, D., Walker, E., Rutter, C., & Katon, W. (1997b). Depressive symptoms and the cost of health services in HMO patients aged 65 years and older. A 4-year prospective study. *Journal of the American Medical Association*, *277*, 1618–1623.
- U.S. Department of Health and Human Services, Administration on Aging, and American Association of Retired Persons (D. G. Fowles, Comp.). (1995). *A profile of older Americans*. Washington, DC: American Association of Retired Persons.
- Vachon, M. L. S. (1979). *Identify change over the first two years of bereavement: Social relationships and social support in widowhood*. Unpublished doctoral dissertation, York University, Toronto, Canada.
- Vachon, M. L., Lyall, W. A., Rogers, J., Freedman-Letofsky, K., & Freeman, S. J. (1980). A controlled study of self-help intervention for widows. *American Journal of Psychiatry*, *137*, 1380–1384.
- Vachon, M. L., Sheldon, A. R., Lancee, W. J., Lyall, W. A., Rogers, J., & Freeman, S. J. (1982). Correlates of enduring distress patterns following bereavement: Social network, life situation and personality. *Psychological Medicine*, *12*, 783–788.
- Wallace, J., & O'Hara, M. W. (1992). Increases in depressive symptomatology in the rural elderly: Results from a cross-sectional and longitudinal study. *Journal of Abnormal Psychology*, *101*, 398–404.
- Ware, J. E., Jr., Bayliss, M. S., Rogers, W. H., Kosinski, M., & Tarlov, A. R. (1996). Differences in 4-year health outcomes for elderly and poor, chronically ill patients treated in HMO and fee-for-service systems. Results from the Medical Outcomes Study. *Journal of the American Medical Association*, *276*, 1039–1047.
- Waters, E. (1995). Let's not wait till it's broke: Interventions to maintain and enhance mental health in late life. In M. Gatz (Ed.), *Emerging issues in the mental health and aging* (pp. 183–209). Washington, DC: American Psychological Association.
- Weinberger, D. R. (1987). Implications of normal brain development for the pathogenesis of schizophrenia. *Archives of General Psychiatry*, *44*, 660–669.
- Weiner, R. D., & Krystal, A. D. (1994). The present use of electroconvulsive therapy. *Annual Review of Medicine*, *45*, 273–281.
- Weissman, M. A., Bruce, M., Leaf, P. J., Florio, L. P., & Holzer C., III. (1991). Affective disorders. In L. N. Robins & D. A. Regier (Eds.), *Psychiatric disorders in America: The Epidemiologic Catchment Area study* (pp. 53–80). New York: Free Press.