

- KNUDSON, R.J., BURROWS, B., LEBOWITZ, M.D. The maximal expiratory flow-volume curve: Its use in the detection of ventilatory abnormalities in a population study. *American Review of Respiratory Disease* 114(3): 871-879, November 1976.
- KNUDSON, R.J., CLARK, D.F., KENNEDY, T.C., KNUDSON, D.W. Effect of aging alone on mechanical properties of the normal adult human lung. *Journal of Applied Physiology: Respiratory, Environmental and Exercise Physiology* 43(6): 1054-1062, December 1977.
- KNUDSON, R.J., LEBOWITZ, M.D. Comparison of flow-volume and closing volume variables in a random population. *American Review of Respiratory Disease* 116(6): 1039-1045, December 1977.
- KNUDSON, R.J., KALTENBORN, W.T. Evaluation of lung elastic recoil by exponential curve analysis. *Respiratory Physiology* 46: 29-42, 1981.
- KNUDSON, R.J., LEBOWITZ, M.D., HOLBERG, C.J., BURROWS, B. Changes in the normal maximal expiratory flow-volume curve with growth and aging. *American Review of Respiratory Disease* 127(6): 725-734, June 1983.
- KRUMHOLZ, R.A., HEDRICK, E.C. Pulmonary function differences in normal smoking and nonsmoking, middle-aged, white-collar workers. *American Review of Respiratory Disease* 107(2): 225-230, February 1973.
- KRZYZANOWSKI, M. Changes of ventilatory capacity in an adult population during a five-year period. *Bulletin Europeen de Physiopathologie Respiratoire* 16(2): 155-170, 1980.
- KUEPPERS, F., BLACK, L.F. Alpha₁-antitrypsin and its deficiency. *American Review of Respiratory Disease* 110(2): 176-194, August 1974.
- KUEPPERS, F., MILLER, R.D., GORDON, H., HEPPEL, N.G., OFFORD, K. Familial prevalence of chronic obstructive pulmonary disease in a matched pair study. *American Journal of Medicine* 63: 336-342, September 1977.
- LAM, S., ABOUD, R.T., CHAN-YEUNG, M., TAN, F. Use of maximal expiratory flow-volume curves with air and helium-oxygen in the detection of ventilatory abnormalities in population surveys. *American Review of Respiratory Disease* 123(2): 234-237, February 1981.
- LARSSON, C. Natural history and life expectancy in severe alpha₁-antitrypsin deficiency, PiZ. *Acta Medica Scandinavica* 204(5): 345-351, May 1978.
- LARSSON, C., ERIKSSON, S., DIRKSEN, H. Smoking and intermediate alpha₁-antitrypsin deficiency and lung function in middle-aged men. *British Medical Journal* 2(6092): 922-925, October 8, 1977.
- LEBOWITZ, M.D. Occupational exposures in relation to symptomatology and lung function in a community population. *Environmental Research* 14(1): 59-67, August 1977a.
- LEBOWITZ, M.D. The relationship of socio-environmental factors to the prevalence of obstructive lung diseases and other chronic conditions. *Journal of Chronic Diseases* 30(9): 599-611, September 1977b.
- LEBOWITZ, M.D. Respiratory symptoms and disease related to alcohol consumption. *American Review of Respiratory Disease* 123(1): 16-18, January 1981.
- LEBOWITZ, M.D., BURROWS, B. Quantitative relationships between cigarette smoking and chronic productive cough. *International Journal of Epidemiology* 6(2): 107-113, June 1977.
- LEBOWITZ, M.D., KNUDSON, R.J., BURROWS, B. Tucson Epidemiologic study of obstructive lung diseases. I: Methodology and prevalence of disease. *American Journal of Epidemiology* 102(2): 137-152, August 1975.
- LEBOWITZ, M.D., KNUDSON, R.J., BURROWS, B. Family aggregation of pulmonary function measurements. *American Review of Respiratory Disease* 129(1): 8-11, 1984.
- LEBOWITZ, M.D., KNUDSON, R.J., BURROWS, B. Risk factors for airways obstructive diseases: A multiple logistics approach. *Chest* 85(6): 11s-12s, June 1984, Supplement.

- LEEDER, S.R., COLLEY, J.R.T., CORKHILL, R., HOLLAND, W.W. Change in respiratory symptom prevalence in adults who alter their smoking habits. *American Journal of Epidemiology* 105(6): 522-529, 1977.
- LEFCOE, N.M., ASHLEY, M.J., PEDERSON, L.L., KEAYS, J.J. The health risks of passive smoking. The growing case for control measures in enclosed environments. *Chest* 84(1): 90-95, 1983.
- LEFCOE, N.M., WONNACOTT, T.H. Chronic respiratory disease in four occupational groups. *Archives of Environmental Health* 29(3): 143-146, September 1974.
- LIANG, A., MACFIE, A.E., HARRIS, E.A., WHITLOCK, R.M.L. Transit-time analysis of the forced expiratory spiogram during clinical remission in juvenile asthma. *Thorax* 34(2): 194-199, April 1979.
- LIARD, R., PEDRIZET, S., COOREMAN, J., BIDOU, S. Smoking and chronic respiratory symptoms: Prevalence in male and female smokers. *American Journal of Public Health* 70(3): 271-273, March 1980.
- LINN, W.W., HACKNEY, J.D., PEDERSEN, E.E., BREISACHER, P., PATTERSON, J.V., MULRY, C.A., COYLE, J.F. Respiratory function and symptoms in urban office workers in relation to oxidant air pollution exposure. *American Review of Respiratory Disease* 114(3): 477-483, September 1976.
- LOVE, R.G., MILLER, B.G., Longitudinal study of lung function in coal-miners. *Thorax* 37(3): 193-197, March 1982.
- MACFIE, A.E., HARRIS, E.A., WHITLOCK, R.M.L. Transit-time analysis of the forced spiogram in healthy children and adults. *Journal of Applied Physiology: Respiratory, Environmental and Exercise Physiology* 46(2): 263-267, February 1979.
- MACKLEM, P.T., MEAD, J. Resistance of central and peripheral airways measured by retrograde catheter. *Journal of Applied Physiology* 22(3): 395-401, March 1967.
- MACNEE, W., POWER, J., INNES, A., DOUGLAS, N.J., SUDLOW, M.F. The dependence of maximal flow in man on the airway gas physical properties. *Clinical Science* 65(3): 273-279, September 1983.
- MADISON, R., MITTMAN, C., AFIFI, A.A., ZELMAN, R. Risk factors for obstructive lung disease. *American Review of Respiratory Disease* 124: 149-153, August 1981.
- MANFREDA, J., NELSON, N., CHERNIACK, R.M. Prevalence of respiratory abnormalities in a rural and an urban community. *American Review of Respiratory Disease* 117(2): 215-226, February 1978.
- MARAZZINI, L., PELOSI, V., VEZZOLI, F., PENNASI, R., LONGHINI, E. Prospective study of airway obstruction in a population with small airway disease. *Bulletin Europeen de Physiopathologie Respiratoire* 13(2): 219-229, March-April 1977.
- MARAZZINI, L., VEZZOLI, F., LONGHINI, E. Respiratory function 8 years after a diagnosis of peripheral airways disease. *Respiration* 42(2): 88-97, 1981.
- MARCQ, M., MINETTE, A. Lung function changes in smokers with normal conventional spirometry. *American Review of Respiratory Disease* 114(4): 723-738, October 1976.
- McCARTHY, D.S., CRAIG, D.B., CHERNIACK, R.M. Effect of modification of the smoking habit on lung function. *American Review of Respiratory Disease* 114(1): 103-113, July 1976.
- McCARTHY, D.S., SPENCER, R., GREENE, R., MICLIC-EMILI, J. Measurement of "closing volume" as a simple and sensitive test for early detection of small airway disease. *American Journal of Medicine* 52: 747-753, June 1972.
- MEAD, J. Problems in interpreting common tests of pulmonary mechanical function. In: Macklem, P.T., Permutt, S. (Editors). *The Lung in the Transition Between Health and Disease*. New York, Marcel Dekker, Inc., 1979, pp. 43-51.
- MEDICAL RESEARCH COUNCIL COMMITTEE ON THE AETIOLOGY OF CHRONIC BRONCHITIS. Standardized questionnaires on respiratory symptoms. *British Medical Journal* 2(5213): 1665, December 3, 1960.

- MENDELSSOHN, A. *Journal of the Russian National Health Society*, September 1897. *Lancet* 2(3867): 952-953, 1897.
- MICHAELS, R., SIGURDSON, M., THURLBECK, S., CHERNIACK, R. Elastic recoil of the lung in cigarette smokers: the effect of nebulized bronchodilator and cessation of smoking. *American Review of Respiratory Disease* 119(5): 707-716, May 1979.
- MILLER, A., THORNTON, J.C., WARSHAW, R., ANDERSON, H., TEIRSTEIN, A.S., SELIKOFF, I.J. Single breath diffusing capacity in a representative sample of the population of Michigan, a large industrial state. Predicted values, lower limits of normal, and frequencies of abnormality by smoking history. *American Review of Respiratory Disease* 127(3): 270-277, March 1983.
- MILNE, J.S. Longitudinal respiratory studies in older people. *Thorax* 33: 547-554, 1978.
- MIMICA, M. *Study of Regional Prevalence and Incidence of Chronic Bronchitis and Asthma in Adults*. Final Report. U.S. Department of Health, Education, and Welfare, Public Health Service, National Institutes of Health, January 1975.
- MINK, S.N., WOOD, L.D.H. How does HeO₂ increase maximum expiratory flow in human lungs? *Journal of Clinical Investigation* 66(4): 720-729, October 1980.
- MITCHELL, R.S., STANFORD, R.E., JOHNSON, J.M., SILVERS, G.W., DART, G., GEORGE, M.S. The morphologic features of the bronchi, bronchioles and alveoli in chronic airway obstruction: A clinicopathologic study. *American Review of Respiratory Disease* 114: 137-145, July 1976.
- MITTMANN, C. The PiMZ phenotype: Is it a significant risk factor for the development of chronic obstructive lung disease? *American Review of Respiratory Disease* 118(4): 649-652, October 1978.
- MONSON, R.R. *Occupational Epidemiology*. Boca Raton, Florida, CRC Press, Inc., 1980.
- MORRIS, J.F., KOSKI, A., JOHNSON, L.C. Spirometric standards for healthy nonsmoking adults. *American Review of Respiratory Disease* 103(1): 57-67, January 1971.
- MORSE, J.O. Alpha₁-antitrypsin deficiency. *New England Journal of Medicine* 299(19): 1045-1048, November 9, 1978 and 299(20): 1099-1105, November 16, 1978.
- MUELLER, R.E., KEBLE, D.L., PLUMMER, J., WALKER, S.H. The prevalence of chronic bronchitis, chronic airway obstruction, and respiratory symptoms in a Colorado city. *American Review of Respiratory Disease* 103(2): 209-228, February 1971.
- MUELLER, W.H., CHAKRABORTY, R., BARTON, S.A., ROTHHAMMER, F., SCHULL, W.J. Genes and epidemiology in anthropological adaptation studies: Familial correlations in lung function in populations residing at different altitudes in Chile. *Medical Anthropology* 4(3): 367-384, Summer 1980.
- MULTIPLE RISK FACTOR INTERVENTION TRIAL RESEARCH GROUP. Risk factor changes and mortality results. *Journal of the American Medical Association* 248(12): 1465-1477, September 24, 1982.
- NADEL, J.A., COMROE, J.H., Jr. Acute effects of inhalation of cigarette smoke on airway conductance. *Journal of Applied Physiology* 16(4): 713-716, July 1961.
- NATIONAL CENTER FOR HEALTH STATISTICS. *Vital and Health Statistics Series No. 136*. DHHS Publication No. (PHS) 81-1564, 1981.
- NATIONAL CENTER FOR HEALTH STATISTICS. Unpublished data.
- NEMERY, B., MOAVERO, N.E., BRASSEUR, L., STANESCU, D.C. Significance of small airway tests in middle-aged smokers. *American Review of Respiratory Disease* 124(3): 232-238, September 1981.
- NERI, L.C., MANDEL, J.S., HEWITT, D., JURKOWSKI, D. Chronic obstructive pulmonary disease in two cities of contrasting air quality. *Canadian Medical Association Journal* 113(11): 1043-1046, December 13, 1975.

- NEUBURGER, N., LEVISON, H., KRUGER, K. Transit time analysis of the forced expiratory vital capacity in cystic fibrosis. *American Review of Respiratory Disease* 114(4): 753-759, October 1976.
- NEUKIRCH, F., LIARD, R., COOREMAN, J., PERDRIZET, S. Prevalence of respiratory symptoms in Parisian teenagers according to smoking habits. *Journal of Epidemiology and Community Health* 36(3): 202-204, September 1982.
- O'DONNELL, T.V., DE HAMEL, F. Objective respiratory screening among Dunedin public servants. *New Zealand Medical Journal* 84: 99-103, August 11, 1976.
- OGILVIE, A.G., STRANG, C., LEGGAT, P.O., NEWELL, D.J. *A Ten-Year Prospective Study of Chronic Bronchitis in the North-East of England*. Edinburgh, Churchill Livingstone, 1973.
- ORELL, S.R., MAZODIER, P. Pathological findings in α_1 -antitrypsin deficiency. In: Mittman, C. (Editor). *Pulmonary Emphysema and Proteolysis*. New York, Academic Press, 1972, pp. 69-88.
- ORIE, N.G., SLUITER, H.J., DeVRIES, K., TAMMELING, G.J., WITKOP, J. The host factor in bronchitis. In: Orie, N.G.M., Sluiter, H.J. (Editors). *Bronchitis, An International Symposium*, April 27-29, 1960. Assen, Netherlands, Charles C Thomas, Royal Vangorcum, 1961, pp. 43-59.
- OSMAN, M., CANTOR, J., ROFFMAN, S., TURINO, G.M., MANDL, I. Tobacco smoke exposure retards elastin repair in experimental emphysema. *American Review of Respiratory Disease* 125(4): 213A, April 1982.
- OXHOJ, H., BAKE, B., WILHELMSEN, L. Spirometry and flow-volume curves in 10-year follow-up in men born 1913. *Scandinavian Journal of Respiratory Diseases* 57(6): 310-312, 1976.
- PARE, P.D., BROOKS, L.A., BATES, J., LAWSON, L.M., NELEMS, J.M.B., WRIGHT, J.L., HOGG, J.C. Exponential analysis of the lung pressure-volume curve as a predictor of pulmonary emphysema. *American Review of Respiratory Disease* 126(1): 54-61, July 1982.
- PARK, S.S., JANIS, M., SHIM, C.S., WILLIAMS, M.H., Jr. Relationship of bronchitis and emphysema to altered pulmonary function. *American Review of Respiratory Disease* 102(6): 927-936, December 1970.
- PARK, S.S., KIKKAWA, Y., GOLDRING, I.P., DALY, M.M., ZELEFSKY, M., SHIM, C., SPIERER, M., MORITA, T. An animal model of cigarette smoking in beagle dogs: Correlative evaluation of effects on pulmonary function, defense, and morphology. *American Review of Respiratory Disease* 115(6): 971-979, June 1977.
- PARK, T.K. Cigarette smoking patterns among students attending a Christian university in Seoul, Korea. *Yonsei Medical Journal* 22(2): 145-154, 1981.
- PERMUTT, S., MENKES, H.A. Spirometry. Analysis of forced expiration within the time domain. In: Macklem, P.T., Permutt, S. (Editors). *The Lung in the Transition Between Health and Disease*. New York, Marcel Dekker, 1979, pp. 113-152.
- PETO, R., SPEIZER, F.E., COCHRANE, A.L., MOORE, F., FLETCHER, C.M., TINKER, C.M., HIGGINS, I.T.T., GRAY, R.G., RICHARDS, S.M., GILLILAND, J., NORMAN-SMITH, B. The relevance in adults of air-flow obstruction, but not of mucus hypersecretion, to mortality from chronic lung disease: Results from 20 years of prospective observation. *American Review of Respiratory Disease* 128(3): 491-500, September 1983.
- PETTY, T.L., PIERSON, D.J., DICK, N.P., HUDSON, L.P., WALKER, S.H. Follow-up evaluation of a prevalence study for chronic bronchitis and chronic airway obstruction. *American Review of Respiratory Disease* 114(3): 881-890, November 1976.
- PETTY, T.L., RYAN, S.F., MITCHELL, R.S. Cigarette smoking and the lungs: Relation to postmortem evidence of emphysema, chronic bronchitis and black lung pigmentation. *Archives of Environmental Health* 14: 172-177, January 1967.

- PETTY, T.L., SILVERS, G.W., STANFORD, R.E. Functional correlations with mild and moderate emphysema in excised human lungs. *American Review of Respiratory Disease* 125(6): 700-704, December 1981.
- POUKKULA, A., HUHTI, E., MAKARAINEN, M. Chronic respiratory disease among workers in a pulp mill. A ten-year follow-up study. *Chest* 81(3): 285-289, March 3, 1982.
- QIZILBASH, A., YOUNG-PONG, O. Alpha₁-antitrypsin liver disease in differential diagnosis of PAS-positive, diastase-resistant globules in liver cells. *American Journal of Clinical Pathology* 79(6): 697-702, June 1983.
- RAWBONE, R.G., KEELING, C.A., JENKINS, A., GUZ, A. Cigarette smoking among secondary school children in 1975. Prevalence of respiratory symptoms, knowledge of health hazards, and attitudes to smoking and health. *Journal of Epidemiology and Community Health* 32(1): 53-58, March 1978.
- REID, L.M. Pathology of chronic bronchitis. *Lancet* 1(6806): 275-278, February 6, 1954.
- ROBERTSON, D.G., WARREL, D.A., NEWTON-HOWES, J.S., FLETCHER, C.M. Bronchial reactivity to cigarette and cigar smoke. *British Medical Journal* 3(5665): 269-271, August 2, 1969.
- ROKAW, S.N., DETELS, R., COULSON, A.H., SAYRE, J.W., TASHKIN, D.P., ALLWRIGHT, S.S., MASSEY, F.J., Jr. The UCLA population studies of chronic obstructive respiratory disease. 3. Comparison of pulmonary function in three communities exposed to photochemical oxidants, multiple primary pollutants, or minimal pollutants. *Chest* 78: 252-262, August 1980.
- RYAN, S.F., VINCENT, T.N., MITCHELL, R.S., FILLEY, G.F., DART, G. Ductectasia: An asymptomatic pulmonary change related to age. *Medicina Thoracalis* 22(2): 181-187, 1965.
- RYDER, R.C., DUNNILL, M.S., ANDERSON, J.A. A quantitative study of bronchial mucous gland volume emphysema and smoking in a necropsy population. *Journal of Pathology* 104(1): 59-71, May 1971.
- RYDER, R.C., THURLBECK, W.M., GOUGH, J. A study of the interobserver variation in the assessment of the amount of pulmonary emphysema in paper-mounted whole lung sections. *American Review of Respiratory Disease* 99: 354-364, March 1969.
- SALMON, R.B., SAIDEL, G.M., INKLEY, S.R., NIEWOEHNER, D.E. Relationship of ventilation inhomogeneity to morphologic variables in excised human lungs. *American Review of Respiratory Disease* 126(4): 686-690, October 1982.
- SAMET, J.M. A historical and epidemiologic perspective on respiratory symptoms questionnaires. *American Journal of Epidemiology* 108(6): 435-446, December 1978.
- SAMET, J.M., SCHRAG, S.D., HOWARD, C.A., KEY, C.R., PATHAK, D.R. Respiratory disease in a New Mexico population sample of Hispanic and non-Hispanic whites. *American Review of Respiratory Disease* 125(2): 152-157, February 1982.
- SAMET, J.M., TAGER, I.B., SPEIZER, F.E. The relationship between respiratory illness in childhood and chronic air-flow obstruction in adulthood. *American Review of Respiratory Disease* 127(4): 508-523, April 1983.
- SARIC, M., LUCIC-PALAIC, S., HORTON, R.J.M. Chronic nonspecific lung disease and alcohol consumption. *Environmental Research* 14(1): 14-21, August 1977.
- SAWICKI, F. *Regression Persistence and Incidence of Chronic Bronchitis in a Sample of an Urban Population Followed Up for 5 Years*. Presented at the 8th International Scientific Meeting of the International Epidemiological Association, Puerto Rico, September 1977.
- SCHENKER, M.B., SAMET, J.M., SPEIZER, F.E. Effect of cigarette tar content and smoking habits on respiratory symptoms in women. *American Review of Respiratory Disease* 125(6): 684-690, June 1982.
- SCHILLING, R.S.F., LETAI, A.D., HUI, S.L., BECK, G.J., SCHOENBERG, J.B., BOUHUY, A. Lung function, respiratory disease, and smoking in families. *American Journal of Epidemiology* 106(4): 274-283, October 1977.

- SCHLESINGER, Z., GOLDBOURT, U., MEDALIE, J.H., RISS, E., NEUFELD, H.N., ORON, D. Pulmonary function and respiratory disease among adult male Israelis. Variations by age and birth place. *Israel Journal of Medical Sciences* 8(7): 957-964, July 1972.
- SCHUYLER, M.R., RYNBRANDT, D.J., KLEINERMAN, J. Physiologic and morphologic observations of the effects of intravenous elastase on the lung. *American Review of Respiratory Disease* 117(1): 97-102, January 1978.
- SEELY, J.E., ZUSKIN, E., BOUHUYS, A. Cigarette smoking: Objective evidence for lung damage in teen-agers. *Science* 172(3983): 741-743, May 14, 1971.
- SHARP, J.T., PAUL, O., LEPPER, M.H., McKEAN, H., SAXON, G.A., Jr. Prevalence of chronic bronchitis in an American male urban industrial population. *American Review of Respiratory Disease* 91(4): 510-520, April 1965.
- SHARP, J.T., PAUL, O., McKEAN, H., BEST, W.R. A longitudinal study of bronchitic symptoms and spirometry in a middle-aged, male, industrial population. *American Review of Respiratory Disease* 108(5): 1066-1077, November 1973.
- SILVERS, G.W., PETTY, T.L., STANFORD, R.E. Elastic recoil changes in early emphysema. *Thorax* 35: 490-495, July 1980.
- SIMONSSON, B. Effect of cigarette smoking in the forced expiratory flow rate. *American Review of Respiratory Disease* 85: 543-539, 1962.
- SLUIS-CREMER, G.K., SICHEL, H.S. Ventilatory function in males in a Witwatersrand town. Comparison between smokers and nonsmokers. *American Review of Respiratory Disease* 98(2): 229-239, August 1968.
- SOBOL, B.J., VAN VOORHIES, L., EMIRGIL, C. Detection of acute effects of cigarette smoking on airway dynamics: A critical and comparative study of pulmonary function tests. *Thorax* 32(3): 312-216, June 1977.
- SPAIN, D.M., SIEGEL, H., BRADESS, V.A. Emphysema in apparently healthy adults. Smoking, age, and sex. *Journal of the American Medical Association* 224(3): 322-325, April 16, 1973.
- SPARROW, D., ROSNER, B., COHEN, M., WEISS, S.T. Alcohol consumption and pulmonary function. A cross-sectional and longitudinal study. *American Review of Respiratory Disease* 127(6): 735-738, June 1983a.
- SPARROW, D., STEFOS, T., BOSSE, R., WEISS, S.T. The relationship of tar content to decline in pulmonary function in cigarette smokers. *American Review of Respiratory Disease* 127(1): 56-58, January 1983b.
- SPEIZER, F.E., TAGER, I.B. Epidemiology of chronic mucus hypersecretion and obstructive airways disease. *Epidemiologic Reviews* 1: 124-142, 1979.
- STANESCU, D.C., VERITER, C., FRANS, A., BRASSEUR, L. Maximal expiratory flow rates and "closing volume" in asymptomatic healthy smokers. *Scandinavian Journal of Respiratory Disease* 54(4): 264-271, 1973.
- STEBBINGS, J.H., Jr. Chronic respiratory disease among nonsmokers in Hagerstown, Maryland. III. Social class and chronic respiratory disease. *Environmental Research* 4(3): 213-232, August 1971.
- STERLING, G.M. Mechanism of bronchoconstriction caused by cigarette smoking. *British Medical Journal* 3(5560): 275-277, July 29, 1967.
- STUART-HARRIS, C.H. Chronic bronchitis (Part I). *Abstracts of World Medicine* 42(9): 649-669, September 1968a.
- STUART-HARRIS, C.H. Chronic bronchitis (Part II). *Abstracts of World Medicine* 42(10): 737-751, October 1968b.
- SUTINEN, S., VAAJALAHTI, P., PAAKKO, P. Prevalence, severity, and types of pulmonary emphysema in a population of deaths in a Finnish city. Correlation with age, sex and smoking. *Scandinavian Journal of Respiratory Disease* 59(2): 101-115, April 1978.
- SYMONDS, G., RENZETTI, A.D., Jr., MITCHELL, M.M. The diffusing capacity in pulmonary emphysema. *American Review of Respiratory Disease* 109(3): 391-393, March 1974.

- TAGER, I.B., ROSNER, B., TISHLER, P.V., SPEIZER, F.E., KASS, E.H. Household aggregation of pulmonary function and chronic bronchitis. *American Review of Respiratory Disease* 114(3): 485-492, September 1976.
- TAGER, I.B., SPEIZER, F.E. Risk estimates for chronic bronchitis in smokers: A study of male-female differences. *American Review of Respiratory Disease* 113(5): 619-625, May 1976.
- TAGER, I., TISHLER, P.V., ROSNER, B., SPEIZER, F.E., LITT, M. Studies of the familial aggregation of chronic bronchitis and obstructive airways disease. *International Journal of Epidemiology* 7(1): 55-62, March 1978.
- TATTERSALL, S.F., BENSON, M.K., HUNTER, D., MANSELL, A., PRIDE, N.B., FLETCHER, C.M. The use of tests of peripheral lung function for predicting future disability from airflow obstruction in middle-aged smokers. *American Review of Respiratory Disease* 118(6): 1035-1050, December 1978.
- TATTERSALL, S.F., PEREIRA, R.P., HUNTER, D., BLUNDELL, G., PRIDE, N.B. Lung distensibility and airway function in intermediate alpha₁-antitrypsin deficiency (PiMZ). *Thorax* 34(5): 637-646, October 1979.
- THURLBECK, W.M. A clinico-pathological study of emphysema in an American hospital. *Thorax* 18(1): 59-67, March 1963.
- THURLBECK, W.M. The diagnosis of emphysema. *Thorax* 19(6): 571-574, November 1964.
- THURLBECK, W.M. Internal surface area and other measurements in emphysema. *Thorax* 22(6): 483-496, November 1967a.
- THURLBECK, W.M. The internal surface area of nonemphysematous lungs. *American Review of Respiratory Disease* 95(5): 765-773, May 1967b.
- THURLBECK, W.M. *Chronic Airflow Obstruction in Lung Disease*. Major Problems in Pathology, Volume 5. Philadelphia, W.B. Saunders Company, 1976, 456 pp.
- THURLBECK, W.M. Aspects of chronic airflow obstruction. *Chest* 72(3): 341-349, September 1977.
- THURLBECK, W.M. Post-mortem lung volumes. *Thorax* 34(6): 735-739, December 1979.
- THURLBECK, W.M. Overview of the pathology of pulmonary emphysema in the human. *Clinics in Chest Medicine* 4(3): 337-350, September 1983.
- THURLBECK, W.M., DUNNILL, M.S., HARTUNG, W., HEARD, B.E., HEPPELSTON, A.G., RYDER, R.C. A comparison of three methods of measuring emphysema. *Human Pathology* 1(2): 215-226, June 1970.
- THURLBECK, W.M., RYDER, R.C., STERNBY, N. A comparative study of the severity of emphysema in necropsy populations in three different countries. *American Review of Respiratory Disease* 109(2): 239-248, February 1974.
- TOBIN, M.J., COOK, P.J.L., HUTCHISON, D.C.S. Alpha₁-antitrypsin deficiency: The clinical features and physiological features of pulmonary emphysema in subjects homozygous for Pi type Z. A survey by the British Thoracic Association. *British Journal of Diseases of the Chest* 77: 14-27, January 1983.
- TOCKMAN, M.S. *Follow-up Study of Pulmonary Function in Nonsmokers*. Doctoral Dissertation, The Johns Hopkins University, Baltimore. Ann Arbor, University Microfilms International, Thesis No. 79-24643, 1979, 650 pp.
- TOCKMAN, M., MENKES, H., COHEN, B., PERMUTT, S., BENJAMIN, J., BALL, W.C., Jr., TONASCIA, J. A comparison of pulmonary function in male smokers and nonsmokers. *American Review of Respiratory Disease* 114(4): 711-720, October 1976.
- U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE. *Smoking and Health: A Report of the Surgeon General*. U.S. Department of Health, Education, and Welfare, Public Health Service, Office of the Assistant Secretary for Health, Office on Smoking and Health, DHEW Publication No. (PHS)79-50066, 1979, 1136 pp.

- U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES. *The Health Consequences of Smoking for Women: A Report of the Surgeon General*. U.S. Department of Health and Human Services, Public Health Service, Office of the Assistant Secretary for Health, Office on Smoking and Health, 1980a, 359 pp.
- U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES. *Report of Task Force on Epidemiology of Respiratory Disease. State of Knowledge, Problems, Needs, July 1979*. Public Health Service, National Institutes of Health, NIH Publication No. 81-2019, October 1980b.
- U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES. *The Health Consequences of Smoking. The Changing Cigarette: A Report of the Surgeon General*. U.S. Department of Health and Human Services, Public Health Service, Office of the Assistant Secretary for Health, Office on Smoking and Health, DHHS Publication No. (PHS)81-50156, 1981, 269 pp.
- U.S. PUBLIC HEALTH SERVICE. *Smoking and Health. Report of the Advisory Committee to the Surgeon General of the Public Health Service*. U.S. Department of Health, Education, and Welfare, Public Health Service, Center for Disease Control, PHS Publication No. 1103, 1964, 387 pp.
- U.S. PUBLIC HEALTH SERVICE. *The Health Consequences of Smoking. A Report of the Surgeon General, 1971*. U.S. Department of Health, Education, and Welfare, Public Health Service, Health Services and Mental Health Administration. DHEW Publication No. (HSM)71-7513, 1971, 458 pp.
- U.S. PUBLIC HEALTH SERVICE. *Prevalence of Selected Chronic Respiratory Conditions, United States, 1970*. Vital and Health Statistics data from the National Health Survey, Series 10, Number 84, September 1973.
- VAN DER LENDE, R., KOK, T.J., PESET REIG, R., QUANJER, Ph.H., SCHOUTEN, J.P., ORIE, N.G.M. Decreases in VC and FEV₁, with time: Indicators for effects of smoking and air pollution. *Bulletin European de Physiopathologie Respiratoire* 17(5): 775-792, September-October 1981.
- WALTER, S., NANCY, N.R., COLLIER, C.R. Changes in the forced expiratory spirogram in young male smokers. *American Review of Respiratory Disease* 119(5): 717-724, May 1979.
- WEBSTER, P.M., LORIMER, E.G., MAN, S.F.P., WOOLF, C.R., ZAMEL, N. Pulmonary function in identical twins: Comparison of nonsmokers and smokers. *American Review of Respiratory Disease* 119(2): 223-228, February 1979.
- WEISS, S.T., TAGER, I.B., SCHENKER, M., SPEIZER, F.E. The health effects of involuntary smoking. *American Review of Respiratory Disease* 128(5): 933-942, November 1983.
- WILHELMSEN, L., ORHA, I., TIBBLIN, G. Decrease in ventilatory capacity between ages 50 and 54 in a representative sample of Swedish men. *British Medical Journal* 3(5670): 553-556, September 6, 1969.
- WOOLCOCK, A.J., COLMAN, M.H., BLACKBURN, C.R.B. Chronic lung disease in Papua New Guinea and Australian populations. *Papua New Guinea Medical Journal* 16(1): 29-35, March 1973.
- WOOLCOCK, A.J., VINCENT, N.J., MACKLEM, P.T. Frequency dependence of compliance as a test for obstruction in the small airways. *Journal of Clinical Investigation* 48(6): 1097-1106, June 1969.
- WOOLF, C.R., SUERO, J.T. The respiratory effects of regular cigarette smoking in women. *American Review of Respiratory Disease* 103(1): 26-37, January 1971.
- WOOLF, C.R., ZAMEL, N. The respiratory effects of regular cigarette smoking in women. A five-year prospective study. *Chest* 78(5): 707-713, November 1980.
- ZAMEL, N., YOUSSEF, H.H., PRIME, F.J. Airway resistance and peak expiratory flow rate in smokers and nonsmokers. *Lancet* 1: 1237-1238, 1963.

**CHAPTER 3. MORTALITY FROM
CHRONIC
OBSTRUCTIVE LUNG
DISEASE DUE TO
CIGARETTE SMOKING**

CONTENTS

Introduction

COLD Mortality Patterns in the United States

Prospective Studies

- The British Doctors Study
 - The American Cancer Society 25-State Study
 - The U.S. Veterans Study
 - The Canadian Veterans Study
 - The American Cancer Society 9-State Study
 - California Men in Various Occupations
 - The Swedish Study
 - The Japanese Study of 29 Health Districts
-

Cigarette Smoking and Overall COLD Mortality

Retrospective Studies

Male and Female Differences in COLD Mortality

Amount Smoked and Mortality From COLD

Inhalational Practice and Mortality From COLD

Age of Initiation and COLD Mortality

Smoking Cessation and COLD Mortality

Pipe and Cigar Smoking Mortality From COLD

International Comparison of COLD Death Rates and
Smoking Habits: The Emigrant Studies

COLD Mortality Among Populations With Low Smoking
Rates

Summary and Conclusions

References

Introduction

The chronic obstructive lung diseases (COLD) that are causally related to cigarette smoking are chronic bronchitis, emphysema, and chronic obstructive pulmonary disease and allied conditions without mention of asthma, bronchitis, or emphysema. The last classification was introduced by the National Center for Health Statistics in response to the changes that occurred in the late 1960s in patterns of reporting causes of death on death certificates. During this period, physicians increasingly recorded deaths as due to "chronic obstructive lung disease" rather than the more specific categories of "emphysema" or "chronic bronchitis" (NCHS 1982). Because of this shift in patterns of reporting, and in recognition of the difficulty of clinically separating these categories from one another as a cause of death, the discussion in this chapter combines all of these categories for analysis, where possible, which should result in a more complete description of death rates from COLD.

COLD Mortality Patterns in the United States

The three chronic obstructive lung diseases related to smoking may account for almost 62,000 deaths in 1983, compared with 56,920 deaths in 1982, according to provisional mortality data recently published by the National Center for Health Statistics. This data is based on a 10 percent sample of all death certificates for the 12-month period ending in November (NCHS 1984). This is a dramatic increase from 1970, when slightly over 33,000 deaths were attributed to COLD.

Complete mortality data are available through 1980, and Table 1 presents the numbers of male and female deaths from COLD for 1970, 1975, and 1980. In addition to the relatively rapid rise in COLD deaths during these years, there was also a shift in the male to female ratio of these deaths. In 1970 male deaths outnumbered female deaths by a ratio of 4.3 to 1. By 1980 this ratio had declined to 2.36.

The age-adjusted death rates for COLD during the years 1960 through 1980 are presented in Figure 1 for white men, white women, and men and women of other races. As described in the previous chapter, however, COLD is a slowly progressive disease, and death from COLD usually occurs only after extensive damage has developed in the diseased lungs. Many individuals with COLD will die with their disease rather than because of it, and even those who do die of COLD are usually symptomatic for an extended period of time prior to death.

Therefore, death rate data may not accurately reflect the true prevalence or incidence of COLD in the U.S. population. In addition, COLD is often not recorded as a cause of death in hospital records

TABLE 1.—Number of and ratio of male to female chronic obstructive lung disease (COLD) deaths for three time periods, United States

Cause of death	1970		1975		1980	
	Men	Women	Men	Women	Men	Women
Chronic bronchitis	4,282	1,564	3,260	1,452	2,380	1,348
Emphysema	18,901	3,820	14,849	3,946	10,133	3,744
COLD and allied conditions	3,601	848	13,411	4,182	24,820	10,734
Total COLD deaths	26,784	6,227	31,520	9,580	37,333	15,826
M:F ratio	4.30		3.29		2.36	

SOURCE: National Center for Health Statistics (1982, and unpublished mortality data).




FIGURE 1.—Age-adjusted COLD mortality rates for whites and nonwhites in the United States, 1960–1980

SOURCE: National Center for Health Statistics (1982, and unpublished data).

(Moriyama et al. 1966) or on death certificates (Mitchell et al. 1968), even though it may have played an important role in a person's death. In a recent prospective study, nearly half of the excess mortality associated with significantly lowered FEV₁ was attributed to other causes (Peto et al. 1983). Relatively advanced lung disease (as judged by pathologic examination) may also exist without clinical

recognition because of the lung's large ventilatory reserve (Mitchell et al. 1968; Hepper et al. 1969). A joint committee of the American College of Chest Physicians and the American Thoracic Society (ACCP-ATS 1975) has developed standardized definitions of these conditions that may improve the accuracy of mortality reporting in the future.

As discussed in the chapter on morbidity in this Report, COLD in an individual is usually a combination of mucus hypersecretion, airway narrowing, and emphysema. The extent of damage represented by each of these three processes can vary substantially from individual to individual, both in the absolute magnitude of the damage and in the proportional contribution of each of these three components. The majority of those with smoking-induced lung damage do not have enough damage to result in clinically significant disease, and only some of those with clinically significant disease have damage to the lung that results in death from COLD. The progressive loss of FEV₁ in smokers described in the preceding chapter is one measure of the extent and progression of lung damage, and individuals with a markedly reduced FEV₁ are far more likely to die of COLD (Peto et al. 1983). These deaths commonly occur secondary to the failure of these severely damaged lungs to carry out the gas exchange required for survival.

Because death from COLD is the end result of lung damage accumulated over many years, these deaths would be expected to occur disproportionately in the older age groups; therefore, the presentation of a single age-adjusted death rate might not reflect a true picture of the changes in this disease with time. Figure 2 presents the age-specific death rates in 1977 for COLD in the different sexes and racial groups. Death rates increase rapidly over the age of 45, and this increase is particularly dramatic over the age of 65. In addition, the bulk of the difference between white men and men of other races, evident in Figure 2, occurs in those over age 65. Indeed, the COLD death rates for nonwhite men are actually higher than that for white men under age 55.

The examination of age-specific death rates over time also presents a somewhat different picture from that presented by the age-adjusted numbers in Figure 1. The age-adjusted rates for white men in Figure 1 seem to have changed only slightly between 1968 and 1980. However, when the age-specific rates for the years 1968 and 1977 are examined (Figure 3), this apparent stability can be seen to be a product of counterbalancing trends in those under and over 65 years of age. The death rates from COLD declined in white men under age 65 between 1968 and 1977, but COLD death rates increased in white men over age 65 during the same years; this increase was particularly dramatic in those over age 75.




FIGURE 2.—Age-specific COLD mortality rates for whites and nonwhites in the United States, 1977

SOURCE: National Center for Health Statistics (1982).

Figure 4 presents the age-specific COLD mortality rates for white women in 1960, 1968, and 1977. As with the male rates, the female COLD death rates rise rapidly with age, but they are substantially lower than the male rates. In contrast with the male rates, however, the white female death rates increased steadily with time from 1960 through 1977 both above and below age 65. In each of the age groups over the age of 45, where significant numbers of COLD deaths would be expected, there was a steady increase in rates from 1960 to 1968 and from 1968 to 1977. As is discussed later in this chapter, these differences between men and women over time are consistent with their differences in smoking behavior.

The effect of the normal aging process on the lung is small, rarely limits maximal exercise, and never results in ventilatory failure. Therefore, death from chronic obstructive lung disease is never a natural part of the aging process; it is the result of an infectious or other disease process or of the cumulative damage of environmental respiratory toxins. The most important of these toxins in the United States is cigarette smoke.




FIGURE 3.—Age-specific COLD mortality rates for white men in the United States, 1960, 1968, and 1977

SOURCE: National Center for Health Statistics (1982).

In spite of the large ventilatory reserve possessed by the lung, death from COLD is a major cause of U.S. mortality. This mortality is closely linked to cigarette smoking and has been examined extensively. Figure 5 shows the differences in COLD death rates for smokers and nonsmokers at different ages. From the rarity of COLD death in nonsmokers and the magnitude of the increased risk associated with smoking, it is clear that the overwhelming importance of cigarette smoking as a determinant of abnormal lung function demonstrated in the previous chapter is matched by the importance of cigarette smoking as a determinant of death from COLD. Examination of the death rates from COLD in smokers and nonsmokers suggests that from 85 to 90 percent of the COLD deaths in the United States can be attributed to cigarette smoking.

Prospective Studies

The relationship between smoking and death from COLD has been evaluated in a large number of prospective mortality studies. There are eight major prospective studies of the disease consequences of smoking. They involve large numbers of smokers and nonsmokers




FIGURE 4.—Age-specific COLD mortality rates for white women in the United States, 1960, 1968, and 1977

SOURCE: National Center for Health Statistics (1982).

and have examined the death rates from COLD in both groups. These studies cumulatively represent more than 17 million person-years of observation and over 330,000 deaths. The size of the populations studied allows a detailed examination of the relationship between smoking and death rates. The characteristics of the populations studied are summarized in Table 2 and are briefly reviewed here.

The British Doctors Study

The British doctors study (Doll and Hill 1954, 1956, 1964a, 1964b, 1966; Doll and Peto 1976, 1977; Doll and Pike 1972; Doll et al. 1980) of 40,000 male and female physicians in Britain was the first prospective study and is the longest running. Deaths from chronic bronchitis and emphysema were combined. Deaths from cor pulmonale (i.e., heart failure secondary to lung disease) were separately analyzed by smoking category and probably include some deaths from chronic bronchitis and emphysema.




FIGURE 5.—Death rate for bronchitis, emphysema, or both, per 100,000 population, by age and smoking status¹, U.S. veterans study, 16-year followup

¹ Smoker is defined as all people who smoke cigarettes and those who have ever smoked other tobacco products
SOURCE: Adapted from Rogot and Murray (1980).

The American Cancer Society 25-State Study

The American Cancer Society 25-State study (Hammond 1965, 1966; Hammond and Garfinkel 1969; Hammond et al. 1976; Lee and Garfinkel 1981) represents the largest investigation. Deaths from emphysema were separately analyzed by smoking habit; deaths from cor pulmonale were also separately recorded.

The U.S. Veterans Study

The mortality experience of approximately 294,000 U.S. veterans who held U.S. Government life insurance policies in December 1953 was examined in the U.S. veterans study (Dorn 1959; Kahn 1966; Rogot 1974a, b; Rogot and Murray 1980). Deaths from COLD were recorded as "bronchitis and/or emphysema"; "bronchitis, underlying or contributory"; and "emphysema without bronchitis."

The Canadian Veterans Study

Initiated in 1955 by the Canadian Department of National Health and Welfare, the Canadian veterans study (Best 1966; Best et al. 1961) included 78,000 men and 14,000 women. Over the next 6 years of followup, there were 9,491 male and 1,794 female deaths. The cause of death in most of these cases was confirmed by autopsy.

TABLE 2.—Outline of eight major prospective studies

Authors	Doll Hill Peto Pike	Hammond	Dorn Kahn Rogot	Hirayama	Best Jose Walker	Hammond Horn	Weir Dunn Linden Breslow	Cederlof Friberg Hrubec Lorich
Subjects	British doctors	Males and females in 25 States	U.S. veterans	Total population of 29 health districts in Japan	Canadian pensioners	White males in nine States	California males in various occupations	Probability sample of the Swedish population
Population size	40,000	1,000,000	290,000	265,000	92,000	187,000	68,000	56,000
Females	6,000	562,671	<1%	142,857	14,000			27,700
Age range	20-85+	35-84	35-84	40 and up	30-90	50-69	33-64	18-69
Year of enrollment	1961	1960	1964 1967	1966	1965	1962	1964	1963
Years of followup reported	20-22 years	12 years	16 years	13 years	6 years	4 years	5-8 years	10 years
Number of deaths	11,166	150,000	107,500	39,100	11,000	12,000	4,700	4,500
Person years of experience	800,000	8,000,000	3,500,000	3,000,000	500,000	670,000	480,000	550,000

The American Cancer Society 9-State Study

In the American Cancer Society 9-State study (Hammond and Horn 1958a, b), 187,783 white men were followed for an average of 44 months by 22,000 American Cancer Society volunteers. All deaths from pulmonary disease (except pulmonary neoplasms) were considered as one group and included deaths from pneumonia, asthma, tuberculosis, lung abscess, pneumoconiosis, bronchiectasis, and emphysema.

California Men in Various Occupations

The study of California men in various occupations (Dunn et al 1960; Weir and Dunn 1970) examined the mortality experience of 68,153 men, aged 35 to 64, drawn from labor union rolls in specified occupations. Deaths from emphysema were separately categorized.

The Swedish Study

The study of a probability sample of 55,000 Swedish men and women (Cederlof et al. 1975), aged 18 to 69, represents a detailed analysis of mortality by smoking status over a period of 10 years. The cause of death was ascertained by death certificates collected by the Central Bureau of Statistics for all of Sweden.

The Japanese Study of 29 Health Districts

In the fall of 1965, a total of 265,118 men and women in 29 health districts in Japan were enrolled in a prospective study (Hirayama 1967, 1970, 1972, 1975a, 1975b, 1977, 1981). Mortality data regarding deaths from asthma and emphysema have recently been reported.

Cigarette Smoking and Overall COLD Mortality

The data from the major prospective studies relating smoking to mortality from COLD in men and women are presented in Table 3. These data demonstrate a uniform increase in death rates from COLD among male and female smokers when compared with nonsmokers of either sex. The mortality ratios for smokers compared with nonsmokers vary markedly, however, from 2.2 in the Japanese study to 24.7 in the study of British doctors. Some of this variability can be attributed to different patterns of certification of cause of death in different countries, but a number of other factors are also important. Perhaps the most important other factor is the age range of the population studied. As described earlier in this chapter, death rates from COLD rise steeply with age, particularly over the age of 65. Studies of populations under age 65 may significantly underestimate the impact of cigarette smoking on COLD because of the long duration of smoking required to damage enough lung to result in

death from COLD. The population under 65 contains large numbers of individuals who have significant airflow obstruction and who will die of COLD, but who have not done so prior to age 65. This effect is demonstrated in the American Cancer Society 25-State study, in which the COLD mortality ratio for male smokers aged 45 to 64 was 6.55, but increased to 11.41 in male smokers aged 65 to 79.

A second reason for differences in mortality ratios is the selection of study populations who are currently employed, particularly if the duration of followup is relatively short. The incremental nature of the lung injury in COLD often results in a prolonged period of disability prior to resulting in death. This disability is usually incompatible with full-time work, particularly in those occupations requiring substantial exertion. Therefore, the study of a working population excludes those with significant existing disability from COLD and underestimates the COLD death rates in the general population. Unless the followup period is long enough to observe the progression of COLD from its asymptomatic stages through the development of disability and finally death, the impact of cigarette smoking on COLD death rates will be underestimated. This effect is particularly important because cigarette smoking is overwhelmingly the major determinant of COLD risk, and therefore an underestimation of the true COLD prevalence leads to an underestimation of the relative risk of smoking. As the followup period is extended for a duration sufficient to allow the full time course of COLD to be observed, the impact of cigarette smoking on COLD death rates also emerges from the small background rate of COLD death certification in nonsmokers (which includes those classified in error and those with disease induced by agents that results in a more rapid progression to death).

This "healthy worker" effect is present to varying extents in all of the prospective studies and is one of the reasons the studies with the longest followup periods also tend to have the largest COLD mortality ratios. This is particularly evident in the study with the longest followup. The British doctors study, with a followup of 20 years, revealed a mortality ratio for male smokers of 24.7.

A final reason for the differences in mortality ratios is the differences in the smoking habits of the various populations. As was discussed in the previous chapter, the extent of lung injury is influenced by both the number of cigarettes smoked per day and the duration of the smoking habit. As is shown in Table 4, some of the variability in mortality ratios among the studies disappears when the mortality ratios are reported by number of cigarettes smoked per day. However, there are also substantial differences in the pattern of cigarette use in different countries, particularly in the use of the milder types of tobacco cigarettes that are more likely to be inhaled and are smoked in the United States. For example, these cigarettes

TABLE 3.—COLD mortality ratios by disease category, eight prospective studies

Study	Size of population	Nonsmoker	Emphysema	Bronchitis	Both	Other	Comments
British physicians							
Men	34,000	1.00					
Women	6,195	1.00			24.7		Ratio for women by amount smoked only; see Table 4
California men in various occupations	68,000	1.00	12.33				
Canadian veterans							
Men	78,000	1.00	5.85	11.42			
American Cancer Society 25-State							
Men	440,500	1.00	45-64 ¹ 6.55	65-79 ¹ 11.41			¹ Age range
Women	562,700	1.00	4.89	7.50			
U.S. veterans							
Men	290,000	1.00	14.82	5.11	12.07		
American Cancer Society 9-State							
Men	188,000	1.00				2.85	All pulmonary diseases other than cancer (pneumonia, influenza, TB, asthma, bronchitis, lung abscess, etc.)

TABLE 3.—Continued

Study	Size of population	Nonsmoker	Emphysema	Bronchitis	Both	Other	Comments
Swedish							
Men	27,000	1.00				•	* Number of deaths too small for statistical analysis; includes deaths due to asthma
Women	28,000	1.00				2.20	
Japanese							
Men	122,000	1.00					Data by amount smoked only; see Table 4
Women	143,000	1.00					