

Pediatricians have considered tobacco smoke exposure in the troubled allergic child an identifiable problem to be faced. McGovern and coworkers (70) emphasized that allergic disease represents a major school health problem because children with hay fever, allergic rhinitis, and asthma account for about one-third of all chronic conditions reported under age 17. A survey is cited in which it was noted that asthma accounted for 11.4 percent of all chronic conditions in children and for 22.9 percent of days lost from school (8). These clinical investigators have, therefore, emphasized the need and value of removing the allergic child from all environmental sources of tobacco smoke exposure as a valid preventive measure.

Since the chances for progression of disease are more likely to occur in the face of continued and uncontrolled presence of causative factors, the potential for chronicity among adults is evident. The magnitude of the problem can be appreciated by noting the large population surveys in the United States which estimate that as many as 15 to 17 percent of the population suffers from asthma or hay fever (97). Thus, to whatever extent tobacco and/or tobacco smoke play a causal or contributory role in allergy, if they are ultimately shown to be allergens, it would be important for allergic patients of all age groups to take appropriate precautions to avoid exposure.

Effects of Cigarette Smoking on the Immune System

That cigarette smoking can affect the immune system has been well documented in both animals and humans. For purposes of discussion, these alterations in immune function can be classified as local and systemic. The local host defense system is comprised of the mucociliary mechanisms and functionally specialized cells, such as the macrophages and lymphocytes. Systemic defense mechanisms divide conveniently along the lines of cellular and humoral immunity.

Microscopic examinations of the respiratory tract mucosa demonstrate that chronic smoking leads to denuding of the ciliated epithelium, an increased number of goblet cells, and squamous metaplasia (89). On the other hand, studies attempting to quantify toxicity of cigarette smoke to cilia have been difficult to evaluate because of variation of mucus transport rates both among and within species studied, differences in techniques used to measure ciliary activity, and variations in methods and periods of exposure employed.

Studies on the short-term effects of smoke on ciliary function *in vitro* and *in vivo* generally show decreased function. Ciliostasis has been produced by *in vitro* exposure of the epithelium of the human respiratory tract to smoke residue passed through an aqueous medium (7) and, along with decreased rates of mucus transport, has also been observed in many animal models (1, 26, 50, 55). However, the effects of short-term smoking on mucociliary function in man have been

contradictory. In studies by Yeates, et al. (128) which measured mucociliary tracheal transport rates, some smokers showed slower bronchial clearance rates, while others showed little or no change over nonsmokers. Camner and coworkers (17), on the other hand, found mucociliary transport to be significantly increased during periods of intensive smoking (to the point of discomfort) compared to non-smoking periods.

Studies of long-term exposure have also been undertaken and, again, both animal and human studies are contradictory. Two studies were carried out in dogs exposed to forced smoke inhalation. One showed no change in tracheobronchial clearance (6) while the second, by different methodology, showed that tracheal mucus velocity was 30 percent of that found in controls (118).

In a study of 10 pairs of identical twins, discordant with regard to smoking (16), five of the smoking twins had decreased clearance rates while the other five demonstrated no differences over controls. Similarly, Albert, et al. (2) found bronchial clearance impaired in 8 out of 14 cigarette smokers tested. Lourenco and coworkers (65) found delayed clearance of particles, particularly in the central airways, at 1 hour after inhalation in nine smokers when compared to controls. On the other hand, Pavia, et al. (82) found no decrease in the efficiency of removal of particulate matter in the lungs of smokers compared to nonsmokers. However, the evidence indicates an adverse effect of long-term smoking on the mucociliary transport mechanisms and mucus composition (58).

It is necessary to understand the functions of alveolar macrophages and lung phagocytic cells as well as the population of immunocompetent lymphocytes in pulmonary tissue in order to appreciate how these elements and their modification can affect the processing of tobacco antigen and the resultant production of antibody and cell-mediated immunity. Since hypersensitivity phenomena are products of the immune system, these cellular elements can serve as determinants of allergic inflammation as well as of immunity.

Alveolar macrophages are important to lung function because of their role as phagocytes, engulfing and digesting particulate matter in the lung. Also, these cells process antigens and interact with lymphocytes in immune and allergic processes.

Many studies have examined the effect of smoking on macrophage function and metabolism. Even though most of these are *in vitro* studies, comparison is difficult because of differences inherent in the human and animal models used. In addition, in some cases, human subjects or animals were exposed to the smoke before the cells were harvested, while in others, cells were exposed directly to the smoke. Other variables included serious differences in amounts and lengths of exposures, filtration of smoke, and different methods of harvesting

cells. Nevertheless, it is clear from these studies that profound alterations in macrophages result from smoke exposure.

One consistent finding concerning the effect of smoking on macrophages is that the total number is increased in smokers. Keast and Holt (57) used a special apparatus simulating human smoking in exposed mice. They found initial and sustained elevations in macrophage populations. Other workers (56) also found increased macrophage numbers after only 2 weeks of cigarette smoking in humans. Studies by Pratt, et al. (88) and Harris, et al. (44) showed that smokers had strikingly increased numbers of macrophages when compared to nonsmokers and, furthermore, that macrophages accounted for 90 to 95 percent of lavaged lung cells found in smokers. The authors (44) speculate that increased alveolar macrophages in smokers might play an important role in pulmonary defense against toxic components of cigarette smoke. Also important is the possibility that macrophage accumulations could contribute to the pathogenesis of chronic pulmonary disease by the release of lysosomal enzyme content.

Changes in ultrastructure of macrophages have also been reported in smokers. Pratt and associates (88) observed that macrophages obtained in lung fluids of smokers were filled with cytoplasmic inclusions, and Martin (67) identified multinucleated giant cells in some smokers but none in nonsmokers. Martin (67) also noted that crystalloid refractile cytoplasmic inclusions were more common among the smokers. Harris, et al. (44) found the most salient feature of the macrophages from smokers to be larger and more numerous lysosomal bodies.

The study by Holt and Keast (47) demonstrated that the immediate toxic effects of tobacco smoke *in vitro* were greater in macrophages than fibroblasts, with surviving macrophages showing an increase in measured protein synthesis. Keast and Holt (57) also found that the macrophages from mice exposed to smoke for many weeks were no longer as susceptible to the untoward effects of smoke and had apparently adapted to the toxic conditions in a fashion similar to that seen in the tissue culture experiments.

Enzyme systems have also been shown to be affected by smoking. Martin (67) demonstrated that increased macrophage acid hydrolase directly correlated with daily cigarette consumption. Meyer, et al. (72) examined the effect of various concentrations of nicotine on the ATPase activity of sheep pulmonary alveolar macrophages and showed significant inhibition of this activity. Additionally, lower concentrations of this alkaloid stimulated cell respiration while higher concentrations were inhibitory. Kasemir and Kerp (56) recorded decreased oxygen uptake in sheep macrophages in contact with tobacco extracts. The *in vitro* studies of Harris and coworkers (44) on human alveolar macrophages demonstrated increased glucose utilization in smokers.

In pertinent studies, macrophage function has been measured by several methods. Green and Carolin (34), using an *in vitro* system to

measure phagocytosis, showed that added cigarette smoke had a depressant effect on the phagocytic activity of alveolar macrophages for *Staphylococcus albus*. The studies by Maxwell, et al. (69) on lung macrophages from guinea pigs exposed to tobacco prior to cell harvest showed that, although these alveolar cells phagocytosed bacteria at normal rates, their capacity for bacterial inactivation was impaired.

Laurenzi, et al. (61) demonstrated a 50 percent reduction in clearance of staphylococci from the lungs of smoke-exposed mice. In two human studies (22, 44) which measured phagocytic properties of alveolar macrophages, no significant differences were found between smokers and nonsmokers. Other studies of *in vitro* function of macrophages after *in vivo* exposure to smoke (employing rat alveolar macrophages) revealed no impairment of bactericidal inactivation of *S. albus* (49).

In the studies of Warr and Martin (119, 120), macrophages of smokers demonstrated an impaired response to an immune effector, MIF, paralleling those situations characterized by the absence of cell-mediated delayed hypersensitivity as well as acquired resistance to aggregate under *in vitro* conditions.

Though more work is needed to define the total qualitative and quantitative influences of tobacco smoking on alveolar macrophages, there is sufficient evidence in these studies to indicate measurable degrees of physiological impairment. Since interference with phagocytosis, endocytosis, and antigen processing can be anticipated as a consequence, there is the potential diminution of specific immune functions by these cells. In turn, the impairment of local immune processes as the first line of host defense exerts its toll on the dependent development of systemic immunity and influences emerging allergic inflammation.

The B and T lymphocytes are involved respectively in the humoral and cell-mediated arms of the immune system that functions both locally and systemically. It is therefore pertinent to examine the effect of smoking on these elements that provide the immunologic basis of hypersensitivity.

Of the immunoglobulins, secretory IgA is known to be predominant in bronchial mucus (29) (although the IgG/IgA ratio is increased in smokers (90)) and presumably plays a role in first-line defense against microbial invasion. Soutar's (101) studies on the distribution of plasma and other immunoglobulin-containing cells in the respiratory tract indicated more IgA-containing cells than those of other immunoglobulin classes. However, the only differential finding between smokers and nonsmokers was localized to the lobar bronchi of smokers where significant increases in IgA-containing cells were identified. Smoking was found to have significant suppressive action on salivary secretory IgA levels in normals, but not in patients with chronic diseases whose IgA levels were already elevated above normal (63). While these

studies show alterations in the expressions of local humoral immunity, the clinical significance of these changes is unknown.

Investigations have also been done to determine the effect of smoking on systemic humoral immunity. An assay which reflects antibody production is the plaque-forming cell (PFC) response. Thomas, et al. (108) examined PFC responses in samples of immunocompetent lung cells from mice exposed to fresh cigarette smoke and found progressive impairment of these responses over the exposure period of up to 10 months. In the studies of Holt, Keast, Nulsen, and Thomas (76,106,108,109,110) concerning the long-term effects of smoking on mice, PFC responses to intratracheally or intraperitoneally introduced antigens were shown to be initially enhanced and then depressed by chronic smoking (108,109). The direct measurement of serum hemolytic and hemagglutinating antibodies also showed depression, but the humoral response to a T cell-independent immunogen was unaffected (109). The secondary PFC response reflecting another aspect of humoral immunity was unaffected by smoking (109). PFC response depression was found to be reversible in a group when smoking was discontinued for 16 weeks (110). Other measurements of humoral immunity in mouse models exposed to tobacco also demonstrated impairment of the production of hemagglutinating antibodies, including those raised in response to the influenza virus (66), although some degree of suppression was reversible (28). Tar content of cigarettes may also play an important role (46).

Roszman, et al. (93, 94, 95), investigating several aspects of smoking and immunity in rabbits, found suppression of mitogen-induced blastogenesis and suppression of the immunoglobulin M and G antibody responses which correlated directly with the concentration either of nicotine or of the water-soluble fraction from cigarette smoke that was added to cultures.

Several surveys have attempted to address the issue of whether smoking influences serum immunoglobulin levels. Vos-Brat and Ruenke (116) found significant depression of IgG in smokers, Kosmider, et al. (59) also found a decreased IgG but increased IgM and IgA, while Wingerd and Sponzilli (127) found a decrease in the entire gamma globulin fraction. A decrease in lymphocytotoxic antibodies among smokers has also been demonstrated in pregnant women (77). On the other hand, no reported differences in mean concentrations of immunoglobulins were found when smokers were compared to nonsmokers by geographic location (71).

While these reports suggest that humoral antibody responses are influenced by cigarette smoke in a variety of ways, critical to this issue is a consideration of possible biologic impact in humans. Whether susceptibility to infection may be the end result of smoking effects on constituent elements of the immune system should be addressed. Thus, especially pertinent are the influenza vaccination studies of Waldman,

et al. (117), indicating that smoking more than one-half pack of cigarettes per day increased the risk of influenza-like illness, although the duration of the illness was unaltered. Finklea and associates (32) showed that the incidence of clinical influenza was 21 percent higher among smokers than nonsmokers. Serological data from this study suggested that smokers also had more frequent subclinical influenza. In pursuing this observation, Finklea, et al. (31) showed that, while serologic response to vaccination did not significantly differ between smokers and nonsmokers, the persistence of antibody titers after either natural infection or vaccination with A₂ antigens was significantly decreased among smokers. Nymand (77), examining histories of pregnant women, found that urinary tract infections and viral illness were observed more often in smokers than nonsmokers.

That elements indicative of immune function appear in the lung is evidenced by the identification of both T and B cells in fluid samples recovered from this site (121). Of interest is the finding of both an increased number of T and B cells and an increase in the T/B ratio in smokers.

Several aspects of cell-mediated immunity have been studied in animal models, including the ability of immunocompetent lymphocytes to proliferate after mitogenic stimulation by phytohemagglutinin (PHA), pokeweed (PW), and Concanavalin A (Con-A). In mice, initial increases of PHA responses in blood and regional lymph node lymphocytes were found after brief exposure to cigarette smoke, but decreases were found after prolonged exposure (107). Another study (18) demonstrated inhibition of proliferation of mouse lymphocytes to both PHA and pokeweed mitogen by an aqueous fraction of tobacco. In the rabbit (94), both nicotine and water-soluble fractions from whole cigarette smoke diminished peripheral lymphocyte blastogenic response to lectin stimulation.

Because of variation in methodology, data from human studies are difficult to compare. While increased numbers of T cells in peripheral blood lymphocytes and enhanced PHA response were noted among younger smokers, responses of older smokers or of those with a history of heavier cigarette consumption did not differ from normals (100). In examining peripheral bloods, Suci-Foca, et al. (103) found no differences in percent of T lymphocytes, PHA responses, or behavior in mixed lymphocyte cultures between smokers and nonsmokers. In another study (125), samples of blood taken from humans after smoking showed no differences in PHA responses even when physiologic levels of nicotine were added directly to the cultures. In contrast, Neher (74) found decreased DNA synthesis in response to PHA in the presence of nicotine. Desplaces, et al. (27) showed that smoke inhibited lymphocyte transformation by PHA yet stimulated lymphocytes in the absence of PHA. The clinical significance of this single aspect of T-cell function has yet to be determined.

Effects on other cellular elements of the immune system have also been described. Vos-Brat and Ruemke (116) and Silverman, et al. (100) demonstrated increased granular leukocytic levels in smokers. Others (54,79,98,129) have shown that smokers have hypereosinophilia. In two studies (79,98) the hypereosinophilia was reversible with abstinence from smoking. Similar lymphocytic and eosinophilic increases among smokers have been noted in patients' post-myocardial infarctions (129).

Serum abnormalities also have been described in smokers, including increased C-reactive (45) protein and an abnormal seroflocculant in smokers. Effects of smoking on manifestations of immune hyperresponsiveness add further evidence to the purported suppressive action of tobacco. Of interest are the reports of diminution of amyloid formation in the hamster model (123) and the inexplicable increase in survival of cardiac transplants in patients who resumed smoking postoperatively (35).

Target Organs of the Allergic Response

Despite the limitations, as previously noted, in appropriate materials and methods to define any possible effects of tobacco and smoking on allergic people, studies dealing with their roles in affecting various organs are noteworthy. A variety of clinical conditions have been ascribed to allergic manifestations to tobacco leaf or smoke, including asthma, rhinitis, hives, dermatitis, migraine headaches, cardiac and other vascular disturbances, as well as gastrointestinal disorders. The respiratory system has been the most widely studied.

Allergic rhinitis, typified by hay fever due to seasonal pollens and molds, is caused by exposure to a wide range of ubiquitous allergens. Apart from investigations of tobacco workers, there are no available studies to date to suggest that tobacco smoke or tobacco allergens are in fact a cause of allergic rhinitis in the general population. Many studies, however, have been reported showing that rhinitis patients suffer exacerbation of symptoms upon exposure to smoke. Speer (102) reported that 67 percent of allergic persons noted aggravation of nasal symptoms upon exposure to smoke, compared to 29 percent of nonallergic persons similarly exposed. Broder, et al. (11) found that most symptoms of allergic rhinitis could be attributed to other definable allergens with smoking or smoke exposure playing only a minor role. Allergic rhinitis believed to be related specifically to hypersensitivity to tobacco leaf products was reported to occur in 14.6 percent of 355 tobacco plantation workers and 8.7 percent of 722 tobacco factory workers (114).

Another study (86) among tobacco workers demonstrated that allergic rhinitis thought to be related to tobacco leaf occurred in approximately 4 percent of cases. However, possible contamination of

tobacco by molds or other allergens or irritants was not excluded in these studies.

It is relevant to note that symptoms of nasal congestion and excess mucous gland secretion, which may mimic those of allergic rhinitis or hay fever, can be caused by the nonspecific irritant or pharmacologic effects of vapor from the constituents of tobacco smoke. Thus, although it is not known whether allergy to tobacco or tobacco smoke plays a primary etiologic role in the usual case of allergic rhinitis, tobacco smoke per se is known to aggravate this condition via an irritant effect.

It is well known (102) that eye irritation manifested by itching, burning, swelling, and lacrimation occurs commonly among both allergic nonsmokers and nonallergic nonsmokers. To date, no studies are available suggesting that this manifestation is due to anything other than the nonspecific irritating effect of cigarette smoke.

Many studies have attempted to assess the relation between tobacco or smoking and asthma. Early investigators, using a variety of skin test materials (64, 91), inferred that allergy to tobacco could be causally related to asthma. Subsequent reports have examined the possible role of passive smoking in asthma. Speer (102) found that wheezing occurred more frequently in allergic people than in nonallergic people upon exposure to smoke. O'Connell and Logan (78), in studying the effects of parental smoking, found that smoke aggravated attacks of asthma in 26 percent of asthmatic children of nonsmoking parents, in contrast to 67 percent of asthmatic children of smoking parents. Importantly, they assessed the effects upon asthmatic children whose parents stopped smoking and reported improvement in 18 of 20 children. In contrast, only 4 of 15 asthmatic children improved when parents continued to smoke. Cameron and coworkers (15) concluded that asthmatic children of smoking parents were more often ill with respiratory disease but that this was related to nonspecific irritation rather than hypersensitivity. On the other hand, Rosen and Levy (92) published a case report of an infant who developed bronchial asthma associated with exposure to smoke. In this study, reaginic antibody to tobacco extract was documented by passive cutaneous transfer. More conclusive studies that tobacco may be causally related to asthma are reported among tobacco workers. Among 286 persons exposed to raw or fermented tobacco, the incidence of allergic manifestations was 8 percent, of which 17 percent had asthma (86). The possible role of tobacco additives has also been considered. Burge, et al. (13) reported the occurrence of occupationally-related asthma in a group of 21 industrial workers where colophony or pine resin, a substance also present in cigarettes as adhesives and filter fillings, was implicated.

The consequences of cigarette smoking in the asthmatic patient have also been examined. Townley and coworkers (112) reported similar

bronchial airway responses to lung function tests by methacholine inhalation in both smoking and nonsmoking asthmatics. Pimm and associates also reported that passive exposure of asthmatics to cigarette smoke resulted in no consistent significant effect on lung volumes and expiratory flow rates when compared with parallel room air exposure (84). On the other hand, Burrows, et al. (14), in a study of smoking and tests of lung function, found that an allergic predisposition, asthma or allergic rhinitis, as defined by positive skin reactivity, were associated with an increased susceptibility to bronchoconstrictor effects of cigarette smoking and to recurrent chest infections. That smoking can adversely effect an asthmatic patient in an indirect manner is illustrated by the finding of Powell, et al. (87) demonstrating interference with normal metabolism of the bronchodilator agent, theophylline, in smokers.

The concept that hyperreactive airways in asthmatics are due to a regulatory dysfunction of the autonomic nervous system is pertinent to this discussion (30). In addition to the effects of specific allergens inducing responsible mediators of bronchoconstriction, it is appreciated that nonspecific irritants (for example odors, temperature extremes, exercise, chemicals) can also act upon the affected cell receptors to precipitate asthmatic attacks.

Thus, apart from any putative allergenic effects of tobacco in a specifically sensitized patient, inhaled tobacco smoke carries the irritant potential to trigger or to aggravate asthmatic symptoms in the patient so affected. Hence, there is further support offered for both cessation of smoking and the following of avoidance procedures of passive exposure in the asthmatic individual.

Allergic effects of tobacco on the cardiovascular system have also received considerable attention. It is well documented that cardiac abnormalities occur in association with allergic phenomena, for example, anaphylaxis or allergic shock (5, 25, 73). However, whether tobacco may play a role in cardiovascular alterations apart from known pharmacologic effects is still not clear. Harkavy's series of observations (36, 37, 38, 39, 40, 42, 43) would support the concept that allergy to tobacco leaf may have important implications in a variety of cardiac and vascular diseases. In these he would include cardiac arrhythmias, intensification of coronary artery insufficiency, thromboangiitis obliterans, migrating phlebitis, and some forms of allergic vasculitis. Although acknowledging the pharmacologic effects of nicotine on the cardiovascular system, Harkavy also suggests that it may act as a hapten in inducing allergic responses. Recent observations by Becker and coworkers (10), using a partially characterized antigenic component of tobacco, led them to hypothesize that circulating tobacco antigens in sensitive individuals might react with corresponding antibody to produce focal injury of blood vessels. If this hypothesis is corroborated, design of further studies of potential adverse conse-

quences of possible tobacco allergy on the cardiovascular system will be possible.

That tobacco may operate through the mechanism of cell-mediated immunity or delayed hypersensitivity is suggested by case reports of contact dermatitis caused by tobacco smoke and tobacco smoke residue (19, 24, 122). Recent surveys among tobacco workers have shown that contact dermatitis related to tobacco was responsible for 14 percent of skin eruptions occurring in this industrial sample (3). By contrast, however, an earlier survey (96) could not implicate tobacco as a cause of dermatitis among cigar factory workers. It has been pointed out that dermatitis among tobacco workers probably represents a nonspecific response due to injury, moisture, or irritants, especially those from the chemicals or other fertilizers used in the growing process (122). To date, therefore, there is little evidence that allergic skin manifestations due to tobacco occur with any significant frequency.

Summary

1. Tobacco and tobacco smoke extracts have been found to act as antigens inducing both precipitating and reaginic antibodies in experimental animals. Tobacco leaf products can also sensitize lymphocytes participating in cell-mediated immune functions.

2. Tobacco and its combustion products are known to be heterogeneous mixtures of particulate and gaseous materials. Additionally, natural contaminants and intentional additives increase the array of components, presenting a complex of toxic, pharmacologic, irritant, and inflammatory effects that can complicate interpretation of a precisely defined role for tobacco in immune and allergic processes.

3. Several tobacco antigens have been isolated by chemical procedures. Of special interest is a glycoprotein common to both tobacco extracts and smoke antigenically corresponding with reaginic antibody in humans.

4. Epidemiologic samplings to define the presence of true allergy to tobacco, either among healthy persons or among those suffering from known allergic conditions, are inconclusive.

5. Tobacco smoking exerts a variety of effects on respiratory tract structures involved in local host defense, and chronic smoking leads to consistent histological changes in the respiratory tract.

(a) There is evidence to indicate an adverse effect of long-term smoking on the mucociliary transport mechanisms and mucus composition.

(b) The number of macrophages isolated from lung fluids of smokers is increased over nonsmokers.

(c) Changes in the ultrastructure of macrophages—most notably the presence of cytoplasmic inclusions—are found in smokers.

- (d) Alveolar macrophages from smokers have altered metabolism and measurable degrees of physiologic impairment.
6. Alterations of indicators of humoral immunity have been demonstrated in the respiratory tracts of smokers, and smoking may impair systemic humoral immunity both *in vitro* and *in vivo*.
7. Alterations in assays of cell-mediated immunity are noted locally and systemically in smokers.
8. Leukocytosis and reversible hypereosinophilia have been seen in smokers.
9. The ability to make a definitive diagnosis of tobacco allergy is complicated by the difficulty of demonstrating a cause and effect relationship between immunologic events and disease manifestations; additional evidence is required to establish whether there is a definitive role for tobacco smoke sensitization in causing allergic diseases.
10. Studies concerned with the adverse consequences of either active or passive smoking have shown that allergic individuals, especially those with rhinitis or asthma, may, in fact, be more sensitive to the nonspecific noxious effects of cigarette smoke than healthy individuals.

Conclusion and Comment

Apart from symptom-relieving drugs, there are no known effective therapeutic measures to prevent or combat the adverse effects of smoking on immune function and on allergy-related problems. It is evident that further studies defining tobacco antigens, determining the clinical incidence of tobacco allergy, further clarifying the nature of immune responses to tobacco, and improving the diagnostic agents and materials should be undertaken. Such studies, however, can not be expected to have an impact on improving the health of individuals subject to tobacco's adverse effects comparable to that which would result from adhering to the mainstay of management of the allergic patient—complete avoidance of the incriminated substance.

Allergy and Immunity: References

- (1) ALBERT, R.E., BERGER, J., SANBORN, K., LIPPMANN, M. Effects of cigarette smoke components on bronchial clearance in the donkey. *Archives of Environmental Health* 29: 96-101, August 1974.
- (2) ALBERT, R.E., LIPPMANN, M. BRISCOE, W. The characteristics of bronchial clearance in humans and the effects of cigarette smoking. *Archives of Environmental Health* 18: 738-755, May 1969.
- (3) ANDREICHUK, I.E. K voprosu o tabochnom dermatite (On tobacco dermatitis). *Vestnik Dermatologii i Venerologii* (4): 57-59, 1975.
- (4) ARMEN, R.N., COHEN, S.G. The effect of forced inhalation of tobacco smoke on the electrocardiogram of normal and tobacco-sensitized rabbits. *Diseases of the Chest* 35: 663-676, June 1959.
- (5) AUER, J., ROBINSON, G.C. An electrocardiographic study of the anaphylactic rabbit. *Journal of Experimental Medicine* 18: 450-460, 1931.
- (6) BAIR, W.J., DILLEY, J.V. Pulmonary clearance of $^{59}\text{Fe}_2\text{O}_3$ and $^{51}\text{Cr}_2\text{O}_3$ in rats and dogs exposed to cigarette smoke. In: Davies, C.N. (Editor). *Inhaled Particles and Vapours II*. New York, Pergamon Press, 1967, pp. 251-268.
- (7) BALLENGER, J.J. Experimental effect of cigarette smoke on human respiratory cilia. *New England Journal of Medicine* 263(17): 832-835, October 27, 1960.
- (8) BARKIN, G.D., MCGOVERN, J.P. What the classroom teacher can do for the asthmatic child. *National Education Association Journal* 56(8): 40-41, November 1967.
- (9) BECKER, C.G., DUBIN, T. Activation of factor XII by tobacco glycoprotein. *Journal of Experimental Medicine* 146: 457-467, 1977.
- (10) BECKER, C.G., DUBIN, T., WIEDEMANN, H.P. Hypersensitivity to tobacco antigen. *Proceedings of the National Academy of Sciences* 73(5): 1712-1716, May 1976.
- (11) BRODER I., HIGGINS M.W., MATHEWS K.P., KELLER J.B. Epidemiology of asthma and allergic rhinitis in a total community, Tecumseh, Michigan, III. Second Survey of the Community. *Journal of Allergy and Clinical Immunology* 53(3): 127-138, March 1974.
- (12) BROWN, A. Present day treatment of asthma. *New York Medical Journal and Medical Review* 98(6): 333-336, September 19, 1923.
- (13) BURGE, P.S., HARRIES, M.G., O'BRIEN, I.M., PEPYS, J. Respiratory disease in workers exposed to solder flux fumes containing colophony (pine resin). *Clinical Allergy* 8: 1-14, 1978.
- (14) BURROWS, B., LEBOWITZ, M.D., BARBEE, R.A. Respiratory disorders and allergy skin-test reactions. *Annals of Internal Medicine* 84(2): 134-139 1976.
- (15) CAMERON, P., KOSTIN, J.S., ZAKS, J.M., WOLFE, J.H., TIGHE, G., OSELETT, B., STOCKER, R., WINTON, J. The health of smokers' and nonsmokers' children. *Journal of Allergy* 43(6): 336-341, June 1969.
- (16) CAMNER, P., PHILIPSON, K. Tracheobronchial clearance in smoking-discordant twins. *Archives of Environmental Health* 25: 60-63, July 1972.
- (17) CAMNER, P., PHILIPSON, K., ARVIDSSON, T. Cigarette smoking in man. Short-term effect on mucociliary transport. *Archives of Environmental Health* 23: 421-426, December 1971.
- (18) CARLENS, E. Smoking and the immune response in the airpassages. *Broncho-Pneumologie* 26(4): 322-323, July/August 1976.
- (19) CHANIAL, G., JOSEPH, J., COLIN, L., DUCLAUX, C. Les dermites chez les travailleurs du tabac (a propos de 9 observations) (Cases of Dermatitis in tobacco workers [reference to nine observations]). *Bulletin de la Societe Francaise de Dermatologie et de Syphiligraphie* 77(2): 281-283, July 1970.
- (20) CHOBOT, R. The significance of tobacco reactions in allergic children. *Journal of Allergy* 6: 383-386, 1934.

- (21) CHU, Y.M., PARLETT, R.C., WRIGHT, G.L., JR. A preliminary investigation of some immunologic aspects of tobacco use. *American Review of Respiratory Disease* 102(1): 118-123, 1970.
- (22) COHEN, A.B., CLINE, M.J. The human alveolar macrophage: Isolation, cultivation in vitro, and studies of morphologic and functional characteristics. *Journal of Clinical Investigation* 50(7): 1390-1398, July 1971.
- (22a) COLLEY, J.R.T., HOLLAND, W.W., CORKHILL, R.T. Influence of passive smoking and parental phlegm on pneumonia and bronchitis in early childhood. *Lancet*:1031-1034, November 2, 1974.
- (23) COOMBS, R.R.A., GELL, P.G.H., Classification of allergic reactions responsible for clinical hypersensitivity and disease. In: Gell, P.G.H., Coombs, R.R.A. (Editors). *Clinical Aspects of Immunology*. Oxford, Blackwell Scientific Publications, 1963, pp. 575-596.
- (24) CORMIA, F.E., DE GARA, P.F. Vesiculobullous dermatitis from tobacco smoke. *Journal of the American Medical Association* 193(5): 141-142, August 2, 1965.
- (25) CRIEP, L.H. Electrocardiographic studies on the effect of anaphylaxis on the cardiac mechanism. *Archives of Internal Medicine* 48(6): 1098-1109, 1931.
- (26) DALHAMN, T. Studies on tracheal ciliary activity. *American Review of Respiratory Diseases* 89(6): 870-877, 1964.
- (27) DESPLACES, A., CHARREIRE, J., IZARD, C. Action de la phase gazeuse de fumee de cigarette sur la transformation lymphoblastique du petit lymphocyte humain (observations preliminaires) (Smoke inhibition of lymphocytes transformation by PHA but stimulation of lymphocytes in the absence of PHA [preliminary observations]). *Revue European Etudes Clinique et Biologique* 16: 822-826, 1971.
- (28) ESBER, H.J., MENNINGER, F.F., JR., BOGDEN, A.E., MASON, M.M. Immunological deficiency associated with cigarette smoke inhalation by mice. Primary and secondary hemagglutinin response. *Archives of Environmental Health* 27(2): 99-104, August 1973.
- (29) FALK, G.A., OKINAKA, A.J., SISKIND, G.W. Immunoglobulins in the bronchial washings of patients with chronic obstructive pulmonary disease. *American Review of Respiratory Disease* 105: 14-21, 1972.
- (30) FARR, R.S. Asthma in adults: The ambulatory patient. *Hospital Practice*: 113-123, April 1978.
- (31) FINKLEA, J.F., HASSELBLAD, V., RIGGAN, W.B., NELSON, W.C. HAMMER, D.I., NEWILL, V.A. Cigarette smoking and hemagglutination inhibition response to influenza after natural disease and immunization. *American Review of Respiratory Disease* 104(3): 368-376, September 1971.
- (32) FINKLEA, J.F., SANDIFER, S.H., SMITH, D.D. Cigarette smoking and epidemic influenza. *American Journal of Epidemiology* 90(1): 390-399, 1969.
- (33) FONTANA, V.J., REDISCH, W., NEMIR, R.L., SMITH, M.K., DECRINIS, K., SULZBERGER, M.B. Studies in tobacco hypersensitivity. III. Reactions to skin tests and peripheral vascular responses. *Journal of Allergy* 30(3): 241-249, May/June 1959.
- (34) GREEN, G.M., CAROLIN, D. The depressant effect of cigarette smoke on the in vitro antibacterial activity of alveolar macrophages. *New England Journal of Medicine* 276(8): 422-427, February 23, 1967.
- (35) GRIEPP, R.B., STINSON, E.B., BIEBER, C.P., REITZ, B.A., COPELAND, J.G., OYER, P.E., SHUMWAY, N.E. Control of graft arteriosclerosis in human heart transplant recipients. *Surgery* 81(3): 262-269, March 1977.
- (36) HARKAVY, J. Cardiac manifestations due to hypersensitivity. *Annals of Allergy* 28(6): 242-251, June 1970.
- (37) HARKAVY, J. Cardiovascular manifestations due to hypersensitivity. *New York State Journal of Medicine* 69(21): 2757-2765, November 1, 1969.

- (38) HARKAVY, J. Hypersensitiveness to tobacco and biopsy studies of skin reactions in vascular disease. *Journal of Allergy* 9: 475-488, 1938.
- (39) HARKAVY, J. Tobacco sensitiveness in angina pectoris and coronary artery disease. *Proceedings of the Society for Experimental Biology and Medicine* 30(104): 683-684, 1932.
- (40) HARKAVY, J. Tobacco sensitiveness in thromboangiitis obliterans, migrating phlebitis and coronary artery disease. *Bulletin of the New York Academy of Medicine* 9: 318-322, 1933. (Abstract)
- (41) HARKAVY, J. Tobacco sensitization in rats. *Proceedings of the Society for Experimental Biology and Medicine* 36(3): 381-383, April 1937.
- (42) HARKAVY, J. Tobacco skin reactions and their clinical significance. *Journal of Investigative Dermatology* 2: 257-279, 1939.
- (43) HARKAVY, J., WITEBSKY, E. Studies of specificity in multiple hypersensitiveness by quantitative titration and absorption of reagins. *Journal of Allergy* 6: 437-447, 1935.
- (43a) HARLAP, S., DAVIES, A.M. Infant admissions to hospital and maternal smoking. *Lancet* :529-532, March 30, 1974.
- (44) HARRIS, J.O., SWENSON, E.W., JOHNSON, J.E., III. Human alveolar macrophages: Comparison of phagocytic ability, glucose utilization, and ultrastructure in smokers and nonsmokers. *Journal of Clinical Investigation* 49: 2086-2096, 1970.
- (45) HEISKELL, C.L., MILLER, J.N., ALDRICH, H.J., CARPENTER, C.M. Smoking and serologic abnormalities. *Journal of the American Medical Association* 181(8): 674-677, August 25, 1962.
- (46) HOLT, P.G., CHALMER, J.E., ROBERTS, L.M., PAPADIMITRIOU, J.M., THOMAS, W.R., KEAST, D. Low-tar and high-tar cigarettes. Comparison of effects in two strains of mice. *Archives of Environmental Health* 31(5): 258-265, September/October 1976.
- (47) HOLT, P.G., KEAST, D. Acute effects of cigarette smoke on murine macrophages. *Archives of Environmental Health* 26(6): 300-304, June 1973.
- (48) HOSEN, H. Letter to the editor. Tobacco Sensitivity. *Annals of Allergy* 29(11): 608-610, November 1971.
- (49) HUBER, G.L., SHEA, J. Does tobacco smoke impair alveolar macrophage function. *Chest* 70(3): 429, September 1971. (Abstract)
- (50) IRAVANI, J. Effects of cigarette smoke on the ciliated respiratory epithelium of rats. *Respiration* 29: 480-487, 1972.
- (51) ISHIZAKA, K., ISHIZAKA, T. Identification of γ E-antibodies as a carrier of reaginic activity. *Journal of Immunology* 99(6): 1187-1198, 1967.
- (52) ISHIZAKA, T., ISHIZAKA, K., ORANGE, R.P., AUSTEN, K.F. The capacity of human immunoglobulin E to mediate the release of histamine and slow reacting substance of anaphylaxis (SRS-A) from monkey lung. *Journal of Immunology* 104(2): 335-343, 1970.
- (53) JUSTUS, D.E., ADAMS, D.A. Evaluation of tobacco hypersensitivity responses in the mouse. A potential animal model for critical study of tobacco allergy. *International Archives of Allergy and Applied Immunology* 51(6): 687-695, 1976.
- (54) KAMESWARAN, L., KANAKAMBAL, K., VIJAYASEKARAN, V. Studies on plasma histaminase levels in normal and allergic individuals. *Indian Journal of Physiology and Pharmacology* 12(4): 159-165, October 1968.
- (55) KAMINSKI, E.J., FANCHER, O.E., CALANDRA, J.C. In vivo studies of the ciliastatic effects of tobacco smoke. *Archives of Environmental Health* 16: 188-193, February 1968.

- (56) KASEMIR, H., KERP, L. Gesamteinwirkungen von luftverunreinigungen auf den menschen aus der sicht der immunologie (Total effects of air pollution on man from an immunological viewpoint). *Praxis Der Pneumologie* 23(special issue): 776-780, December 1974.
- (57) KEAST, D., HOLT, P. Smoking and the immune response. *New Scientist* 61(891): 806-807, March 28, 1974.
- (58) KOLLERSTROM, N., LORD, P.W., WHIMSTER, W.F. A difference in the composition of bronchial mucus between smokers and non-smokers. *Thorax* 32(2): 155-159, 1977.
- (59) KOSMIDER, S., FELUS, E., WYSOCKI, J. Ocena niektorych wyznacznikow odpornosci humoralnej palaczy papierosow (Evaluation of some humoral resistance determinants in smokers). *Polski Tygodnik Lekarski* 28(2): 47-50, January 8, 1973.
- (60) KREIS, B., PELTIER, A., FOURNAUD, S., DUPIN-GIROD, S. Reaction de precipitation entre certains serums humains et des extraits solubles de tabac (Precipitation reaction between some human serums and soluble tobacco extracts). *Annales de Medecine Interne* 121(4): 437-440, April 1970.
- (61) LAURENZI, G.A., GUARNERI, J.J., ENDRIGA, R.B., CAREY, J.P. Clearance of bacteria by the lower respiratory tract. *Science* 142: 1572-1573, 1963.
- (62) LEHRER, S., WILSON, M., SALVAGGIO, J.E. Immunogenic properties of tobacco smoke. *Journal of Allergy and Clinical Immunology* 62: 368-370, 1978.
- (63) LEWIS, D.M., LAPP, N. LeR., BURRELL, R. Quantitation of secretory immunoglobulin A in chronic pulmonary disease. *American Review of Respiratory Disease* 101(1): 55-61, January 1970.
- (64) LIMA, A.O., ROCHA, G. Cutaneous reactions to tobacco antigen in allergic and nonallergic children. *Annals of Allergy* 7(4): 528-531, July/August 1949.
- (65) LOURENCO, R.V., KLIMEK, M.F., BOROWSKI, C.J. Deposition and clearance of 2 μ particles in the tracheobronchial tree of normal subjects—smokers and nonsmokers. *Journal of Clinical Investigation* 50: 1411-1420, 1971.
- (66) MACKENZIE, J.S. The effect of cigarette smoke on influenza virus infection: A murine model system. *Life Sciences* 19(3): 409-412, August 1, 1976.
- (67) MARTIN, R.R. Altered morphology and increased acid hydrolase content of pulmonary macrophages from cigarette smokers. *American Review of Respiratory Disease* 107: 596-601, 1973.
- (68) MATSUSHITA, H., NOGUCHI, M., TAMAKI, E. Conjugate of bovine serum albumin with nicotine. *Biochemical and Biophysical Research Communications* 57(4): 1006-1010, 1974.
- (69) MAXWELL, K.W., MARCUS, S., RENZETTI, A.D., JR. Effect of tobacco smoke on the phagocytic and cytopeptic activity of guinea pig alveolar macrophages. *American Review of Respiratory Disease* 96(1): 156, 1967.
- (70) MCGOVERN, J.P., PEIRCE, K.E., LEE R.E., JR. The allergic child and his challenge to the school. *Clinical Pediatrics* 10(11): 636-644, November 1971.
- (71) MELOY LABORATORIES, INC. Air pollution exposure and immunoglobulin levels. Springfield, Virginia, Meloy Laboratories, Inc., Biological Products Division, March 1974, pp. 1-15.
- (72) MEYER, D.H., CROSS, C.E., IBRAHIM, A.B., MUSTAFA, M.G. Nicotine effects on alveolar macrophage respiration and adenosine triphosphatase activity. *Archives of Environmental Health* 22(3): 362-365, March 1971.
- (73) MIKULICICH, G. Electrocardiographic changes in experimental anaphylactic reactions. *Journal of Allergy* 22(3): 249-263, May 1951.
- (74) NEHER, G.H. Nicotine-induced depression of lymphocyte growth. *Toxicology and Applied Pharmacology* 27(2): 253-258, February 1974.

- (75) NORMAN, V. An overview of the vapor phase, semivolatile and nonvolatile components of cigarette smoke. In: *Recent Advances in Tobacco Science*. Volume 3. Tobacco Smoke: Its Formation and Composition. Thirty-first Tobacco Chemists Research Conference. Greensboro, North Carolina, October, 5-7, 1977, pp. 27-58.
- (75a) NORMAN-TAYLOR, W., DICKINSON, V.A. Dangers for children in smoking families. *Community Medicine*: 32-33, April 21, 1972.
- (76) NULSEN, A., HOLT, P.G., KEAST, D. Cigarette smoking, air pollution, and immunity: A model system. *Infection and Immunity* 10(6): 1226-1229, December 1974.
- (77) NYMAND, G. Maternal smoking and immunity. *Lancet* 2(7893): 1379-1380, December 7, 1974. (Letter)
- (78) O'CONNELL, E.J., LOGAN, G.B. Parental smoking in childhood asthma. *Annals of Allergy* 32: 142-145, March 1974.
- (79) PAINTAL, I.S., MININA, R.J. Tobacco smoking—a probable cause of eosinophilia. *Indian Practitioner* 28(5): 243-245, May 1975.
- (80) PANAYOTOPOULOS, S., GOTSIS, N., PAPAZOGLU, N., CONCOURIS, L. Antigenic study of nicotiana tabacum and research on precipitins against tobacco antigens in the serum of smokers and nonsmokers. *Allergologia et Immunopathologia* 2(1): 111-114, January/February 1974.
- (81) PATON, W.D.M. The mechanism of histamine release. In: Wolstenholme, G.E.W., O'Connor, C.M. (Editors). *Histamine*. Boston, Little, Brown and Co., 1956, pp. 59-73.
- (82) PAVIA, D., SHORT, M.D., THOMSON, M.L. No demonstrable long term effects of cigarette smoking on the mucociliary mechanisms of the human lung. *Nature* 226: 1228-1231, June 27, 1970.
- (83) PESHKIN, M.M., LANDAY, L.H. Cutaneous reactions to tobacco antigen in allergic and nonallergic children with the direct and indirect (local passive transfer) methods of testing. *Journal of Allergy* 10: 241-245, 1939.
- (84) PIMM, P., SHEPHARD, R.J., SILVERMAN, F. Physiological effects of acute passive exposure to cigarette smoke in asthmatics. *Proceedings of the Federation of American Societies for Experimental Biology*, 61st Annual Meeting, Chicago, Illinois, April 1-8, 1977, 36: 606, 1977. (Abstract)
- (85) PIPES, D.M. Allergy to tobacco smoke. *Annals of Allergy* 28(3): 277-282, July/August 1945.
- (86) POPESCU, I. GR., PAUN, R., MOLNER, C., OLARU, C., GHEORGHIU, T., IOTA, C.G. Contributions to the study of tobacco allergy. *Revue Roumaine de Medecine Interna* 1(5): 427-436, 1964.
- (87) POWELL, J.R., THIERCELIN, J.-F., VOZEH, S., SANSOM, L., RIEGELMAN, S. The influence of cigarette smoking and sex on theophylline disposition. *American Review of Respiratory Disease* 116: 17-23, 1977.
- (88) PRATT, S.A., FINLEY, T.N., SMITH, M.H., LADMAN, A.J. A comparison of alveolar macrophages and pulmonary surfactant (?) obtained from the lungs of human smokers and nonsmokers by endobronchial lavage. *Anatomical Record* 163: 497-508, 1969.
- (89) REGLAND, B., CAJANDER, S., WIMAN, L.-G., FALKMER, S. Scanning electron microscopy of the bronchial mucosa in some lung diseases using bronchoscopy specimens. *Scandinavian Journal of Respiratory Diseases* 57: 171-182, 1976.
- (90) REYNOLDS, H.Y., NEWBALL, H.H. Analysis of proteins and respiratory cells obtained from human lungs by bronchial lavage. *Journal of Laboratory and Clinical Medicine* 84(4): 559-573, October 1974.
- (91) ROSEN, F.L. Studies in tobacco allergy. *Journal of the Medical Society of New Jersey* 51(3): 109-114, March 1954.

- (92) ROSEN, F.L., LEVY, A. Bronchial asthma due to allergy to tobacco smoke in an infant. *Journal of the American Medical Association* 144: 620-621, 1950.
- (93) ROSZMAN, T.L. Effect of nicotine and water soluble condensate from whole smoke in the *in vitro* secondary antibody response. *Proceedings of the University of Kentucky Tobacco and Health Research Institute, Tobacco and Health Workshop Conference, Lexington, Kentucky, March 26-28, 1973*, pp. 530-541.
- (94) ROSZMAN, T.L., ELLIOTT, L.H., ROGERS, A.S. Suppression of lymphocyte function by products derived from cigarette smoke. *American Review of Respiratory Disease* 111(4): 453-457, 1975.
- (95) ROSZMAN, T.L., ROGERS, A.S. The immunosuppressive potential of products derived from cigarette smoke. *American Review of Respiratory Disease* 108(5): 1158-1163, November 1973.
- (96) SAMITZ, M.H., MORI, P., LONG, C.-F. Dermatological hazards in the cigar industry. *Industrial Medicine and Surgery* 18(10): 434-439, October 1949.
- (97) SCHACHTER, J., *Allergy Statistics: Allergic Conditions in the United States*. U.S. Department of Health, Education and Welfare, Public Health Service, National Institutes of Health, National Institute of Allergy and Infectious Diseases, DHEW Publication No. (NIH) 75-757, 1975, 2 pp.
- (98) SCHOEN, I., PIZER, M. Eosinophilia apparently related to cigarette smoking. *New England Journal of Medicine* 270(25): 1344-1347, June 18, 1964.
- (99) SELYE, H. *The Mast Cells*. Washington, Butterworths, 1965, pp. 132-216.
- (100) SILVERMAN, N.A., POTVIN, C., ALEXANDER, J.C., JR., CHRETIEN, P.B. *In vitro* lymphocyte reactivity and T-cell levels in chronic cigarette smokers. *Clinical and Experimental Immunology* 22(2): 285-292, November 1975.
- (101) SOUTAR, C.A. Distribution of plasma cells and other cells containing immunoglobulin in the respiratory tract of normal man and class of immunoglobulin contained therein. *Thorax* 31(2): 158-166, April 1976.
- (102) SPEER, F. Tobacco and the nonsmoker. A study of subjective symptoms. *Archives of Environmental Health* 16(3): 443-446, March 1968.
- (103) SUCIU-FOCA, N., MOLINARO, A., BUDA, J., REEMTSMA, K. Cellular immune responsiveness in cigarette smokers. *Lancet* 1: 1062, May 25, 1974. (Letter)
- (104) SULZBERGER, M.B. Studies in tobacco hypersensitivity. I. A comparison between reactions to nicotine and to denicotinized tobacco extract. *Journal of Immunology* 24(1): 85-91, 1933.
- (105) SULZBERGER, M.B., FEIT, E. Studies in tobacco hypersensitivity. II. Thromboangiitis obliterans with positive urticarial skin reactions and negative reagin findings. *Journal of Immunology* 24(5): 425-432, 1933.
- (106) THOMAS, W.R., HOLT, P.G., KEAST, D. Antibody production in mice chronically exposed to fresh cigarette smoke. *Experientia* 30(12): 1469-1470, December 15, 1974.
- (107) THOMAS, W.R., HOLT, P.G., KEAST, D. Cellular immunity in mice chronically exposed to fresh cigarette smoke. *Archives of Environmental Health* 27(6): 372-375, December 1973.
- (108) THOMAS, W.R., HOLT, P.G., KEAST, D. Development of alterations in the primary immune response of mice by exposure to fresh cigarette smoke. *International Archives of Allergy and Applied Immunology* 46(4): 481-486, 1974.
- (109) THOMAS, W.R., HOLT, P.G., KEAST, D. Humoral immune response of mice with long-term exposure to cigarette smoke. *Archives of Environmental Health* 30(2): 78-80, February 1975.
- (110) THOMAS, W.R., HOLT, P.G., KEAST, D. Recovery of immune system after cigarette smoking. *Nature* 248(5446): 358-359, March 22, 1974.

- (111) TOMIOKA, H., ISHIZAKA, K. Mechanisms of passive sensitization. II. Presence of receptors for IgE on monkey mast cells. *Journal of Immunology* 107(4): 971-978, October 1971.
- (112) TOWNLEY, R.G., RYO, U.Y., KOLOTKIN, B.M., KANG, B. Bronchial sensitivity to methacholine in current and former asthmatic and allergic rhinitis patients and control subjects. *Journal of Allergy and Clinical Immunology* 56(6): 429-442, December 1975.
- (113) TRASOFF, A., BLUMSTEIN, G., MARKS, M. The immunologic aspect of tobacco in thromboangiitis obliterans and coronary artery disease. *Journal of Allergy* 7: 250-253, 1936.
- (114) TULEBAEV, R.K., MOSHKEVICH, V.S. Allergicheskie zabolevaniya verkhnikh dykhatel, nykh pytei u rabochikh tabachnogo proizvodstva (Allergic diseases of the upper respiratory tract in tobacco production workers). *Zhurnal Ushnykh Nosovykh I Gorlovykh Bolezney* (4): 12-14, 1975.
- (115) VON PIRQUET, C. Allergie, *Muenchener Medizinische Wochenschrift*. 53: 1457, 1906. In: Gell, P.G.H., Coombs, R.R.A. (Editors). *Clinical Aspects of Immunology*. Philadelphia, F.A. Davis Co., 1963, pp. 40-41.
- (116) VOS-BRAT, L.C., RUEMKE, P. Immunoglobuline concentraties, PHA reacties van lymfocytten in vitro (Immunoglobulin concentrations, PHA reactions of lymphocytes in vitro, and some antibody titers of healthy smokers). *Jaarboek Kankeronderzoek Kankerbestruiding* 19: 49-53, 1969.
- (117) WALDMAN, R.H., BOND, J.O., LEVITT, L.P., HARTWIG, E.C., PRATHER, E.C., BARATTA, R.L., NEILL, J.S., SMALL, P.A., JR. An evaluation of influenza immunization. Influence of route of administration and vaccine strain. *Bulletin of the World Health Organization* 41: 543-548, 1969.
- (118) WANNER, A., HIRSCH, J.A., GREENELTCH, D.E., SWENSON, E.W., FORE, T. Tracheal mucous velocity in beagles after chronic exposure to cigarette smoke. *Archives of Environmental Health* 27: 370-371, December 1973.
- (119) WARR, G.A., MARTIN, R.R. Concanavalin-A agglutination of human pulmonary alveolar macrophages from nonsmokers and cigarette smokers: A comparison. *Life Sciences* 18(10): 1177-1184, May 15, 1976.
- (120) WARR, G.A., MARTIN, R.R. In vitro migration of human alveolar macrophages: Effects of cigarette smoking. *Infection and Immunology* 8(2): 222-227, August 1973.
- (121) WARR, G.A., MARTIN, R.R., HOLLEMAN, C.L., CRISWELL, B.S. Classification of bronchial lymphocytes from nonsmokers and smokers. *American Review of Respiratory Disease* 113(1): 96-100, January 1976.
- (122) WEARY, P.E., WOOD, B.T. Allergic contact dermatitis from tobacco smoke residues. *Journal of the American Medical Association* 208(10): 1905-1906, June 9, 1969.
- (123) WEHNER, A.P., OLSON, R.J., BUSCH, R.H. Increased life span and decreased weight in hamsters exposed to cigarette smoke. *Archives of Environmental Health* 31(3): 146-153, May/June 1976.
- (124) WESTCOTT, F.H., WRIGHT, I.S. Tobacco allergy and thromboangiitis obliterans. *Journal of Allergy* 9: 555-564, 1938.
- (125) WHITEHEAD, R.H., HOOPER, B.E., GRIMSHAW, D.A., HUGHES, L.E. Cellular immune responsiveness in cigarette smokers. *Lancet* 1(7868): 1232-1233, June 15, 1974.
- (126) WIDE, L., BENNICHT, H., JOHANSSON, S.G.O. Diagnosis of allergy by an in vitro test for allergen antibodies. *Lancet* 2: 1105-1107, November 25, 1967.
- (127) WINGERD, J., SPONZILLI, E.E. Concentrations of serum protein fractions in white women: Effects of age, weight, smoking, tonsillectomy, and other factors. *Clinical Chemistry* 23(7): 1310-1317, 1977.

- (128) YEATES, D.B., ASPIN, N., LEVISON, H., JONES, M.T., BRYAN, A.C. Mucociliary tracheal transport rates in man. *Journal of Applied Physiology* 39(3): 487-495, September 1975.
- (129) ZDICHYNEC, B. Zmeny v bilem krevnim obraze u dlouhodobych kuraku. (Na Zaklade srovnani souboru nemocnych s transmuralnim srdecnim infarktem a nemocnych bez priznaku koronarni choroby v predchorobi) (The change of the white blood cells count in long-term smokers. [Basis for the comparison of patients with transmural myocardial infarct and patients without symptoms of coronary disease]). *Vnitri Lekarstvi* 17(4): 327-330, April 1971.
- (130) ZUSSMAN, B.M. Atopic symptoms caused by tobacco hypersensitivity. *Southern Medical Journal* 61(11): 1175-1179, 1968.
- (131) ZUSSMAN, B.M. Tobacco sensitivity in the allergic population. *Journal of Asthma Research* 11(4): 159-168, 1974.

11. INVOLUNTARY SMOKING.

CONTENTS

Introduction	5
<hr/>	
Constituents of Tobacco Smoke and Their Absorption by the Nonsmoker	6
Carbon Monoxide.....	15
Nicotine.....	24
Other Substances.....	24
<hr/>	
Effects of Tobacco Smoke on the Nonsmoker.....	25
General Population	25
Effects of Carbon Monoxide in Psychomotor Tests ...	28
Special Populations	29
Cardiovascular Disease	29
Chronic Obstructive Lung Disease.....	31
Hypersensitivity	31
Children.....	31
<hr/>	
Summary.....	33
<hr/>	
Recommendations	35
<hr/>	
References	36

LIST OF FIGURES

Figure 1.—Calculated buildup of CO under varying conditions of ventilation and smoking.....	22
--	----

LIST OF TABLES

Table 1.—Constituents of cigarette smoke: ratio of sidestream smoke to mainstream smoke.....	6
Table 2.—Measurement of constituents of tobacco smoke in experimental conditions	7
Table 3.—Measurement of constituents of tobacco smoke under natural conditions.....	16
Table 4.—Median percent carboxyhemoglobin (COHb) saturation and 90 percent range for nonsmokers by location.....	23
Table 5.—Admission rates (per 100 infants) by diagnosis, birth weight, and maternal smoking.....	33
Table 6.—Pneumonia and bronchitis in the first 5 years of life, by parents' smoking habit and morning phlegm....	34

Introduction

The effects of smoking on the smoker have been extensively documented in other chapters of this report. This chapter will review the effects of tobacco smoke on the nonsmoker, an area in which there has been increasing concern in the past several years (66a, 76, 77). This topic has been referred to as "passive smoking" or "secondhand" smoking as well as "involuntary smoking." The term involuntary smoking will be used to mean the inhalation by the nonsmoker of tobacco combustion products from smoke-filled atmospheres. This type of exposure is, in a sense, "smoking" because it provides exposure to many of the same constituents of tobacco smoke that voluntary smokers experience. It is also "involuntary" because the exposure occurs as an unavoidable consequence of breathing in a smoke-filled environment.

The chemical constituents found in an atmosphere filled with tobacco smoke are derived from two sources—mainstream and sidestream smoke. Mainstream smoke emerges from the tobacco product while being drawn through the tobacco during puffing. Sidestream smoke rises from the burning cone of tobacco. For several reasons, mainstream and sidestream smoke contribute different concentrations of many substances to the atmosphere: different amounts of tobacco are consumed in the production of mainstream and sidestream smoke; the temperature of combustion for tobacco is different during puffing than while smouldering; and certain substances are partially absorbed from the mainstream smoke by the smoker. The amount of a substance absorbed by the smoker depends on the characteristics of the substance and the depth of inhalation by the smoker.

When the smoker does not inhale the smoke into his lungs, the smoke he exhales contains less than half its original amount of water-soluble volatile compounds, four-fifths of the original nonwater-soluble compounds and particulate matter, and almost all of the carbon monoxide (25). When the smoker inhales the mainstream smoke, he exhales into the atmosphere less than one-seventh of the amount of volatile and particulate substances that were originally present in the smoke, and he also reduces the exhaled CO to less than half its original concentration (26). As a result, different concentrations of substances are found in exhaled mainstream smoke depending on the tobacco product, composition of the tobacco, and degree of inhalation by the smoker.

The effects of cigarette smoke on the environment and on the nonsmoker in the environment will be examined by reviewing data on the constituents of cigarette smoke measured under various conditions and on the absorption of these constituents by the nonsmoker. The physiologic effects of this "involuntary smoking" will then be considered.

Constituents of Tobacco Smoke and Their Absorption by the Nonsmoker

Brunnemann, et al. (14) have recently presented a compilation of the levels of some of the important substances in mainstream cigarette smoke and the ratio of sidestream to mainstream levels for these substances (Table 1). The actual amount of the substance and the mainstream-to-sidestream ratio will vary with different types of tobacco tested and the method used to burn the cigarette, but Table 1 gives values generally consistent with those found by others (23, 45, 50). Many of the substances, including nicotine, carbon monoxide, and ammonia, are found in much higher concentrations in sidestream smoke than in mainstream smoke. Thus, the total smoke exposure of nonsmokers is quantitatively much smaller than the exposure of smokers, but the smoke nonsmokers inhale may be qualitatively richer in certain compounds than mainstream smoke. This qualitative

TABLE 1.—Constituents of Cigarette Smoke.¹ Ratio of sidestream smoke (SS) to mainstream smoke (MS)

A. GAS PHASE	MS	SS/MS		MS	SS/MS
Carbon Dioxide	20-60 mg	8.1	Nitrogen Oxides (NO _x)		
Carbon Monoxide	10-20 mg	2.5	Ammonia	80 µg	73
Methane	1.3 mg	3.1	Hydrogen cyanide	430 µg	0.2 ²
Acetylene	27 µg	0.8	Acetonitrile	120 µg	3.9
Propane Propene	0.5 mg	4.1	Pyridine	32 µg	10
Methylchloride	0.65 mg	2.1	3-Picoline	24 µg	13
Methylfuran	20 µg	3.4	3-Vinylpyridine	23 µg	28
Propionaldehyde	40 µg	2.4	Dimethylnitrosamine	10-65 µg	52
2-Butanone	80-250 µg	2.9	Nitrosopyrrolidine	10-35 µg	27
Acetone	100-600 µg				
B. PARTICULATE PHASE	MS	SS/MS		MS	SS/MS
"Tar"	1-40 mg	1.7	Quinoline	1.7 µg	11
Water	1-4 mg	2.4	Methylquinolines	0.7 µg	11
Toluene	108 µg	5.6	Aniline	360 ng	30
Stigmasterol	53 µg	0.8	2-Naphthylamine	2 ng	39
Total Phytosterols	130 µg	0.8	4-Aminobiphenyl	5 ng	31
Phenol	20-150 µg	2.6	Hydrazine	32 ng	3
Catechol	130-280 µg	0.7	N'-Nitrosornicotine	100-500 ng	5
Napthalene	2.8 µg	16	NNK ²	80-220 ng	10
Methylnapthalene	2.2 µg	28	Nicotine	1-2.5 mg	2
Pyrene	50-200 µg	3.6			
Benzo(a)pyrene	20-40 µg	3.4			

¹Nonfilter cigarette

²NNK = 4-(N-methyl-N-nitrosamino)-1-(3-pyridyl)-1-butanone (tobacco specific carcinogenic nitrosamine)

SOURCE: Adapted from Brunnemann (14).

TABLE 2.—Measurement of constituents of tobacco smoke in experimental conditions.¹

Reference, location, and dimensions	Ventilation	Amount of tobacco burned	Level of constituent	Measure of absorption
Anderson and Dalhamn (3). Room 80 m ³	6.4 air changes per hour	46 cig & 3 pipefuls	4.5 ppm CO 377 mg/m ³ nicotine	COHb .6%
Bridge and Corn (13). Party room 145 m ³	7.0 air changes per hour	50 cig & 17 cigars in 1.5 hr	7.0 ppm CO	
Party room 101 m ³	10.6 air changes per hour	63 cig & 10 cigars in 1.5 hr	9.0 ppm CO	
Brunnemann, et al. (16). Box .4 m ³	none 1.5 liters/min	10 cig in 1 hr 10 cig in 1 hr	2.7 ng/l dimethylnitrosamine 2.9 ng/l dimethylnitrosamine	
Small room 20 m ³	none none some	100 cig in 1 hr 100 cig in 1 hr 100 cig in 1 hr	.33 ng/l dimethylnitrosamine .23 ng/l dimethylnitrosamine 1.85 ng/l dimethylnitrosamine	