

Parvin (Theo.)

Three Cases of Rupture
of the Uterus

BY

THEOPHILUS PARVIN, M. D.


INDIANAPOLIS, IND.



REPRINT FROM VOLUME III.

Gynecological Transactions

1879



THREE CASES OF RUPTURE OF THE UTERUS.

BY THEOPHILUS PARVIN, M. D.,

Indianapolis, Ind.

THE three cases of uterine rupture which I have grouped, occurred in parturition. They illustrate three of the causes of this accident, and they also repeat the common proofs of its terrible mortality.

As to the frequency of rupture of the parturient uterus, statistics and statements have given various results, as one in 940,¹ one in 1,300 or 1,400,² one in 1,331,³ one in 2,433,⁴ one in 3,403,⁵ and less than one in 4,887.⁶ Johnson and Sinclair⁷ state that in 13,748 deliveries, the accident occurred seventeen times, or once in a little over 800.

The accident is probably much more frequent than the larger of these proportions would lead us to believe. The number of cases of rupture would, I am persuaded, be greatly increased if, first, known cases were published, and second, if unknown cases were discovered. The first statement is so obvious that it needs neither proof nor comment. As to the second, the testimony of Baudelocque was that he rarely made post-mortem examinations after craniotomies, without finding either ruptures or severe contusions of the uterus.

CASE I. *Rupture from mal-presentation.*—The history of the first case, for which I am indebted to my friend Dr. I. C. Walker, illustrates mal-presentation as a cause of uterine rupture, and is briefly this.

When I was called to Mrs. G., a stout, healthy German

¹ Burns.

² Ingleby.

³ Churchill.

⁴ Lehman.

⁵ Jolly.

⁶ Ramsbotham.

⁷ *Practical Midwifery*, London, 1858.

woman, about forty years of age, the mother of six living children, the physician in attendance stated that she had been in labor forty-eight hours, that the pains had become quite active, but had abruptly ceased six hours previously, and that there was a shoulder presentation which he had vainly endeavored to correct. The pulse was 150, the respiration hurried, and the countenance expressive of great distress and prostration. The rupture was upon the right side of the uterus, and the child had partly escaped into the abdominal cavity. There was no difficulty in turning and delivering. The child of course was dead, and the patient died five hours after delivery.

CASE II. *Rupture from fetal deformity—Hydrocephalus.*—Mrs. C., a healthy woman, thirty-five years of age, who had given birth to seven children and had one miscarriage, was taken in labor on the evening of April 9, 1878. The pains were tormenting but inefficient, and dilatation exceedingly slow, so that it was not until twelve o'clock the next day that a satisfactory digital examination could be made, when the head was found unusually large. By 1 P. M. the pains had become quite severe, but after a terribly intense pain ceased altogether. They did not return, although the patient made frequent voluntary efforts. A thorough examination convinced Dr. McShane¹ that the infant was hydrocephalic. Dr. G. N. Duzan, one of the ablest practitioners of the State, was summoned. Upon his arrival at 5 P. M. he perforated the head, evacuating about a quart of fluid, and delivered. Immediately after removing the placenta he was struck by the woman's remarkable pallor and evidences of prostration.

At once thinking of concealed hemorrhage, he introduced his hand into the uterus until it came in contact with the liver,² there being a rent of not less than six inches in the

¹ I am indebted to Dr. J. T. McShane, of Carmel, Ind., who had charge of this patient, and to Dr. G. N. Duzan, the consultant, for this report.

² Dr. Duzan's experience reminds one of a passage in Blundell's *Lectures on Midwifery*, where this eminent obstetrician describes his pass-

fundus and right side of the uterus. Nevertheless, the patient lived for thirty-eight hours.

CASE III. *Rupture from improper administration of ergot.*—On Wednesday, July 3, at 1 P. M., my friend, Dr. T. N. Bryan, requested me to visit with him a case of ruptured uterus to which he had been called in the morning. The patient, a robust Irish woman, was thirty-five years of age. She had been taken in labor at 9 on the previous evening. Two former pregnancies had occurred, but the children were still-born. In this third labor a German midwife, at 11 P. M., two hours after the commencement of labor, was sent for. The midwife at 1 A. M. called in a doctor who was associated with her in obstetrical practice.

She represented both then, and subsequently, that the presentation was of the vertex, and that the labor was making favorable progress. Nevertheless two doses, each of a drachm, of the fluid extract of ergot were given at half an hour's interval. The pains were frequent and severe until 3 A. M., and then ceased.

Dr. Bryan found her at 6 A. M. with a feeble, rapid pulse, and complaining of great abdominal distress. She was vomiting every few minutes, and there was some external hemorrhage. The doctor recognized the nature¹ of the accident, and the patient rallying somewhat, and her friends consenting to have efforts made for her delivery, I was summoned.

I found the patient lying upon her left side, her greatest distress being referred to the left iliac region, and its vicinity. The utero-fetal tumor was directed somewhat obliquely from this to the right. Introducing my right hand gently and gradually into the vagina, it first came in contact with a fold of intestine; then passing it on and direct-

ing his hand through a rent in the uterus—for the purpose of reaching the feet to turn—when “he perceived the intestines, felt the beat of the large abdominal arteries, touched the edge of the liver,” etc.

¹ When Dr. Bryan suggested to the midwife and her assistant that there was a rupture of the womb, the latter naïvely replied: “It was quite likely. He had seen several cases of the kind.”

ing it towards the left side of the false basin, I reached the child's right hand, and further up the head. A few minutes' careful examination satisfied me that there was a lateral rupture of the uterus and also of the vagina, that the head and shoulders had escaped through this aperture, but that it was ample for their return without the least violence. Then turning my hand into the uterus, and carrying it upward and to the right, I touched the right knee of the child, and hooking a finger over it, soon succeeded in bringing the foot to the vulva.

Version was then readily completed, and with Dr. Bryan's skillful assistance the delivery of the child, which evidently had been dead some hours, was soon accomplished. Dr. Bryan removed the placenta without difficulty, and but trifling hemorrhage ensued; no prolapsed intestine could now be discovered, and the uterus seemed fairly contracted. The very night after the accident, the patient's attendants permitted her to get out of bed and sit in a chair for half an hour! In the treatment of this case, opium and quinia were used, ice-bags to the abdomen, and injections of a solution of carbolic acid. Her temperature only once rose as high as 103° , and most of the time varied between 101° and 102° . The pulse for one day was less than 100, and the bladder and bowels were evacuated spontaneously. Early Friday night she was seized with violent vomiting, and it continued through the night. Most carelessly and unfortunately neither Dr. Bryan nor I was sent for. From that night she commenced to sink, so that all hope of her recovery was lost, and she died a little more than four days after the rupture occurred.

In reviewing the history of the case, my belief is that this patient would probably have recovered had she been properly nursed.

According to Velpeau,¹ Albucasis first mentioned a case of rupture of the uterus; then Plater² one in 1584, and

¹ *Traité Complet de l'Art des Accouchemens.*

² Plater's case was the concubine of a priest.

Fabricius de Hildanus cites an analogous case observed in 1593, and adduces two others from Cornarius; "but Guillemeau seems first to have comprehended the nature of the phenomenon which his predecessors had only indicated."

In regard to the etiology of uterine rupture, may we not sum up the intrinsic causes — of course all violent manipulation and all use of instruments are excluded — in this statement: want of correspondence between the uterine force and the resistance it attempts to vanquish. Thus there may be weakened muscular fibre in some part of the uterus; uterine force is exerted, resistance is greater and the weak fibre gives way. Or again, the pelvic entrance is blocked by a hydrocephalic head, or the pelvic cavity packed by a shoulder, and the uterus, roused by this resistance impossible of removal, is ruptured in the vain effort. Or the uterus, equal to the gradual propulsion of the descending head or the slow dilatation of the mouth, is called upon, when goaded to unnatural activity by ergot, to accomplish in a few minutes that which nature designed should occupy hours, and rupture is the result.

Duparcque in his classic work¹ has placed this first among his "conclusions" as to rupture of the body of the uterus in labor: such ruptures have for their determining cause uterine contractions.

And our own eminent countryman and fellow-member, Dr. Trask, who has studied the subject so thoroughly, observes: "Unless caused by direct violence, rupture must, in almost every case, be the result of the contraction of the uterine fibres, whether the uterus be healthy or diseased."²

Dr. Tyler Smith³ remarked, in referring to the causes of rupture of the uterus: "I do not think sufficient prominence has been given to uterine motor action, which, in many cases, is the sole cause of the mischief, and which plays an important part in all." And, again, in the same lecture:

¹ *Histoire Complète des Ruptures et des Déchirures de l'Utérus, du Vagin et du Périnée.* Paris, 1839.

² *American Journal of the Medical Sciences*, vol. xv., 1848.

³ *Lancet*, October 28, 1848.

"Undoubtedly cases of rupture of the uterus do occur which are dependent upon softening of the uterus from inflammatory action, either during or before labor, or upon malignant disease of the uterus; but such cases are rare when compared with rupture from self-contraction of the uterus."

Playfair,¹ by the way, in referring to Tyler Smith's views, attributes to him the assertion "that ruptures are relatively as common in first as subsequent pregnancies." On the contrary, I read Tyler Smith as saying,² "It is an interesting and remarkable fact that ruptures of the uterus seldom happen to primiparous women." Be this as it may, and to return, confirmation of Tyler Smith's views as to rupture occurring in certain cases without there being any structural change in the uterine walls to explain the accident, was given by Dr. Robert Barnes when he asserted³ "that he had carefully examined the tissues in three cases of rupture, and he had found no more degeneration than that normal amount of granular change of the fibre-cells which always existed toward the end of pregnancy as a preparation for solution of the tissues about to become superfluous; certainly, then, although degeneration of tissue might sometimes be present, it was not a constant or necessary condition."

On the other hand, Dr. Angus McDonald makes a statement in almost direct contradiction. He says,⁴ "For, as rupture of the cervical portion forms nearly the whole of the cases of uterine rupture, seeing that both experiment and clinical observation agree in leading us to the conviction that spontaneous rupture of the body or fundus is impossible if the uterine tissue is healthy, if we have fully mastered the mechanism by which cervical rupture is brought about, we have been able to explain the great bulk of uterine ruptures."

¹ *System of Midwifery.*

² *Op. cit.*

³ *Trans. Obst. Soc., London, vol. x., p. 45.*

⁴ *Trans. Edin. Obst. Soc., vol. iv., p. 431.*

Certainly the first part of Dr. McDonald's statement, viz. : "rupture of the cervical canal forms nearly the whole of the cases of uterine rupture," finds no support in the statistics of Dr. Trask. Those statistics show that of 148 cases of rupture occurring (21) in gestation, and (127) in parturition, seventy-six were of the fundus and the body, and only seventy-two of the cervix, involving more or less of the body and of the vagina.

It is not necessary to enlarge upon the special causes — mal-presentation, fetal deformity, and ergot improperly given — that produced rupture in the cases I have reported ; for obstetric literature has often had similar testimonies.

In regard to the treatment of this accident, but few words will be said. That treatment has varied from doing nothing to gastrotomy. Each of these two can boast some extraordinary results. Thus, Lambron,¹ of Orleans, in 1775, performed gastrotomy in a case seventeen hours after the rupture ; in 1779 he repeated the operation upon the same patient, and in 1781 she gave birth to a living child *per vias naturales*. On the other hand, Dubois² narrates a case where rupture occurred, and the escaped infant lodged in the right hypochondrium, the patient recovering. A new pregnancy was followed by rupture, the second child finding a lodgment in the left hypochondrium. One was finally removed by an abscess, the other by incision.

Just now the advocates for gastrotomy in this accident are presenting their views with signal ability. The teaching of Levret and Baudelocque is revived, revived too when the large experience of ovariologists in abdominal surgery has given priceless wisdom to guide and sustain it.

Nevertheless, while gastrotomy must be admitted to be the best course in case the fetus has entirely escaped into the abdominal cavity, and where it has partially thus es-

¹ Duparcque, *op. cit.* The first successful gastrotomy, on account of uterine rupture, given by Duparcque, was Thibault's, in October, 1767.

² Velpeau, *op. cit.*, ii., 199.

caped and cannot be returned to the uterus without violence; yet if the rent involve merely cervix and vagina and thus present abundant opportunity for drainage, is it not best to deliver by forceps, by craniotomy, or by podalic version? On the other hand, let the rupture be large and high up, may not gastrotomy be advisable to remove effused blood, etc., from the peritoneal cavity because there can be drainage?

The researches of Dr. Robert P. Harris certainly present gastrotomy in a more favorable light than would have been supposed. That seventy-five per cent. of the mothers may be thus saved is indeed a most encouraging statement.

Finally in utterance, though first in practice, the prevention of the terrible accident is chiefly attainable, I believe, holding the creed of Tyler Smith, by the golden rule of moderating excessive reflex motor action; when these turbulent waves of contraction are beating in vain against unnatural barriers — of calming, if need be, with chloroform or other anesthetic, with opium or other anodyne, with bleeding or other sedative — and of removing those barriers.

NOW READY:

Gynecological Transactions.

VOLUME III.

*Being the Transactions of the American Gynecological
Society for the Year 1878.*

CONTENTS.

I. Annual Address on the Relation of Neurasthenia to Diseases of the Womb. By the first Vice-President, Dr. William Goodell.

II. A Case of Rupture of the Perineum without Implication of the Vulva. By Dr. J. C. Reeve.

III. On the Surgical Treatment of Stenosis of the Cervix Uteri. By Dr. J. Marion Sims.

IV. A Case of Extra-uterine Pregnancy, with Discharge of the Fetal Bones through the Bladder. By Dr. J. P. White.

V. A Case of Foot and Head Presentation; Fracture of the Spine in Utero. By Dr. J. T. Johnson.

VI. The Necessity of Early Delivery, as demonstrated by the Analysis of One Hundred and Sixty-one Cases of Vesico-vaginal Fistula. By Dr. T. A. Emmet.

VII. The Hand as a Curette in Post Partum Hemorrhage. By Dr. H. P. C. Wilson.

VIII. The Treatment of Post Partum Hemorrhage. By Dr. R. A. F. Penrose.

IX. Dermoid Tumors of the Ovary. By Dr. W. H. Byford.

X. A Contribution to the Study of the Treatment of the Acute Parenchymatous Nephritis of Pregnancy. By Dr. W. L. Richardson.

XI. Alternating Anterior and Posterior Version of the Uterus. By Dr. S. C. Busey.

XII. Remarks on Gastro-elytrotomy. By Dr. H. J. Garrigues.

XIII. The Pêndulum Leverage of the Obstetric Forceps. By Dr. A. H. Smith.

XIV. Rectal Alimentation in the Nausea and Inanition of Pregnancy; Intestinal Inanition, an Important Factor and the True Solution of its Efficiency. By Dr. H. F. Campbell.

XV. Three Cases of Rupture of the Uterus. By Dr. T. Parvin.

XVI. On the Early Delivery of the Placenta when Previa; with the Relation of a Case of Spontaneous Detachment of the Placenta without hemorrhage. By Dr. I. E. Taylor.

XVII. Unexpected Narcotism induced suddenly on the Third Day of the Administration of Three-grain Suppositories of Opium. By Dr. J. P. White.

XVIII. Treatment of Pelvic Indurations and Adhesions. By Dr. E. Van de Warker.

XIX. On some Points in Connection with the Treatment of Sterility. By Dr. A. Reeves Jackson.

XX. A Case of Extreme Anteversion and Ante flexion of the Uterus at the Full Term of Pregnancy. By Dr. I. E. Taylor, of New York.

XXI. Memoir of Edmund Randolph Peaslee, M. D., LL. D. By Dr. Fordyce Barker.

XXII. In Memoriam: Washington Lemuel Atlee. By Dr. T. M. Drysdale.

XXIII. The Mechanism of Retroversion and Prolapsus of the Uterus considered in Relation to the Simple Lacerations of the Cervix Uteri and their Treatment by Bloody Operations. By Dr. Nathan Bozeman.

Index of Obstetric and Gynecological Literature of all Countries for the Year 1877.

Price of Vol. III., \$5.00; Vols. I. and II., contents of which will be found on the following pages, will be sent to one address by mail or express, prepaid, on receipt of \$10.00 by the publishers. The three volumes supplied for \$15.00.

Address

HOUGHTON, OSGOOD AND COMPANY,
WINTHROP SQUARE, BOSTON.

TRANSACTIONS
OF
The American Gynecological Society.

VOL. I., FOR THE YEAR 1876.

400 pages. Many illustrations.

The Annual Address by the President, Dr. Fordyce Barker, of New York, and the following papers:—

I. Etiology of Uterine Flexures, with the proper mode of treatment indicated. By Thomas Addis Emmet, New York.

II. Cicatrices of the Cervix Uteri and Vagina. By Alexander J. C. Skene, M. D., New York.

III. Extirpation of the Functionally Active Ovaries for the Remedy of otherwise Incurable Diseases. By Robert Battey, Rome, Ga.

IV. On Central Rupture of the Perineum. By J. Matthews Duncan, M. D., Edinburgh, Scotland.

V. *Viburnum Prunifolium* (Black Haw); its Uses in the Treatment of the Diseases of Women. By Edward W. Jenks, M. D., Detroit, Mich.

VI. An Illustration of Xenomenia. By Theophilus Parvin, M. D., Indianapolis, Indiana.

VII. Relations of Pregnancy to General Pathology. By Robert Barnes, M. D., London, Eng.

VIII. The Spontaneous and Artificial Destruction and Expulsion of Fibrous Tumors of the Uterus. By W. H. Byford, Chicago, Ill.

IX. Case of Abdominal Pregnancy, treated by Laparotomy. By T. Gaillard Thomas, M. D., New York.

X. Pneumatic Self-Replacement in Dislocations of the Gravid and Non-gravid Uterus. By Henry F. Campbell, M. D., Augusta, Ga.

XI. Hydrate of Chloral in Obstetrical Practice. By W. L. Richardson, M. D., Boston, Mass.

XII. Labor Complicated with Uterine Fibroids and Placenta Previa. By James R. Chadwick, M. D., Boston, Mass.

XIII. Latent Gonorrhœa, with Regard to its Influence on Fertility in Women. By Emil Noeggerath, M. D., New York.

XIV. Death from Urinemia in Certain Cases of Malignant Diseases of the Uterus. By Alfred Wiltshire, M. D., London, England.

XV. Clinical Memoir on some of the Genital Lesions of Childbirth. By William Goodell, M. D., Philadelphia, Pa.

XVI. Hermaphroditism. By Lawson Tait, F. R. C. S., Birmingham, England.

XVII. Cases of Cystic Tumors of the Abdomen and Pelvis. By George H. Bixby, M. D., Boston, Mass.

XVIII. Case of Solid Uterus Bipartitus; both ovaries removed for the relief of epileptic seizures ascribed to ovarian irritation. By E. Randolph Peaslee, M. D., New York.

XIX. The Origin and History of Calculi found in the Bladder after the Cure of Vesico-Vaginal Fistula by Operation. By Henry F. Campbell, Augusta, Ga.

XX. Rare Forms of Umbilical Hernia in the Fetus. By James R. Chadwick, M. D., Boston, Mass.

XXI. In Memoriam Gustav Simon. By Paul F. Mundé, M. D., New York

Price of Volume I., \$5.00.

As contributions to advanced gynecology they exceed in value anything which we have ever before seen collected in any one volume.—*N. Y. Medical Record.*

This volume of Transactions is certainly superior to any book of the kind that has been issued by the American Press.—*St. Louis Clinical Record.*

Now we have the first of, we hope, a long row of volumes which will bear comparison with the Transactions of European Societies. Indeed, these must look to their laurels in this noble emulation.—*Edinburg Medical Journal.*

This volume is one of the best collections of gynecological papers that has been published this year.—*Annales de Gynécologie.*

VOLUME II., FOR 1877.

650 pages, 12 chromo-lithographs and many other illustrations.

Contains the following papers, with the discussions to which they gave rise:—

I. Annual Address, on Medical Gynecology. By the President, Dr. Fordyce Barker, of New York.

II. The Functions of the Anal Sphincters, so called. With two wood-cuts. By Dr. James R. Chadwick, of Boston, Mass.

III. Excision of the Cervix Uteri; its Indications and Methods. With one wood-cut. By Dr. John Byrne, of Brooklyn, N. Y.

IV. Report on the Corpus Luteum. With twelve chromo-lithographic plates. By Dr. John C. Dalton, of New York.

V. The Pathology and Treatment of Puerperal Eclampsia. By Professor Otto Spiegelberg, of Breslau, Prussia.

VI. Dilatation of the Cervix Uteri for the Arrest of Uterine Hemorrhage. By Dr. George H. Lyman, of Boston.

VII. The Principles of Gynecological Surgery applied to Obstetric Operations. By Dr. A. J. C. Skene, of Brooklyn, N. Y.

VIII. On the Necessity of Caution in the Employment of Chloroform during Labor. By Dr. William T. Lusk, of New York.

IX. The Present Status of the Intra-Uterine Stem in the Treatment of Flexions of the Uterus. By Dr. Ely Van de Warker, of Syracuse, N. Y.

X. Supplement to the report of a case of Xenomenia made at the first Annual Meeting in 1876. By Dr. Theophilus Parvin, Indianapolis, Ind.

XI. A Case of Vaginal Ovariectomy. By Dr. William Goodell, of Philadelphia, Penn.

XII. Is there a Proper Field for Battey's Operation? By Dr. Robert Battey, of Rome, Ga.

XIII. Subsulphate of Iron as an Antiseptic in the Surgery of the Pelvis. By Dr. H. P. C. Wilson, of Baltimore, Md.

XIV. Tetanus after Ovariectomy. By Dr. Theophilus Parvin, of Indianapolis, Ind.

XV. Sarcoma of the Ovaries. By Dr. Washington L. Atlee, of Philadelphia, Penn.

XVI. The Value of Electrolysis in the Treatment of Ovarian Tumors, as seen in the Light of Recent Experience. By Dr. Paul F. Mundé, of New York.

XVII. Congenital Absence and Accidental Atresia of the Vagina; Mode of Operation to Establish the Canal, and to Evacuate retained Menstrual Blood. With three wood-cuts. By Dr. Thomas Addis Emmet, of New York.

XVIII. Case of Sarcoma of the Kidney in a negro child. With Heliotype. By Dr. W. H. Geddings, Aiken, S. C.

XIX. The Hystero-Neuroses. By Dr. George J. Engelmann, of St. Louis, Mo.

XX. Cases illustrating Important Points connected with Ovariectomy. By Dr. Gilman Kimball, of Lowell, Mass.

XXI. The Radical Treatment of Dysmenorrhea and Sterility by Rapid Dilatation of the Canal of the Neck of the Uterus. With one wood-cut. By Dr. Ellwood Wilson, of Philadelphia, Penn.

XXII. Dr. Uvedale West's Views of Rotation, as illustrated by the Contrast between the Mechanism of Simple Occipito-Posterior Positions and those of the Bregmato-Cotyloid Variety. By Dr. John P. Reynolds, of Boston, Mass.

XXIII. Vascular Tumors of the Female Urethra; with the Description of an Instrument devised to Facilitate their Removal. By Dr. A. Reeves Jackson, of Chicago, Ill.

XXIV. Lying-in Hospitals; especially those in New York. By Dr. Henry J. Garrigues, of Brooklyn, N. Y.

XXV. The Simpler Varieties of Perineal Laceration; their Consequences and Treatment. By Dr. Thaddeus A. Reamy, of Cincinnati, Ohio.

XXVI. The Cyclical Theory of Menstruation. By Dr. John Goodman, of Louisville, Ky.

XXVII. In Memoriam Charles E. Buckingham. With a Heliotype Portrait. By Dr. George H. Lyman, of Boston, Mass.

Index of Obstetric and Gynecological Literature of all Countries, from July 1, 1876, to January 1, 1877.

The price of Volume II. is \$6.50. Vols. I. and II. will be sent by mail or express, post-paid, for \$10.00. Address,

HOUGHTON, OSGOOD & CO., BOSTON.

OPINIONS OF THE PRESS.

As contributions to advanced gynecology they exceed in value anything which we have ever before seen collected in any one volume. Without exception, every paper has an interest and value attached to it which is intrinsic, and which reflects the best efforts of the respective authors. . . . The work is elegantly printed, neatly and tastefully bound. — *N. Y. Medical Record*.

This volume of Transactions is certainly superior to any book of the kind that has been issued by the American press. The papers read and here presented are well written and well digested; and the discussions reported have almost as much value as the papers themselves. — *St. Louis Clinical Record*.

We think that never before was such a number of excellent monographs collected together in one volume of Transactions. It is a model in every respect. It is a volume that should be in the hands of every general practitioner as well as specialist, as in it may be found most useful suggestions for every day practice. We cannot commend the work too highly, and it is our earnest hope that future meetings of the society will give origin to volumes of Transactions equal to the one before us. — *The Hospital Gazette*.

In conclusion, we cannot but regard this volume as a fitting monument of the progress of this most progressive department of medicine. The discussions are rich, and are full of such epitomized good sense well expressed, as is rarely found in any volume of Transactions, while the whole contents exhibit the most careful supervision of the editor. — *Toledo Medical and Surgical Journal*.

Organized amid the whirl and bustle of last summer, the American Gynecological Society bids fair to realize the brightest hopes of its founders. Its first volume of Transactions, now before us, will bear the closest scrutiny, and safely challenge the severest criticisms, so superior is it in its breadth of thought and observation, scientific worth, and literary excellence; the writers of the papers in these Transactions being no mere novices, but men skilled in the profession they represent, and experts in the branch of study and practice which they especially call their own. — *Philadelphia Medical Times*.

It is not a trifling matter to be able to say of this volume that it is the handsomest of the kind anywhere produced. . . . But this is not only an elegant volume, it is a good one; creditable to the authors, creditable to the great republic. . . . Now we have the first of, we hope, a long row of volumes which will bear comparison with the Transactions of European Societies. Indeed, these must look to their laurels in this noble emulation. — *Edinburgh Medical Journal*.

This book of nearly 400 pages contains a series of excellent articles upon gynecological topics, written by the most prominent gynecologists of America and England. — *Allg. Wiener Med. Zeitung*.

This volume is one of the best collections of gynecological papers that has been published this year. . . . We do not doubt that it will confer the greatest honor upon the medical literature of America. — *Annales de Gynécologie*.

Few volumes that have recently come under our notice have been so extremely interesting and instructive. The papers reach a very high order of excellence, and some indeed are superlatively good. — *American Journal of Obstetrics*.

