

PATRICK (H. T.)

The Bryson Symptom  
in Exophthalmic Goitre.

BY

HUGH T. PATRICK, M. D.,  
Professor of Neurology in the Chicago Polyclinic.

REPRINTED FROM THE  
New York Medical Journal  
*for February 9, 1895.*





Reprinted from the *New York Medical Journal*  
for February 9, 1895.



THE BRYSON SYMPTOM  
IN EXOPHTHALMIC GOÏTRE.\*

WITH A REPORT OF FORTY CASES.

By HUGH T. PATRICK, M. D.,  
PROFESSOR OF NEUROLOGY IN THE CHICAGO POLYCLINIC.

FIVE years ago Dr. Louise Fiske Bryson † first called attention to a condition in Graves's disease that has since been called the Bryson symptom—viz., diminished chest expansion or vital capacity. She cites it as one of the "distinctive, fixed conditions" prevailing in this disease which "afford the only rational basis for prognosis." Other fixed conditions are not specified and no measurements or cases are given. Dr. Græme M. Hammond ‡ considers it "of the greatest importance in regard to the prognosis" and "also of assistance in locating the seat of the lesion." He says further: "Dr. Bryson states that where the expansion is found to be reduced to half an inch or less the termination of the case is invariably fatal." He found the symptom present in the eight cases reported in

\* Read before the Chicago Medical Society, October 1, 1894.

† A Preliminary Note on the Study of Exophthalmic Goître. *N. Y. Med. Jour.*, December 14, 1889.

‡ A Contribution to the Study of Exophthalmic Goître. *N. Y. Med. Jour.*, January 25, 1890.

his paper, and as recovery took place the expansion increased. Dr. A. B. Pope\* has reported a case of exophthalmic goitre in a man with chest expansion of an inch and a half, and in the discussion Dr. E. Le Fevre said the symptom was "owing to some nervous influence similar to that which gave rise to the rapid heart action."

Three years after her first paper Dr. Bryson † resumes the theme, giving particular prominence to the respiratory symptoms, and emphasizing especially diminished chest expansion, but she seems to have receded somewhat from her former positive position as to its pathognomonic importance, as of twenty cases the Bryson symptom was present in only thirteen. She considers the disease, however, to be "a disorder of nutrition *respiratory in its first manifestations*, ‡ and even makes this assumption the foundation of a therapy consisting of a series of respiratory gymnastics executed by means of Taylor's respirator. These views have been quoted at home and abroad, § and the Bryson symptom would seem to be about to take its place among the important signs of Graves's disease as bearing upon the diagnosis, prognosis, pathology, and treatment. Ham-

\* *N. Y. Med. Jour.*, April 19, 1890.

† Exophthalmic Goitre: A View of Thirty Cases. *The Post-graduate*, July, 1892.

‡ Italics mine.

§ Frank S. Johnson, *Trans. Ill. State Med. Soc.*, 1893, p. 193; J. Madison Taylor, *Med. News*, 1893, pp. 673, 711; William Oliver Moore, *Internat. Clinics*, 1893, i, p. 92; Edmund Lee Tompkins, *Amer. Jour. of Obstet.*, November, 1893; W. H. Thomson, *N. Y. Med. Jour.*, June 3, 1893; William A. Hammond, *Dis. of the Nerv. Syst.*, 1891, p. 790; Landon Carter Gray, *Treat. on Nerv. and Ment. Dis.*, p. 551; Mannheim, *Der Morbus Gravesii*, Berlin, 1894, p. 33; Oppenheim, *Lehrb. der Nervenkrank.*, Berlin, 1894, p. 829; Grasset et Rauzier, *Mal. du syst. nerv.*, Montpellier and Paris, 1894, ii, p. 309; Debove et Achard, *Manuel de méd.*, Paris, 1894, a-t. by Létienne, vol. iv, p. 397; Blocq, *Études sur les mal. nerv.*, Paris, 1894, p. 173.



mond,\* indeed, distinctly calls it the fourth cardinal symptom, a distinction, by the way, which belongs to Marie's symptom—tremor.

At the time when Dr. Bryson's first article appeared I had under observation a case of exophthalmic goître which I measured several times, finding the chest expansion not materially diminished. After this I measured a number of cases, finding the expansion sometimes diminished and sometimes about normal, and I presently noticed that diminished expansion seemed to go with a diminution in general vitality and muscular strength. I finally, then, began comparing the chest expansion with the hand grasp as recorded by the dynamometer, this affording the most convenient, if only approximately accurate, index of the general muscular condition. It is the result of these measurements that I wish to present. I may premise that only well-developed, in a measure typical, cases are comprised in this report, no *formes frustes* being included, and that the diagnosis was in every instance confirmed by some well-known neurologist. I have therefore deemed it unnecessary to lengthen my paper by giving the symptoms of each case in detail. With the exception just noted, the cases have not been selected, but embrace all that I was able to measure from November 1, 1892, to May 1, 1894.

The patients were all women, and it may be worthy of note that of some sixty cases seen since 1891 only two were in men.

It is a matter of common remark that no two observers take the chest expansion exactly alike, so that figures obtained by different persons can not be compared with confidence. To form a safe basis for comparison, therefore, I took the chest expansion and dynamometric measurement

\* Græme M. Hammond, *loc. cit.*

4 THE BRYSON SYMPTOM IN EXOPHTHALMIC GOÏTRE.

of twenty-eight women who came to the dispensary for various ailments, selecting those whose troubles would not, in my opinion, affect the result. These are presented in Table II.

TABLE II.—*For Comparison.*

No.	Age.	Affection.	General condition.	Chest expansion.	DYNAMOMETER.		
					R.	L.	Average.
1	52	Tabes.	Fair.	<small>Ctm.</small> 4	30	25	27·5
2	24	Hysteria.	Very good.	7	80	70	75
3	31	"	Good.	5	45	45	45
4	51	Chronic rheumatism.	Fair.	4	40	25	32·5
5	41	Incipient tabes.	Good.	5½	55	55	55
6	29	Cerebro-spinal syphilis; dement. paral. (?)	Good; very muscular.	5	100	75	87·5
7	30	Epilepsy.	Good.	3½	50	50	50
8	46	Ophthalmoplegia.	"	7	47	47	47
9	23	Brach. neuralgia.	Very good.	4½	80	70	75
10	43	"	Fair.	6	72	70	71
11	39	Chronic rheumatism.	Very fair.	5½	60	40	50
12	18	Sciatica.	Fair.	4½	55	50	52·5
13	26	Neurasthenia.	Very fair.	3¾	65	50	57·5
14	38	Hysteria and neuralgic pains.	Fair.	4½	58	42	50
15	17	Hystero-epilepsy.	Good.	5	55	47	51
16	28	Lumbago.	Very good.	6	95	95	95
17	36	Hysteria.	Fair.	4	30	30	30
18	22	"	Good.	4¾	46	46	46
19	34	Hysteria and neuralgia.	"	6	75	70	72·5
20	22	Hysteria and organic heart disease.	Fair.	4½	37	50	43·5
21	36	Headache.	Anæmic.	4	75	70	72·5
22	30	Tabes.	Very good.	5½	80	67	73·5
23	15	Hysteria.	Good; well developed.	4½	45	35	40
24	19	Valvular heart disease.	Anæmic.	4½	51	55	53
25	27	Headache.	.....	4½	50	40	45
26	41	Hysteria.	Good.	4½	61	54	57·5
27	23	Hysteria and lumbago.	Very good.	4	47	47	47
28	26	Sciatica.	Good.	6	77	75	76
				4·8	..	..	56·36

I am well aware also of the varying results given by the dynamometer, and I do not allege absolute accuracy for my figures or wish to make too sweeping deductions from them, but all measurements were taken by myself in as nearly a uniform manner as possible, and sources of error excluded as well as might be.

It will be seen that the average chest expansion in the forty cases of exophthalmic goitre is 4·3 centimetres and the hand grasp 43·75 kilogrammes; in the twenty eight other cases, 4·8 centimetres and 56·36 kilogrammes respectively—that is, the expansion in Graves's disease is diminished half a centimetre, or ten and a half per cent., the hand grasp 12·61 kilogrammes, or twenty two and two fifths per cent. In other words, the grasp is diminished more than twice as much as the expansion. It may be contended that the average expansion of the twenty-eight women used for comparison is not up to the normal, and I think this may possibly be true, but I also think they make a far better basis for comparison than statistics of measurements taken by some one else whose methods would doubtless differ in some degree from mine, and consequently whose results, other things being equal, would differ from mine. Further, these measurements were made in the same dispensary which furnished the majority of the cases of Graves's disease,\* and the subjects are fairly representative of the class of patients furnishing the material for this report, and, as before mentioned, the pathological conditions in these women were not such as to materially affect chest expansion or hand grasp. But even supposing that the expansion of the twenty-eight might be materially below the normal, we could not suppose in all these various cases a condition which would affect chest expansion to the exclusion of other muscular action, in-

\* Professor Mendel's *Poliklinik*, Berlin.

cluding hand grasp, and as we find that the hand grasp in Graves's disease is diminished twenty-two and two fifths per cent. and the expansion only ten and a half per cent., the natural conclusion would be that if one of these two is pathognomonic of this malady it would be the weakened grasp and not the diminished chest expansion, an assumption I would not for a moment entertain. Again, we find of the forty cases twenty-six with an expansion below the average of the twenty-eight, while thirty show a diminished grasp.

Could we take accurate dynamometric measurements of the lower extremities and of the pelvic muscles, I have no doubt we should find an equal falling off. Many of these patients complain of a sudden "giving way of the legs," and placed in the recumbent posture show their muscular weakness in the manner of rising.

A woman now under observation, not included in this report, is a case in point. It is not what could be called a very severe case; she does her own housework, gets about very well, hand grasp and chest expansion only slightly diminished, and yet she rises from the supine position much like a patient with idiopathic muscular atrophy; that is, she first raises the trunk by means of the arms till she rests on her elbows; then, again by help of the arms and with a peculiar wriggle or writhing movement to bring accessory muscles into play, she brings the trunk to the perpendicular; the erect position is then attained, with manifest effort, by turning round and pushing herself up with her hands first on the floor and then on her thighs or adjacent objects.

I think the diminished power of convergence often observed, as well as the occasional affection of the laryngeal muscles, is quite analogous to these other findings, and is simply a part of a general myasthenia, sometimes affecting one set of muscles more, sometimes another.



I find, further, that many of these cases show a rapid and marked falling-off in both chest expansion and hand grasp on repeated effort. Thus either may diminish one half after three or four trials in rapid succession, and in a general way the decrease affects both and is proportional to the debility.

The idea that the diminished chest expansion is simply part of a diminished vitality or energy is borne out by the history of individual cases. I have generally found that as the patient improves in general tone the expansion increases (and *vice versa*), although not necessarily in the same ratio, and that the grasp goes with it hand in hand. But as the diminished hand grasp shows the larger percentage in the table, so here it is apt to show the larger fluctuations, which would seem again to indicate that it is the more delicate index of the two. These facts are well illustrated by Case III. The rule, however, is not absolute, and Case XXXVII is a striking exception in the first two examinations.

The only one of my patients (Case III) who ever showed an expansion as low as half an inch (1.25 ctm.), indicating, according to Dr. Bryson, an absolutely fatal termination, had, four months later, gained over twenty pounds in weight, and was better than at any time during the six months she was under observation.

I think, then, we may conclude that the Bryson symptom, although present in many cases of exophthalmic goître, is in no wise pathognomonic of this affection, or even an important sign; that it has no special significance in relation to the prognosis, pathology, seat of the lesion, or treatment, and should be relegated back to the comparative obscurity of an individual in a large community of manifestations which all depend alike upon the general state; a state which makes the French designation of the

## 8 THE BRYSON SYMPTOM IN EXOPHTHALMIC GOÏTRE.

disease, exophthalmic cachexia, quite as appropriate as any other.\*

### VENETIAN BUILDING.

\* I wish to express my great gratitude to Professor Mendel, Professor Jolly, Professor Oppenheim, Professor Bernhardt, Professor Eulenburg, and Dr. Goldscheider of Berlin, Professor Brissaud of Paris, and Dr. J. Hughlings Jackson of London, for permission to examine cases entering into this report.



TABLE I.—GRAVES'S DISEASE.

No.	Age.	Duration of disease.	Date of examination.	Chest expansion. Centimetres.	Average.	DYNAMOMETER.			Remarks.
						R.	L.	Average.	
1	29	4½ years.	Nov. 1, 1892. Dec. 8, 1892. Feb. 25, 1893.	4½ 4 4	..... ..... .....	40 47 46	30 40 37	..... ..... .....	Following fright.
2	30	2 years.	Mar. 2, 1893. Nov. 1, 1892.	5½ 5	4·31 .....	50 55	45 45	41·8 .....	
3	32	4 years.	Feb. 17, 1893. Nov. 30, 1892. Dec. 14, 1892.	4 1½ 1½	4·75 ..... .....	60 57 12	57 2½ 16	54·2 ..... .....	At fourth examination she had been two weeks confined to bed, following mental worry. May 17th, much better; had gained over twenty pounds.
4	37	1 year.	Dec. 21, 1892. Jan. 28, 1893. May 17, 1893.	2½ 1½ 2	..... ..... 1·8	26 16 16	16 16 16	..... ..... 10·7	Is right-handed, but has always been stronger in left. Much worse since a confinement nine weeks ago.
5	31	Probably 4 years.	Nov. 9, 1892. Nov. 30, 1892. Oct. 15, 1892.	3 3½ 7	..... 3·25 .....	37 40 67	47 50 47	..... 43·5 .....	No treatment the past year, as treatment the two preceding years almost cured her. Much worse since influenza two years ago. Developed after hemiplegia. Cerebral apoplexy.
6	23	1 year.	Dec. 2, 1892.	5½	6·25	49	48	52·75	
7	17	1½ year.	Feb. 20, 1893.	4½	4·5	22	15	18·5	
8	40	12 years.	Nov. 19, 1892. Nov. 1, 1892.	5 4	5 3·75	65 30	55 25	60 25·75	Goitre from youth.
9	55	8 years.	Jan. 5, 1893. Nov. 2, 1892.	3½ 3½	3·5 3·5	37 30	37 30	33·5	
10	35	8 years.	Nov. 3, 1892.	2	2	20	10	15	
11	30	2 years.	Nov. 9, 1892.	6	6	67	63	65	
12	21	3 years.	Nov. 12, 1892.	7	7	60	50	55	
13	18	2 years.	Nov. 14, 1892.	6½	6·5	50	50	50	
14	28	4 months.	Nov. 12, 1892.	5½	5·5	44	44	44	
15	51	4½ years.	Nov. 16, 1892.	5½	.....	45	55	.....	Locomotor ataxia for the last ten years.
16	24	1 year.	Mar. 2, 1893. Nov. 19, 1892. Dec. 14, 1892.	5 4 4	5·25 ..... 4·5	42 30 3	40 45 62	45·5 ..... 45·6	Stigmata of hysteria; right hand excluded from estimate, as low figure is due to psychic inhibition.
17	19	3 years.	Nov. 22, 1892.	3½	3·5	45	45	45	
18	26	9 months.	Nov. 29, 1892.	4	.....	62	57	.....	
19	35	1 year.	Dec. 14, 1892. Jan. 10, 1893. Dec. 7, 1892.	3½ 4 4	3·83 ..... 4	67 67 62	53 53 56	59·33 ..... 59	Was much worse eight months ago; all symptoms were more pronounced.
20	48	2 years.	Dec. 20, 1892.	5	5	15	10	12·5	
21	20	9 months.	Dec. 20, 1892.	4	4	47	37	42	
22	21	Uncertain, 1 to 4 years.	Jan. 5, 1893. Jan. 20, 1893. Feb. 16, 1893.	4½ 3½ 5	..... ..... .....	48 50 47	47 35 37	..... ..... .....	Patient thinks an operation on the nose for catarrh aggravated the disease.
23	39	8 years.	Mar. 2, 1893. Apr. 27, 1893. July 10, 1893.	5 4½ 4½	..... ..... 4·45	42 40 35	40 30 35	..... ..... 40·5	
24	25	.....	Jan. 10, 1893. Feb. 17, 1893.	3½ 3½	..... 3·5	60 50	60 50	..... 55	Has had goitre ten years. Nervous and irritable four years. All symptoms of Graves's disease five weeks, following failure in business.
25	38	4 years (?).	Jan. 13, 1893. Feb. 16, 1893.	3 4	3 4	16 37	16 33	16 35	Has <i>imperative ideas</i> . March 2d, feels much better. Symptoms aggravated by operation on uterus two years ago.
26	21	3 years.	Feb. 18, 1893.	6½	6·5	30	20	25	
27	30	6 months.	Feb. 22, 1893.	4½	.....	67	67	.....	Difficult labor seven months ago, since which time is nervous and anamic; is now two months pregnant; all symptoms of Graves's disease six weeks.
28	29	4 years.	Mar. 2, 1893. Feb. 24, 1893.	4½ 5	4·5 5	85 71	85 59	76 65	Complicated with hysteria.
29	37	7 months (?).	Mar. 1, 1893.	5½	5·5	65	60	62·5	Symptoms aggravated by operation on uterus two years ago.
30	..	2 years.	Mar. 3, 1893.	3	3	60	57	58·5	Following confinement eleven months ago.
31	34	11 months.	Mar. 9, 1893. Mar. 20, 1893.	4½ 5½	..... 5	60 50	57 50	..... 56·75	
32	23	2 months.	Mar. 15, 1893.	4	.....	37	30	.....	
33	38	4 years.	June 13, 1893. May 17, 1893.	3½ 3½	3·75 3·5	32 45	30 45	32·25 45	Following a fright; sister has same disease, and is much worse. Following influenza; thyroidectomy thirteen months ago, with great improvement.
34	38	3 years.	May 12, 1893.	5½	.....	43	43	.....	Following fright.
35	29	4 to 5 years.	May 22, 1893. May 12, 1893.	4½ 4½	5 4·75	42 55	45 65	43·25 .....	
36	46	2 years.	July 10, 1893.	5	4·75	55	65	60	
37	18	2 years.	Apr. 23, 1893. Nov. 17, 1893.	4½ 4	4·5 .....	50 50	47 40	48·5 .....	
38	26	2 years.	Jan. 18, 1894. Jan. 30, 1894.	2 3½	..... 3·16	66 32	40 37	..... 45	
39	39	3 years.	Dec. 5, 1893. Jan. 19, 1894.	2 2½	..... .....	55 55	55 55	..... .....	
40	29	2 years (?).	Jan. 30, 1894. Jan. 18, 1894. Jan. 30, 1894.	3 3 3	..... ..... .....	50 40 32	45 38 27	..... ..... 38·66	Epilepsy; much worse the last three years.
			Feb. 12, 1894. Feb. 24, 1894. Mar. 6, 1894.	4 4 4½	..... ..... .....	30 35 40	25 23 32	..... ..... .....	After inception cured (?) by rest, etc.; recurrence one year ago.
			Mar. 31, 1894. Apr. 13, 1894.	4½ 4	..... 4·2	40 35	32 25	..... 28·5	
				4·3	.....	.....	.....	43·75	
				0·5	.....	.....	.....	12·61	
				10½	.....	.....	.....	22½	

Diminution.....

Diminution, per cent.....





# The New York Medical Journal.

A WEEKLY REVIEW OF MEDICINE.

EDITED BY

FRANK P. FOSTER, M.D.

---

THE PHYSICIAN who would keep abreast with the advances in medical science must read a *live* weekly medical journal, in which scientific facts are presented in a clear manner; one for which the articles are written by men of learning, and by those who are good and accurate observers; a journal that is stripped of every feature irrelevant to medical science, and gives evidence of being carefully and conscientiously edited; one that bears upon every page the stamp of desire to elevate the standard of the profession of medicine. Such a journal fulfills its mission—that of educator—to the highest degree, for not only does it inform its readers of all that is new in theory and practice, but, by means of its correct editing, instructs them in the very important yet much-neglected art of expressing their thoughts and ideas in a clear and correct manner. Too much stress can not be laid upon this feature, so utterly ignored by the “average” medical periodical.

Without making invidious comparisons, it can be truthfully stated that no medical journal in this country occupies the place, in these particulars, that is held by THE NEW YORK MEDICAL JOURNAL. No other journal is edited with the care that is bestowed on this; none contains articles of such high scientific value, coming as they do from the pens of the brightest and most learned medical men of America. A glance at the list of contributors to any volume, or an examination of any issue of the JOURNAL, will attest the truth of these statements. It is a journal for the masses of the profession, for the country as well as for the city practitioner; it covers the entire range of medicine and surgery. A very important feature of the JOURNAL is the number and character of its illustrations, which are unequalled by those of any other journal in the world. They appear in frequent issues, whenever called for by the article which they accompany, and no expense is spared to make them of superior excellence.

---

Subscription price, \$5.00 per annum. Volumes begin in January and July.

---

PUBLISHED BY

D. APPLETON & CO., 72 Fifth Avenue, New York.

