




## 

# Dr. Douglass's Practical HISTORY

OF

A New Epidemical Eruptive Milary Forer, with an Angina Ulcufculofa, Ec.

\$ .



### THE

## Practical HISTORY

#### OF

A New Epidemical Eruptive Miliary Fever, with an Angina Ulcufculofa

#### Which

Prevailed in Boston New-England in the Years 1735 and 1736.


By William Douglass, M. D.



### BOSTON, N.E.

Printed and fold by Thomas Fleet, at the Sign of the Heart and Crown in Cornhill. 1736.

GEON GENERA



## [i]

#### TO

## A Medical Society

### in Boston.

Gentlemen,

HIS Piece of Medical Hiltory does naturally addrefs it felf to you, confidering that I have the pleafure of being one of your number, that you have been fellow labourers in the management of this diftemper, and therefore competent judges of this performance, and that where difficult or extraordinary Cafes have occurred in any of your private practice, I was favoured to vifit the Patients in order to make a minute clinical enquiry: in flort, without your offifance this piece would have been lefs perfect, and not fo well vouched.

As this diftemper continues to spread and prevail in several Towns of this and the neighbouring Provinces, I thought it might prove a piece of humanity and benevolence, if after many months diligent observations made in most of the varieties which occur in this Illness, I did endeavour to reduce them to some easy distinct Historical and Practical Method. The vanity of appearing as an Author or writer was no inducement, because we all know that in a plantation life neither honour nor credit are to be acquired by writing. It is not published by way of

400104

of a Quack bill to procure Patients and their money, as has been the practice of some void of modesty and truth; because the Distemper is almost over in Boston, and while it prevailed here I could not well have attended more patients than what I had from time to time under my care, and make with attention the proper observations at the same time.

A fecondary reason for my writing is, to induce fome Gentlemen of the profession in our other Provinces and Colonies, where this distemper does or may prevail; to give some account of its appearance with them, in order to discover what influence, progress of time, varieties of climate and Soil may have in the phænomina of this disease. This method, of taking things originally that is from the life, if pursued (but by abler bands) in the Epidemical Distempers which may from time to time happen among st us, may be of considerable advantage in Physick.

A Speculation that is a novelle might have been composed sooner, but not a real History: for as amongst Naturalists, many repeated observations and experiments are requisite to form established truths or conclusions; so it ought certainly to be in the practice of Medicine, where no affair of Speculation or curiosity, but the life and death of a fellow Citizen is the object of our enquiry.

Yours, &c.

William Douglass.

## The Practical Hiftory of a new Epidemical Miliary Fever with an Angina Ulcusculosa.

(1)

HIS Diftemper did emerge 20th. May, 1735. in Kingston Township 50 Miles Eastward from Boston ; it was no foreign importation, Kingston being an inland place, of no Trade or confiderable communication. The first feized was a child who died in three Days Illness; about a Week thereafter in another Family at four Miles diftance, three Children were feized fucceffively, and died alfo the third Day; it continues fpreading gradually in that Township, feizing here and there particular Families with that degree of violence, that of the first circiter forty decumbents none recovered as we were informed. It was vulgarly called the Throat Illness, or a Plague in the Throat, and alarmed the Provinces of New-England very much. Some died of a sudden or acute Necrofis; but most of them by a Symptomatick affection of the Fauces or Neck; that is by Sphacelations or corresive Ulcerations in the Fauces, or by an infiltration and tumefaction in the Chops and fore part of the Neck, to turged, as to bring all upon a level between the chin and sternum, occasioning a strangulation of the Patient in a very fhort time.

After a few Weeks it fpreads into the neighbouring Townships, but with more mildnels. The first

first appearance that we can recollect of it in Bofton, was 20th August on a Child of Capt. Stannys at the North End; having white Specks in the Throat, and a cutaneous efflorescence : A few more in the fame Neighbourhood were feized in like manner, about the fame time. Towards the end of September it appeared in feveral parts of the Town, with a complaint of forenels in the Throat, Tonfils fwelled and speckt, Uvula relaxed, flight Fever, flush in the Face, and an Erysipelas like efflorescence on the neck, cheft and extremities; but being of no fatal or bad confequence, nothing more than a common cold was fuspected. Our first alarm was from a young Man How At. 20. in the beginning of Oftober : His Hiftory runs thus; He was lately arrived from Exeter to the Eastward, where his Brother died of this Illnefs; his Symptoms were great profiration of Strength, a speck in one of the Tonfils, colliquative Sweats, Pulse not high and full, but low, hard, firingy, unequal and more frequent than natural, deglutition good to the laft, no Sphacelation in the Throat, no eruption; from a rash inconfiderate opinion of forcibly quelling the Malignity, he was thrice let Blood, had fome Emeticks and Catharticks adminstred, and by profuse evacuations was gradually reduced, so as to die of a gentle decay of natural Strength, the 6th Day of Illnefs.

Beginning of November it fpread confiderably in Boston, especially amongft Children, with more violent Symptoms, and severals die of it in various Periods: it seemed to be at the hight, as to Numbers ailing and quantity of Deaths, the second Week of March; that Week there were 24 Burials, whereas communibus annis in that Season they are only 9 or 10 per Week.

Ic

It is generally in fo confiderable a Degree more favourable in Bofton, than in the Townships where it first prevailed; that many can scarce be persuaded of its being the fame Diffemper : It is neverthe lefs effentially the fame, there is no Symptom, even the most malignant that has appeared in News Hampfhire, but what the like has occured in Bofton. Perhaps Boston dry healthy air, good feeding, con-flitutions less Pforick, and the better management of the Sick, favoured us; the reasons for its proving more mortal in the other Towns, may be, the Country woodland and fresh water damps, (the Sheep in fenny lands are most susceptible of and fuffer most by the Rot) their coarse Food, salt Pork diet, Pforick Constitutions, (which is one of the principal Endemial distemperatures in New-England) bad Lodgings, and that mischievous Practice of using this Diftemper with protule evacuations, whereby the laudable and falutary cuticular eruption has been fo perverted as to be noticeable only in a few, and in these it was called a Scarlet Fever; the great profiration of Strength effential to this Diftemper is so much increased, as to render Nature an under match for the affaults of this Illnefs and its confequences. In fact to the Eaftward in forme Country Towns, at certain times have died 1 in 3 of the Sick, in others I in 4, in scarce any fewer than I in 6, whereas in Bofton not above I in 35 have died.

As in most Epidemical acute Illneffes, especially eruptive Fevers, (witness the Small Pox) fo in this, are very many varities or degrees, from the moft gentle and benign to the most malignant. Symptoms did vary chiefly from fomething inferutable in the Constitutions of Families and Persons ; the Scropholous and Plorick were the most susceptible of it, and did did fuffer moft by it; the Regimen had a confiderable influence, here fome who might have furvived the natural Symptoms did fuccumb by profufe U. S. and other evacuations, one of the moft effential Symptoms of this Diftemper (as before hinted) being great profiration of Strength. In fo great variety it is not poffible to give any concife fcholastick description, which may comprehend all: We shall therefore, as a Standard first describe the most frequent fort, as it appeared in good conflictuions. A previous liftlesness and languishing counte-

nance for a Day or two, or some other pranuncia as u. g. wet Nurses loofing their Milk. The first attack is fomewhat of a chill or fhivering; foon after follows Head ake or fome other versatile spasmodick pains, as pain in the back, joints, fide, Ge; a vomit. ing or nausea, or in some constitutions which are not eafily provoked to vomit, only a certain uneafinefs or ficknefs at Stomach; at the fame time the Uvula but chiefly the Tonfils were tumified, inflam; ed and painful, with fome white Specks; then follows a flush in the Face and fome miliary cruptions there, with a benign mild Fever, the fame efflorescence foon after appears on the neck, cheft and extremities; the 3d or 4th Day, Eruption is at the hight and well defined with fair intervals; the flushing goes off gradually, with a general *itching*; and in a Day or two more the *cuticle* fcales or peels off, especially in the extremities : At the same time the cream coloured floughs or fpecks in the Fauces become loofe and caft off, and tumefactions there do fubfide. The Tongue from the beginning is fur'd as in a Mercurial ptyali(m, urine high coloured, Blood by U.S. more florid than natural, in the whole course of the Diftemper a very great proftration

tion of Strength, and faintnefs upon recovery, nervous pains and weaknefs in the joints, particularly in the neck, wrifts and ancles; univerfal tendernefs to the touch; a tickling guttural cough, fome fhort Hectick flufhings, and lofs of enbon point. As in the Meafles there is a peculiar fmell, fo in our Diftemper the effluvia from the Patient have a proper fmell; in Children as if troubled with Worms, in grown Perfons the rancid fmell of foul Bed Linnen. The alvine excrement is of a dark caft and very fetid.

This Standard kind when left to nature, with a warm foft Regimen, had generally an eafy and falutary courfe in fix or feven Days; but when by a hot cordial method, or on the other extreme, by being too much exposed to the cold, or by officious profuse evacuations Nature was diffurbed in her Work; the Diffemper was protracted, or fome confequential ails from an imperfect defection enfued.

Where Nature required any affistance, the principal intentions were with regard to the cuticular eruption and the ulcuscula in the Throat. Any Affection of the Throat does frequently produce a natural ptyalism: Mercurials used with diferetion are a kind of specifick in fuch like ulcers & ulcuscula, and in fact here they moifined the Throat and Mouth, ftopt the fpreading of the ulcufcula, and promoted the caffing off of the floughs; and as an acceffory advantage (the Patients being moftly Children) destroy'd Worms: amongst all its pre-parations Calomel answered best, the gentle vomiting or few stools that it occasioned in some, did not confound the natural courfe of the Diftemper ; Turbith proves generally too ftrong a revultion, and the Eruption is thereby too much diverted ; this Diftem B 2

(5)

Diftemper did not well bear any other evacuations but Mercurials. Any detergent Gargle, with an addition of the Tinsture of Myrrb and Aloes, was of good use, especially for the Ulcuscula, and did promote the discharge of a ropy Phlegm lodged in the Fauces. As to the cuticular efflorescence, it was not a scarlet suffusion, but a miliary palpable eruption, or in lieu thereof in fome conflitutions a continued gentle breathing Sweat; and in a very few, who have naturally a liberior transitus by the Pores than is ufual, no fenfible cuticular excretion; in all the morbid effluvia discovered themselves by their peculiar imeli: Thefe were with good effect follicited by Snake-root Tea; or (as in fome Perfons) where this did occasion an ardor or burning heat, instead of a breathing mellow Sweat, Sp. C. C. or any other volatile Spirit in small Herb Teas anfwered well. Blifters and Suppedanea, in the beginning where Symptoms were not violent, occafioned a protracted Eruption; in fome immediately upon their application, the Eruptions vanished or became lefs vivid. When the Eruption began to decline a few loofe Stools were very refreshing. The Pati-ent being up, and having recovered a competent degree of Strength, is to be purged once or twice, to carry off any feculency that may have remained in the Blood and Juices.

For a more diffinct conception of the varities in this Diftemper, they may be reduced to three Classes.

I Those who die the first, second and third day of Illness, by an irremediable Necrosis of the Occonomy: in such the Seizure is generally sudden, a finking pain at the Stomach, an extreme prostration of Strength, a titubating low pulse, in some a flupor, in others a delirium, in some children

con-

convultions, and all of them generally die dozie : they are attended with fome colliquation, as continued vomiting, purging, profuse Sweats, bloatednefs of the habit, an infiltration like that of the Mumps vulgarly fo called, one or more of these: in general the texture of their blood and juices is much deftroy'd and rendred an incoherent puddle of corruption; in fact immediately upon (fometimes before) their exit, they have an intolerable fator. In this Class U. S, and other evacuations did accelerate death.

(7)

II. Those where the diffemper has its common or ordinary course; here the 6 or 7 day seems to be critical, and the Symptoms of death or recovery do generally then begin to manifest themselves. Some by peculiarity of conflitution, and from improper administrations do die or have an incipicent recovery fooner: others for the like reafons or fome particular accidents (u.g. if about the time of regular menstruation, the complicated fret occasions worfe Symcoms, and of longer continuance) have this period protracted and in fuch (where death is inevitable) the Symptoms of death may continue a day or two longer, that is the Patient may die the eighth or ninth day. All who continue Ill after that period, belong to the third Clafs, that is of confequential ails.

The Symtoms of bad Omen in this Clafs, are very great profiration of firength, dejection and despondency of mind, titubating low pulle, inceffant vomitings purgings or sweats, Tonfils much inflamed endangering firangulation, the specks in the Fauces of a brownish or leaden colour, or ragged and jagged, a continued jactantia in some, in others a fupor, retural of affumenda even of diluting common

mon drinks, a dry parched skin, Eruptions appears ing and difappearing alternately, Eruption univerfal of a dark redish cast continuing crude many days (because in this as in all eruptive Fevers, the darker or more livid the efflorescence, the more malignant) where the miliary pufiles are large, diffinct and pale like a chrystaline Small. Pox; where ftrong Cordials and Alexipharmicks have been used, the face, eye-lids, arms, hands, legs, feet fwell, and are of a dark red complexion, as in the most malignant Small-Pox; in children if the velum Palati be much affected, with an ichorous discharge by the Nose; where many mucous linings are expectorated, refembling the cuticle raifed by Vefications; when pus was brought up, where no floughs or exulcerations could be feen in the Fauces; where without any difficulty in fwallowing, this affection has reached down the Bronchia unto the Lungs with the Symptoms of a New-England Quinfey, and was erroniously deemed fuch : the deeper in the Thorax the complaint the greater the danger : in fome young children with fcarce any appearance in the Throat, fpreading Ulcers did form behind the Ears in the place where Infants have a natural Iffue or running. In fome the Tongue did throw off a flough or Exuvia, retaining the impreffions of the Papillæ; being a Mucus infpiffated, and of the fame nature with those mucous linings expectorated from the Bronchia or Oefophagus. Some have had impostumations in the Fauces, with a fatal ftrangulation, while others have escaped by the discharge of Ichorous curdly matter. Some especially of the adult female kind, have had Hysterical or Nervous Suffocations; but of no bad confequence, unless officiously and ignorantly treated with U. S. and other evacuations. The

The Fever is feldom too high, fometimes it is too low for a thorough laudable Eruption. If the Fever is too high, if the patient is plethorick or accustomed to U.S.; take away fome Blood but with diferetion; if the Tonfils are much inflamed with great pain and difficulty in fwallowing, ufe U.S. in the Jugulars, Epispafficks ad Nucham, encourage the Eruption, or its fuccedaneum a breath-ing fweat; a profuse fudor is equally to be avoided as a continued Diarrhea, either of them confound the diftemper in its natural courfe. In cafe of colliquations give ol. Cinamomi, decott. Alb. Elixir Vitriol, torrified Rhubarb and the like. As to the fpecks or floughs in the Fauces (they caft of in course in the benign kind) Mercurials inwardly, and the Gargles before mentioned topically, are useful; the practice in fome Country places of feparating them forcibly by spatulas is hurtful, because the irritation occafioned thereby induces a further flux upon the part, and the floughs form again worfe conditioned than before. Where the Brain is affected as in Vigilia, jastantia, delirium, Coma, fupor, &c; glyfters, Veficatories & Suppedanea are to be Where faintness or great profiration of ufed. ftrength, give toafted Bread foaked in fome generous Wine and Water, or volatile Spirits in their common drinks; Bezoars, Testacea and the like are of no use, the Shop Cordial Juleps and mixtures are only fugar'd drams. To enumerate all the other accidental Symptoms which do happen here, in common with other acute difeafes, would be trifling.

III. Confequential ails, which may be enumerated as in the following articles.

1. The natural Effects of an intense corrosive scorbusick

butick like colliquation of the Blood and Juices. Anafarcous fwelling or blotedness of the face, in some to that degree as to fhut up the Eyes; the fame Oedemetous swellings in the extremities; in a few an Infiltration in the Scrotum; in fome Petechia, Purple spots, scorbutick like fugillations upon the leaft icratch or bruise; hamorrhages of all forts, by the Nose, from the Lungs in expectoration, by Stool, by Urine, Proflufiums in Women tempore non debito ; these are difmal phanomina in the flate of any acute Fever, u. g. Small-Pox, and fcarce any recover ; but in our diftemper being only fhort tem. porary confequential ails, fcarce any of them proved mortal, but gave way to a foft milk diet, in fome to Cortex Peruv. or Elixir Vitrioli in others; a Girle æt. 14. with hæmorrhages of feveral forts, with Purple ipots, and fcorbutick like fugillations, recovered, notwithftanding of a very loofe Regimen. N. B. Thefe were not to be attributed to the Mercurial administrations, because they equally happened to those who had taken no Mercury.

2. Where the defecation has not been compleat, from want of natural firength, or from catching cold, or from undue evacuations: the reliquia were thrown off by Uritations, by Vesscations in feveral parts of the Body, by serpiginous eruptions chiefly in the face, by purulent Pussues, by Boils, by swellings and impostumations in the groin, armpits and other parts of the Body. The most frequent confequential ail of this kind is, when from cold received, the glands and cellulary tegument called the panicula adiposa in the fore part of the neck becomes infiltrated and obstructed; if not foon refolved by the continued fotus of warm woollens and hot animating applications; the induration increases creafes and fpreads every way, to as to fuffocate fome, in others they fphacelate and become Ulcers mortal or of difficult cure: thus a few have died with us in Bofton, but many in the Country. By catching cold likeways the Tonfils have afterwards inflamed and come to fuppuration: In a young Woman the Tonfils and Uvula being much ulcerated, did unite and coalefce into one mafs and remain fo; this might have been prevented by frequent gargling.

While these indurations are only in the form of Kernels as they are vulgarly called, woollen mufflers, Empl. de R anis cum Mercurio and the like, with gentle Catharticks, foon refolves them. Cataplasms in this case have done much mischief; because so foon as they are become cold, they act as a chilling damp upon the part, and deftroy its vitality. When they arrive to the state of putrid flaccid Ulcerations, digestives and soft fomentations intenerate the part and occasion the Ulcer to spread; spirituous animating deficcative dressings have done better. Exposing the part to the cold, either in flate of Tumesaction only, or in the subsequent exulcerations aggravates the ail.

3. From the violence which the Nerves have fuffered in this Illnefs; even where the Symptoms were apparently mild, they all complain of great faintnefs and Universal weaknefs, particularly in the joints. Some Women have Hyfterick affections, in a few upon recovery imbecility of mind or fillinefs, in fome flammering or lots of Speech for a few days, fome have had fhort fits of Melancholy, fome were feized with Epileptick fits, but not fo as to become habitual. All these diforders foon vanished, as the Patient recovered his Strength in course of time, and and by the help of a reftorative cordial Regimen and diet.

4. Other confequential ails in common with other fevers; particularly where the Strength of nature has been much impared by the diffemper it felf, or by immoderate evacuations, the Patient is left in a languifhing weaknefs. Where the Eruption has been impeded by being expoled to the cold, or by unfeafonable  $\mathcal{O}$ . S. or Catharticks; the patient falls into Heltical wasfings, fatal to fome in a very short time. All who underwent immoderate evacuations, were a long time in recovering of their Strength.

SCHOLIA or Some general remarks upon the whole. 1. This seems to be a new kind of Epidemical difeafe. It is not the fame with the Aphtha which have at times prevailed in Holland, as defcribed by Foreftus, and mentioned by Boerhaave in his Colleges. Tournefort says there is a diffemper not uncommon in the Levant, viz. a Carbuncle or plague fore in the bottom of the Throat; it carries off children in a few days, but does not affect grown People as does ours. Capt. Morton of late Ply-mouth Colony, who wrote many years ago his New England Memorial, fays that an. 1650. a difeafe in the Mouth and Throat prevailed, which proved mortal to many in a fhort time; but he does not defcribe it, and mentions nothing of a Fever. In Bofton November 1719. a flight milary fever chiefly with children, but was over in two or three days, unlefs by catching cold it continued appearing and difappearing alternately for fome days longer; there was no complaint of the Throat, and no deaths enfued. It is not the fame with the fore Throats which are observed from time

(12)

time to time in some of our Country Towns, especially in the Winter feason: these are Endemial and conflicutional, being tumefactious and exulceratious with fluxion in the Fauces and Neck ; proceeding from an intenfe fcropholous, fcorbutick, or Plorick habit (in fuch lubjects vefications by Cantharides did putrifie) without any Eruptive fever: ours have generally an Eruptive fever or tendency that way, fo that of those who have died in Bofton, not above one in feven died of any Throat ail, but of this fever. It is however observable that the Scrophulous and Pforick, are most fusceptible of this diftemper, and fuffer more remarkably.

2. This Epidemical distemper is no creature of the Seafons, it having prevailed from May 1735, when it first emerged, the whole year or all the Sealons round. It is no produce of peculiar climates and foils, becaufe it hath made its appearance in various places from Pemaquia in 44 N. Lat. to Carolina Southward, and as we are lately informed, it is in our West India Islands. It is remarkable that in damp places, as near large Ponds, trefh water Rivers, woodlands, and the like, it has done the greatest execution, as does the Rot amongst Sheep in fenny Lands.

It is not perfonally infecting after the rate of the Plague, Small Pox, &c. where every Perfon is fufceptible, excepting a very few anomolous conftitutions. Children are the most obnoxious to any intection; but with us feveral Children in the family, where the diffemper appeared, have escaped ; it is true where it happens in a family, it frequently feizeth feverals, as is the cafe with our Country Peripneumonick Fevers, and our Autumnal remitting flow Fevers, which cannot be faid to be CONT contagious. The diftance in time of Infection to be supposed received from a fick Person, to the time of the diffemper's appearing in the fuppofed infected, could never, with any reafonable allowance of latitude, be reduced to any rule, as in Small-Pox, Meafles, &c. We have Anatomically infpected Perfons who died of it with fo intense a fator from the violence of the difease, that some Practioners could not continue in the room ; but without being infected our felves or carrying it into families. Many children without referve, frequent the houfes and chambers of the fick, and escape. It does sherefore proceed from some undiscovered quality of the air, affecting only peculiar conflications of perfons and families: notwithftanding of its being generally favourable, it proves fatal to certain families; many families for this reason have buried all or most of their children, u. g. Boynton of Newbury-Falls left his eight children, at Hampton. Falls in 5 families died 27 Perfons:

3. This is a very anomalous Illnefs, fome complain a day or two before they are confined, fome are feized as it were inflantaneoufly, it is generally moft fevere with these laft. In some a foreness of the Throat and darting pain there, reaching the Ears, is previous to all other Symptoms; in others the common Symptoms of a fever appear, before any inflamation or specks are perceivable in the Fauces. Some have a fore Throat without any perceivable eruption, only a gentle breathing continued Sweat, or an increased infensible perspiration with the peculiar smell of the morbid effluvia. Some (but very sew) have the cuticular eruptions without any sloughs in the Throat; only the Tonfils, Uyula, and yelum Palati, tumified and inflamed; and in a few, a purulent discharge from some parts deeper than the Fauces, that is lower than the fight can reach, these are not without danger. Many of those who died early of a Necrosis, had no tumesaction, inflamation or specks in the Throat.

The time of Eruption is very uncertain; in a very few it preceeds the forenefs of the Throat, in a few it goes pari paffu with the affection of the Fauces; but generally it is (not much) later than the first complaints of the Throat, in a young Woman it was later by 14 days.

In ruddy complexions the efflorefcence is very difcernable; it is not fo diffinctly perceivable in Brunets, Indians, and Negroes; unlefs the miliary Eruption have a confiderable *Relievo* as in fome, they generally fcale and peal notwithflanding. Sometimes it appears only in the cheeks, fometimes only a few clufters in the extremities. Sometimes the *fuffufion* was fcarce miliary and vanifhed infenfibly by becoming gradually paler without fcaling. Where the Miliary Eruptions were confiderable, the extremities peel in fcraps or ftrips like *Exuvia*; in one or two the nails of the fingers and toes did caft off. The period or continuance of Eruption is fometimes prolonged by weaknefs of nature, by undue evacuations, or by the Patients being expoied to the cold.

4. In fome who were very flightly affected, their Illnefs was of a much fhorter continuance, than is defcribed in the Standard kind. Most of those who died of the Physician died by immoderate evacuations. As to the deaths, only a few were occasioned immediately by any diffemperature of the Throat; they were generally the effect of the Fever, either by an immediate Necrofis at first feizure, or by the or-

ordinary fatality of Fevers, or by confequential ails. In Bofton at a medium of the laft eight healthy years (1723. 1724. 1725. 1726. 1727. 1728. 1732. and 1733) in the Months of October, November, December, January, February, March, April to 18th May, died pr. an. 268 Whites and Slaves; this year in the fame space of time died 382, is 114 extra deaths, and may be refonably charged to this Illnefs, it being otherways a healthy time : of these 114. about 71 cafes came to my knowledge, whereof in the first period died 35, in the fecond period 28, and of confequential ails, 8. Of these 71, only about 10 can be faid to have died of fore Throats. Of these 71 only 9 were upwards of 14. et. According to the nearest estimate I can make in round numbers, about I in 35 have died, that is about 4000 Perfons in Boffon have had this diftemper, which is about one 4th part of the Inhabitants.

5. The Summer 1735 was unufually wet and chilly with many Easterly winds, in the Summer & Autumn it prevail'd and was very mortal in feveral Country Towns. In Boston it began in Autumn, but did not prevail until Winter, which was not rigid with hard frosts as is usual, but with a very dilagreeable chill in the air, especially in the Month of March last, in which Month was our greatest Mortality.

6. Most Malignant distempers affect to throw off their malignancy by some Emunctory. The desputation of this acrid inquation of the juices in our distemper, that is, its natural Criss, seems to be by the patent and falutary Emunctories of the Fauces and skin. In corrosive taints, u. g. Venereal and others, a Mercurial ptyalism and sudoristick decoction of the woods, answer best; this gave us the hint of promoting the the tendency of nature in our Illnefs, by Mercurials, and gentle breathing Sweats a bed; which with good management feldom fail'd, excepting where the Necrofis was irremediable from the beginning.

Some affection of the Throat ferms to attend most kinds of Eruptive Fevers. In the Small-Pox (even where the pufules and other Symptoms were in the fmalleft degree) they all complain of a foreness of the Throat, but without ulcerating. In the Measles there is a hoarsness, and foreness of the Throat. In ours besides the foreness, tumesaction, and inflammation in the Fauces; there are specks or floughs of a mellow white or Cream colour, like those on the infide of the cheeks in a Mercurial ptyalism; the Scrophulous and Venereal ulcers in the Throat are yellow; Aphthæ are more of the nature of phyctenæ; many of our Patients complain of a copperish tafte or peppery smart in the Throat, as they express it.

7. As in all other diftempers fo in this there do femetimes happen violent Symptoms, meerly from the Regimen and Medicines used; which on that account are not of that bad confequence, as if they had proceeded from the diftemper in its natural course u.g. in some confficutions a Turbith bolus operates with violence, so as to occasion shiverings, torsions of the Bowels, and Spasms, as if the Patient were moribund: Calomet even in very small doses feizeth the Mouth of some to a very confiderable degree of inconveniency.

8 We did not observe any genuine second seizures. It is true, being Winter Season, many common fore Throats, that is, relaxations of the Uvula and inflammations of the Tonfils; have passed with the less observing practitioners, for the genuine Epedemick

demick and were used accordingly; fuch have afterwards had this Illnefs, and was erroneoully called a second seizure. N. B. Our Epidemick is attended with no cough, unlefs when complicated with a cold or some old habitual Tuffis : upon recovery, it leaves frequently a small catarrhous colliquation or cough. but of short continuance.

In some after being well, upon catching cold, the Tonfils have been inflamed even to suppuration; in others the Uvula and Velum Palati infiltrated and some phlystene or common Aphthe, have appeared. Such have allo by fome been deem'd as fecond feizures, and used as such.

After a long continuance of cold chilly Weather. there fet in fuddenly warm Weather hot as mid Summer. May 25th, 26th, Ge. feveral children, who formerly had this Eruptive fever, have an efflorescence or miliary eruption by the heat, as is not unufual with children in hot weather : this was by miltake of some practitioners and others, called a fecond feizure.

9. No conditions of Mankind were exempted (in our Epidemical Autumnal dysentery A. 1734. The Ne-groes escaped) Europeans, West-India Islanders, Indians and Negroes, of all ages, were equally fubject to it: but, as in most Epidemical diseases, it affected Children and the younger Perfons more generally:

10. This is a Real History of the diftemper as it appeared in Boston New-England, taken clinically from the life and not copied. There is no ftroak or claufe, but what I can vouch by real not imaginary cafes. It is founded only upon observations or phanomina, that is upon the Symptoms that appeared in the courfe of this Epidemical disease; it must therefore be of permanent truth.

FINIS



