

TABLE 15.—Grand total for curvature of spine with ratio per 1,000 men.

State.	Number of cases.	Ratio per 1,000.	State.	Number of cases.	Ratio per 1,000.
Vermont.....	76	9.03	North Dakota.....	94	5.26
Wisconsin.....	538	8.51	Michigan.....	508	5.13
Maine.....	165	8.32	New Jersey.....	386	5.08
Missouri.....	698	8.15	Georgia.....	316	4.95
New Hampshire.....	65	7.63	Florida.....	119	4.93
Colorado.....	162	7.20	Kentucky.....	314	4.91
Tennessee.....	411	6.72	Washington.....	173	4.87
Utah.....	79	6.71	Nevada.....	16	4.86
Virginia.....	371	6.67	District of Columbia.....	56	4.85
Connecticut.....	236	6.57	Mississippi.....	179	4.75
Massachusetts.....	600	6.57	Oklahoma.....	279	4.59
Rhode Island.....	94	6.48	South Dakota.....	93	4.48
Oregon.....	101	6.31	Montana.....	122	4.45
New York.....	1,614	6.26	Alabama.....	221	4.26
Illinois.....	1,119	6.24	Kansas.....	159	4.14
Minnesota.....	444	6.17	Nebraska.....	111	3.77
Ohio.....	860	6.02	Louisiana.....	189	3.65
Indiana.....	430	6.01	Texas.....	426	3.59
Delaware.....	30	5.93	New Mexico.....	36	3.52
California.....	470	5.83	Arkansas.....	150	3.51
Iowa.....	389	5.67	Wyoming.....	41	3.16
West Virginia.....	220	5.65	Alaska.....	3	2.46
North Carolina.....	312	5.63	Arizona.....	16	1.93
South Carolina.....	215	5.46	State not specified.....	187	1.97
Idaho.....	68	5.45			
Maryland.....	201	5.40	Total.....	15,231	5.53
Pennsylvania.....	1,089	5.38			

16. *Diabetes mellitus*.—(See Table 16.)

TABLE 16.—Grand total for diabetes mellitus with ratio per 1,000 men.

State.	Number of cases.	Ratio per 1,000.	State.	Number of cases.	Ratio per 1,000.
Connecticut.....	44	1.23	Ohio.....	33	0.23
Vermont.....	6	.71	Montana.....	6	.22
Oregon.....	11	.69	North Dakota.....	4	.22
South Dakota.....	13	.63	Pennsylvania.....	45	.22
Maine.....	11	.55	Colorado.....	4	.18
Wisconsin.....	35	.55	Indiana.....	13	.18
Iowa.....	34	.50	Kansas.....	7	.18
Arizona.....	4	.48	Idaho.....	2	.16
Utah.....	5	.42	North Carolina.....	9	.16
Nebraska.....	12	.41	Tennessee.....	9	.15
Maryland.....	15	.40	Wyoming.....	1	.15
New York.....	101	.39	Alabama.....	7	.14
California.....	31	.38	Oklahoma.....	7	.12
District of Columbia.....	4	.35	Texas.....	12	.10
New Hampshire.....	3	.35	Arkansas.....	4	.09
Rhode Island.....	5	.34	Kentucky.....	6	.09
Illinois.....	57	.32	Florida.....	2	.08
Massachusetts.....	29	.32	South Carolina.....	3	.08
Nevada.....	1	.30	Georgia.....	4	.06
New Jersey.....	22	.29	Louisiana.....	1	.02
West Virginia.....	11	.28	Alaska.....		
Michigan.....	27	.27	Delaware.....		
Mississippi.....	10	.27	New Mexico.....		
Virginia.....	14	.25	State not specified.....	10	.11
Washington.....	9	.25			
Minnesota.....	17	.24	Total.....	740	.27
Missouri.....	20	.23			

17. *Exophthalmic goiter*.—This disease ranged in frequency from 0.3 to 9.4 per mille in the different States. The distribution as shown in Table 17 and Plate II, figure 3, is extremely suggestive. The State of Washington has a larger proportion than any other State, nearly 1 per cent of the men examined. This is followed by Wisconsin